

532 h 26
8560 h 4

Astronomia Accurata;

OR THE
ROYAL ASTRONOMER and NAVIGATOR.
CONTAINING NEW IMPROVEMENTS IN
Astronomy, Chronology, and Navigation.

PARTICULARLY
New and correct SOLAR and LUNAR TABLES;
With PRECEPTS and EXAMPLES of their Use, according to Old or New STYLE.

The ELEMENTS of the *Radical*
Mean Places and Motions of the
PLANETS and SATELLITES.

A New and Correct CATALOGUE
of STARS.

With their *Right Ascensions, Declinations, and Annual Variations,*
(supplying the Place of URANOGRAPHIA BRITANNICA.)

And ANSWERS to Fifty-three
Astronomical QUESTIONS.



CHRONOLOGICAL RULES and
TABLES.

REDUCTION of the DATES
of ÆRAS.

With *Universal and Perpetual*
TIME-TABLES; serving for
any Year, *past, present, or to*
come.

AND
ANSWERS to many *Chronological*
QUESTIONS.

Male inv et del. J. L. Müller sc.

The SEAMAN'S READY COMPUTER, Or New and EASY NAVIGATION:

Shewing how to keep an *exact* Ship's Reckoning; with *Examples*, according to *all* the Methods
of Sailing in PRACTICE, by only *summing up a few Figures*, taken out of *Tables*.
Tabular Answers to all the CASES in *Plain and Spherical TRIANGLES*; and to all the
CASES of SAILING.

An *alphabetical* TIDE-TABLE: An *universal* SEA-COAST TABLE, of the *Latitudes and Longitudes*
from *Greenwich*, and also the *Island of Ferro*, of the most *remarkable* Sea-Coast Places, *Ports*, and *Har-*
bours, lying *contiguous* round the WORLD.

To which are added
An *explanatory* COMMENT on the MOON's *Theory*: An *easy* Method, with *Examples*, of *computing* the
LUNATIONS, and of *Solar and Lunar* ECLIPSES. And how to determine the *Difference* of LONGITUDE,
from the *Royal Observatory*, at *Greenwich*, by observing the *Moon's Place*, under any *distant Meridian*.

With *some useful Remarks and Improvements* on the Motion and Theory of COMETS.
The *whole* interspersed with a Variety of *necessary* Tables, and *Articles*, subservient to the *three* principal SUBJECTS.

By ROBERT HEATH, a *Military* OFFICER.

*Now Heav'n in all her Glory shone, and roll'd
Her Motions, as the Great FIRST MOVER's Hand
First wheel'd their Course.*

Milton's Par. Lost. B. VIII. L. 499.

LONDON: Printed for the AUTHOR. Price in Sheets. M.DCC.LX.

Those Books not having the *Copper-Plate* of the *Palladium* (*Time and Minerva*) affixed to the *Title*, or *Frontispiece*, and signed with the
Author's own Hand, and sealed with his Seal, (*a Coat of Arms, and Lion rampant*) or any with the *Frontispiece-space* left *Blank*, are *spurious*,
counterfeit, pilfer'd, or pirated. And a handsome *Reward* is hereby offered to any *Person* who shall be pleased to discover to the *Author* any such
Books, published, or distributed, against his *Knowledge or Consent*. — All *Purchasers* whatsoever may be supplied at *Mr. Cole's, Mathematical*
Instrument-maker, in *Fleetstreet, London*, Successor to the eminent *Mr. Wright*, with proper Allowance of *Seven* for the Price of *Six* Books,
to all *Booksellers* and their *Country Correspondents*. Where *only the Trade*, in *Town*, may be supplied at the *proper Prices*, for *ready Money*,
with Allowance of the *Quarter-of-Hundred Book*, besides the *Seven* for *Six* Books, taken together; by those *Gentlemen* signing their *Order*
and *Place of Abode*; or subscribing for what *Number* they please.

TO THE
HONOURABLE and REVEREND
SPENCER COWPER, D.D.
DEAN of DURHAM.

SIR,

HOW illustrious and ennobled is SCIENCE! when Persons of high Birth, Honour, and Distinction, become her Votaries, and worship at her Altar; while other humble Admirers are obliged to have Recourse to the Favour and Interest of their SUPERIORS to be admitted to her Smiles!

Had not you, Sir, vouchsafed me the Honour to compose the Comment on the lunar Theory, to model the lunar Equation-Tables, and to lend me your Aid, in bringing the ROYAL ASTRONOMER to Light, that useful Design must have still lain dormant in Oblivion: Which being matured by your actuating Genius, naturally seeks your Patronage and Protection.

The incomparable Sir ISAAC NEWTON laid the Foundation of a perfect lunar Theory and Computation, and pointed out the Means to his indefatigable Successors for accomplishing so desirable an Attainment; to which the illustrious Halley, of our own Nation, the celebrated Mayer, M. De la Caille, and other ingenious Foreigners, have amazingly contributed. But you are the first Englishman who so easily resolved the difficult CENTRAL EQUATION OF THE MOON, in two Parts, and introduced such other new and easy lunar EQUATIONS, deduced from the Newtonian Principles of Gravity, as have, at the greatest Difference, rendered the final Computation of the Moon within two Minutes of Truth; which the great Newton himself left at a Difference of above five Minutes, and Halley's Computation left at above eight Minutes; as appears by his observed and computed lunar Places, compared for a Plinian Period of 18 Years; in the noble (and before unequalled) Correction he has given us of the lunar Theory. But your Computation, at a mean Difference from Truth, is not above three Fourths of a Minute; as appears by your Table of one Hundred lunar Places computed. Which great Accuracy is shewn, in the following Work, to be of prodigious Advantage in observing the Longitude from Greenwich Observatory, in long Voyages, at Sea.

The celebrated Mayer of Gottingen, undertook the arduous Task of correcting the lunar Theory about the same Time with yourself. You have since adopted his concise Method of Computation in the Construction of your Equation Tables, and so far improved upon it, as to perform that, with equal Accuracy, by ten easy lunar Equations, and one correcting Table, which that celebrated Foreigner has constructed thirteen different Equations, and as many different Arguments, to perform. And as you have as usefully and elegantly improved and adorned the lunar Theory and Computation, you are, among other celebrated Improvers of the celestial Science, (of which KINGS and PRINCES have been Patrons and Promoters) justly entitled to your merited Honours!

Your great Example, Sir, in encouraging and improving Science, will not only do Service, but redound Honour to this Nation. And, as your Motive, in your indefatigable Pursuit of new Acquisitions in Astronomy, appears to be the public Utility more than a View to your own Reputation, your excellent Improvements will reflect Honour, as long as the SUN gives, and the MOON reflects, Light.



But

a/a

DEDICATION.

But it will not seem strange to those who are the least acquainted with your NOBLE FAMILY, and your Affection for the belles Lettres, (which you seem to inherit as much by Nature as Acquisition) that you are a Lover and Promoter of the LIBERAL ARTS.

The noble Earl the LORD CHANCELLOR, your FATHER, was the Ornament of the Age in which he lived. His manly Ciceronian Eloquence, in debating the weighty Affairs of State; his Patriot Zeal, for the Liberties and Constitution of his Country, when attacked by perfidious and implacable Enemies; his uncorrupted Fidelity, in the Discharge of Offices of the highest Trust; his indefatigable Labours, in the Discovery of Truth; and unbiassed Integrity, in the Administration of public Justice; are so many Monuments of the Greatness of his Mind, the Honesty of his Heart, his vast Capacity, and unexampled Worth. The Rebel Lords, on whom he, as Lord High Steward, pronounced Sentence of Death, were no less edified and convicted by the Force of his Reasoning, than they were instructed to embrace their Fate with a becoming Resignation.

But you, Sir, were not educated to adorn the Profession in which that great Man, your Father, was early and eminently distinguished, but you were destined to improve the COELESTIAL SCIENCE, and to instruct Mankind in the Knowledge of their true Happiness; to encounter Vice, and reform an immoral Age, by the benevolent Spirit of CHRISTIANITY. Wherein the Dignity of your Birth and noble Alliance, (being Brother to an Earl and married to the Daughter of a late noble Lord) are of Advantage to the Doctrines you inculcate; in making the greater Impression on the Minds of your Auditors.

But I must intreat your Pardon, Sir, for having thus far trespassed on your Patience; occasioned by the Dignity and Lustre of the Subject, as well as by the Respect and Gratitude I owe you, for the singular Obligations and Honours I have frequently received at your Hands.

Who am, with the highest Regard,

S I R,

Your most devoted,

Most obedient,

Upnor-Castle, September 29,
1760.

and most humble Servant,

ROBERT HEATH.

PREFACE.

P R E F A C E.

AFTER long Expectation, we have published our *Astronomical Tables* of the Sun and Moon; which being connected with the Subjects of *Chronology* and *Navigation*, we are obliged, on Account of the Alteration and Enlargement of our original Plan and Design, to omit our *Astronomy* of the Planets and Satellites; referring our Reader to our *Astronomia Perfecta*, or Supplemental *Astronomer and Navigator*, for Satisfaction in those Subjects, and in all others promised and expected, to complete our Undertaking. Which Work will be printed as soon as due Encouragement is given us; but without proper Encouragement (as all Subscribers may, conditionally, depend on our Honour of strictly fulfilling our Engagement) no such Work will be printed, solely, at our own Hazard of Expence; though great Part of the Copy is actually finished.

To get the correct mean Places and Motions of the Planets and Satellites (employing the utmost Skill and Industry of the ablest Astronomers, for many Years, since the Improvement of accurate Instruments to observe them) is no easy Attainment. But to settle the Variation of these several Motions, by an Acceleration, and determine the Forms and Equations of the several Orbits, so that the computed Places may correspond nearly with the Places observed, for the present Century and remote Ages, is a Task requiring the greatest Judgment and Assiduity to perform. In which we have not wanted proper Assistance from our ablest mathematical Astronomers, and for perfecting which we are beholden to foreign Observations more than those made at Home, the more difficult to acquire; and not obtainable at all but by the deepest Artifice. One British Observer brooded, like a Miser, over his Observations, depriving this Nation, as 'tis said, of a perfect lunar Theory, by secreting some, and falsifying other, of his Observations, ordered Sir Isaac Newton; instead of being usefully and generously communicable of his Discoveries, like the late SCIENTIFIC and ILLUSTRIOUS HALLEY!

As to what is pretended of the Discoveries of the INDEFATIGABLE and ACCURATE Dr. BRADLEY, neither the Motion of the Earth's Aphelion (could it be exactly known) the Aberration of the Light of the fixed Stars, nor the Equation of the Precession of the equinoctial Points, nor that of the Nutation of the Earth's Axis, (all but imperfectly known, though pretended of such Importance by the great Halley's Preface Writer) are sufficient to account for the several Variations, in the planetary Orbits, where the computed Errors, in each, partly depend on those of the Earth's Orbit. And had the late Mr. Flamsteed been acquainted with all these refined Matters, as we are at present, they, probably, would not have reconciled the apparent Errors in his Observations.

The celebrated Halley's Skill and Contrivance, in supplying the Defects of the lunar Equations, by a Correction of Errors of the lunar Computation for a Plinian Period, as recurring, after which, of the same Magnitude, are no less to be admired for the Invention thereof, than for the Utility and Importance of the Improvement. Which Method can only be exceeded by some such direct Correction of the lunar Theory and Computation, as in our TABLES is attempted.

As to what is given us for Halley's Computation, by his Precept-Writer, seems to carry a Mistake. For otherwise, what Reason can be assigned for the sagacious Halley placing the Fourth (or Newton's 6th) before the central Equation; if, as the Precept-Writer represents, it was designed to be used after the central Equation. Using it before requires an awkward Correction of its Argument; without which Correction he pretends there will come out near half a Minute Difference, more or less, in the Error of Halley's Abacus of computed and observed Places, compared in his TABULÆ Lunæ Meridianæ. Whereas if this fourth Equation were designed by Halley to be used after the central Equation, it had been properly and naturally placed after it, immediately before his Variation Equation; whereby the present troublesome Correction of its Argument, and confounding the Computer how this Equation should be used, had been avoided. But this Precept Writer palpably mistakes, (in p. A. 8. Halley's Tables) where he applies the Excess of the Sun's Equation, and Sum of the Moon's Equations (reduced to Time in Mean Motion of the Moon from the Sun) to the Time of the Mean for that of the True Syzygy; without considering the Sign of the greater Quantity, in the said Excess; and therefore mistakes, in his Account of the true Syzygy preceding the Mean, or the Mean preceding the True: Which intirely depend on the Negative or Affirmative Sign of the Sum of the Sun's Equation with its proper Sign, and of the Moon's Equations with a contrary Sign; which Sum, with its proper Sign, is universally the Equation of the Interval between the Mean and True Syzygy, and not as Halley's Precept Writer describes.

We have given, what was much wanted, the correctest Catalogue of eminent fixed Stars of any extant, from M. De la Caille's Observations, with their Right Ascensions, Declinations, and annual Variations; by which the Moon's Right Ascension or Place on the Meridian, or at other Times, may be readily determined; as also the Right Ascension or Place of a Planet or Comet, at any Time, moving in its Orbit. This correct Catalogue, so far as it extends, may be said to more than supply the Place of the once promised and expected *Uranographia Britannica*. And it were to be wished, for the Honour of this Nation, that we had a *Britannic Catalogue* published, of all the most faithful, diligent, and correct Observations of the fixed Stars, that have been made by our most able and celebrated Astronomers. Since it is a Matter of high Importance to this Nation, that all the Observations made at the Royal Observatory at Greenwich, should be published at certain Times, for the Correction of our Theories, and Improvement of Astronomy; on which our Navigation so much depends.

P R E F A C E.

In the Preface to *Halley's Tables*, it is remarked, by his *Preface-Writer*, "That the Publication of astronomical Observations is of very great Importance, as they never become obsolete like Tables; but, on the contrary, the Usefulness of such Observations as are carefully made and faithfully delivered is greatly enhanced by their Antiquity. That the more diligent an Observer is, the more is the Public concerned in the Preservation (and he might have said Publication) of his Labours, and the less suitable is the Expence" (of such Publication) "to the Circumstances of a private Person." — Yet we find that no *British* Observer, since the communicative *Halley*, has followed the Example of the above Precept; though the *Precept-Writer* has invidiously spoken of that great Astronomer and Mathematician in another Part of his Preface, as not having published his Observations. Whose Pretence, that the *Greenwich*-Observations are best printed at the public Expence, "as less suitable to the Expence of a private Person" (while the *Palladium* Author, or any other Promoter of Science, would print them, were they open to Access) seems calculated for a prudent Augmentation of the present Salary, or else is proposed for a *Perquisite* to some Favourite Assistant-Observers.

As to communicating to the Ladies, the Method of making celestial Observations, from *Greenwich-Hill*, shewing them the rising Moon, and other Objects of Pleasure and Improvement, from that delightful View, we greatly approve. And we wish, those Ladies, vainly attending on the ORACLE in the *Old Bailey*, London, to consult their Destinies, and other future Events, would henceforth, more wisely, attend the celestial Observers; it being not above an Hour's Travel further, in their Coaches, to have infallible Proof of the secret Transactions among all the bright Rulers in the Heavens! There they may see the celebrated Mr. *Gael Morris*, enamoured of the Goddess of Science, walking Hand in Hand, and the renowned Mr. T. S. F. R. S. Usurper of the Ladies Diary, holding up her Train!

The Change of Declination of the fixed Stars, according to Succession of Years, happening with the Variation of the *Ecliptic* Obliquity, or Distance of the Pole of the *Ecliptic* from the Pole of the World, is no more to be wondered at than their annual Progression in the Signs of the *Ecliptic*, by the Precession of the equinoctial Points. By which Means, the Distances of the fixed Stars are found to increase, or decrease, from the Vertex of any Place, according to their annual Variations, as seen in our Catalogue. So that the Variation of the Declination of the polar Star, and of all other Stars, inserted in our Catalogue, for any Number of Years, past or to come, and thence the present Declination of a given Star, may be readily and truly determined.

Our supplemental EQUATION-TABLES, in the present Work, used with the Eccentricity of the Moon's Orbit, and second Equation of the Moon's Apogee, are designed for making Trial and Improvement of Computation, by the different Kind and Number of lunar Equations to be used. Mayer's lunar Equations are inserted for the same Purpose; to compare their Result with that from the lunar Equations marked C; and also with the Result from our other Equations.

The Rest of our Work we leave to speak for itself, or the Critics for it. On which we observe, that we shall be so far from taking it amiss from the judicious Critics, that we shall be obliged to them to mark our Defects, for the Improvement of Truth and Science. But if Cavillers, instead of Critics, should presume to find Fault with what they do not understand, or, according to the Custom of some Monthly Reviewers, use Contempt, Derision, and Abuse, for Criticism, on Subjects they are not Masters of, we shall have Recourse to Words with the Publisher, in our Vindication, unless he delivers up his Emisseries; as all such Scoffers, and Pretenders to Science, are very easily answered in their own Way; being no more than so many Flies under the Brush of a Flapper, — one being now sick of a *Sarcasm* — another maimed by a Stroke of *Ridicule*; — and a third disabled by a Strain given to his Attention, &c. — Some have ventured, without Ideas on the Subject, to characterize Mr. *Emerson's* Treatises of Fluxions, Mechanics, &c. and some to criticize and correct *Newton's Principia*; but alas! how vain and insignificant must all these Pretenders to Criticism appear, when it is considered that they attempt to criticize such unequalled Productions; of which Subjects we have but very few Persons, in this Nation, (now appearing) eminently qualified to judge. Who then, but Men of equal Genius and Capacity, shall be able to judge of the difficult yet certain Truths and Demonstrations those few Persons of Ability write? It is true their Method, or Language, may be cavilled at, to which all Method and Language are liable; but this is Nothing to the Truth of their Conclusions, deduced from a Series, or Chain, of demonstrative mathematical Arguments; whose right or wrong Inferences are the principal Part of the Subject to be criticised. And though many speculative Truths are not reducible to Practice, yet the Demonstrations of those Truths, on assumed Principles, are not the less certain; as will be found, where the Critic has Ideas and Judgment to pursue the intricate Chain of Reasoning. — Nevertheless there are many speculative Truths that are not necessary to be introduced in Practice; such as building Bridges on circular Arches instead of on elliptic or other Arches, full as sufficient, as circular Arches, to answer the End proposed, and to last out the Materials: Which Truth is in the Province of the practical Engineer, more than with the speculative Mathematician. There are other speculative Truths, in Astronomy, not reducible to Observation; as we find in the *Miscellaneous Tracts*. — And we hereby advertise the judicious Critics, that we are sensible of the Breaches of Connexion and Method in our Work, in our promiscuous Way of writing it, while the Press was employed; which will be all rectified in our second Edition: TRUTH (which we have endeavoured to preserve through the whole) being our principal OBJECT!

Our Royal Astronomer was not begun on the Plan whereby it is now finished. It was originally designed to contain only a few solar and lunar Tables, for accurately computing the Sun and Moon's true Places, to vie with, and make Improvement on, the Plan and Method of the celebrated Mayer's Tables. But finding this Accuracy of Computation would not have its Use unless applied to Navigation, (in observing the Moon's Place under a distant Meridian, for determining the Difference of Longitude, at Sea, by the Difference of that Time, and the Time when the Moon has the same computed Place at *Greenwich*, by our Tables) we farther extended our Plan, to the Royal Astronomer and Navigator. To which we found, that Chronology, and some other subservient Subjects being added, would render the

Work

P R E F A C E.

Work still the more useful and acceptable to the Public. But, in this progressive Augmentation, our Encouragement from Subscribers, we are sorry to say, was not proportionable to our Endeavours to oblige; and we find but few Encouragers of Science in this Kingdom.

Mr. Emerson, in his Preface to his *Mechanics*, observes to the following Purpose. — *That no Mæcenæ, in this Age, appears to patronise and protect Science; that the Decline of Arts and Science (he thinks) is in a great Measure owing to the false Ambition and extreme Avarice of the present Age. Where Man (unlike the antient Sons of Science and Encouragers of Arts and Literature) not being able to lift their Eyes above this Earth, make Nothing so much their Care as raking together the Dress it affords; striving, like the Mole, who shall die with the most Earth in his Possession. (Where some find more Plague to take Care of their Abundance, than others to supply their natural Wants.) That natural Knowledge finds Nothing but Contempt; and that Minerva yields to Pluto. That, if the general Disposition of Mankind had been in all Ages, as in the present, he doubts, whether we should have had either a Mill to grind Corn for our Bread; or a Pump to draw us Water. That, it is a trifling Excuse for Men in exalted Station to plead their Unacquaintance with Science; since Learning, of all Sorts, has been ever deemed under the Care and Superintendency of the GREAT: Who ought to protect and encourage Science and its Professors, for the Promotion of public Good. That Friendship and Generosity among the Great are hard to be found.*

Scire volunt omnes, mercedem solvere nemo.

Juv.

And the Reason the World has so few Aristotles is because there are so few Alexanders.

And we have observed, that invented *Histories*, corrupt *Novels*, and other false Amusements, find much more Encouragement than a Treatise on real and useful Knowledge.

But, that we may account to our worthy Subscribers and Encouragers of Science for the extreme Delay in the Delivery of our present *Work*, besides the Time required to select, vary, and new model our several Materials, the most for Advantage in our extended *Plan*, the Nature of Printing the *Work* has been such, and the Time required to convey and correct the Proofs, that, on Account of these several unexpected Hindrances, it could not be executed and delivered sooner. — And we hope that our Subscribers will excuse us (having shared in their respective Uneasiness) for these unforeseen Occasions of Delay.

☞ To prevent all such Uneasiness and Disappointment, on all Sides for the future, we have provided sufficient Means of Dispatch, for printing our second Edition; also our Supplemental Astronomer and Navigator, &c. as soon, as future Subscribers shall be pleased sufficiently to encourage those separate Works, on the Terms proposed in our *Palladium* for the Year 1761.

TO the ENCOURAGERS and IMPROVERS of ARTS and SCIENCES.

Vincit Amor Patriæ.

Virg. Æn. VI. v. 823.

WHATE'ER in Egypt, Greece, or Rome was known,
England can boast a GENIUS of her own;
Who, form'd with Passions Science to befriend,
And vers'd in Arts, Encouragement shall lend.
While the less noble stoop to abject Ways,
His Works a lasting Monument shall raise!
His Principles — as noble as his Blood —
Who acts in private for the public Good —
Descended from a loyal Patriot-Race,
Whom Trust still honours, and no Arts debase;
To Statesmen, Heroes, Gen'als, allied,
Their PRINCE's Glory, and the Nation's Pride!
In Fame's imperial Temple highly plac'd,
And some with laurel'd Honours doubly grac'd —
O Sons of Science, Merit still pursue,
And, by Example, keep the Prize in View;
Exert your Talents still to bless Mankind,
And a Reward your Services shall find;
Though Loss of sacred Science is deplor'd,
That Treasure in a COWPER is restor'd!

R. HEATH.


*The NEW absurd SYSTEM, lately published in Latin, (in the Imp. Magazine, September 1760)
for the true SYSTEM of the WORLD. By F. B.*

THE true System of the World is not that of Ptolemy, so much to be admired for its harmonious Epicycles, nor that of Copernicus, so famous for explaining the several Phenomena; nor yet that of Tycho, representing a noble, and, as it were, a Gothic, Structure. Among the Ancients, Ecphantus the Pythagorean, and Heraclides of Pontus; among the Moderns, Origanus, and Longomontanus were not unacquainted with this System. But I find none, in our Age, but William Comins, an Englishman, Vicar of Uffenden, in Rutlandshire, who has writ Something in Favour of this System. Plato, after he had begun to make a Figure in Philosophy, was pleased with it: I own, indeed, that he afterwards altered his Opinion of it, because the true Theory of this System, as one may guess, did not occur to his Thoughts. The whole System is thus explained on Principles consentaneous both to natural Philosophy and Geometry. — *Crede quod habes, et habes.* —

"The Earth is situate in the Center of the World, and revolves round its Axis from West to East, whereby Light and Darkeness succeed each other daily, and the Stars appear to make their daily Revolutions from East to West. The Moon is

The New Absurd SYSTEM of the WORLD. By F. B.

FIRMAMENT.



"carried round the Earth from West to East in the Space of one Month. Mercury and Venus revolve by a particular Motion about the Sun; the former in about 3 Months, the latter in about 7 $\frac{1}{2}$ Months, and at the same Time are carried round the Earth, every Year, with the Sun. Mars, Jupiter, and Saturn, run through their respective Orbits, (near the Center of which the Earth is fixed) the first in 2 Years, the second, with his Satellites, in 12 Years, and the third, or Saturn, with his, in 30. The fixed Stars are at Rest; whence the starry Heaven is called, in sacred Writ, the Firmament.

"The Stations and Retrogradations of Venus and Mercury, in the true System of the World, are in the Copernican Hypothesis; and in Tycho's System are particularly explained; but the Stations and Retrogradations, which appear to the Eye in Mars, Jupiter, and Saturn, proceed solely from the Circulation of Light from West to East, which is greater or less, according as the Planet is in Apogee or Perigee, or at a greater or less Distance from the Earth.

"The Manner of this Circulation will clearly appear from the following Example. If a Stone fall from the Firmament to the Earth, beside that Motion, by which it is precipitately carried toward the Earth's Center, it must, necessarily, partake of that Motion also, by which all Things are carried from

"West to East, according to the general LAW of the World. What we have said of the Stone, is to be understood likewise of the Globules or Atoms of Light; as all other Things are carried along from West to East, while they pass from the lucid Body to the Earth. And the nearer the Planet is to the Earth, the less will be the Circulation of LIGHT from West to East, when it reaches the Eye. The Circulation of Light toward the East in one Hour, for Instance, will be less than in two; whence the Planets appear to move more swiftly or more slowly, or to stand in the same Point of the Heaven, or to return toward the West, though they do really always continue advancing to the East."

F. B.

ON which We QUERE. How will the Comets shift for themselves, in their regular Orbits; or what Center or Law must they obey, in their respective Motions, to correspond with Observation, according to the above confused, or absurdly conceived, System? And how consistent this System with the Harmony of a Multitude of Systems, each having a fixed Star, or new Sun, for its Center? And how will all the Planets, according to this new Hypothesis, describe equal Areas in equal Times, about the Sun, as observed in the Copernican System.

CONFUTATION

CONFUTATION of the foregoing SYSTEM.

Magna est Veritas, et prevalebit. —

Observation 1. The Squares of the periodical Times are observed to be as the Cubes of the mean Distances from the Sun, of all Bodies revolving round the Sun; considered as the Fountain of Light and Heat, irradiated through the whole Copernican System: Which universal Law observed, is, at once, destroyed, if the Earth is considered at Rest. Therefore F. B.'s System is absurd.

2. The Times now observed of the Conjunctions, Oppositions, and Retrogradations of the Planets, are not such as would be, if they revolved, in Orbits, round the Earth at Rest.

3. The Bodies of Mercury and Venus in the lower Conjunction with the Sun, are hid behind the Sun's Body; and in the upper Conjunction, (in the Copernican System) are seen to pass over the Sun's Body, in Form of a dark round Spot, which is impossible to happen with the Earth at Rest.

4. Mercury and Venus are observed to have two Conjunctions with the Sun, but no Opposition; which cannot happen unless those Orbits are within the Earth's Orbit. And therefore cannot happen in F. B.'s System.

5. The Earth being at Rest, in the Center of F. B.'s System, will not account for the different Brightness and apparent Diameters of the planetary Bodies. Venus's apparent Diameter, when greatest, is nearly 66'; but when least not above 9' $\frac{1}{2}$. Mars's apparent Diameter, when greatest is about 21'; but when least not above 2' $\frac{1}{2}$; which F. B.'s System, with the Earth at Rest, will not account for.

6. The greatest Elongation or Distance of Mercury from the Sun is about 28 Degrees, and that of Venus about 47 Degrees, answering exactly to the Copernican System; but cannot agree with F. B.'s System, with the Earth at Rest.

7. Mars, Jupiter, and Saturn, have each their Conjunctions and Oppositions to the Sun alternate and successive; which cannot happen in F. B.'s System.

8. The Earth being placed in the Center of F. B.'s System, with the Sun and Planets revolving round it, the Planets must then, like the Comets in the Copernican System, be scorched with Heat, when nearest to the Sun; and frozen with Cold, at their greatest Distances; contrary to the Harmony of the Copernican System; proving F. B.'s System absurd.

* * Men of Ignorance and Ostentation endeavouring to raise Doubts and absurd Opinion, concerning settled and demonstrative Knowledge, founded on Experiment and Observation, are Enemies to Truth, useful Science, and the Public. Newtonienfis.

☞ The Dates of the Eclipses in the Palladium for 1761, subservient to this Work, are all according to New Style, since the Gregorian Æra commenced, October 5, 1582.

REMARKS on the said SYSTEM. By our ingenious and learned Correspondent,
ARGUS BRIAREUS. (See Palladium for 1761, P. 76.)

Clarius è tenebris depressa resurgit veritas.

THE preceding System, according to my View of it, makes a very absurd and irrational Appearance, more likely to be rejected, than received as Truth, by the Astronomers of this Age.

The Author whereof would do well to assign a Reason, why the Earth should be fixed in the Center of the whole System, rather than one of the other Planets. To say that it contains on it more noble Creatures, Beings of more exalted Faculties and Perfections, and therefore is the most illustrious of them all, would be a bold Assertion, and strongly favouring of a Mind overmuch attached to this little Ball, and its Inhabitants. And yet, if this be not allowed a substantial Reason, I presume it will be hard to find another that is so. — It may be asked, why the Sun should revolve round the Earth and not rather round one of the other superior or inferior Planets; seeing he is allowed to be the great Source of Light and Heat to all those dark Bodies. And also, why the Sun be made the Center of the Orbits of Venus and Mercury, while the larger and superior Planets, Mars, Jupiter, and Saturn, perform their Revolutions round an inferior Planet, the Earth, as their Center. Clear, rational Answers to these Questions may go a good Way to remove Objections to this System. If the Answer be made, that these Things must necessarily be so, to agree with the Appearances of the planetary Bodies, it will follow that this System cannot agree with Appearances; though the present Copernican System fully and clearly solves all those Appearances. The Copernican or Pythagorean Hypothesis not only effectually answers that End, in agreeing with the most accurate Observations, which the absurd System does not; but is infinitely more simple, uniform, and harmonious in its Construction, and consequently is infinitely more worthy the Wisdom and Contrivance of the divine and matchless Architect.

Burfordiæ, Cal. Nov. 1760.

ARGUS BRIAREUS.

EXPLANATION of TERMS.

A.

Achernar. A bright Star of the 1st Magnitude in Eridanus; observed by Dr. Halley at St. Helena, in 1677; never rising at Greenwich.

Achronical Rising and Setting of the Stars. When they rise in the Eastern Horizon, in the Evening, as the Sun sets; and set in the Western Horizon, in the Evening, with the Sun.

Æras. Points of Time; whence Chronologers begin to compute their Years. Of which Accounts of Time there are many Kinds; as the Æra of Creation; the Christian, Turkish, &c. Æra; some more famous than others.

Aldebaran.

EXPLANATION of TERMS.

- Aldebaran.** An Arabic Name for a fixed Star of the 1st Magnitude, called the Bull's South Eye.
- Algenib.** A Star of the 2d Magnitude, in the right Side of *Perseus*.
- Algol.** A Star of the 3d Magnitude in the right Side of *Perseus*, called *Medusa's Head*.
- Allioth.** A Star in the Tail of the great Bear.
- Almanac.** A Succession of all the Week-days, Month-days, and Holydays, correspondent, throughout the Year: Nothing without those Requisites being *truly* and *literally* an Almanac. Which fixed, and not variable or astronomic Computations, are its chief Properties.
- Alramech.** The same as *Arcturus*.
- Amplitude.** An Arch of the Horizon between the Rising or Setting of the *Sun*, *Moon*, or *Star*, and the East or West Points, of the same Name (N. or S.) with their Declination.
- Analemma.** A Projection of the Sphere, *orthographically*, on the Plane of the Meridian, the Eye being at an infinite Distance, placed in a Line perpendicular to the Middle of the said Plane.
- Analogy.** The same as Proportion.
- Angle of Direction.** The Angle made by the *Axis* of the Moon's Orbit and *Axis* of the Earth, at the Center of the Earth's Disk. Of great Use in constructing solar Eclipses. In which Projection, if
 Sun in { ϖ , Ω , $\cap$, $\triangle$, $\cap$, $\dagger$, } Earth's *Axis* to the { Right } of the *Axis* of the Ecliptic.
 { ϖ , $\cap$, $\times$, ∇ , $\cap$, Π , } { Left }
- Angle of Evection.** The Difference between the mean and true central Equation of the Moon, in the *new Astronomy*, the greatest Evection Angle being about $1^{\circ} 20'$ in Quadrature. Or the Difference between the least and true central Equation, according to *Street's* Computation; the greatest Angle of Evection, this Way of computing being about $2^{\circ} 37'$, in Quadrature.
- Angle of Reflection.** The same as *Variation* Equation of the Moon; being greatest in *Orbits* of the Moon from the Sun, or $37^{\circ} 33''$, according to *Street's Astronomy*.
- Anomaly.** A Planet's angular Distance from the Aphelion Point, or farthest Distance from the Sun.
- Anses or Anse.** The Ring of *Saturn*; appearing like Handles to the Body of the Planet.
- Antares.** A fixed Star of the 1st Magnitude, or the *Scorpion's Heart*.
- Antecedentia.** According to a preceding or contrary Order of Signs; the same as *Retrogradation*.
- Apsis, or Apfides.** The Extreme, or Extremes of an Orbit's Transverse Diameter. Signifying also the *Apogee* and *Perigee*, of the Moon's Orbit.
- Argument of Latitude.** The Distance of the Moon from the North or ascending *Node*, in Degrees, &c. by which the Moon's Latitude is determined.
- Ascensional Difference.** The Difference between the right and oblique Ascension or Descension; or the Time the Sun rises and sets before and after Six.
- Ascendent.** The rising Point of the Ecliptic.
- Azimuths.** Vertical Circles passing through the *Zenith* and *Nadir*: Which Terms see.
- B.
- Babylon.** A famous and antient City in *Egypt*. It stood upon a Square of 120 Furlongs long, of 15 Miles each Way. Circumference 480 Furlongs, or 60 Miles.
- Walls { 587 } Feet { Thick } In the Walls were 100 Gates, 25 on each Side, with 25 Streets each Way,
 { 1350 } { High } crossing one another at right Angles; making 676 Squares, each of which was
 { 480 Furlongs Round } $4\frac{1}{2}$ Furlongs, on the Side, and $2\frac{1}{4}$ Miles round.
- In the Middle over *Euphrates*, was a Bridge 220 Yards long and 10 broad. The Temple of *Belus* was a Furlong Square, consisting of 8 Towers, one above another, each 75 Feet high. King *Nebuchadnezzar* enlarged it to 2 Furlongs on each Side, making a Mile he enlarged it round. The old Palace was 4 Miles in Circumference; the new 8 Miles. The Hanging Gardens, 400 Feet Square, by several Terrasses, one above another, till the highest became 350 Feet. The Ascent was from Terrass to Terrass, by Stairs of 10 Feet wide, built by Arches upon Arches, and a Wall 22 Feet thick. This wonderful City like the famous and antient City of *Troy*, is vanished out of the World, where so many great Things were once transacted! And as magnificent Cities and Empires successively vanish in Oblivion; so shall Systems of the celestial Bodies, and all their Inhabitants, pass away, like *Shakespeare's* baseless Fabrick of a Vision, leaving not a Wreck behind!
- C.
- Calendar.** The same with *Almanac*: Which see.
- Center of the Equant.** The same as the upper Focus of an Ellipsis, round which the Planets uniformly move.
- Conjunction true.** When the Centers of two Planets (as of the SUN and MOON) are seen conjoined in a right Line, from the Earth's Center.
- Conjunction apparent or visible.** When two Planets are seen conjoined in a right Line from the Earth's Surface.
- Consequentia.** According to a following Order of Signs.
- Copernicus Nicholas.** A Native of *Thom*, in *Polish Prussia*, born in 1473, and died in 1543. A great Astronomer and Reviver of the *Pythagorean System*.
- Cosmical Rising and Setting of Stars.** Rising in the Morning with the Sun, and Setting as the Sun rises.
- Cycle of the Moon.** A Revolution of 19 Years, after which Time the mean Lunations return on the same Month-days. The Number of this Cycle is the Golden Number.
- Cycle of Indiction.** A Revolution of 15 Years, substituted in the Room of the *Greek Olympiads*; denoting the Year the Tribute was to be paid to *Constantine the Great*; established 312 since Christ, on September 24.
- D.

EXPLANATION of TERMS.

D.

Day Artificial. Day of the Sun's Light, from Sun-rising to Sun-setting: Being the Distinction of Days according to Custom, which is the greatest Authority in the proper Use of Words.

Day Natural. Day of the Year, or 24 Hours. The Time of an apparent Revolution of the Sun, from any Hour Circle to the same.

Descendent. Setting Point of the Ecliptic.

Digit. The twelfth Part of the Sun's Diameter, in solar Eclipses. In lunar Eclipses, the Moon's Digits eclipsed may be 23; all above 12 Digits shewing how much the Earth's Shadow more than covers the Moon's nearest Edge to the Middle of that Shadow, or Eclipse.

Direct. A Planet's Motion is direct, moving according to the following Order of Signs; as from *Aries* to *Taurus*.

Disk of the Sun and Moon. Their round flat Faces, as they appear at a Distance.

Disk of the Earth. Her Face, as seen from the Sun or Moon; or from other Planets; being equal to the Difference between the Sun and Moon's horizontal Parallax.

E.

Earth. The Globe we inhabit; carried round its Orbit, called the Ecliptic, in 365 D. 5 H. 48 M. 55 S. but to the same fixed Star in 365 D. 6 H. 9 M. 24 S. keeping its Axis always parallel to itself. Whose swift Rotation causes its Figure to be flattened next its Poles, and swelled out about its Equator. Its equatorial exceeds its polar Diameter by about 63 Miles; Sir Isaac Newton having proved the equatorial to the polar Diameter in the Ratio of 698 to 692. — Whence, according to Norwood's Measure of the Miles in a Degree of a great Circle, the Earth's

Circumference 25035.83
Diameter 7969.15 } English Miles.

Height of the Atmosphere 47.11

Eccentricity. The Distance from the Center to the Focus of a Planet's Orbit.

Elongation. A Planet's farthest angular Distance from the Sun, as seen from the Earth. Mercury's Elongation is never more than $28^{\circ} 21' 8''$, nor less than $17^{\circ} 35' 42''$. Venus's never more than $47^{\circ} 38' 35''$, nor less than $44^{\circ} 56' 14''$.

Emerfion. When a Planet, or Satellite, eclipsed, begins to recover its Light.

Epacl. The Difference between the solar and lunar Year, of $365^d 5^h 49^m$ and $354^d 8^h 49^m$, or $10^d 21^h 0^m$; being reckoned 11 whole Days, the Days of the Month sooner the Moon changes in the following than in a former Year.

Ephemeris. A yearly and daily Account of the Places of the celestial Bodies.

Equation. The Difference between one Quantity and another to which it is to be equated. As the Difference between the Planet's mean and true Places, Anomalies, &c. or Equation of mean to apparent solar Time.

Eridanus. A Southern Constellation, or the River.

F.

Fomabaut. A Star of the first Magnitude in the Mouth of the Southern Fish.

G.

Golden Number. The Number of the Moon's Cycle.

H.

Heliacal Rising. When a Star recovers from the Sun's Light to the Westward, so as to be seen again in the Morning before the Sun.

Heliacal Setting. When a Star first becomes invisible, by the Sun's near Approach. The Moon being large, can be seen 17° from the Sun, when the other Planets cannot be seen till they are near 30° distant from him.

Heliocentric Place. The Place of a Planet as seen from the Sun; from whence all the Planets appear to move direct, according to following Order of Signs; and the heliocentric Latitude is the same with the Inclination of the Planet's Orbit with the Ecliptic.

Horizon. The great Circle of the Heavens, bounding our Sight; dividing the upper from the lower Hemisphere. It is distinguished into sensible, as what we see from the Earth's Surface; and rational Horizon, as what it would truly appear, if we beheld it from the Earth's Center.

Hour Circles. The same as Meridians; or great Circles cutting the equinoctial right Angles, at any Hour of the Day or Night.

Hypothesis. A System, Theory, or Conjecture, to explain Appearances by; which is the more consistent with Truth and Reality, the more it corresponds, in general, with Observation.

I.

Jewish Hours, otherwise called *Planetary Hours.* The twelfth Part of the Day (of the Sun's Light) from Sun Rising to Setting; and the twelfth Part of the Night, from Setting to Rising of the Sun: Being the antient Jewish Account of Time.

Illuminative Month. The Space of Time the Moon is visible from Conjunction to Conjunction.

Immersion. When a Planet or Satellite first begins to be eclipsed.

Jupiter. Marked Υ . One of the primary Planets, next to the highest, or Saturn. He moves round the Sun in 11 Yrs. 313 D. 5 H. 35 M. 4 S. Inclination of his Orbit, $1^{\circ} 20'$. Diurnal Motion, $4' 59''$. The Law of his Motion about the Sun is such, that the Square of his periodical Revolution is as the Cube of his mean Distance from the Sun, compared with that immutable Law, observed in the Motion of the Rest of the Planets; first discovered by Kepler, and since demonstrated by Sir Isaac Newton. Mr. Flamsteed and Dr. Halley observed that Jupiter moved too slow by the Tables, which were corrected accordingly.

K.

EXPLANATION of TERMS.

K.

Kepler, John. Of Wittemberg, in Germany, flourished in 1620; he was Mathematician and Astronomer to three Emperors; and was the first who discovered the *elliptical* Orbits of the Planets, and that their *periodical* Times were as the *Cubes* of the Distances from the Sun. And explained the general *Phænomena* of solar Eclipses. In the Year 1618 he published his Epitome, intitled *Astronomiæ Copernicæ: Ephemerides de Harmoniâ Mundi, Mysterium Cosmographium*: Likewise, *De Motibus Stellæ Martis*, and other Pieces of Astronomy.

L.

Latitude of a Star or Planet. The Distance of either from the *Ecliptic*; measured on the Arch of a Circle of Latitude, passing through the Center of either, and the *Poles* of the *Ecliptic*.

Libration of the Moon. 1. In *Longitude*, being a Motion arising from the Plane of that Meridian of the Moon in general, nearly turned towards us; not directed towards the Earth; but towards the other *Focus* of the Moon's *elliptic* Orbit; whereby to an Eye at the Earth she appears to *librate* to and fro. 2. In *Latitude* arising from her *Axis* not being perpendicular, but inclined to the Plane of her Orbit, whereby sometimes one of her *Poles* and sometimes the other dips a little towards the Earth. 3. A *Libration*, whereby one Part of the Moon is not really turned towards the Earth as in the former Manner; but another is illuminated by the Sun. For her *Axis* being nearly perpendicular to the Plane of the *Ecliptic*, when she is at her greatest Southern Limit of the Parts adjacent to her North Pole, will be illuminated by the Sun, while the contrary, or South Pole, will be involved in Darkness.— These *Librations* are completed in her synodical Month. See *Newton's Principia*.

Limits of Eclipses. Sun or Moon's Distances from the Node at new or full Moon, within which Distances all *solar* or *lunar* Eclipses must happen.

		Fr. Node.	Lat. Moon.	Greater than which Distances from the Node, or Latitude of the
Greatest Limit	{ Moon }	12° 2' 5"	{ 1° 2' 23" }	Moon, no Eclipse of the Moon or Sun can happen, at nearest Distances of the Sun and Moon from the Earth.
	{ Sun }	18 20 4	{ 1 34 14 }	
		Fr. Node.	Lat. Moon.	Greater than which Distances from the Node, or Latitude of the
Least Limit	{ Moon }	10 19 13	{ 0 53 2 }	Moon, no Eclipse of the Moon or Sun can happen, at farthest Distances of the Sun and Moon from the Earth.
	{ Sun }	16 35 1	{ 1 25 30 }	

M.

Mars. Marked δ , One of the *primary* Planets; moving round the Sun, in an Orbit between the Earth and *Jupiter*, in 1 Yr. 321 D. 22 H. 18 M. 18 S. His *diurnal* Motion, 31' 27". The *Inclination* of his Orbit to the *Ecliptic*, 1° 52'; being 15 Times less than our Earth. His Colour is ruddy like that of the Star *Aldebaran*; and is *retrograde* once in two Years.

Medium Cæli. The Degree or Point of the *Ecliptic* on the Meridian at any Time of the Day.

Mercury. Marked ξ . One of the *primary* Planets; moving through his Orb next the Sun, in 87 D. 23 H. 14 M. 34 S. His mean heliocentric *diurnal* Motion 4° 5' 32". The *Inclination* of his Orbit with the *Ecliptic*, 6° 59'; being never more elongated from the Sun than 28° 21' 28", and therefore seldom seen. — He appears *retrograde* 4 or 5 Times every Year; and is 27 Times less than our Earth.

Meridional Parts. Tables fitted for the Use of *Navigation*, from a Projection of the Earth's Surface, where the *Meridians* increase as the *Parallels* of Latitude decrease. For as all the *Parallels* end in a Point at the Pole, so the *Parallel* Meridians, being infinitely continued, never meet.

Micrometer. An Instrument invented by Mr. *Townly*, fitted to a Telescope, for taking the *Diameters* of the Planets.

Month, lunar or periodical. The Time the Moon is going through her Orbit, 27^d 7^h 43^m.

Synodical. From her Conjunction to the next Conjunction with the Sun, in 29^d 12^h 44^m.

Solar. Time of the Sun going through one of the Signs in the *Ecliptic* about 30 $\frac{1}{2}$ Days, but all of different Length.

— of twenty eight Days. Nearly equal to the periodical Month.

— of the Calendar. More or less Days as in the Calendar.

Moon. Marked Δ , a *secondary* Planet; moving round the Earth, to the same fixed Star, in 27 D. 7 H. 43 M. 5 S. but returns not in Conjunction with the Sun, till 29 D. 12 H. 44 M. 3 S. &c. While the Earth moves round the Sun in 365 D. 5 H. 48 M. 55 S. Her Orbit intersects the *Ecliptic* in two opposite Points, called *Nodes*. She is 50 Times less than our Earth.

Moon's Inclination of her Orbit with the Ecliptic. 4° 59' 35" in Conjunction and Opposition, 5 17 20 in Quadrature with the Sun.

N.

Nadir. The Point of the Heavens under the Earth; diametrically opposite to the *Zenith* above it.

Nodes. The Points where the several Orbits of the Planets intersect the Earth's Orbit, or *Ecliptic*; having a slow *progressive* Motion, as well as the *Aphelion* Points.

Nonagesimal Degree. The 90th Degree or highest Point of the *Ecliptic* at any Time of the Day or Night; whose *Altitude* is always equal to the Angle that the *Ecliptic* makes with the *Horizon*; which is also equal to the Distance between the Pole of the *Ecliptic* and Vertex or *Zenith* of the Place.

Nychthemeron. A natural Day in the Planets; including a Day of Light, and Night, where the Sun sets in a *diurnal* Revolution of the Planet; being with us 24 Hours.

O.

Oblique Ascension. The Degree and Minute of the *Equinoctial*, rising with the Sun, Moon, or Planets, in an oblique Sphere.

Oblique Descension. The Degree and Minute of the *Equinoctial*, setting with the Sun, Moon, or Planets, in an oblique Sphere.

Occultation.

EXPLANATION of TERMS.

Occultation. The Time that a *Star* or *Planet* is hid from us; when eclipsed by the Interposition of the Body of the Moon, or some other Planet, betwixt it and our Sight.

Orthographic Projection of the Sphere. Drawing the Superficies of a Sphere on a Plane, cutting it in the Middle, the Eye being placed at an infinite Distance vertical to one of the Hemispheres; in which all the Hour Circles become Ellipses: The same with *Analemma*.

P.

Parallax. The Difference between the true Place of a Planet, as seen from the Earth's Center, and its apparent Place as seen from the Earth's Surface.

Parallax in the annual Orbit. The Angle the Earth would appear under to the Eye, at each Planet, to be elongated from the Sun; being greatest and least at extreme Positions.

Path of the Vertex. A Circle described by any Point of the Earth's Surface, by the Rotation about its *Axis*. This Point is considered as vertical to the Earth's Center; being the same with *Vertex* or *Zenith*. The Semidiameter of the *Path* of the *Vertex* is equal to that of the Complement of Latitude of the Place describing it.

Penumbra. A faint Shadow, or outmost Edge of the dark Shadow of the Earth, in an Eclipse of the Moon: It being difficult to determine where the dark Shadow begins, and where the Light ends, in (*solar*) Eclipses of the Earth, by the Moon's Shadow. The Penumbra's Semidiameter is equal to the Sum of the apparent Semidiameters of the Sun and Moon. For if, at any Time of the true Conjunction of the Sun and Moon, none of the *Penumbra* falls within the Earth's Disk, the Sun will then be *nowhere* seen eclipsed.

Perigeon. A Planet's nearest Distance to the Earth. When the Moon's mean Anomaly is 6 Signs she is then in *Perigeon*, and her diurnal Motion about 15° . This Point is opposite to the *Apogee*, both at the Extremes of the transverse Diameter of her elliptic Orbit.

Perihelion. A Planet's nearest Distance to the Sun; the *heliocentric* Motion being here the *swiftest*, with 6 Signs Planet's mean Anomaly.

Place of the Sun or Planet. The same with Longitude.

Place true and apparent. As seen from the Earth's Center and Surface, respectively.

Poetical Rising and Setting of Stars. *Achronical*, *cosmical*, and *heliacal*, before described.

Precession of the Equinoxes. The fixed Stars are immoveable; the Earth travels round the Sun in its annual Orbit, with its *Axis* carried $66^{\circ} 31' 30''$, inclined to the Plane of the *Ecliptic*: And by the Earth's diurnal Motion East, round its *Axis*, in 24 Hours, the Equinoctial Points are moved contrary at about $50''$ yearly; whereby the fixed Stars are made to go forward in a following Order of Signs, yearly, as much.

Projection of the Sphere. A geometrical Delineation of the Circles of the Sphere, or any assigned Part, on the Plane of any great Circle, as on the *Horizon*, *Meridian*, *Equinoctial*, *Ecliptic*, *Colures*, or on the *Tropics*, &c. being either *stereographic*, supposing the Eye 90° distant from, and perpendicular to the Place of Projection; or *orthographic*, the Eye being at an infinite Distance, in the Center of Projection.

Ptolomy. *Claudius Ptolomeus*. A Native of *Pelusium*, a City of *Africa*, in *Egypt*; who flourished 135 Years after *Christ*; and was Author of the *System* known by that Name.

R.

Radius. The Semidiameter of a Circle. A mean Proportional between the *Tangent* and *Cotangent* of any Arch.

A mean Proportional between *Sine* and *Secant* Complement of any Arch.

Ratio. The particular Relation, or Habitude, of two Numbers. The Similitude of Ratios signifies Proportion, or Analogy.

Simple, duplicate, triplicate, quadruplicate, &c. Ratio. Is the single, square, cube, 4th Power, &c. of two Quantities.

Subduplicate, Subtriplicate, &c. Ratio. The square, cube Root, &c. of two Quantities, &c.

The sesquiplicate Ratio. Is the one and $\frac{1}{2}$ or $\frac{3}{2}$ Power, consequently *subsesquiplicate Ratio*, the $\frac{2}{3}$ Root, and the subquintuplicate Ratio, the $\frac{1}{5}$ Root, of any two Quantities; and so of the Rest. *Leadbetter*, in his *Astronomy*, has misrepresented what a Ratio is, in every Case and Circumstance; giving a wrong Idea thereof.

Ratios.	$\left\{ \begin{array}{l} a \text{ to } b \text{ the simple.} \\ a^2 \text{ to } b^2 \text{ the duplicate.} \\ a^3 \text{ to } b^3 \text{ the triplicate.} \\ a^4 \text{ to } b^4 \text{ the biquadrate.} \\ \text{\&c.} \end{array} \right.$	$\left\{ \begin{array}{l} a^{\frac{1}{2}} \text{ to } b^{\frac{1}{2}} \text{ the subduplicate.} \\ a^{\frac{1}{3}} \text{ to } b^{\frac{1}{3}} \text{ the subtriplicate.} \\ a^{\frac{2}{3}} \text{ to } b^{\frac{2}{3}} \text{ the sesquiplicate.} \\ a^{\frac{2}{5}} \text{ to } b^{\frac{2}{5}} \text{ the subsesquiplicate.} \\ a^{\frac{1}{5}} \text{ to } b^{\frac{1}{5}} \text{ the subquintuplicate, \&c.} \end{array} \right.$	Ratios.
---------	---	---	---------

Recession of the Equinoxes. Going back of the Equinoctial Points, about $50''$ in each Year; caused by the Earth's spheroidal Figure, in its diurnal Motion.

Reduction of the Orbit to the Ecliptic Place. The Angle between the *Axis* of the *Ecliptic* and *Axis* of the Planet's Orbit; being equal to the *Ecliptic's* Arch between the two *Axes*.

Reflexion. The Pole's Distance from the *Horizon* of the *Disk*. — The same as the Sun's Declination.

Refraction. Produced by the Atmosphere, making a Star appear more elevated than it really is.

Retrogradation. Going back, in a contrary Order of Signs: as the Moon's North Node moves retrograde upon the whole of its Motions.

Right Ascension. The Degrees of the *Equinoctial*, reckoned from the Beginning of *Aries*, coming to the *Meridian* with a Star or Planet; or to any Hour Circle, at right Angles with the *Equinoctial*.

S.

EXPLANATION of TERMS.

S.

Saros, Chaldean. Of 18 Yrs. 11^d 7^h 43^m, when 4 Leap-Years intervene — but only 18 Yrs. 10^d 43^m when 5 Leap-Years intervene. — The same as *Plinian Period*.

Satellites. Secondary Planets, revolving round their respective Primaries, as the Moon round the Earth; the Satellites of *Jupiter* and *Saturn* round those Bodies.

Saturn. Marked ♄, One of the *primary* Planets; the highest in all our System; moving round the Sun in 29 Yrs. 158 D. 13 H. 14 M. 4 S. His diurnal Motion 2'. Inclination of his Orbit 2° 33'. He is retrograde once every Year.

Scenographic Projection. The same as *Perspective*.

Sidereal Day. The Time between any *fixed Star* departing from, to its Return to, the same Meridian; being 23^h 56^m 4^s 6th, Parts of a mean solar Day.

Sidereal Year. The Time of the Sun going from a *fixed Star* to his Return to the same, 365^d 6^h 9^m 24^s, &c.

Solar Year. The Time of the Sun's going from, to his Return to, the same Point of the *Ecliptic*; called also *tropical Year*, of about 365^d 5^h 48^m 55^s.

Street's Computation. 1. Having found the Moon's mean Anomaly, Apogee, and Node; you find the *eccentric Equation* answering to her mean Anomaly, which being added to, or subtracted from, the said Anomaly, according to its Title, you have thence the 1. *equated Anomaly* of the Moon; to which adding the Moon's Apogee, you have 1. *equated Place* of the Moon.

2. From the 1st *equated Place* of the Moon take the Sun's true Place, and the *Residue* is the Distance of the Moon from the Sun, by which you may take out the Equation of Variation (as at P. 74, according to *Horrox*) which added to or subtracted from the 1st *equated Anomaly* (before found) will give the 2. *equated Anomaly* of the Moon.

3. The *synodical Anomaly* being found, as under that Term, take its Half, if less than 6 Signs, but the Half of its Supplement to 12 Signs, if above 6 Signs.

4. To the *Logarithm* 3.640432, of the Diameter of the Circle of *Evection*, add the Log. Sine of the Moon from the Sun (before found) which Sum, rejecting Log. of Radius, will be the Log. of the Chord of *Evection*. This *Logarithm* to be taken from the *Logarithm* of the Moon's Distance from the Earth (found in a Table) adding the *Logarithm* of Radius to the Remainder, will be the *Tangent* of an Arch, from which reject 45°.

Now, say, As Radius to the *Tangent* of this Remainder, in Degrees, so the *Tangent* of half the *synodical Anomaly*, or *Tangent* of Half its Supplement to 12°, (before found) to the *Tangent* of an Arch, whose Difference from that Half will be the *Evection-Equation* to be added or subtracted according as the whole *synodical Anomaly* is more or less than 6 Signs.

— With which *Evection-Equation* you must connect the Variation, or *Reflexion*, according to its Sign, (before found) for an Equation; which being connected with the Moon's Place 1. *equated*, will give her *Orbit-Place*; which is reduced to her *Ecliptic Place*, by the *Reduction-Equation*, found as usual.

The true Place of the Node (2. *equated*) being taken from the Moon's *Orbit-Place*, the Remainder will be the *Argument of Latitude*.

By the Distance of the Moon from the Sun, you find the *Excess* of the greatest Latitude above the present Latitude. 5. To find the present Latitude. — As Radius to the Sine of the *Argument of Latitude*, so is the Sine of the greatest Latitude, to the Sine of the present Latitude.

To find the *Reduction*. — As Radius to *Cofine* of the greatest Latitude, so is *Tangent* of the *Argument of Latitude* to *Tangent* of an Angle, whose Difference from the former is the *Reduction*.

* * Leadbetter, in his *Astronomy*, has used *Street's Logarithms* of the Distance of the Moon from her Earth, according to her mean Anomaly, as he has also used *Street's Diameter* of the Circle of *Evection*, and also *Variation Equation*, yet has deviated from this Computation, in using *Newton's central Equation* to the greatest *Eccentricity*, instead of *Street's Eccentric*; the Maximum of the latter being 5° 1' 43" and of the former but 4° 57' 56"; being 3' 47" different. Leadbetter's *Logarithm* of the Chord of *Evection* computed to every Degree of the Moon à Sun saves no Labour; being more difficult and less certain to proportion the *Logarithm* of the Chord of *Evection* to the odd Minutes and Seconds of Moon à Sun, then to gain that *Logarithm* by adding the *Logarithm Sine* of Moon à Sun to the given *Logarithm* of the Diameter of *Evection Circle*, viz. 3.640432.

Sun. The Globe of Fire placed in the Center of our System, revolving round its *Axis* in 25 $\frac{1}{2}$ Days; and giving Light and Heat to the whole.

<i>Symbols.</i> ' (cross a trigon. Fig.) given.	× Multiplication.	√ Square Root.	T. <i>Tangent</i> .	∴ Therefore.
○ Required.	÷ Division.	S. Sine.	Cot. Cotangent or ^t .	: To.
R. Radius.	= Equal.	Cof. Cofine or S ^t .	° Degrees.	: So is.
+ More.	∠ Angle.	Sec. Secant.	' Minute.	Δ Triangle.
— Less.	∠s Angles.	Cofec. Cofecant or s ^t .	" Second. " Third, &c.	□ Square.

Synodical Anomaly. From *Syzygies* to *Quadratures* is the Moon from Sun less than 3 or 9 Signs added to the Moon's mean Anomaly 2. *equated*, by her *Eccentricity* and *Variation*. — From the *Quadratures* to the *Syzygies* is the Moon from the Sun more than 3 or 9 Signs subtracted from the Moon's mean Anomaly 2. *equated*, as before. This is according to *Street's Lunar Theory*; who determines the Moon's true Place, from her mean Place, at any Time, by Means of 3 Equations. The first of her Center, for the greatest *Eccentricity* (called the *eccentric Equation*) the *Evection-Equation* correspondent to the Difference of present *Eccentricity* from the greatest (making together the whole central Equation) and the third, the *Variation-Equation* (otherwise called the *Reflexion*) used before to equate the 1. *equated Anomaly*. Which three Equations give the Moon's Place sometimes very near the Observation. However this Method of Computation seems to want Improvement, in a new Circle of *Evection*, new *eccentric Equation*, and new *Variation*.

EXPLANATION of TERMS.

Synygies. The Conjunction or Opposition of the Sun and Moon.

T.

Time of Incidence. The Time from the Beginning to the Middle of an Eclipse. In the Moon's Eclipse it is equal to Half the Time of Duration.

Time of Repletion. The Time from the Middle to the End of a solar Eclipse.

Transit. The Passing of one Planet by, or over, another.

V.

Vector. A Line drawn from any Planet moving round the Sun, or Focus of its Orbit, describing equal Areas in equal Times.

Venus. Marked ♀, One of the primary Planets; the brightest of all. When she is West, at her greatest Elongation from the Sun, she sometimes shines extremely bright. She has her Increase and Decrease of Light from the Sun, like the Moon; and moves in an Orb, between the Earth and Mercury, round the Sun in 224 D. 16 H. 49 M. 24 S. and is never farther from the Sun than $47^{\circ} 38' 35''$. She has the least Eccentricity, but greatest geocentric Latitude. — For being in π , retrograde, her Latitude is above 8° N. and being in Υ retrograde, she has about 9° Latitude S. In every 8 Years she is seen in the same Place of the Heavens.

Vortex. A System of Particles of Matter, according to the Cartesian Philosophy, rapidly moving round like a Whirlpool; by which the French Mathematicians endeavoured to solve the Motions of the celestial Bodies: But being proved absurd, and contrary to Experience, by Sir Isaac Newton, it was exploded.

X.

Xiphias. The Sword-Fish; a Southern Constellation.

Y.

Year. The Time of the Earth's Revolution through the Ecliptic, in 365 D. 5 H. 48 M. 55 S. called tropical Year, or its Return to the same fixed Star, in 365 D. 6 H. 9 M. 24 S. called sidereal Year.

Z.

Zenith. The Point of the Heavens directly over our Heads, otherwise called the Vertex; diametrically opposite to the Nadir.

Zodiac. A Zone or Girdle surrounding the Heavens, crossing the Equinoctial, in Aries and Libra, at an Angle of $23^{\circ} 28' 30''$, equal to the Sun's greatest Declination. The Middle of which Zodiac is the Ecliptic. The Breadth of the Zodiac is $18^{\circ} 30'$; taking in the Latitude of all the Planets. It is equally divided into 12 Parts, called Signs, denominated Aries, Taurus, Gemini, Cancer, Leo, Virgo, Libra, Scorpio, Sagittary, Capricorn, Aquarius, and Pisces; being Constellations of those Names in the Zodiac.

All the foregoing Circles and Properties of the Earth and Heavens, are best seen on Mr. Cole's new improved terrestrial and celestial Globes, (which no Astronomer should be without) sold at his House in Fleetstreet, London, being Successor to the eminent Mr. Wright, Mathematical Instrument Maker to his late Majesty. Where likewise may be had all Sorts of Mathematical Instruments, Optical Glasses, &c. according to the most correct and newest Improvements.

DEFINITIONS of QUANTITY.

Def. I. Whatever Being, or Object, consists of Parts, and is capable of being augmented or diminished, is called Quantity.

Corollary 1. Hence there is Quantity of Extension, Number, Weight, Measure, Motion, Time, &c. taken more or less, heavier or lighter, longer or shorter, swifter or slower, &c. relative to the same Kind: great and small being but comparative Terms for Things of the same Kind.

2. And as the principal Property of Quantity is in its being capable of more or less, therefore Quantities may be added to, subtracted from, and multiplied by, one another, or divided into the Parts they contain.

Def. II. All Quantities have their Parts united or separated.

Def. III. Quantity having its Parts separated is called Multitude, or Number, being the Subject of ARITHMETIC.

Def. IV. Quantity having its Parts united is called Magnitude, being the Subject of GEOMETRY.

Def. V. The mutual Relation of two Quantities of the same Kind compared together, is called a Ratio; and the Similitude of Ratios of Quantities is called Proportion.

Def. VI. The Knowledge of the Comparisons, Properties, and Relations of Quantities (Ratios and Similitude of Ratios) is the Subject of MATHEMATICS; including Fluxions, Algebra, Arithmetic, and all subordinate Methods of Computation.

Corol. 1. Hence, a Part of Quantity is only considered in Relation to the whole, and may be taken as an indivisible Component of it, or for that whose farther Divisibility is not considered.

2. Unit signifies any Quantity, considered as indivisible.

3. Number is a Name of Quantity consisting of a Collection or Sum of Units, whether considered as Parts of a whole, or of several whole Quantities together.

4. Integer is a Name of Quantity considered as consisting of whole Units.

5. Fraction is a Name of Quantity considered as consisting of Parts of an Unit, or whole Quantity.

Some

Some Ancient OBSERVATIONS of Solar and Lunar ECLIPSES compared with the Computations from the Tables in the Royal Astronomer, corrected by the new Table of ACCELERATION, at P. 374. According to D. C.

1. A Total Eclipse of the Sun, June 16, Anno Chr. 364. According to Mr. Dunthorne. (See Ph. Transact. Vol. XLVI. P. 168.) at app. T. Under the Meridian of Greenwich the Moon was in Conseq. with the Sun . . .	Computation. 2 ^h 4 ^m 20 ^s 0° 0' 39" 41" II 25 9 49 II 25 49 30 Error — II 25 45 54 3' 36"
The Sun's Ecliptic Longitude by the Tables was then . . .	
Therefore the Moon's true Ecliptic Longitude was . . .	
But by the Tables thus corrected . . .	
2. A partial solar Eclipse, December 12, Anno Chr. 977, at . . .	19 ^h 12 ^m 30 ^s App. Time.
Under the same Meridian the Moon was in Anteced. to the Sun . . .	— 43' 39"
The Sun's Ecliptic Longitude by the Tables being then . . .	‡ 27 8 53
The Moon's Ecliptic Longitude observed was . . .	‡ 26 25 14 Error +
But by the Tables was . . .	‡ 26 26 12 0' 58"
3. Another partial Eclipse of the Sun, on June 8, Anno Chr. 978, at . . .	1 ^h 16 ^m 10 ^s App. Time.
Under the same Meridian the Moon was in Consequentia . . .	+ 29' 3"
The Sun's Longitude by the Tables was then . . .	II 21 53 1
The Moon's observed Ecliptic Longitude was therefore . . .	II 22 22 4 Error +
By the Tables was . . .	II 22 22 34 0' 30"
N. B. The first of these was observed at Alexandria, in Egypt, by Theon. The two last at Grand Cairo; and are the same as are quoted by Mayer in the Prolegomena to his Tables.	

The antient Astronomers were so negligent in their Observations, that hardly any can be depended on, as Dunthorne remarks, for establishing the Certainty, much less the Quantity of the Moon's Acceleration, at different Times. And yet there are a few attended with such lucky Circumstances, that they serve, as he observes, not only to establish the Hypothesis, but to settle the Quantity of the Acceleration of the Moon. I shall therefore add these, as they lie in Dunthorne's Letter to Mr. Mason, and compare the Computations from the corrected Tables with them.

1. The first is said by Hipparchus, to have been observed at Babylon, in the 366th Year of Nabonassar, the Night between the 26th and 27th Days of Thoth; when a small Part of the Moon's Disk was eclipsed from the North-East, Half an Hour before the End of the Night, and the Moon set eclipsed. This was in the Year before Christ 383, (falsely printed 313) December 22. The Sun rose then at 7 ^h 12 ^m ; and as the Moon then had South Latitude, her Setting must have been something sooner.	Computation. Beg. 7 ^h 11 ^m 30 ^s Mid. 7 58 0 Dig. Lat. 2° 12'	The Beg. by Comp. just at Sun-rise: therefore the Time by Tables about 15 ^m too late.
2. It is also said, by Hipparchus, to be observed in the 54th Year of the second Callipic Period, at Alexandria, the 16th Day of Mifson; when, he says, the Moon began to be eclipsed Half an Hour before her Rising, and was wholly clear again in the Middle of the third Hour of the Night. This was in the Year before Christ 201, September 22. The Sun set about 7 ^m after six; and the Moon being in Anteced. with N. Latitude, her true Rising had been somewhat sooner; but the Depression by Parallax, over and above Refraction, brings the Times to a near Equality.	Computation. Beg. 5 ^h 34 ^m 42 ^s End 8 42 0 Dig. Ecl. 9° 0'	These Times correspond as near as can be expected, and being not 2 Centuries after the former: the Error of that may partly arise from the computed Duration being too small, partly from the assumed Longitude of Babylon being too great.
3. The most antient Eclipse recorded, is related by Ptolemy, to have been observed at Babylon, the first Year of Mardokempad, in the Night, between the 29th and 30th Days of Thoth, in which the Moon began to be eclipsed, when one Hour after her Rising was fully passed. This Eclipse was in the Year before Christ 721, on March 19. The apparent Rising of the Moon was at 5 ^h 46 ^m . The Sun-Setting at 5 ^h 48 ^m .	Computation. Beg. 6 ^h 57 ^m 12 ^s Mid. 8 51 0 End 10 45 0	The Beginning of this also agrees very nearly with the Computation: The Words admitting of Latitude. But if Babylon should be 40', and not 50' East of Alexandria, both Computations will correspond; and the Longitude of Babylon has not been determined since Ptolemy's Days.

*. In regard to the Acceleration of the Apogee and Node, supposed by Mr. Morris, if it had been founded on Observation, the solar Eclipses in 977, and 978 were, of all others, the most likely to have discovered the Quantity, as well as Reality of it: As they happened one near the Perigeon, the other near the Apogee of the Moon's Orbit. But we see as near a Conformity of the Error of Computation as can be expected. And the Acceleration of the Moon's Anomaly being then nearly equal to 40'; it being the Sum of the Correction of the Moon's mean Motion, and twice the Acceleration of the Moon, the Equation of Longitude thence arising would be about 4' in the Perigeon New Moon; and 3' in the Apogee. Which would give the Errors of the above Eclipses, in December 977 . . . + 5' 0"

in June 978 . . . — 3 30

I do not doubt but Mayer omitted a Correction of the Apogee from the observed Disagreement with Observation.

As to the Acceleration of the Moon's Nodes, I can say Nothing to it from a Comparison of ancient Observations; but I think there is as little Reason to suppose any. S. C. — December 9, 1760.

LUNAR

LUNAR ECLIPSES.

Yrs	M.	D.	Errors	Yrs	Diff.
1628	January	10	1' 54" +	54	- 1' 17"
1682	February	11	0 37 +	54	- 0 27
1736	March	15	0 10 +	54	
1638	Decem.	10	1 36 +	54	- 0 52
1693	January	11	0 54 +	54	- 1 14
1747	February	13	0 20 -	54	
1642	April	4	1 26 +	54	- 0 52
1696	May	6	0 34 +	54	- 0 24
1750	June	2	0 10 +	54	
1681	August	18	1 11 +	54	- 0 51
1735	Septemb.	20	0 20 +	54	
1685	Novemb.	30	0 46 +	54	- 0 53
1740	January	2	0 7 -	54	

N. B. The Moon's Places are corrected by the Reduction of 6. and 7. Equations. Total 410" Divided by 8 = 51 1/8 In one Century = 95"

SOLAR ECLIPSES.

1661 D. March 19	0 8 +	54	- 0 42
1715 L. April 21	0 34 -	18	+ 0 42
1733 L. May 2	0 8 +		
1676 D. May 31	0 38 -	72	+ 0 57
1748 Gr. July 14	0 19 +		
1699 O. Sept. 12	0 9 -	54	- 0 3
1753 Co. Oct. 14	0 12 -		

Places of Observation, Dantzick, London, Oxford, Greenwich, and Gottingen.

If we take a Mean between the Differences in lunar Eclipses, and those made from Transits, the Acceleration of mean Motion for a Century, above our Tables, will be 2' 16". — The Computations from solar Eclipses prove Nothing; which I wonder at, as the Times of Beginning and End are more easily to be determined than in lunar Eclipses. I suppose the Errors of Latitude or Parallax may account, in some Measure, for it. It is surprizing too, that the Seven should agree so much nearer with Observation than the same Number of lunar ones. — But, by comparing the ancient observed lunar Eclipses, with the Tables, and fixing the Acceleration at 10" X by Sq. of the Centuries, the mean Motion is to be accelerated 2' 20" in a Century, according to the Table of Correction in P. 374. S. C.

Correction of the Index in last Page. D. Cowper, from the foregoing Observations and Remarks, made since the printing off the last Page, is doubtful whether the Apogee and Node have been accelerated, since the earliest Ages; or rather, from the near Agreement in the Errors of Computation, in the two famous solar Eclipses of 977 and 978, (the one when the Moon was near the Perigee, the other her Apogee) is inclined to think they have none at all. But is of Opinion, if the Apogee and Node be accelerated as well as the Moon, that the Theory of Gravity requires that their Acceleration should not be in the Ratio as is given by Mr. Gael Morris, (in the Astronomia Arcana, or Greenwich Tables, he has secretly printed for the Use of the PRIVATE) but in the Ratio of the annual Equations to each other, as 6, 10, and 5, very nearly. In which Proportions of Acceleration of the Moon, Apogee, and Node, the following COMPUTATIONS are made.

Two EXAMPLES of computing the MOON'S PLACE. The Mean Motions corrected by P. 374. C.

Long.	M. Anom.	Ap.	Arg.	M. Ev.
1733	8 1 3 58	6 20 25 40	8 8 27 8	7 2 35
May 13	10 12 27 38	14 49 2	7 2 35	1 12
Hours 9	4 56 28	2 30		
Mo. Sec. 33	. . . 18			
M. Motions.	6 18 28 22	7 5 17 12	8 1 23 21	7 3 47
Ann. Equat.	- 6 38	+ 15 12	- 5 28	
D, Ap. & 1 Eq.	6 18 21 44	7 5 32 24	8 1 17 53	6 20 36 50
6 Equat.	- 9 18	6 18 21 44		
D 7 Equat.	6 18 12 26	11 12 49 20	10 19 18 57	6 Eq. [D & 8]
Elliptic Equat.	+ 1 45 47	- 9 18		
D 8 Equat.	6 19 58 13	11 12 40 2	D cor. Anom.	
Evection	+ 1 16 17	+ 0 52 53	1/2 M. Ell. Equat.	
D 9 Equat.	6 21 14 30	11 13 32 55	D true Anom.	
Variation	- 37 40	1 25 54 -	2. Ann. Arg.	
D Orb. Long.	6 20 36 50	9 17 38 55	Arg. Evect.	
Reduct. + 20"	+ 7 13			
D in Eclipt.	20 44 3	No. Arguments	Equations.	
Observed	20 43 41	2 6 27 57	4 24	
Error of Computation	+ 0 22	3 6 2	3	
Halley's Error	+ 1 49	4 9 21	56	
		5 8 13	1 20	
		6 8 10	2 35	
		7 10 24		
			9 18	

The above is an Improvement in the Form of the lunar Computation by D. C. who has commodiously placed the Argument of Mean Evecton, at the Top of the Page, for taking out the left Equations. And has corrected the Mean Motions from Tab. p. 374.

63 21° 14' D
2 3 29 C
4 17 45 D & C

LUNAR COMPUTATIONS for 1690 and 1744, from Mayer's Tables, compared with OBSERVATIONS, by Flamsteed, and Bradley, at the Royal Observatory. According to D. C.

Yrs	M.	D.	Errors	Yrs	Diff.
1689	Decem.	12	0' 13" +	54	- 0' 11"
1744	January	14	0 2 +		
1690	January	12	1 2 +		- 1 49
1744	February	13	0 47 -		
1690	February	7	1 53 +		- 2 0
1744	March	11	0 7 -		
1690	February	8	2 23 +		- 2 51
1744	March	12	0 28 -		
1690	February	14	0 22 -		- 0 19
1744	March	18	0 41 -		
1690	February	19	0 6 +		- 1 6
1744	March	23	1 0 -		
1690	March	6	1 24 +		- 2 5
1744	April	7	0 41 -		
1690	April	8	2 10 +		- 2 12
1744	May	10	0 2 -		

N. B. The mean Radixes are reduced to our Radixes, in these Computations. Total - 753" ÷ 8, = - 94 1/8 In one Century = 2' 54"

The Mean of Differences gives, at an Average, 3' nearly, for a Century; but they are so very unequal that one can only lament the Carelessness of the 1st Observers.

Long.	M. Anom.	Ap.	Arg.	M. Ev.
1733	8 1 3 58	6 20 25 40	8 8 28 18	7 47 4
May 27	4 16 55 48	16 22 37	7 47 4	2 39
Hours 20	10 58 49	5 34	2 39	4
Mot. 27 S. 33	15 18	8		
M. Motions	0 29 13 53	7 6 53 59	8 0 38 31	- 7 49 47
Ann. Equat.	- 4 4	+ 9 23	- 3 22	
D Ap. & 1 Eq.	0 29 9 49	7 7 3 22	8 0 35 9	0 26 19 8
6 Equat.	- 0 19	0 29 9 49		
D 7 Equat.	0 29 9 30	5 22 6 27	4 25 43 59	6 Eq. [D & 8]
Elliptic Equat.	- 0 55 47	- 0 19		
D 8 Equat.	0 28 13 43	5 22 6 8	D cor. Anom.	
Evection	- 1 20 26	- 0 27 53	1/2 M. Ell. Eq.	
D 9 Equat.	0 26 53 17	5 21 38 15	D true Anom.	
Variation	- 34 9	2 20 42 -	2. Ann. Arg.	
D Orb. Long.	0 26 19 8	3 0 56 15	Arg. Evecton	
Reduc. + 20"	+ 6 49			
D in Eclipt.	26 25 57	No. Arguments	Equations.	
Observed	26 26 16	2 7 10 21	5 13	
Error of Computation	- 0 19	3 6 16 50	26	
Halley's Error	- 8 20	4 4 5 2	. . . 0 49	
		5 2 16 7	. . . 1 22	
		6 2 9 48	. . . 3 9	
		7 3 23 0		
			5 39	5 20
			5 20	+
			0 19	

The Moon's Latitude and Parallax are taken out of Tables, p. 52 and 53, by the proper Arguments, as shown in Examples, p. 54. Of which these two last printed Examples are an Improvement by D. C. to whom no Task is insuperable!

THE CONTENTS.

GEOGRAPHICAL TABLE.

TABLE of correct Latitudes and Longitudes of remarkable Places — from Page 1 to 2

ASTRONOMICAL TABLES OF THE SUN.

Table of annual radical mean Places of the Sun and first Star of <i>Aries</i> , O. S.	3
— annual R. M. Places of Sun and $1^{\circ} \star \gamma$, N. S.	4
— Monthly M. Motions of Sun and $1^{\circ} \star \gamma$ from 5 to 7	
— Radical mean Places of the Sun and first Star of <i>Aries</i> for Centuries	8
— Mean Motion of the Sun and fixed Stars for 99 <i>Julian</i> Years	9
— Mean Places of the Sun and fixed Stars for Years and Months, O. and N. S.	10
— Mean Motions of the Sun and fixed Stars for Months and Days	11
— Equation of the Sun's Center	12
— Decimals for proportioning the Dif. of Equat.	13
— Examples of proportioning the Dif. of Equat.	14
— Sun's proportional Distances from the Earth. Eccentricity 1685	15
— Logarithms of the Sun's proportional Distances	16
— Sun's true Place, or Anomaly, to the mean	17
— Examples of its Use	18
— Absolute Equation of Time, for 58 Years	19
— Equation of Time for Sun's true Place only	20
— Equation of Time for Sun's M. Anomaly	
— Refraction of Light for <i>Greenwich</i> Observatory	21
— Hourly Motion and Sun's app. Diameter, and Acceleration of fixed Stars	
— Decimals of a Degree for proportioning Sexages.	22
— Sun's Declination and meridian Angle	23
— Sun's right Ascension to Sun's Place	fr. 24 to 25
— Sun's right Ascension in Time	26 & 27
— Reducing Parts of Equator to Time	28
— Reducing Time into Parts of Equator	
— Refraction and Parallax, for <i>Paris</i>	29
— Reducing Parts of Equator from Noon to Noon into mean Time	
— Reducing M. solar Hours into Parts of Equator	30
— Difference of mean solar above sydereal Hours	
— Dif. between <i>Julian</i> , <i>Gregorian</i> , and <i>solar</i> Years	31
— Construction of the above	32
— Decimals of a Day	33
— M. Mot. Sun for Months and Days to Thirds	
— Equation of Points of the Equinoctial. Equat. of Ecliptic's Obliquity. Annual Precession of Equinox, according to Longitude of Moon's Node.	

ASTRONOMICAL TABLES OF THE MOON.

Table of annual rad. M. Places of Moon, O. S.	Page 34
— Annual radical M. Places of the Moon, N. S.	35
— Annual monthly M. Mot. of the Moon	fr. 36 to 39
— Radical mean Places and Motions of the Moon for Centuries, O. and N. Style	40
— Mean Motion of the Moon for 99 <i>Julian</i> Years	41
— Mean Places of the Moon for Years and Months, O. and N. S.	42
— Moon's mean Motions for Months and Days	43
— I. Equation of the Moon. C.	from 44 to 45
— II. and III. Equations of the Moon. C.	46
— IV. V. VI. and VII. Equations of the Moon. C.	47
— VIII. Equation of the Moon. C.	48
— IX. Equation of the Moon. C.	49
— X. Equation of the Moon. C.	50
— Correction of the Moon's mean Variation P. 73, with Examples. C.	51
— XI. Equation of Moon, or Reduction to Ecliptic	52
— Moon's mean Latitude. C.	
— Equation of the Moon's mean to her true Lat.	53
— Moon's mean horizontal Parallax	
— I. Equat. of Moon's M. horizontal Parallax	54
— II. Equat. of Moon's M. horizontal Parallax. C.	
— Examples of computing the Sun and Moon's true Places. C.	55
Remarks on Mr. <i>Mayer's</i> Acceleration of the Moon	56
Account of our new solar and lunar Tables	
Precepts for computing the Sun and Moon's true Places accurately	

SUPPLEMENTAL EQUATION-TABLES OF THE SUN.

Table of Equation of the Sun's Center, for Eccentricity 1692	57
— Logarithms of the Sun's proportional Distances, for Eccentricity 1692	58
— Equat. Sun's Center, to Eccentricity 1680	59
— Sun's proportional Distances from the Earth, for Eccentricity 1680	60

SUPPLEMENTAL EQUATION-TABLES OF THE MOON.

Table of the I. Equation of the Moon, Apogee, and Node	fr. 61 to 62
— II. Equation of the Moon	63
— Decimal Multipliers to the foregoing Table	
— Greatest Equation (Moon in Octants of Sun)	63
— III. Equation of the Moon	

Table

C O N T E N T S.

Table of second Equation of the Moon's Apogee, according to <i>Newton</i> . With decimal and sexagesimal Multipliers for greatest Dif. under and above the Moon's central Equation, for present central Equation	Page 64
— Second Equation of the Moon's Apogee according to <i>Horrox</i> . With decimal and sexagesimal Multipliers for the greatest Difference under and above the Moon's mean central Equation, for present central Equation	65
— IV. or mean central Equation of the Moon, according to <i>Newton</i> . With the greatest Difference of Equation (under and above the mean central Equation) corresponding to the greatest and least Eccentricities	from 66 to 68
— IV. or mean central (or elliptic) Equation of the Moon, according to <i>Horrox</i> . With the greatest Diff. of Equation (under and above the mean central Equation) corresponding to the greatest and least Eccentricities	from 69 to 71
— V. Equation of the Moon, or Moon's greatest and least Variation, according to <i>Newton</i>	72
— Decimal Multipliers for the foregoing Equation	
— Moon's greatest Variation in Octants of $\Delta\alpha\odot$	73
— V. Mean Variation Equation of the Moon	
— Logistical Logarithms for Correction of the foregoing Equation	74
— V. Entire Variation-Equation of the Moon, according to <i>Horrox</i> or <i>Flamsteed</i>	
— VI. Equation of the Moon, or second Equation of her Center. According to <i>Newton</i>	75
— VII. Equation of the Moon, or second mean Variation. According to <i>Gregory</i> .	
— Correction of <i>Gregory's</i> VII. or second Variation	76
— VII. Equation of the Moon, or second mean Variation. According to <i>Newton</i>	
— Correction of VII. Equation foregoing, or second mean Variation	77
— Farther Correct. of foregoing 2d M. Variation. According to Diff. of Δ 's Dist. fr. Earth and Sun	
— Second Equation of the Moon's Node, according to <i>Newton</i> . With decimal Multipliers and Logistical Logarithms to be applied to the greatest Excesses of Reduction and Moon's Latitude. For finding the present Reduction and Latitude. Also present Excesses above the least Inclination of the Moon's Orbit and Ecliptic, for finding the present Inclination	76
— Reduction-Equation of the Moon from her Orbit to Ecliptic	77
— Latitude of the Moon, for least and greatest Inclination of her Orbit. With greatest Excess of Latitude for finding the present Lat. Moon	
— Mean horizontal Parallax of the Moon in Conjunction and Opposition of the Sun and Moon. With greatest Difference of horizontal Parallax	78
— Decimal Multipliers for greatest Difference of horizontal Parallax. To find the horizontal Parallax for present Eccentricity	
— Correction of foregoing horizontal Parallax, out of Conjunction and Opposition of the Sun and Moon	79
— Eccentricities of the Moon's Orbit. Difference of those Eccentricities from the mean, according to <i>Horrox</i> and <i>Flamsteed</i> . With the Difference of Eccentricities from the mean, according to <i>Newton</i>	

Table of Equation of mean elliptic Equation of the Moon's Center, to the mean elliptic Equation. According to <i>Ward's</i> Hypothesis	80
Construction of constant Logarithms in following Table, for finding the Moon's central Equation	
— Constant Logarithms correspondent to the different Eccentricities of Moon's Orbit. For finding the true elliptic Equation of the Moon's Center	81
— Equation or Correction of Half Moon's equated Anomaly. For expeditiously finding (at one Operation) the true Equat. of Moon's Center	fr. 82 to 83
— Elliptic Equation of the Moon's Center in all States of her variable Orbit, according to her different Eccentricity. With the Increase or Decrease of Equation from mean Eccentricity for every 5000, Difference of Eccentricity	fr. 84 to 89
— Equation of the mean to the present true elliptic Equation of the Moon's Center. According to equated Anom. Moon, and Sun fr. Moon's Ap.	fr. 90 to 95
Use of the foregoing Equation	96
Of finding the true from the mean Anomaly universally and correctly	96

MAYER'S LUNAR EQUATIONS.

Table of Equation of Node and Moon's mean Anom.	97
— I. II. III. IV. V. Equations of the Moon	98
— VI. VII. VIII. IX. X. Equations of the Moon	99
— XI. Equation, or Equation of Moon's Center	100
— XII. Equation, of Moon's Evection	101
— XIII. Equation, of the Moon's Variation	102
— XIV. Reduction-Equation to the Ecliptic	
MAYER'S LATITUDE OF THE MOON.	
— I. Moon's mean Latitude	103
— II. Equation of the Moon's mean to her true Latitude	

MAYER'S HORIZONTAL PARALLAX OF THE MOON.

— Mean equatorial Parallax of the Moon	104
— I. Equation of Moon's horizontal Parallax	
— II. Equation of Moon's horizontal Parallax	

M. EULER'S EQUATIONS OF THE MOON.

Table of Equation of the mean to eccentric Anomaly of the Sun and Moon	105
— Sun's mean Motion and Acceleration	
— I. II. III. Equations of the Moon	106
— IV. Equation of the Moon	107
— V. VI. VII. and VIII. Equations of the Moon, with Remarks	108

DIFFERENT METHODS of solving the *Keplerian* PROBLEM.

A limited Correction of <i>Ward's</i> Hypothesis, determining the true Angle at the upper Focus	109
Examples of several Solutions, by different Methods, according to <i>Halley</i> , <i>Bullialdus</i> , and the foregoing Method, compared together	fr. 110 to 114
A Remark on the miscellaneous Tracts	114
Universal and short Method of solving the <i>Keplerian</i> Problem	96

RULES and EXAMPLES for determining other REQUISITES of ELLIPTIC ORBITS.

<i>Prop. I.</i> To determine the Distance of a Planet, or Comet, from the Sun, from the true Anomaly and Eccentricity of the Orbit given	115
<i>Prop.</i>	

C O N T E N T S.

Prop. II. To determine the eccentric Anomaly of a Planet, or Comet, from the true Anomaly and Eccentricity given. ————— 115

Prop. III. To determine the mean Anomaly and Equation correspondent, of a Planet, or Comet, from the eccentric Anomaly and Eccentricity of the Orbit given ————— 115

Prop. IV. To determine the Distance of a Planet, or Comet, from the Sun, from the eccentric Anomaly, true Anomaly, and Eccentricity of the Orbit given. ————— 116

Prop. V. To determine the greatest Equation of a Planet, or Comet, or greatest Difference between its mean and true Anomalies ————— 116

Prop. VI. To determine the Eccentricity and other Dimensions of a Planet's or Comet's Orbit from the greatest Equation given ————— fr. 117 to 118

Remarks on the greatest Equat. of Mercury fr. 118 to 119

PRINCIPLES of the PLANETS mean PLACES and MOTIONS.

Decimal Table of the Sun's mean radical Places and Motions: With Equations of the Sun's mean Places in astronomical Tables to those in others fr. 120 to 121

———— The Moon's mean radical Places and Motions. With Equations of the Moon's mean Places in astronomical Tables to those in others ————— 121

———— The mean radical Places and Motions of the Moon from the Sun.

———— Mean radical Places and Motions of the five planetary Bodies, Mercury, Venus, Mars, Jupiter, and Saturn ————— fr. 122 to 123

———— Planets mean periodical Revolutions.

———— Planets mean synodical Revolutions.

How to determine a Planet's mean Place at any Time, by its periodical Revolution, and the mean Place of that Planet for some one Time being given ————— 123

Comparison of the Elements of the Planets mean Places and Motions, according to late Improvements.

Law of the five primary Planets Motion ————— 124

PRINCIPLES of the SATELLITES Mean PLACES and MOTIONS.

Decimal Table of the radical mean Places and Motions of Jupiter's four Satellites ————— 124

Law of Motion of Jupiter's four Satellites.

Periodical Times, Distances, Magnitudes, and variable Motion of Jupiter's four Satellites. With Observations and Remarks by Dr. Pound and Dr. Bradley on the Phenomena and Properties of these Bodies ————— 125

Decimal Table of the radical mean Places and Motions of the five Satellites of Saturn. According to late Improvements.

———— Of the radical mean Places and Motions of the five Satellites of Saturn deduced from their periodic Times.

A Comparison of the Orbits, Revolutions, and Velocities of Jupiter and Saturn, with the Orbits, Revolutions, and Velocities of their Satellites ————— 126

Law of Motion of Saturn's five Satellites.

A Comparison of the periodic Times and Distances of Saturn's five Satellites from Saturn's Center, in Radii of his Ring ————— 127

b 2

Remarks, by different Astronomers, on the Phenomena and Properties of Saturn's five Satellites; and on Saturn and his Ring.

A general View of the Elements of the Theory of our planetary System ————— fr. 127 to 128

Astronomical and Chronological DEDUCTIONS and DEMONSTRATIONS.

A Table shewing, at Sight, the Number of Days from the 1st of January to any Month-day of the Year following, in a common or Leap-Year.

———— For reducing Julian Years to Days, and the contrary.

With Examples of the Use of the foregoing Tables — 129

Table of the Julian, solar, Gregorian, and syderal Years compared.

With Examples of the Use of the foregoing Table — 130

Of the Precession of the Equinoxes.

The mean annual Quantity deduced from Observation.

Of the Progression of the Aphelions and Nodes of the Orbits of the Planets in respect of the fixed Stars. And of the Progression of the Aphelion of the Earth's Orbit.

The Quantity of that Progression deduced ————— 131

Of the Inequalities of the Earth's Motion, caused by the Gravitation of the Moon.

Rules for determining the Quantity of Precession, Nutation, and Changes thereby of Declination and right Ascension, in respect of the fixed Stars.

Present Obliquity of the Ecliptic deduced from Observation ————— 132 to 133

Of the MEASURE of TIME.

The syderal and solar Day.

An astronomical or solar Day.

The syderal Day correctly deduced.

To regulate a Clock or Watch ————— 134 to 135

EQUATION of TIME.

Mean and true Days defined.

Examples of the Equation of Time ————— 136 to 137

NATURE and DIVISION of TIME.

Time measured by Motion.

Variable solar Year.

Mean astronomical or solar Year, called also the tropical Year.

Near syderal Year.

Near anomalistical Year.

Correct syderal Year.

Correct anomalistical Year.

Mean lunar Year.

Civil Year.

Mean Julian and mean solar Year compared.

The falling back of the Seasons before the Style was altered.

The Quantity deduced. ————— 138 to 139

Alteration of Style, how ————— 140

Difference

C O N T E N T S.

Difference of Years.	
Civil, solar, and lunar, complete, vacant, and <i>Roman</i>	
Year, described.	
Difference of Months.	
Astronomical, civil, periodical, synodical, and antient.	
Division of the Year by the Antients into	
<i>Jewish, Egyptian, Arabic or Turkish, and Grecian</i> Months.	
A common Month. A Week.	140 to 141
Day, natural, artificial, astronomical.	
Hours, equal, unequal.	
Minutes, Seconds, Scruples.	
Cycles and Periods.	
Sun's Cycle, Moon's Cycle, and Indiction	142

Of the FIXED STARS.

Appear bigger to the naked Sight than through a Tele-	
scope.	
Made not on Purpose to give Light in the Night.	
Readily known.	
Constellations, a Table of their Names	143 to 144
Different Properties of fixed Stars.	
Milky Way, lucid and cloudy Stars, <i>Megalennic</i> Clouds.	
Changes of, and periodical Stars.	
Changes in the Heavens.	
Ecliptic Obliquity changes	144 to 145

TABLE of Chronological *ÆRAS* of great EVENTS;

According to the Year of <i>Julian</i> Period of Creation, and	
of <i>Christ</i> .	
An Account, and Examples of, the Use of <i>Æras</i> , Cycles,	
and Periods.	
The Birth and Death of <i>Christ</i> .	
Sir <i>Isaac Newton</i> 's Method of fixing the <i>Æras</i> of Chrono-	
logy, and particularly the Time of the <i>Argonautic</i> <i>Æra</i> ,	
by the advanced Places of the Constellations	146 to 147
Chronological Tables, shewing, at Sight, the dominical	
Letter, or Letters, for any Year before <i>Christ</i> , O. S.	
For any Year since <i>Christ</i> , O. S.	148
For any Year before <i>Christ</i> , according to N. S.	
For any Year since <i>Christ</i> , according to N. S.	
Shewing by the dominical or <i>Sunday</i> Letter of ei-	
ther Style, the Day of the Week answering to any Day	
of the Month in that Year; for ever	149
Shewing the Golden Number, at Sight, for any	
Year before or since <i>Christ</i>	150
Shewing, at Sight, the near Place or Sign of the	
Moon, by the Golden Number for the Year and Day	
of the Month given, N. S.	151
Shewing, at Sight, the Moon's Age, for any Day	
of the Month given, N. S.	152
The Sun's Place through the Year, by which the	
Moon's Place may be found, and her Rising and Setting	
nearly for <i>London</i> , or any other Part of the World.	
A Table of the Time of high Water at several noted Pla-	
ces before or after high Water at <i>London</i>	153
A chronological Table, to find, at Sight, the Time of the	
<i>Paschal</i> , or <i>Easter</i> Full Moon, and consequently the	
Time of <i>Easter</i> for ever, by the Golden Number and	
Days that the lunar Cycle of 19 Years, goes forward	
in any given Period, New Style.	154

A Table of the Number of Days, at Sight, the New,	
Full, or Quarter Moons, in the <i>Gregorian</i> Account, go	
forward in the Months; or how much the lunar Cycle	
of 19 Years advances in the Month-day, from 1600	
Beginning	154
Shewing, at Sight, the moveable Feasts by the	
dominical Letter and Golden Number.	
Shewing, at Sight, the moveable Terms, by the	
dominical Letter and Golden Number	155
Shewing, at Sight, the moveable Feasts and	
Terms for ever, by the dominical Letter and Golden	
Number, for old Style — Of Use in antient History	
and Chronology.	
A general Table of the Sun's Rising and Setting in <i>Eng-</i>	
<i>land</i> and <i>Ireland</i> , according to new Style	156
Remarks on <i>Æras</i> , or Points of Time.	
Controversies concerning <i>Æras</i> .	
Remarks on the Nature and Measure of Time	157
Of Rest and Motion of Bodies	
Of antient and modern Hypotheses	158

PRACTICAL QUESTIONS in CHRONOLOGY,

Of sundry Kinds, answered	159 to 166
To turn Time into Degrees and Parts of the Equator, and	
the contrary, arithmetically	167
A Table to find, at Sight, the Number of Days advanced	
and retreated by the <i>Epaets</i> and <i>Lunations</i> , in the	
Month-days, (from 1600 before to 6000 Years since	
<i>Christ</i>) from old to new Style from 1800, and the con-	
trary.	
For determining the <i>Epaet</i> and Fall of <i>Easter</i> , according	
to new Style, as settled by Pope <i>Gregory</i> . Also to find	
the <i>Epaet</i> , O. S. for the same Centuries	168
Examples of the Use of the foregoing Tables	168 to 169
Chronological Remarks and Improvements.	
Chronological Scale.	
Chronological Rules, arithmetically applied to Practice in	
answering many Kinds of chronological Questions;	
such as finding the dominical Letter, Week-day answer-	
ing to a given Month-day, the Golden Number, Cycle	
of the Sun, <i>Roman</i> Indiction, <i>Epaet</i> , &c. for new and	
old Style, before and since <i>Christ</i> , &c. With memo-	
rial Couplets to assist the Memory; and small chrono-	
logical Tables added for the same Purpose	170 to 177
A Table of Calends, Ides, and Nones, according to the	
<i>Roman</i> Year. Of Use in History.	
An Account of the old and new ecclesiastical and civil	
Calendar.	
With several other chronological Matters decided	178
	to 179
PERPETUAL TIME-TABLES, or GENERAL	
RULES for CHRONOLOGISTS, HISTORIANS,	
NAVIGATORS, &c.	
Determining, at Sight, the Points of Time for any one	
Season of the Year, past, present, or to come, particular-	
ly useful for all Persons writing Almanacs for the <i>Sta-</i>	
<i>tioners</i> Company.	
Rules for finding the moveable Feasts and Holidays in	
any Year	180
	Names

C O N T E N T S.

Names of all the fixed Feast-days, observed in the Church of England.

Names of all the Vigils and Fast-days.

The Days of Fasting and Abstinence.

Solemn Days kept, for which particular Church-Service is appointed — — — 180

Table of moveable Feasts for ever, according to the Fall of *Easter*, N. S.

— Moveable Terms for ever according to the Fall of *Easter*, N. S. — — — 181

— Moveable Feasts from 1759 to 1800, N. S. 182
Perpetual Tables, of the Month-day and Week-day correspondent, according to the dominical Letter; with the fixed Feasts, Holidays, remarkable Days, &c. according to N. S. for any Month in the Year.

Month-day of New, Full, and Quarter Moons for any Month in the Year, according to the Golden Number 183 to 188

RULES for the proper USE of the RIGHT ASCENSION and DECLINATION of the SUN and STARS, their Meridian Altitudes and Height of the Pole, in *astronomical* OBSERVATIONS.

Particularly useful in Navigation 189 to 191

Table of the apparent Semidiameter of the Sun, Time of his passing the Meridian, and his Distance from the Earth for every 10th Day of the Leap-Year, 1756, N. S. — — — 191

A Solar Ephemeris, commencing 1756, and serving, by Means of small Equations, till 1800.

Being a monthly Table of the Time of the Day by a regulated Clock when it is Noon by the Sun: Shewing the Equation of natural Days, and Advance and Retreat of Clock-Time for Leap-Year.

— The Sun's Place, Declination, and Right Ascension, for each Day of the Month in the Year 1756; by which, and small Equations, his Place, Declination, and Right Ascension are easily determined for any future Day of the Year, till 1800. 192 to 196

TABLES in ASTRONOMY.

I. A Catalogue of 140 remarkable FIXED STARS, with their Magnitudes, Longitudes, and Latitudes, N. S. According to Dr. *Halley* — — — 197

II. Catalogue of 69 remarkable fixed Stars, with their Right Ascension in Time, Degrees, and also Declination N. or S. According to the Foreigners 198

Equation-Table of the middle Time in Seconds, between the Times (by a Clock or Watch) of Forenoon and Afternoon, equal Altitudes of the Sun. Correspondent to the Increase or Decrease of the Sun's Declination during the Interval of Observation — 199

Table of semi-diurnal or semi-duration Arcs, for finding the Rising and Setting of the Sun and Stars 200 to 203

Table of the Sun's Amplitude of Rising and Setting; for finding the Variation of the Compass 204 to 207

III. Catalogue of 307 eminent fixed Stars, in the Northern and Southern Hemisphere: With their Places in the

Constellations, correct Right Ascension, Declination for the Year 1750, and their annual Variation. According to the last improved Observations 208 to 213

Small Equations of Clock-Time, and of the Sun's Place and Declination, for every 4th Year from 1750, (or from any common, or Leap-year after) to 1800, N. S. Which will serve, with contrary Sign, back to 1700, N. or O. Style. According to M. *De la Caille's* Motions — — — 214

Equations of the Sun's Right Ascension in Time for every 4 Years.

Sun's apparent Diameter for every 5th Day in the Year 1760.

Equation of the horizontal Semi-diameter of the Moon, to her apparent Semi-diameter in Altitude.

Ecliptic Obliquity for given Years — 215

Table of Clock-Time at Noon by the Sun.

Sun's Right Ascension, Complement of Right Ascension, or Distance of *Aries* from the Meridian, in Time. For each Month and Day in the Year. Reduced to the Meridian of *Greenwich* Observatory 216 to 217

LIII. Principal ASTRONOMICAL QUESTIONS applied to NAVIGATION; practically answered by spherical Proportion and Logarithms 218 to 250

The Nature and Cause of Twilight — 224

XXXIX. *Question.* To find, the apparent Place of the Moon, from her Longitude, or true Place, and Latitude given.

Wherein D. *Cowper's* new and short Method of finding the **Parallactic Angle** is exemplified. 231

Rules of applying the Moon's Parallaxes 233

Equation-Table of the Ecliptic Right Ascension to the Right Ascension with Latitude.

Equation to be subtracted from the Latitude of a Planet, for the Equation of the ecliptic Declination to the Declination with the Planet's Latitude — 251

Table of the Longitude and Right Ascension of the Mid-heaven, with the near Longitude of the nonagesimal or 90° Degree from the Horizon, correspondent in Latitude of *London* — — — 252

Table for determining the near Time in any Day, when the Sun, Moon, a Planet, or Star, is in the highest Point of the ecliptic Latitude of *London* 253

Rules and Examples to find when the Sun, Moon, Planets, or Stars, will be in the nonagesimal Degree 254 to 255

Sexagesimal Tables for proportioning the Difference of Equations at Sight — 256 to 259

Table of the visible Eclipses of the Sun, and Phenomena, for 21 principal Cities of *Europe*.

The visible Eclipses of the Moon, and Phenomena for *Greenwich* Observatory.

Likewise the invisible Eclipses of the Sun and Moon 260

The New, Full, and Quarter Moons or Lunations for *Greenwich* Observatory to the Year 1765 261

Table of the semi-diurnal Arcs of the Sun to every Degree of his Longitude in the Ecliptic, for *London*.

— Semi-diurnal Arcs of the Moon to every Degree of her Longitude in the Ecliptic, at Southings, without Latitude, for *London* — — — 262

Table

C O N T E N T S.

Table shewing, by the Moon's Age, the near Time of her Rising and Setting in <i>England</i> or <i>Ireland</i> ; when she is in the Beginning, Middle, or End of each Sign in the <i>Ecliptic</i> (supposing her without Latitude) her mean Place and Age being known at Sight, by <i>Tab. P.</i>	241, 252	262
Equation of the true horizontal Diameter of the Moon, to the apparent horizontal Diameter, according to the Moon's Declination.		
Equation of the horizontal Diameter of the Moon to the apparent in Altitude.		
Reduction of <i>Paris</i> mean Places to <i>Greenwich</i> Places	263	
A Table shewing the proportional Difference to be added to or subtracted from the Sun, Moon, or Planet's Places, Declination, Right Ascension, &c. at a first Meridian for the respective Place, Declination, Right Ascension, &c. under a distant Meridian	264	
Equation of the Difference of Right Ascension, in Time, at Noon, between the Sun, Moon, Planets, and Stars, to the nearer Time of their respective Southings at <i>Greenwich</i>	265	

LOGISTICAL LOGARITHMS.

Their Nature and Use	266
Table of Logistical Logarithms	267 to 272

The SEAMAN'S READY COMPUTER.

An Improvement in Tabular Navigation	273 to 280
Use of the Seaman's Computer exemplified	281 to 287
Our new Method of Computation, without Scale, Compasses, Chart, or Canon, exemplified in Answers to the five Cases of Sailing, proposed by Mr. <i>Martin</i> , in his new Principles of Navigation; according to Sailing by the Spheroid	283 to 287
Table of meridional Parts on <i>Mercator's</i> Chart, according to the Sphere.	
Table of meridional Parts according to the Spheroid	288 to 293
The Length of a Degree, or Number of Degrees, being elliptic Arcs of the Earth's Meridian in Minutes of the Equator, and Hundredth Parts for any Latitude. According to Don <i>Juan</i> .	
Length of the Meridian Arcs of the Sphere by Mr. <i>Murdoch</i> , compared with the foregoing Arcs of Don <i>Juan's</i> Spheroid	294
Dimensions of the Earth as a Spheroid considered, from	295 to 301
Properties of right-angled plane Triangles.	
Properties of Sines, Tangents, &c.	302
Answers to all the Cases of plane Triangles in Symbols and Words	303
Rules and Proportions for answering all the Cases of Sailing	304
Tabular Rules, in one View, to answer all the Cases in Sailing	305
Examples in answering the principal Cases of <i>Mercator</i> Sailing	306 to 307

Of correcting a Ship's Course steered by the Compass.	
Correction of a Ship's Course by the magnetic Variation.	
Magnetic Variation from 1580 to 1759.	
Farther Correction by allowing for Lee-Way	307
General Rules of Lee-Way from the Sails set or furled.	
Correction of the Course and Distance, by trying the Swiftness and Setting of a Current	308
Of Measuring the Ship's Way.	
Of the Seaman's Compass, Log-Line, Log, Half-minute Glass, and Log-board	309
Example of a Log-board.	
Seaman's Compass.	
Day's Reckoning on Board the <i>Dreadnought</i>	310
Rules for correcting a Ship's daily Reckoning	311
Remark on the foregoing Method of correcting a Ship's dead Reckoning.	
Remarks on Mr. <i>B. Martin's</i> new Navigation, in a second Letter from a Correspondent	312
Dimensions and Weight of Shot.	
Place of Turning to Windward.	
To compute the Distance of a Head-Land, or Cape, at Sea	313
To measure the Tonnage, or Burthen of a Ship.	
Best Position of the Sails, and Rudder, in working a Ship	314
The Properties of Magnetic Variation.	
Rules for determining the Quality and Quantity of Variation	315
Practical Rules for finding the Latitude of a Ship's Place of Observation	316
Geographical Rules for finding the tabular Difference of Longitude.	
Remarks on the different Beginnings of terrestrial Longitudes	317
A Journal of a Voyage in the <i>Dreadnought</i> Man of War. From <i>England</i> to <i>Martinico</i>	318 to 319
Nearness of Sailing to the Wind; and Rules for resolving a Ship's Traverse geometrically; by the Line of Chords, with Scale and Compass	320
Of Trade Winds and Monsoons	321 to 322
Variation of the Compass, as observed at different Times and Places.	
Remarks on the vulgar Error in Books of Navigation of supposing meridional Distance and Departure to be alike, and of fixed Quantity	323
Table shewing the Miles and Minutes of the Equator, making a Degree of Longitude in any Latitude.	
Division of the Earth and Ocean	324
General Affection of spherical Triangles	325
Affections of right angled spherical Triangles.	
Some Properties of Sines, Tangents, &c.	326
Answers to all the Cases of right-angled spherical Triangles, in Symbols and Words	327
Answers to all the Cases of oblique spherical Triangles	328 to 329
The Properties of spherical Triangles	330 to 331
Some general Observations on the Tides.	
To find the Time of High Water, at a given Time and Place	332
An alphabetical Tide-Table from the best Authorities: shewing the Time of high Water at the most remarkable Sea-Ports, and Sea-Coast Places, at New and Full Moon	333 to 334
Observations	

C O N T E N T S.

Observations on the different Beginnings of terrestrial Longitudes.

An astronomical Question of Use in Navigation.

Spherical Questions answered — 335

Observations on the Answers to *Martin's* or *Murdoch's* Examples.

Variation of Case 6th of *Mercator-Sailing*.

Analytically answered — 336

The Mariner's Universal Sea-Coast Table. Of the principal Ports, Harbours, Head-Lands, Bays, Rivers-Mouths, and Islands, lying contiguous, or nearly so, round the World.

With their Latitudes, North and South, and Longitudes East from *Ferro* Island. Also their Difference of Longitude East or West from *Greenwich* Observatory 337 to 344

An alphabetical Table of the Latitudes of promiscuous and remarkable Places, with their Longitudes, from the Island of *Ferro*, and Difference of Longitude from *Greenwich* Observatory — 345

Captain *Brown's* Directions for the Safety of homeward bound Ships, coming into the *English* Channel. 346

REDUCTION of CHRONOLOGY.

Commencement of the most famous *Aeras* in Years current and complete, before and since *Christ*, *Julian* Style — — — 347

Table for Reduction of Chronology.

Table for reducing *Julian* Years into Days, and the contrary.

— Reducing *Turkish* or *Arabic* Years into Days and the contrary.

Several Examples of the Reduction of Chronology. 348 to 355

INTRODUCTION to the COMMENT on the MOON'S THEORY.

An explanatory Comment on the Moon's Theory. According to which Principles the astronomical Tables at the Beginning of our Work (preceding the Supplemental Tables) are constructed 356 to 357

Of determining the Quantity of Evection-Equation 371

Table of the Moon's Longitude computed from the Tables to a 100 Places, and compared with the respective Observations — — 372 to 373

Total Correction of the Moon's mean Motion. According to the Correction of her mean Motion, and Acceleration allowed — — — 374

Table for correcting the Sum of the VI. and VII. Equations of the Moon, at the Beginning of the Tables. C.

Comparison of the Number, Arguments, and greatest Equations of the Moon — — 375

Method of determining the *Maxima* of the several lunar Equations (C.) at the Beginning of our Work 376 to 377

Astronomical Tables equated to others 377

Explanation of *Newton's* lunar Theory — 378

Explanation of *Machin's* lunar Theory — 379

Of *Machin's* Theory of the Moon 379 to 383

Of *De la Caille's* Theory and physical Rules of the Moon's Motion — — 383 to 384

OF LUNATIONS and ECLIPSES.

Table of the mean Time of the first mean New Moon in *January*, from 1700 to 1800, O. S.

Places of the Sun's mean Anomaly, annual Argument, and Sun from Moon's Apogee, corresponding.

Monthly Motions of the same — 385

Revolutions of the Moon to the Sun, next after *Julian* Centuries — — — 386

Plinian Period of solar and lunar Eclipses.

Equation of Time for a *Plinian* Period 387

Radical mean Places and Motions of the Moon from the Sun; for determining the mean Time (by Means of Equations) of the true Orbit or Ecliptic New and Full Moon — — — 388

Mean Motions of the Moon from the Sun, &c. for Months, Days, Hours, Minutes, and Seconds 389

Equation of the Moon's Center (and second Equation supplemental) compounded, at New and Full.

The Moon's Latitude, Inclination, and Reduction, at New and Full — — — 390

The true hourly Motions, Diameters, and Parallaxes of the Sun and Moon in Eclipses.

Times between the true Ecliptic Conjunction or Opposition of the Sun and Moon, and Middle of Eclipse 391

Angles which the Way of the Moon from the Sun makes with a Circle of Latitude, in Eclipses — 392

Examples of computing the Times and Places of the New and Full Moons in Eclipses — 392 to 397

Precepts for computing the Times and Places of the Orbit New and Full Moons — 393

What an Eclipse of the Moon is — 397

Precepts for computing an Eclipse of the Moon.

What an Eclipse of the Sun is.

Precepts for computing an Eclipse of the Sun, by Parallaxes — — — 398

To find the Places, on the Earth, of the principal Appearances of solar Eclipses, by Projection 399

Method of computing the greatest Number of Eclipses of the Sun and Moon that can happen in one Year.

Of the *Plinian* Period of Eclipses, and their Returns.

Of antient Eclipses.

Plurality of Eclipses.

Historical Eclipses of the Sun and Moon, at great Events. 400

Reduction of astronomical Tables, with Examples 401

Geometrical Construction of a lunar Eclipse 402

Table of the Angle of the Hour Circle (passing through the Sun and Moon) with the Ecliptic 403

Examples of computing the parallactic Angle, according to different Methods.

Explanation of different parallactic Angles 404

Precepts of the lunar Computation.

To find the Moon's true from her apparent Place.

Remark on Dr. *Halley's* Tables — 405

Of *Halley's* Method of determining the Longitude by Observation.

Of *De la Caille's* Method. 406 Of

C O N T E N T S.

Of observing the Longitude at Sea by the Moon's Theory	406
Account of a Pamphlet pretending a Discovery of the Longitude.	
Example of computing the Difference of Longitude from Greenwich	407
A new Method of making a lunar Ephemeris.	
Of the Motion of Comets	408
Example of lunar Computation, according to D. C.	409
Example of Computation. According to Mayer.	410

Correction of the Longitude of the Sun, and Sun's Distance from the Earth. According to Mayer	410
A Table and Example for the Reduction of a Comet's Longitude from the Perihelion in a parabolic, to that from the Perihelion in an elliptic Orbit, by the Difference of Motion in those Orbits	411
Computation of an Opposition of the Sun and Moon, with their Places.	
CONCLUSION, OF LAND	412

CORRECTIONS of literal and typographical ERRORS observed; to be made with the Pen, before our WORK is used.

N. B. *Most of the following Errors are obvious and trifling; but are all recited to prevent any being overlooked.*
Where b is placed after the Number of a Line, it signifies the Number of that Line, from the Bottom of the Page.

- ✓ **PAGE 1.** Line 6. b. For $0^h 11^m 16^s$ and $20^s 49' 4''$.
Read $0^h 7^m 16^s$ and $1^s 49' 4''$.
- ✓ L. 23. b. For Lima $86^{\circ} 49' 0''$ r. $76^{\circ} 49' 0''$.
- ✓ P. 2. L. 1. For Nantz $0^h 24^m 55^s$ and $60^s 13' 48''$ E.
Read $0^h 6^m 56^s$ and $1^s 44^s 0''$ W.
- ✓ L. 14. For Porto Bello $89^{\circ} 50' 6''$ r. $79^{\circ} 50' 6''$.
- ✓ P. 11. L. 16. For October 1 r. October 5.
- ✓ P. 23. L. 28. For $67^{\circ} 30' 31''$ r. $67^{\circ} 40' 31''$.
- ✓ P. 24 to 27. Right Hand Col. For $\odot$ Pl. in Degr. r. its Comp. to 30° ,
or invert in writing the Degrees in that Column by Paper pasted
over its Length and Breadth.
- ✓ P. 72. L. 5. b. For i. e. $7, 24 \times$ any L. Eq. = G. D. r. i. e. $1206 \times$ any L. Eq. = G. D.
- ✓ P. 73. L. 8. b. For 2. $\text{D} \hat{=}$ $\odot$ (from next Quadrature or Syzygy) &c. r. S. 2. $\text{D} \hat{=}$ $\odot$ (from next Quadrature or Syzygy) : S. Pref. M. Variation, &c.
- ✓ P. 78. L. 24. b. under Col. 7. For $+$ r. $+$.
- ✓ P. 97. L. 18. Col. 2. For $0^{\circ} 38' 35''$ r. $0^{\circ} 28' 15''$.
- ✓ P. 100. L. 31. Col. 8. For $60^{\circ} 59' 38''$ r. $50^{\circ} 59' 38''$.
- ✓ P. 102. L. 7. Col. 8, 10, 12. For $+$, $+$, $+$, (at Top) r. $-$, $-$, $-$.
- ✓ P. 119. L. 2. For elliptic Trajectories, r. elliptic Trajectories.
- ✓ L. 10. For Where the mean and true Motion is equal, r. Where the mean and true Anomalies are equal.
- ✓ P. 120. For Com. $\left\{ \begin{array}{l} 1. +14 19 45 \\ 2. +28 39 31 \\ 3. +42 59 17 \end{array} \right\}$ r. Com. $\left\{ \begin{array}{l} 1. -14 19 45 \\ 2. -28 39 31 \\ 3. -42 59 17 \end{array} \right\}$ Yrs.
- ✓ L. 34. b. Yrs. $\left\{ \begin{array}{l} 1. -14 19 45 \\ 2. -28 39 31 \\ 3. -42 59 17 \end{array} \right\}$ Yrs.
- ✓ P. 121. L. 8. Col. 3. For $+97017925237$, r. $+9701786292$, &c.
 $29^{\circ} 6' 19''$ in a Lunat.
- ✓ L. 28. Col. 1. For, from 1600 and 1700 reduce to, r. from 1700 reduce to.
- ✓ L. 22. Col. last. For $+4' 3''$, r. $4' 1''$.
- ✓ L. 33. Col. last. For, from 1600 and 1700, r. from 1700.
- ✓ **P. 122.** L. 1. b. Col. 3. For $21^m 36iv$, r. $21iv 36v$.
- ✓ 4. b. same Col. $8^m 38^m 24iv$ $8^m 38iv 24v$.
- ✓ 1. b. Col. 5. $4^m 6iv$, &c. $4^m 0iv$, &c.
- ✓ 1. b. Col. 6. $23^m 15iv$ $23iv 15v$.
- ✓ 4. b. same Col. $9^m 18^m$ $9^m 18iv$.
- ✓ 1. b. Col. 7. $12^m 45iv$ $12iv 45v$.
- ✓ 4. b. same Col. $5^m 6^m$ $5^m 6iv$.
- ✓ 10. b. same Col. 31^m 31^m .
- ✓ **P. 123.** L. 21. Col. 3. $11^m 31^m 12iv$ $11^m 31iv 12v$.
- ✓ 24. same Col. $28^m 48iv$ $28iv 48v$.
- ✓ 21. Col. 4. $6^m 14^m 24iv$ $6^m 14iv 24v$.
- ✓ 24. same Col. $15^m 35iv$ $15iv 35v$.
- ✓ 21. Col. 6. $11^m 47^m 12iv$ $11^m 47iv 12v$.
- ✓ 24. same Col. $29^m 32iv$, &c. $29iv 32v$, &c.
- ✓ 21. Col. 7. $8^m 13^m 12iv$ $8^m 13iv 12v$.
- ✓ 24. same Col. $20^m 33iv$ $20iv 33v$.
- ✓ 1. b. Col. 3. $32^m 57iv$ $32iv 57v$.
- ✓ 3. b. same Col. $12^m 8^m$ $12^m 8iv$.
- ✓ 1. b. Col. 4. $7^m 20iv$ $7iv 20v$.
- ✓ 3. b. same Col. $2^m 56^m 24iv$ $2^m 56iv 24v$.
- ✓ **P. 124.** L. 2. b. Col. 2. For $1^d 12^h 27^m$, &c. r. $1^d 18^h 27^m$, &c.
- ✓ P. 127. L. 7. b. Col. last. For $8 16^{\circ} 6' 29''$, r. $8 14^{\circ} 37' 27''$.
- ✓ P. 146. L. 11. Col. last. For 44 Yrs bef. Chr. r. 45 Yrs bef. Chr.
- ✓ L. 1. b. For, from 4713, r. from 4714.
- ✓ P. 147. L. 4. For 4713 Year Jul. Period, r. 4714, &c.
- ✓ L. 5. For Dif. 2954, r. Dif. 2955.
- ✓ P. 165. L. 10. b. For divisible by 4, r. divisible by 2.
- ✓ P. 170. L. 16. Col. 1. For requires, r. require.
- ✓ P. 172. L. 16. Col. 2. For Years since Christ, r. Years before Christ.
- ✓ L. 29. same Col. For Periods so happened, r. Period so happened.
- ✓ P. 193. L. 1. below the Tables. For correcting or beginning, r. commencing or beginning.
- ✓ P. 231. L. 18. b. Col. 2. For, from Cosine Declination for 5th Arc, r. from polar Distance for the 5th Arc, &c.
- ✓ P. 234. L. 29. Col. 2. For, To the No. of Earth's Diameters, r. To the No. of the Earth's Semi-Diameters.
- ✓ P. 237. Midd. Col. 2. For 1.1287, r. 1.2187.
- ✓ L. 7. lower. For $3^h 12$, r. $3^h 67$.
- ✓ P. 263. L. 1. under Tab. Col. 1. For true Semi-Diam. r. true Diam.
- ✓ P. 281. L. 28. Col. 1. For, Middle Latitude be less than 45° , r. be greater than 45° .
- ✓ L. 29. — For, if greater than 45° , r. if less than 45° .
- ✓ P. 297. L. 11. b. For, as a Means for the Defect, r. as Amenis made for the Defect.
- ✓ P. 312. L. 9. For 1 Mile in 10 less, r. 1 Mile in 10 more.
- ✓ P. 317. Third of 3d Col. Rule 2. Diff. Long. $5^{\circ} 54' W$. r. Dif. Long. $5^{\circ} 14' W$.
- ✓ P. 323. L. 3. Col. 1. For Buffin's Bay, r. Baffin's Bay.
- ✓ P. 324. L. 25. Col. 2. For, and some Islands, r. and some other Islands.
- ✓ L. 3. b. For, From the Divisions, r. From the Division.
- ✓ P. 326. L. 24. For less } than 180° . r. less } greater } than 90° .
- ✓ L. 29. b. For Rad. = 10, r. Rad. = 1.
- ✓ P. 328. L. 6. Col. 7. For, Sum and Dif. r. Sum or Dif.
- ✓ P. 359. L. 12. b. For 1,000 to 178725 r. 1000 to 178725.
- ✓ L. 9. b. For 178725, r. 178,725.
- ✓ L. 5. b. For, at the same Time, r. in the same Time.
- ✓ P. 360. L. 5. b. For, gravitate more towards, &c. r. gravitate more or less towards the Sun than the Earth.
- ✓ L. 3. b. After KT — introduce, by which the Moon is all attracted towards the Earth, in the former Case, &c.
- ✓ P. 372. Obs. 16. For $6^s 29^{\circ} 29'$, r. $5^s 29^{\circ} 29'$ $\text{D} \hat{=}$ $\odot$.
- ✓ P. 376. L. 12. b. Col. 2. Page. For 357,45, r. 356,50.
- ✓ L. 4. b. Marg. For 112,26, r. 119,26.
- ✓ P. 390. Col. 8. $3^s 16^{\circ}$. . for $3^{\circ} 40' 21''$, r. $4^{\circ} 40' 21''$.
- ✓ P. 397. L. 18. Col. 1. For, Ecciptic Conj. r. Ecciptic Opposition.
- ✓ L. 18. Col. 2. For, Ecciptic Conj. r. Ecciptic Opposition.
- ✓ L. 21. same Col. For, Ecciptic $10^m 43^s$ Dif. r. $10^m 43^s$ Dif. blot Ecciptic.
- ✓ L. 3. For Examp. xi, r. Examp. ix.
- ✓ L. 10. b. For 10 6 41 21, r. 10 7 41 21.

Any material Errors contained, escaping our Observation, we shall be obliged to the Men of Judgment and Critics to point out. So far from our taking such Notice amiss, that we hereby offer our Thanks to any, who shall be pleased to detect our Mistakes, (paying the Postage of their Letters franking them, or publicly informing us) that TRUTH (which is our principal View) may be promoted.

THE ROYAL ASTRONOMER AND NAVIGATOR.

A GEOGRAPHICAL TABLE of many eminent and remarkable PLACES in the WORLD, with their correct temporary Difference of Meridians and Longitudes from GREENWICH OBSERVATORY; and their correct Latitudes.

Places Names.	Dif. Time. After or Bef.	Dif. Longit. East or West.	Latitude North or Sou.	Places Names.	Dif. Time. After or Bef.	Dif. Longit. East or West.	Latitude North or Sou.
h m s	o / //	o / //	o / //	h m s	o / //	o / //	o / //
Athens, in Greece,	1 35 3 A	23 52 30 E	38 5 0 N	Dantzick, Poland,	1 15 12 A	18 48 0 E	54 22 0 N
Alexandria, Egypt,	2 1 6 A	30 16 30 E	31 7 0 N	Dunkirk,	0 9 30 A	2 22 23 E	51 2 4 N
Alexander, Syria,	2 25 20 A	36 20 0 E	36 35 10 N	Edinburgb,	0 12 21 B	3 5 15 W	55 58 0 N
Araçta, or Racca,	2 35 20 A	38 50 0 E	36 1 0 N	Florence,	0 45 18 A	11 19 30 E	43 46 30 N
Agra, in Mogul,	5 6 56 A	76 44 0 E	26 43 0 N	Francfort,	0 34 20 A	8 35 0 E	49 55 0 N
Aleppo, Syria,	2 29 20 A	37 20 0 E	35 45 23 N	Goa, East-Indies,	4 55 0 A	73 45 0 E	15 31 0 N
Algiers,	0 8 51 A	2 12 45 E	36 49 30 N	Geneva,	0 36 20 A	6 20 0 E	46 12 0 N
Amsterdam,	0 19 56 A	4 59 0 E	52 22 45 N	GREENWICH Observ.	0 0 0 A	0 0 0 E	51 28 30 N
Antibes,	0 28 34 A	7 8 33 E	43 34 50 N	Jerusalem,	2 21 20 A	35 20 0 E	31 55 0 N
Avignon,	0 19 14 A	4 48 33 E	43 57 25 N	Ingoldstadt,	0 45 30 A	11 22 30 E	48 46 0 N
Babylon, Asia,	2 51 6 A	42 46 30 E	33 0 0 N	Ispahan, Persia,	3 31 20 A	52 50 0 E	32 25 0 N
Babylon, Egypt, G. Cairo,	2 5 45 A	31 26 15 E	30 2 30 N	Kébec, or Quebec, Cana.	4 58 12 B	73 33 0 W	46 55 0 N
Berlin, Prussia,	0 53 50 A	13 27 30 E	52 33 0 N	Lima, Peru,	5 7 18 B	76 49 0 W	12 1 15 S
Batavia, East-Indies,	5 44 0 A	86 0 0 E	21 20 0 N	London,	0 0 20 B	0 5 0 W	51 30 40 N
Bagdad,	2 55 6 A	43 46 30 E	33 21 0 N	Lisbon,	0 36 30 B	8 57 30 W	38 42 20 N
Barcelona,	0 8 52 A	2 13 0 E	41 26 0 N	Lizard Point,	0 19 0 B	4 45 0 W	49 55 0 N
Bayonne,	0 8 19 A	2 4 42 E	43 29 21 N	Landau,	0 32 30 A	8 7 30 E	49 11 40 N
Brest,	0 18 3 B	4 30 45 W	48 23 0 N	Leipsick,	0 49 20 A	12 20 0 E	51 19 14 N
Bologna, Italy,	0 46 28 A	11 37 0 E	44 30 0 N	Lisle, Flanders,	0 12 17 A	3 4 16 E	50 37 50 N
Buenos Ayres, America,	3 52 20 B	58 5 0 W	34 35 0 S	Moscow, Russia,	2 41 20 A	40 20 0 E	55 36 0 N
Bordeaux,	0 2 19 B	0 34 19 W	44 50 18 N	Marfeilles,	0 21 29 A	5 22 15 E	43 17 45 N
Bologne, France,	0 6 27 A	1 36 44 E	50 43 31 N	Madrid,	0 14 58 B	6 2 10 W	40 25 0 N
Breslaw, Silesia,	1 8 35 A	17 17 30 E	51 3 0 N	Malacca, Indies,	6 48 20 A	102 5 0 E	1 15 50 N
Brussels,	0 17 27 A	4 21 43 E	50 51 0 N	Malta,	0 57 50 A	14 29 30 E	35 54 0 N
Cape of Good Hope,	1 8 0 A	17 0 0 E	34 15 0 S	Manila, Islands, Indies,	8 1 20 A	110 20 0 E	14 30 0 N
Carthagina, America,	5 1 46 B	75 26 0 W	10 26 35 N	Martinico, West-Indies,	4 3 55 B	60 58 45 W	14 43 9 N
Cayann Is. America,	3 51 20 B	57 50 0 W	4 56 0 N	Macao, China,	7 35 5 A	113 46 15 E	22 12 44 N
Cadix, Spain,	0 24 28 B	6 7 0 W	36 33 30 N	Mexico, America,	6 54 40 B	103 40 0 W	20 0 0 N
Copenhagen, Denmark,	0 51 0 A	12 45 0 E	55 40 45 N	Milan,	0 36 20 A	9 20 0 E	45 25 0 N
Calais,	0 7 16 A	4 49 4 E	50 57 31 N	Modena,	0 44 50 A	11 12 30 E	44 34 0 N
Cambray,	0 5 55 A	3 13 42 E	50 10 30 N	Montpellier,	0 15 31 A	3 52 44 E	43 36 33 N
Cape Verd,	1 8 40 B	17 10 0 W	14 43 0 N	Mons,	0 15 49 A	3 57 10 E	50 27 10 N
Cologne,	0 38 20 A	7 5 0 E	50 55 0 N	Munich,	0 46 20 A	11 35 0 E	48 2 0 N
Constantinople,	1 55 34 A	28 53 30 E	41 0 0 N	Mentz,	0 24 44 A	6 11 0 E	49 7 5 N
Cracow,	1 19 20 A	19 50 0 E	50 10 0 N	Namur,	0 19 26 A	4 51 37 E	50 28 0 N

GEOGRAPHICAL TABLE Continued

Places Names.	Dif. Time.			Dif. Longitud.	East or West.	Latitude	North or Sou.	Places Names.	Dif. Time.			Dif. Longit.	East or West.	Latitude.	North or Sou.
	h	m	s			°	'		h	m	s			°	'
Nantz,	0	55	A	63	W	47	13	Smyrna,	1	49	19	A	27	19	45
Naples,	0	58	A	14	0	40	50	Stockholm,	1	17	40	A	19	25	0
Nuremberg, Germany,	0	44	A	11	4	49	26	Straßburg,	0	31	5	A	7	46	18
Nice,	0	29	A	7	17	43	41	Surat,	4	49	20	A	72	20	0
New Orleans,	6	18	35	A	94	38	45	SHERBURN CASTLE,	0	4	0	B	1	0	0
Olinda, Brasil,	2	10	40	B	35	10	0	Toledo, Spain,	0	12	12	B	3	20	0
Orleans, France,	0	7	37	A	1	54	22	Toulon,	0	23	36	A	5	56	35
Ostend,	0	11	40	A	2	55	2	Tripoli, Barbary,	0	52	21	A	13	5	15
Oxford,	0	5	4	B	1	16	0	Turin,	0	30	40	A	7	40	0
PARIS Observatory,	0	9	20	A	2	20	0	Tours,	0	2	45	A	0	41	11
Pekin, China,	7	45	30	A	116	22	30	Torneo, Lapland,	1	36	50	A	24	22	30
Petersburg,	2	1	20	A	30	20	0	Touloufe,	0	5	45	A	1	26	13
Pike of Teneriff,	1	6	11	B	16	32	47	Terra del Gada, Madaga.	2	58	0	A	44	30	0
Porto Bello, America,	5	19	20	B	73	50	6	Tbeffalonica, Turkey,	1	32	32	A	23	8	0
Pondicherry, Indies,	5	20	50	A	80	12	50	Valparais, Cbili,	4	49	17	B	72	19	15
Prague, Bobemia,	0	59	0	A	14	45	0	Venice,	0	48	18	A	12	4	30
Poitiers,	0	1	20	A	0	20	5	Verona,	0	45	14	A	11	18	30
Padua, in Italy,	0	47	42	A	11	55	30	Versailles,	0	8	29	A	1	7	10
Pike of Azores,	1	52	40	B	28	10	0	Vienna,	1	5	30	A	16	22	30
Quitto, City, America,	5	11	40	B	77	55	0	UPNOR CASTLE,	0	1	48	A	0	27	0
Quanton, or Canton, China	7	32	13	A	113	3	15	Upsal, Sweden,	1	11	0	A	17	45	0
Quimper,	0	16	30	B	4	7	5	URANIBURG,	0	51	30	A	12	52	30
Rome,	0	49	57	A	12	29	15	Wirtemburg, Saxony,	0	50	14	A	12	33	30
Rochelle,	0	5	3	B	1	13	44	Ylo, Peru,	4	44	52	B	71	13	0
Rhodes,	0	10	17	A	2	34	20	Ypres,	0	11	32	A	3	2	55
Rheims,	0	16	12	A	4	2	53								
Rouen,	0	4	21	A	1	5	20								
St. Malo,	0	8	9	B	2	2	22								
St. Omer,	0	9	0	A	1	14	57								
Siam, Indies,	6	43	20	A	100	10	0								

USE of the foregoing TABLE.

When it is 12 at Noon, Mean Time, at Greenwich Observatory, what is the Hour, or Mean Time, at Porto Bello, in America?

EXAMPLE I.—Against Porto Bello, in the Table you find the Time is 5^h 19^m 20^s Before the Time at Greenwich.

Therefore, From 12^h 0^m 0^s at Greenwich,

Deduct — — 5 19 20

There remains 6 40 40 Time in the Morning at Porto Bello, req^d.

And, in general, you deduct the Time Before from the Time at Greenwich, or add it thereto, when the Time is after that at Greenwich, for the Time at any Place required; and you are to observe the contrary, when the Time is given at any Place, to find the Time at Greenwich Observatory, in Reference to the former Time and Place.

UNIVERSALLY.—The Difference of Degrees in West or East Longitude, between the Meridians of two Places, converted into Time, allowing 15° to an Hour, is the Time or Hour at one Place before or after the Mean Time at another; (the West Longitude before, and East Longitude after) therefore, the Difference or Sum of the Times of any two Places from the Time at Greenwich Observatory, as they are both B or A, or one B, and one A, Greenwich-Time (using After and Before to Greenwich, instead of Before and After from thence) will be the Time at one Place in Reference to the Time at another.

EXAMPLE II.—To find the Hour at Babylon, or Grand Cairo, when it is 10 in the Forenoon at Upnor Castle?

From the Time at Greenwich { Upnor Castle 0^h 1^m 48^s A
Grand Cairo 2 5 45 A

Time at Upnor Castle B. that at Grand Cairo, or Difference 2 3 57 A, at Grand Cairo, the Time at Upnor Castle.

Time at Grand Cairo — 12 3 57 Afternoon required.

EXAMPLE III.—To find the Time at Oxford when it is Noon at Rome?

From the Time at Greenwich { Rome 0^h 49^m 57^s A
Oxford 0 5 4 B

Time at Oxford B. Time at Rome. Sum 0 55 1 B—

Time at Oxford referred to Rome 11 4 59 Forenoon req^d: being all the Cases that can happen.

Mean

Mean Radical PLACES of the SUN, and first Star of Aries, from 1652 to 1752, Old STYLE.

Julian Style.	M. Pla. O.	M. Pla. Ap. O.	M. Pl. 1*V.	Julian Style.	M. Pla. O.	M. Pla. Ap. O.	M. Pl. 1*V.
B. 1652	9 20 36 12	3 6 54 17	0 28 18 56	B. 1700	9 20 58 3	3 7 43 29	0 28 59 20
53	9 20 21 52	3 6 55 18	0 28 19 46	1	9 20 43 43	3 7 44 30	0 29 0 10
54	9 20 7 32	3 6 56 20	0 28 20 37	2	9 20 29 23	3 7 45 32	0 29 1 1
55	9 19 53 13	3 6 57 21	0 28 21 27	3	9 20 15 4	3 7 46 33	0 29 1 51
B. 1656	9 20 38 1	3 6 58 23	0 28 22 18	B. 1704	9 20 59 53	3 7 47 35	0 29 2 42
57	9 20 23 41	3 6 59 24	0 28 23 8	5	9 20 45 33	3 7 48 36	0 29 3 32
58	9 20 9 21	3 7 0 26	0 28 23 59	6	9 20 31 13	3 7 49 38	0 29 4 23
59	9 19 55 2	3 7 1 27	0 28 24 49	7	9 20 16 54	3 7 50 39	0 29 5 13
B. 1660	9 20 39 51	3 7 2 29	0 28 25 40	B. 1708	9 21 1 42	3 7 51 41	0 29 6 4
61	9 20 25 31	3 7 3 30	0 28 26 30	9	9 20 47 22	3 7 52 42	0 29 6 54
62	9 20 11 11	3 7 4 32	0 28 27 21	10	9 20 33 2	3 7 53 44	0 29 7 45
63	9 19 56 52	3 7 5 33	0 28 28 11	11	9 20 18 43	3 7 54 45	0 29 8 35
B. 1664	9 20 41 40	3 7 6 35	0 28 29 2	B. 1712	9 21 3 31	3 7 55 47	0 29 9 26
65	9 20 27 20	3 7 7 36	0 28 29 52	13	9 20 49 11	3 7 56 48	0 29 10 16
66	9 20 13 0	3 7 8 38	0 28 30 43	14	9 20 34 51	3 7 57 50	0 29 11 7
67	9 19 58 41	3 7 9 39	0 28 31 33	15	9 20 20 32	3 7 58 51	0 29 11 57
B. 1668	9 20 43 29	3 7 10 41	0 28 32 24	B. 1716	9 21 5 20	3 7 59 53	0 29 12 48
69	9 20 29 9	3 7 11 42	0 28 33 14	17	9 20 51 0	3 8 0 54	0 29 13 38
70	9 20 14 49	3 7 12 44	0 28 34 5	18	9 20 36 40	3 8 1 56	0 29 14 29
71	9 20 0 30	3 7 13 45	0 28 34 55	19	9 20 22 21	3 8 2 57	0 29 15 19
B. 1672	9 20 45 18	3 7 14 47	0 28 35 46	B. 1720	9 21 7 10	3 8 3 59	0 29 16 10
73	9 20 30 58	3 7 15 48	0 28 36 36	21	9 20 52 50	3 8 5 0	0 29 17 0
74	9 20 16 38	3 7 16 50	0 28 37 27	22	9 20 38 30	3 8 6 2	0 29 17 51
75	9 20 2 19	3 7 17 51	0 28 38 17	23	9 20 24 11	3 8 7 3	0 29 18 41
B. 1676	9 20 47 8	3 7 18 53	0 28 39 8	B. 1724	9 21 8 59	3 8 8 5	0 29 19 32
77	9 20 32 48	3 7 19 54	0 28 39 58	25	9 20 54 39	3 8 9 6	0 29 20 22
78	9 20 18 28	3 7 20 56	0 28 40 49	26	9 20 40 19	3 8 10 8	0 29 21 13
79	9 20 4 9	3 7 21 57	0 28 41 39	27	9 20 26 0	3 8 11 9	0 29 22 3
B. 1680	9 20 48 57	3 7 22 59	0 28 42 30	B. 1728	9 21 10 48	3 8 12 11	0 29 22 54
81	9 20 34 37	3 7 24 0	0 28 43 20	29	9 20 56 28	3 8 13 12	0 29 23 44
82	9 20 20 17	3 7 25 2	0 28 44 11	30	9 20 42 8	3 8 14 14	0 29 24 35
83	9 20 5 58	3 7 26 3	0 28 45 1	31	9 20 27 49	3 8 15 15	0 29 25 25
B. 1684	9 20 50 46	3 7 27 5	0 28 45 52	B. 1732	9 21 12 38	3 8 16 17	0 29 26 16
85	9 20 36 26	3 7 28 6	0 28 46 42	33	9 20 58 18	3 8 17 18	0 29 27 6
86	9 20 22 6	3 7 29 8	0 28 47 33	34	9 20 43 58	3 8 18 20	0 29 27 57
87	9 20 7 47	3 7 30 9	0 28 48 23	35	9 20 29 39	3 8 19 21	0 29 28 47
B. 1688	9 20 52 35	3 7 31 11	0 28 49 14	B. 1736	9 21 14 27	3 8 20 23	0 29 29 38
89	9 20 38 15	3 7 32 12	0 28 50 4	37	9 21 0 7	3 8 21 24	0 29 30 28
90	9 20 23 55	3 7 33 14	0 28 50 55	38	9 20 45 47	3 8 22 26	0 29 31 19
91	9 20 9 36	3 7 34 15	0 28 51 45	39	9 20 31 28	3 8 23 27	0 29 32 9
B. 1692	9 20 54 25	3 7 35 17	0 28 52 36	B. 1740	9 21 16 16	3 8 24 29	0 29 33 0
93	9 20 40 5	3 7 36 18	0 28 53 26	41	9 21 1 56	3 8 25 30	0 29 33 50
94	9 20 25 45	3 7 37 20	0 28 54 17	42	9 20 47 36	3 8 26 32	0 29 34 41
95	9 20 11 26	3 7 38 21	0 28 55 7	43	9 20 33 17	3 8 27 33	0 29 35 31
B. 1696	9 20 56 14	3 7 39 23	0 28 55 58	B. 1744	9 21 18 5	3 8 28 35	0 29 36 22
97	9 20 41 54	3 7 40 24	0 28 56 48	45	9 21 3 45	3 8 29 36	0 29 37 12
98	9 20 27 34	3 7 41 26	0 28 57 39	46	9 20 49 25	3 8 30 38	0 29 38 3
99	9 20 13 15	3 7 42 27	0 28 58 29	47	9 20 35 6	3 8 31 39	0 29 38 53
				B. 1748	9 21 19 55	3 8 32 41	0 29 39 44
				49	9 21 5 35	3 8 33 42	0 29 40 34
				50	9 20 51 15	3 8 34 44	0 29 41 25
				51	9 20 36 56	3 8 35 45	0 29 42 15
				B. 1752	9 21 21 44	3 8 36 47	0 29 43 6

SOLAR TABLES.

Mean Radical PLACES of the SUN, and first Star of Aries, from 1752 to 1852, New STYLE.

Gregorian Style.	M. Pla. $\odot$.	M. Pl. Ap. $\odot$.	M. Pla. 1^{st} γ .	Gregorian Style.	M. Pla. $\odot$.	M. Pl. Ap. $\odot$.	M. Pla. 1^{st} γ .
s o / //	s o / //	s o / //	s o / //	s o / //	s o / //	s o / //	s o / //
B. 1752	9 10 31 13	3 8 36 45	0 29 43 5	1800	9 9 53 55	3 9 25 57	1 0 23 29
53	9 10 16 53	3 8 37 46	0 29 43 55	1	9 9 39 35	3 9 26 58	1 0 24 19
54	9 10 2 33	3 8 38 48	0 29 44 46	2	9 9 25 15	3 9 28 0	1 0 25 10
55	9 9 48 14	3 8 39 49	0 29 45 36	3	9 9 10 56	3 9 29 1	1 0 26 0
B. 1756	9 10 33 2	3 8 40 51	0 29 46 27	B. 1804	9 9 55 44	3 9 30 3	1 0 26 51
57	9 10 18 42	3 8 41 52	0 29 47 10	5	9 9 41 24	3 9 31 4	1 0 27 41
58	9 10 4 22	3 8 42 54	0 29 48 8	6	9 9 27 4	3 9 32 6	1 0 28 32
59	9 9 50 3	3 8 43 55	0 29 48 58	7	9 9 12 45	3 9 33 7	1 0 29 22
B. 1760	9 10 34 51	3 8 44 57	0 29 49 49	B. 1808	9 9 57 33	3 9 34 9	1 0 30 13
61	9 10 20 31	3 8 45 58	0 29 50 39	9	9 9 43 13	3 9 35 10	1 0 31 3
62	9 10 6 11	3 8 47 0	0 29 51 30	10	9 9 28 53	3 9 36 12	1 0 31 54
63	9 9 51 52	3 8 48 1	0 29 52 20	11	9 9 14 34	3 9 37 13	1 0 32 44
B. 1764	9 10 36 41	3 8 49 3	0 29 53 11	B. 1812	9 9 59 23	3 9 38 15	1 0 33 35
65	9 10 22 21	3 8 50 4	0 29 54 1	13	9 9 45 3	3 9 39 16	1 0 34 25
66	9 10 8 1	3 8 51 6	0 29 54 52	14	9 9 30 43	3 9 40 18	1 0 35 16
67	9 9 53 42	3 8 52 7	0 29 55 42	15	9 9 16 24	3 9 41 19	1 0 36 6
B. 1768	9 10 38 30	3 8 53 9	0 29 56 33	B. 1816	9 10 1 12	3 9 42 21	1 0 36 57
69	9 10 24 10	3 8 54 10	0 29 57 23	17	9 9 46 52	3 9 43 22	1 0 37 47
70	9 10 9 50	3 8 55 12	0 29 58 14	18	9 9 32 32	3 9 44 24	1 0 38 38
71	9 9 55 31	3 8 56 13	0 29 59 4	19	9 9 18 13	3 9 45 25	1 0 39 28
B. 1772	9 10 40 19	3 8 57 15	0 29 59 55	B. 1820	9 10 3 1	3 9 46 27	1 0 40 19
73	9 10 25 59	3 8 58 16	1 0 0 45	21	9 9 48 41	3 9 47 28	1 0 41 9
74	9 10 11 39	3 8 59 18	1 0 1 36	22	9 9 34 21	3 9 48 30	1 0 42 0
75	9 9 57 20	3 9 0 19	1 0 2 26	23	9 9 20 2	3 9 49 31	1 0 42 50
B. 1776	9 10 42 9	3 9 1 21	1 0 3 17	B. 1824	9 10 4 51	3 9 50 33	1 0 43 41
77	9 10 27 49	3 9 2 22	1 0 4 7	25	9 9 50 31	3 9 51 34	1 0 44 31
78	9 10 13 29	3 9 3 24	1 0 4 58	26	9 9 36 11	3 9 52 36	1 0 45 22
79	9 9 59 10	3 9 4 25	1 0 5 48	27	9 9 21 52	3 9 53 37	1 0 46 12
B. 1780	9 10 43 57	3 9 5 27	1 0 6 39	B. 1828	9 10 6 40	3 9 54 39	1 0 47 3
81	9 10 29 37	3 9 6 28	1 0 7 29	29	9 9 52 20	3 9 55 40	1 0 47 53
82	9 10 15 17	3 9 7 30	1 0 8 20	30	9 9 38 0	3 9 56 42	1 0 48 44
83	9 10 0 58	3 9 8 31	1 0 9 10	31	9 9 23 41	3 9 57 43	1 0 49 34
B. 1784	9 10 45 47	3 9 9 33	1 0 10 1	B. 1832	9 10 8 29	3 9 58 45	1 0 50 25
85	9 10 31 27	3 9 10 34	1 0 10 51	33	9 9 54 9	3 9 59 46	1 0 51 15
86	9 10 17 7	3 9 11 36	1 0 11 42	34	9 9 39 49	3 10 0 48	1 0 52 6
87	9 10 2 48	3 9 12 37	1 0 12 32	35	9 9 25 30	3 10 1 49	1 0 52 56
B. 1788	9 10 47 36	3 9 13 39	1 0 13 23	B. 1836	9 10 10 18	3 10 2 51	1 0 53 47
89	9 10 33 16	3 9 14 40	1 0 14 13	37	9 9 55 58	3 10 3 52	1 0 54 37
90	9 10 18 56	3 9 15 42	1 0 15 4	38	9 9 41 38	3 10 4 54	1 0 55 28
91	9 10 4 37	3 9 16 43	1 0 15 54	39	9 9 27 19	3 10 5 55	1 0 56 18
B. 1792	9 10 49 26	3 9 17 45	1 0 16 45	B. 1840	9 10 12 7	3 10 6 57	1 0 57 9
93	9 10 35 6	3 9 18 46	1 0 17 35	41	9 9 57 47	3 10 7 58	1 0 57 59
94	9 10 20 46	3 9 19 48	1 0 18 26	42	9 9 43 27	3 10 9 0	1 0 58 50
95	9 10 6 27	3 9 20 49	1 0 19 16	43	9 9 29 8	3 10 10 1	1 0 59 40
B. 1796	9 10 51 15	3 9 21 51	1 0 20 7	B. 1844	9 10 13 57	3 10 11 3	1 1 0 31
97	9 10 36 55	3 9 22 52	1 0 20 57	45	9 9 59 37	3 10 12 4	1 1 1 21
98	9 10 22 35	3 9 23 54	1 0 21 48	46	9 9 45 17	3 10 13 6	1 1 2 12
99	9 10 8 16	3 9 24 55	1 0 22 38	47	9 9 30 58	3 10 14 7	1 1 3 2
For Advantage of taking out the Numbers after January and February, Bissextile Years Places are advanced a Day's Motion, in both Styles, correspondent to the mean Noon of January the 1 st .				B. 1848	9 10 15 46	3 10 15 9	1 1 3 53
				49	9 10 1 26	3 10 16 10	1 1 4 43
				50	9 9 47 6	3 10 17 12	1 1 5 34
				51	9 9 32 47	3 10 18 13	1 1 6 24
				B. 1852	9 10 17 36	3 10 19 15	1 1 7 15

Mean

Mean MOTIONS of the SUN and fixed STARS for MONTHS and DAYS.

Days.	JANUARY.				FEBRUARY.				MARCH.				APRIL.			
	M. Mot. ☉		Ap.	*s.	M. Mot. ☉		Ap.	*s.	M. Mot. ☉		Ap.	*s.	M. Mot. ☉		Ap.	*s.
	s	o	/'	''	s	o	/'	''	s	o	/'	''	s	o	/'	''
1	0	0	59	8	1	1	32	27	1	29	8	19	2	29	41	38
2	0	1	58	17	1	2	31	35	2	0	7	28	3	0	40	47
3	0	2	57	25	1	3	30	43	2	1	6	36	3	1	39	55
4	0	3	56	33	1	4	29	52	2	2	5	45	3	2	39	3
5	0	4	55	42	1	5	29	0	2	3	4	53	3	3	38	11
6	0	5	54	50	1	6	28	8	2	4	4	1	3	4	37	20
7	0	6	53	58	1	7	27	17	2	5	3	10	3	5	36	28
8	0	7	53	7	1	8	26	25	2	6	2	18	3	6	35	36
9	0	8	52	15	1	9	25	33	2	7	1	26	3	7	34	45
10	0	9	51	23	1	10	24	42	2	8	0	35	3	8	33	53
11	0	10	50	32	1	11	23	50	2	8	59	43	3	9	33	1
12	0	11	49	40	1	12	22	58	2	9	58	51	3	10	32	10
13	0	12	48	48	1	13	22	7	2	10	58	0	3	11	31	18
14	0	13	47	57	1	14	21	15	2	11	57	8	3	12	30	26
15	0	14	47	5	1	15	20	23	2	12	56	16	3	13	29	34
16	0	15	46	13	1	16	19	32	2	13	55	25	3	14	28	43
17	0	16	45	22	1	17	18	40	2	14	54	33	3	15	27	51
18	0	17	44	30	1	18	17	48	2	15	53	41	3	16	27	0
19	0	18	43	38	1	19	16	57	2	16	52	50	3	17	26	8
20	0	19	42	47	1	20	16	5	2	17	51	58	3	18	25	16
21	0	20	41	55	1	21	15	13	2	18	51	6	3	19	24	25
22	0	21	41	3	1	22	14	22	2	19	50	14	3	20	23	33
23	0	22	40	12	1	23	13	30	2	20	49	23	3	21	22	41
24	0	23	39	20	1	24	12	38	2	21	48	31	3	22	21	50
25	0	24	38	28	1	25	11	47	2	22	47	40	3	23	20	58
26	0	25	37	37	1	26	10	55	2	23	46	48	3	24	20	6
27	0	26	36	45	1	27	10	3	2	24	45	56	3	25	19	15
28	0	27	35	53	1	28	9	11	2	25	45	5	3	26	18	23
29	0	28	35	2	In Leap-Year for Jan. & Feb. take out for a Day sooner, both Styles.				2	26	44	13	3	27	17	31
30	0	29	34	10					2	27	43	21	3	28	16	40
31	1	0	33	18					2	28	42	30				

To find the Mean Place of the Sun, his Apogee, and of the γ , on March, 27, 1758, at Noon, New Style.

EXAMPLE I.

	☉				Ap. ☉				γ			
	s	o	/'	''	s	o	/'	''	s	o	/'	''
N. S. Year 1758.	9	10	4	22	3	8	42	54	0	29	48	8
March 27	2	24	45	56			+	14			+	12
Answer -	0	4	50	18	3	8	43	8	0	29	48	20

To find the Mean Place of the Sun, Apogee, and γ on January 20, 1852, at Noon, Gregorian Style.

EXAMPLE II.

	☉				Ap. ☉				γ			
	s	o	/'	''	s	o	/'	''	s	o	/'	''
N. S. Year 1852	9	10	17	36	3	10	19	15	1	1	7	15
B. January 19	0	18	43	38			+	3			+	3
Answer -	9	29	1	14	3	10	19	18	1	1	7	18

Mean

Mean MOTIONS of the SUN and fixed STARS for MONTHS and DAYS.

Days.	MAY.						JUNE.						JULY.						AUGUST.					
	M. Mot. ☉.			Ap.			M. Mot. ☉.			Ap.			M. Mot. ☉.			Ap.			M. Mot. ☉.			Ap.		
	s	o	l	l	l	l	s	o	l	l	l	l	s	o	l	l	l	l	s	o	l	l	l	l
1	3	29	15	48	20	17	4	29	49	6	25	21	5	29	23	16	31	25	6	29	56	34	36	29
2	4	0	14	56	20	17	5	0	48	15	26	21	6	0	22	24	31	25	7	0	55	42	36	29
3	4	1	14	5	21	17	5	1	47	23	26	21	6	1	21	33	31	25	7	1	54	51	36	30
4	4	2	13	13	21	17	5	2	46	31	26	21	6	2	20	41	31	25	7	2	53	59	36	30
5	4	3	12	21	21	17	5	3	45	39	26	21	6	3	19	49	31	26	7	3	53	7	36	30
6	4	4	11	30	21	17	5	4	44	48	26	22	6	4	18	58	31	26	7	4	52	16	37	30
7	4	5	10	38	21	17	5	5	43	56	26	22	6	5	18	6	32	26	7	5	51	24	37	30
8	4	6	9	46	21	18	5	6	43	4	27	22	6	6	17	14	32	26	7	6	50	32	37	30
9	4	7	8	55	22	18	5	7	42	13	27	22	6	7	16	22	32	26	7	7	49	41	37	30
10	4	8	8	3	22	18	5	8	41	21	27	22	6	8	15	31	32	26	7	8	48	49	37	31
11	4	9	7	11	22	18	5	9	40	29	27	22	6	9	14	39	32	26	7	9	47	57	37	31
12	4	10	6	20	22	18	5	10	39	38	27	22	6	10	13	47	32	27	7	10	47	6	38	31
13	4	11	5	28	22	18	5	11	38	46	27	23	6	11	12	56	33	27	7	11	46	14	38	31
14	4	12	4	36	22	18	5	12	37	55	28	23	6	12	12	4	33	27	7	12	45	22	38	31
15	4	13	3	45	23	19	5	13	37	3	28	23	6	13	11	12	33	27	7	13	44	31	38	31
16	4	14	2	53	23	19	5	14	36	11	28	23	6	14	10	21	33	27	7	14	43	39	38	31
17	4	15	2	1	23	19	5	15	35	20	28	23	6	15	9	29	33	27	7	15	42	47	38	32
18	4	16	1	10	23	19	5	16	34	28	28	23	6	16	8	37	33	27	7	16	41	56	39	32
19	4	17	0	18	23	19	5	17	33	36	28	23	6	17	7	46	34	28	7	17	41	4	39	32
20	4	17	59	26	23	19	5	18	32	45	29	24	6	18	6	54	34	28	7	18	40	12	39	32
21	4	18	58	35	24	19	5	19	31	53	29	24	6	19	6	2	34	28	7	19	39	21	39	32
22	4	19	57	43	24	20	5	20	31	1	29	24	6	20	5	11	34	28	7	20	38	29	39	32
23	4	20	56	51	24	20	5	21	30	10	29	24	6	21	4	19	34	28	7	21	37	37	39	32
24	4	21	56	0	24	20	5	22	29	18	29	24	6	22	3	28	34	28	7	22	36	46	40	33
25	4	22	55	8	24	20	5	23	28	26	30	24	6	23	2	36	35	28	7	23	35	54	40	33
26	4	23	54	16	24	20	5	24	27	35	30	24	6	24	1	44	35	29	7	24	35	2	40	33
27	4	24	53	25	25	20	5	25	26	43	30	25	6	25	0	52	35	29	7	25	34	11	40	33
28	4	25	52	33	25	20	5	26	25	51	30	25	6	26	0	1	35	29	7	26	33	19	40	33
29	4	26	51	41	25	20	5	27	25	0	30	25	6	26	59	9	35	29	7	27	32	27	40	33
30	4	27	50	50	25	21	5	28	24	8	30	25	6	27	58	17	35	29	7	28	31	36	41	33
31	4	28	49	58	25	21							6	28	57	26	36	29	7	29	30	44	41	33

To find the Mean Place of the Sun, his Apogee, and $1^{\circ} \gamma$, on March 16, 1758, at Noon, Old Style.

EXAMPLE III.

	☉				Ap. ☉				$1^{\circ} \gamma$			
	s	o	l	l	s	o	l	l	s	o	l	l
O. S. Year 1658	9	20	9	21	3	7	0	26	0	28	23	59
Motion Years 100			45	32		1	42	30		1	24	10
16 March	2	13	55	25				13				10
O. S. 16th March 1758.	0	4	50	18	3	8	43	9	0	29	48	19

To find the Mean Place of the Sun, Apogee, and $1^{\circ} \gamma$, on January 8, 1852, at Noon, O. S.

EXAMPLE IV.

	☉				Ap. ☉				$1^{\circ} \gamma$			
	s	o	l	l	s	o	l	l	s	o	l	l
O. S. Year 1752	9	21	21	44	3	8	36	47	0	29	43	6
Motion Years 100			45	32		1	42	30		1	24	10
B. 7th January	0	6	53	58				1				1
O. S. 8th Jan. 1852	9	29	1	14	3	10	19	18	1	1	7	17

Being a complete Proof of the Truth of the Tables, from what is done on the other Side.

Mean

Mean MOTIONS of the SUN, and Fixed STARS, for MONTHS and DAYS.

Days.	SEPTEMBER.			OCTOBER.			NOVEMBER.			DECEMBER.		
	M. Mot. ☉.	Ap.	*s.	M. Mot. ☉.	Ap.	*s.	M. Mot. ☉.	Ap.	*s.	M. Mot. ☉.	Ap.	*s.
	s o / //	//	//	s o / //	//	//	s o / //	//	//	s o / //	//	//
1	8 0 29 52	41	34	9 0 4 2	46	38	10 0 37 20	51	42	11 0 11 30	56	46
2	8 1 29 1	41	34	9 1 3 10	46	38	10 1 36 29	51	42	11 1 10 38	56	46
3	8 2 28 9	41	34	9 2 2 19	46	38	10 2 35 37	52	42	11 2 9 47	57	46
4	8 3 27 17	41	34	9 3 1 27	47	38	10 3 34 45	52	42	11 3 8 55	57	47
5	8 4 26 26	42	34	9 4 0 35	47	38	10 4 33 54	52	42	11 4 8 3	57	47
6	8 5 25 34	42	34	9 4 59 44	47	38	10 5 33 2	52	43	11 5 7 12	57	47
7	8 6 24 42	42	34	9 5 58 52	47	39	10 6 32 10	52	43	11 6 6 20	57	47
8	8 7 23 51	42	35	9 6 58 0	47	39	10 7 31 19	52	43	11 7 5 28	57	47
9	8 8 22 59	42	35	9 7 57 8	47	39	10 8 30 27	53	43	11 8 4 37	58	47
10	8 9 22 7	42	35	9 8 56 17	48	39	10 9 29 35	53	43	11 9 3 45	58	47
11	8 10 21 16	43	35	9 9 55 25	48	39	10 10 28 44	53	43	11 10 2 53	58	48
12	8 11 20 24	43	35	9 10 54 34	48	39	10 11 27 52	53	44	11 11 2 2	58	48
13	8 12 19 32	43	35	9 11 53 42	48	39	10 12 27 0	53	44	11 12 1 10	58	48
14	8 13 18 41	43	35	9 12 52 50	48	40	10 13 26 9	53	44	11 13 0 18	58	48
15	8 14 17 49	43	36	9 13 51 59	48	40	10 14 25 17	54	44	11 13 59 27	59	48
16	8 15 16 57	43	36	9 14 51 7	49	40	10 15 24 25	54	44	11 14 58 35	59	48
17	8 16 16 5	44	36	9 15 50 15	49	40	10 16 23 34	54	44	11 15 57 43	59	48
18	8 17 15 14	44	36	9 16 49 24	49	40	10 17 22 42	54	44	11 16 56 52	59	49
19	8 18 14 22	44	36	9 17 48 32	49	40	10 18 21 50	54	45	11 17 56 0	59	49
20	8 19 13 30	44	36	9 18 47 40	49	40	10 19 20 59	54	45	11 18 55 8	59	49
21	8 20 12 39	44	36	9 19 46 49	49	41	10 20 20 7	55	45	11 19 54 17	60	49
22	8 21 11 47	45	37	9 20 45 57	50	41	10 21 19 15	55	45	11 20 53 25	60	49
23	8 22 10 55	45	37	9 21 45 5	50	41	10 22 18 24	55	45	11 21 52 33	60	49
24	8 23 10 4	45	37	9 22 44 14	50	41	10 23 17 32	55	45	11 22 51 42	60	49
25	8 24 9 12	45	37	9 23 43 22	50	41	10 24 16 40	55	45	11 23 50 50	60	50
26	8 25 8 20	45	37	9 24 42 30	50	41	10 25 15 49	55	46	11 24 49 58	60	50
27	8 26 7 29	45	37	9 25 41 39	50	41	10 26 14 57	56	46	11 25 49 7	60	50
28	8 27 6 37	46	37	9 26 40 47	51	41	10 27 14 5	56	46	11 26 48 15	61	50
29	8 28 5 45	46	37	9 27 39 55	51	42	10 28 13 13	56	46	11 27 47 23	61	50
30	8 29 4 54	46	38	9 28 39 4	51	42	10 29 12 22	56	46	11 28 46 32	61	50
31				9 29 38 12	51	42				11 29 45 40	61	50

RADICAL MEAN PLACES and MOTIONS of the SUN, from several AUTHORITIES.

M. Pl. ☉.	M. Pl. Ap. ☉.	M. Mot. ☉.	M. Mot. Ap.	1 mean Solar	Eccentry	Greatest Eqn.	According to
Julian Style.	Julian Style.	in 100 Julian	in 100 Julian	Year, 365	to M. Diff.	☉ Orb.	
Jan. o. 1701.	Jan. o. 1701.	Years.	Years.	Days.	100000.		
s o / //	s o / //	s o / //	s o / //	h m s th		s o / //	
9 20 43 40	3 7 44 30	0 0 45 20 100 Revol.	0 1 41 0 Ap. a * s 17 40	5 48 57 41 &c.	1692	0 1 56 20	Sir Isaac Newton's last Edit. Principia. 1 ^o . 23'. 20". * s a Eq ^x in 100 J. Yrs.
9 20 43 33	3 7 39 10	0 0 45 32	0 1 41 7	5 48 54 41	same	same	Dr. Ed. Halley's Tables. 1701, 1 * 9
9 20 43 53	3 7 46 30	Same [20 th]	Same [20 th]	&c.			Gael Morris Aft Afr. [0 ^s . 29 ^o . 0'. 10".
9 20 43 35	3 7 45 30	0 0 45 47	0 1 45 0	5 48 51 7	1680	0 1 55 30	Tobias Mayer, of Gottingen, Germany, from his Tables.
9 20 41 31	3 8 5 41	0 0 46 0	0 1 25 50	5 48 47 57	1676	0 1 55 15	Leonard Euler, of Berlin, Prussia, F. R. S. from his Tables.
9 20 40 21	3 7 16 0	0 0 43 25	0 1 20 0	5 49 25 41	1732	0 1 59 6	Thomas Street's Caroline Tables.
9 20 43 43	3 7 44 30	0 0 45 32	0 1 42 30	5 48 54 46	1685	0 1 55 50	Our own Tables.
		* 0 1 24 10	&c.				

Hence, From a Comparison of These, with other Observations, we give the following correct

RADICAL

RADICAL MEAN PLACES and MOTIONS of the SUN, his APOGEE, and of the FIXED STARS, for
Julian and Gregorian YEARS.

Julian Style.		M. Places○.				M.Pl.Ap.○.				M. Pl. 1*γ.				Gregori- an Style.		M. Places ○.				M.Pl.Ap.○.				M. Pl. 1*γ.			
Years before and since Christ.		s	o	i	h	s	o	i	h	s	o	i	h	Years of Christ.		s	o	i	h	s	o	i	h	s	o	i	h
B.	1000	9	0	28	39	1	21	35	59	11	21	6	50	B.	1600	9	10	21	8	3	6	0	57	0	27	35	9
B.	900	9	1	14	11	1	23	18	29	11	22	31	0	1700	9	10	7	32	3	7	43	27	0	28	59	19	
B.	800	9	1	59	43	1	25	0	59	11	23	55	10	1800	9	9	53	55	3	9	25	57	1	0	23	28	
B.	700	9	2	45	15	1	26	43	29	11	25	19	20	1900	9	9	40	19	3	11	8	27	1	1	47	38	
B.	600	9	3	30	47	1	28	25	59	11	20	43	30	B.	2000	9	10	25	51	3	12	50	57	1	3	11	48
B.	500	9	4	16	19	2	0	8	29	11	28	7	40	2100	9	10	12	15	3	14	33	27	1	4	35	58	
B.	400	9	5	1	51	2	1	50	59	11	29	31	50	2200	9	9	58	38	3	16	15	57	1	6	0	8	
B.	300	9	5	47	23	2	3	33	29	0	0	56	0	2300	9	9	45	2	3	17	58	27	1	7	24	18	
B.	200	9	6	32	55	2	5	15	59	0	2	20	10	B.	2400	9	10	30	34	3	19	40	56	1	8	48	28
B. Jul. Period.	100	9	7	18	27	2	6	58	29	0	3	44	20	2500	9	10	16	58	3	21	23	26	1	10	12	38	
B. 4713. B. Chr.	0	9	8	3	59	2	8	40	59	0	5	8	30	2600	9	10	3	21	3	23	5	56	1	11	36	48	
B.	100	9	8	49	31	2	10	23	29	0	6	32	40	2700	9	9	49	45	3	24	48	26	1	13	0	57	
B.	200	9	9	35	3	2	12	5	59	0	7	56	50	B.	2800	9	10	35	17	3	26	30	56	1	14	25	7
B.	300	9	10	20	35	2	13	48	29	0	9	21	0	2900	9	10	21	41	3	28	13	26	1	15	49	17	
B.	400	9	11	6	7	2	15	30	59	0	10	45	10	3000	9	10	8	4	3	29	55	56	1	17	13	27	
B.	500	9	11	51	39	2	17	13	29	0	12	9	20	3100	9	9	54	28	4	1	38	25	1	18	37	37	
B.	600	9	12	37	11	2	18	55	59	0	13	33	30	B.	3200	9	10	40	0	4	3	20	55	1	20	1	47
B.	700	9	13	22	43	2	20	38	29	0	14	57	40	3300	9	10	26	24	4	5	3	25	1	21	25	57	
B.	800	9	14	8	15	2	22	20	59	0	16	21	50	3400	9	10	12	47	4	6	45	55	1	22	50	7	
B.	900	9	14	53	47	2	24	3	29	0	17	46	0	Bissextile Years Places are advanced a Day's Motion.													
B.	1000	9	15	39	19	2	25	45	59	0	19	10	10	From Hundreds of Years <i>Gregorian</i> deduct their <i>Fourth</i> (re- jecting Fractions) and the <i>Remainder</i> will be equal to the No. of Days there are left in any No. of <i>Gregorian</i> Years than there are in the same No. of Years <i>Julian</i> , from 1600. Whence is had the Diff. of Motion between <i>Julian</i> and <i>Gregorian</i> Years.													
B.	1100	9	16	24	51	2	27	28	29	0	20	34	20	EXAMPLES.													
B.	1200	9	17	10	23	2	29	10	59	0	21	58	30	In the Year 3200. In the Year 3756													
B.	1300	9	17	55	55	3	0	53	29	0	23	22	40	—1600 1600													
B.	1400	9	18	41	27	3	2	35	59	0	24	46	50	— 1600													
B.	1500	9	19	26	59	3	4	18	29	0	26	11	0	1/4 16 00 1/4 21 56													
B.	1600	9	20	12	31	3	6	0	59	0	27	35	10	— 4 — 5													
B.	1700	9	20	58	3	3	7	43	29	0	28	59	20	12 Days less 16 Days less than in 2													
B.	1800	9	21	43	35	3	9	25	59	1	0	23	30	[than in 16 Hund. Yrs <i>Jul.</i> [Hundred Years <i>Julian</i>													
B.	1900	9	22	29	7	3	11	8	29	1	1	47	40														
B.	2000	9	23	14	35	3	12	50	59	1	3	11	50														
Bissextile Years Places are advanced a Day's Motion.																											
Julian Years.		M. Mot. ○.				Mean Mot. Ap. ○.				Mean Mot. fixed Stars.				Gregori- an Years		M. Mot. ○.				Mean Mot. Ap. ○.				Mean Mot. fixed Stars.			
		s	o	i	h	s	o	i	h	s	o	i	h	from 1600		s	o	i	h	s	o	i	h	s	o	i	h
B.	100	0	0	45	32	0	1	42	30	0	1	24	10	100	11	29	46	24	0	1	42	30	0	1	24	10	
B.	200	0	1	31	4	0	3	25	0	0	2	48	20	200	11	29	32	47	0	3	25	0	0	2	48	20	
B.	300	0	2	16	36	0	5	7	30	0	4	12	30	300	11	29	19	11	0	5	7	30	0	4	12	30	
B.	400	0	3	2	8	0	6	50	0	0	5	36	40	B.	400	0	0	4	43	0	6	50	0	0	5	36	40
B.	500	0	3	47	40	0	8	32	30	0	7	0	50	B.	800	0	0	9	26	0	13	39	59	0	11	13	10
B.	600	0	4	33	12	0	10	15	0	0	8	2	0	B.	1200	0	0	14	9	0	20	29	59	0	16	49	5
B.	700	0	5	18	44	0	11	57	30	0	9	49	10	B.	1600	0	0	18	52	0	27	19	58	0	22	26	3
B.	800	0	6	4	16	0	13	40	0	0	11	13	20	B.	2000	0	0	23	35	1	4	9	58	0	28	3	1
B.	900	0	6	49	48	0	15	22	30	0	12	37	30	B.	2400	0	0	28	18	1	10	59	57	1	3	39	5
B.	1000	0	7	35	20	0	17	5	0	0	14	1	40	B.	2800	0	0	33	1	1	17	49	57	1	9	16	3
B.	2000	0	15	10	40	1	4	10	0	0	28	3	20	B.	3200	0	0	37	44	1	24	39	56	1	14	53	1
B.	3000	0	22	46	0	1	21	15	0	1	12	5	0	B.	3600	0	0	42	27	2	1	29	56	1	20	29	5
B.	4000	1	0	21	20	2	8	20	0	1	26	6	40	B.	4000	0	0	47	10	2	8	19	55	1	26	6	3
B.	5000	1	7	56	40	2	25	25	0	2	10	8	20	B.	6000	0	1	10	45	3	12	29	53	2	24	9	5
B.	6000	1	15	32	0	3	12	30	0	2	24	10	0														

Mean

Mean MOTION of the SUN, and of the FIXED STARS, for 99 JULIAN YEARS.

Julian Years.	M. Mot. ☉.	Comp.	Mot. ☉ Apogee.	Mean Mot. fixed Stars.	Julian Years.	M. Mot. ☉.	Comp.	Mot. ☉ Apogee.	Mean Mot. fixed Stars.
s o i //	1 2 3 4 5 6 7 8 9 10 11 12	1 2 3 4 5 6 7 8 9 10 11 12	s o i //	s o i //	s o i //	s o i //	1 2 3 4 5 6 7 8 9 10 11 12	s o i //	s o i //
1	11 29 45 40	14 20	1 1	0 50	49	0 0 7 31		50 13	41 14
2	11 29 31 20	28 40	2 3	1 41	50	11 29 53 11	6 49	51 15	42 5
3	11 29 17 1	42 59	3 4	2 31	51	11 29 38 52	21 8	52 16	42 55
B. 4	0 0 1 49		4 6	3 22	52	0 0 23 41		53 18	43 46
5	11 29 47 29	12 31	5 7	4 12	53	0 0 9 21		54 19	44 36
6	11 29 33 10	26 50	6 9	5 3	54	11 29 55 1	4 59	55 21	45 27
7	11 29 18 50	41 10	7 10	5 53	55	11 29 40 42	19 18	56 22	46 17
B. 8	0 0 3 39		8 12	6 44	56	0 0 25 30		57 24	47 8
9	11 29 49 19	10 41	9 13	7 34	57	0 0 11 10		58 25	47 58
10	11 29 34 59	25 1	10 15	8 25	58	11 29 56 50	3 10	59 27	48 49
11	11 29 20 39	39 21	11 16	9 15	59	11 29 42 31	17 29	1 0 28	49 39
B. 12	0 0 5 28		12 18	10 6	60	0 0 27 19		1 1 30	50 30
13	11 29 51 8	8 52	13 19	10 56	61	0 0 12 59		1 2 31	51 20
14	11 29 36 48	23 12	14 21	11 47	62	11 29 58 39	1 21	1 3 33	52 11
15	11 29 22 29	37 31	15 22	12 37	63	11 29 44 20	15 40	1 4 34	53 1
B. 16	0 0 7 17		16 24	13 28	64	0 0 29 8		1 5 36	53 52
17	11 29 52 57	7 3	17 25	14 18	65	0 0 14 48		1 6 37	54 42
18	11 29 38 38	21 22	18 27	15 9	66	0 0 0 28		1 7 39	55 33
19	11 29 24 18	35 42	19 28	15 59	67	11 29 46 9	13 51	1 8 40	56 23
B. 20	0 0 9 6		20 30	16 50	68	0 0 30 58		1 9 42	57 14
21	11 29 54 46	5 14	21 31	17 40	69	0 0 16 38		1 10 43	58 4
22	11 29 40 26	19 34	22 33	18 30	70	0 0 2 18		1 11 45	58 55
23	11 29 26 7	33 53	23 34	19 21	71	11 29 47 59	12 1	1 12 46	59 45
B. 24	0 0 10 56		24 36	20 12	72	0 0 32 47		1 13 48	1 0 36
25	11 29 56 36	3 24	25 37	21 2	73	0 0 18 27		1 14 49	1 1 26
26	11 29 42 16	17 44	26 39	21 53	74	0 0 4 7		1 15 51	1 2 17
27	11 29 27 57	32 3	27 40	22 43	75	11 29 49 48	10 12	1 16 52	1 3 7
B. 28	0 0 12 45		28 42	23 34	76	0 0 34 36		1 17 54	1 3 53
29	11 29 58 25	1 35	29 43	24 24	77	0 0 20 16		1 18 55	1 4 48
30	11 29 44 5	15 55	30 45	25 15	78	0 0 5 56		1 19 57	1 5 39
31	11 29 29 46	30 14	31 46	26 5	79	11 29 51 37	8 23	1 20 58	1 6 29
B. 32	0 0 14 34		32 48	26 56	80	0 0 36 26		1 22 0	1 7 20
33	0 0 0 14		33 49	27 46	81	0 0 22 6		1 23 1	1 8 10
34	11 29 45 54	14 6	34 51	28 37	82	0 0 7 46		1 24 3	1 9 1
35	11 29 31 35	28 25	35 52	29 27	83	11 29 53 27	6 33	1 25 4	1 9 51
B. 36	0 0 16 24		36 54	30 18	84	0 0 38 15		1 26 6	1 10 42
37	0 0 2 4		37 55	31 8	85	0 0 23 55		1 27 7	1 11 32
38	11 29 47 44	12 16	38 57	31 59	86	0 0 9 35		1 28 9	1 12 23
39	11 29 33 25	26 35	39 58	32 49	87	11 29 55 16	4 44	1 29 10	1 13 13
B. 40	0 0 18 13		41 0	33 40	88	0 0 40 4		1 30 12	1 14 4
41	0 0 3 53		42 1	34 30	89	0 0 25 44		1 31 13	1 14 54
42	11 29 49 33	10 27	43 3	35 21	90	0 0 11 24		1 32 15	1 15 45
43	11 29 35 14	24 46	44 4	36 11	91	11 29 57 5	2 55	1 33 16	1 16 35
B. 44	0 0 20 2		45 6	37 2	92	0 0 41 53		1 34 18	1 17 26
45	0 0 5 42		46 7	37 52	93	0 0 27 33		1 35 19	1 18 16
46	11 29 51 22	8 38	47 9	38 43	94	0 0 13 13		1 36 21	1 19 7
47	11 29 37 3	22 57	48 10	39 33	95	11 29 58 54	1 6	1 37 22	1 19 57
B. 48	0 0 21 51		49 12	40 24	96	0 0 43 43		1 38 24	1 20 48
					97	0 0 29 23		1 39 25	1 21 38
					98	0 0 15 3		1 40 27	1 22 29
					99	0 0 0 44		1 41 28	1 23 10

Connect these Motions with Places for whole Hundreds, in either Style, for the present Year's Places.

C

Mean

Mean PLACES of the SUN and Fixed STARS for Years and Months, Old and New Style.

1600		Julian Style.											
Months.		Mean Pl. $\odot$.				M. Pl. Ap. $\odot$.				M. Pl. $1^{\circ} \gamma$.			
D ^s .		s	o	i	''	s	o	i	''	s	o	i	''
January	1	9	20	12	31	3	6	0	59	0	27	35	10
February	1	10	20	45	49	3	6	1	4	0	27	35	14
March	0	11	18	21	42	3	6	1	9	0	27	35	18
April	0	0	18	55	1	3	6	1	14	0	27	35	22
May	0	1	18	29	11	3	6	1	19	0	27	35	27
June	0	2	19	2	29	3	6	1	24	0	27	35	31
July	0	3	18	36	39	3	6	1	29	0	27	35	35
August	0	4	19	9	57	3	6	1	35	0	27	35	39
September	0	5	19	43	15	3	6	1	40	0	27	35	44
October	0	6	19	17	25	3	6	1	45	0	27	35	48
November	0	7	19	50	43	3	6	1	50	0	27	35	52
December	0	8	19	24	53	3	6	1	55	0	27	35	56

1600		Gregorian Style.											
Months.		Mean Pl. $\odot$.				M. Pl. Ap. $\odot$.				M. Pl. $1^{\circ} \gamma$.			
D ^s .		s	o	i	''	s	o	i	''	s	o	i	''
January	1	9	10	21	8	3	6	0	57	0	27	35	9
February	1	10	10	54	26	3	6	1	2	0	27	35	13
March	0	11	8	30	19	3	6	1	7	0	27	35	17
April	0	0	9	3	38	3	6	1	12	0	27	35	21
May	0	1	8	37	47	3	6	1	17	0	27	35	26
June	0	2	9	11	6	3	6	1	22	0	27	35	30
July	0	3	8	45	16	3	6	1	27	0	27	35	34
August	0	4	9	18	34	3	6	1	33	0	27	35	38
September	0	5	9	51	52	3	6	1	38	0	27	35	43
October	0	6	9	26	2	3	6	1	43	0	27	35	47
November	0	7	9	59	20	3	6	1	48	0	27	35	51
December	0	8	9	33	30	3	6	1	53	0	27	35	55

1700		Julian Style.											
Months.		Mean Pl. $\odot$.				M. Pl. Ap. $\odot$.				M. Pl. $1^{\circ} \gamma$.			
D ^s .		s	o	i	''	s	o	i	''	s	o	i	''
January	1	9	20	58	3	3	7	43	29	0	28	59	20
February	1	10	21	31	21	3	7	43	34	0	28	59	24
March	0	11	19	7	14	3	7	43	39	0	28	59	28
April	0	0	19	40	33	3	7	43	44	0	28	59	32
May	0	1	19	14	43	3	7	43	49	0	28	59	37
June	0	2	19	48	1	3	7	43	54	0	28	59	41
July	0	3	19	22	11	3	7	43	59	0	28	59	45
August	0	4	19	55	29	3	7	44	5	0	28	59	49
September	0	5	20	28	47	3	7	44	10	0	28	59	54
October	0	6	20	2	57	3	7	44	15	0	28	59	58
November	0	7	20	36	15	3	7	44	20	0	29	0	2
December	0	8	20	10	25	3	7	44	25	0	29	0	6

1700		Gregorian Style.											
Months.		Mean Pl. $\odot$.				M. Pl. Ap. $\odot$.				M. Pl. $1^{\circ} \gamma$.			
D ^s .		s	o	i	''	s	o	i	''	s	o	i	''
January	0	9	10	7	32	3	7	43	27	0	28	59	19
February	0	10	10	40	50	3	7	43	32	0	28	59	23
March	0	11	8	16	43	3	7	43	37	0	28	59	27
April	0	0	8	50	2	3	7	43	42	0	28	59	31
May	0	1	8	24	12	3	7	43	47	0	28	59	36
June	0	2	8	57	30	3	7	43	52	0	28	59	40
July	0	3	8	31	40	3	7	43	57	0	28	59	44
August	0	4	9	4	58	3	7	44	3	0	28	59	48
September	0	5	9	38	16	3	7	44	8	0	28	59	53
October	0	6	9	12	26	3	7	44	13	0	28	59	57
November	0	7	9	45	44	3	7	44	18	0	29	0	1
December	0	8	9	19	54	3	7	44	23	0	29	0	5

1800		Julian Style.											
Months.		Mean Pl. $\odot$.				M. Pl. Ap. $\odot$.				M. Pl. $1^{\circ} \gamma$.			
D ^s .		s	o	i	''	s	o	i	''	s	o	i	''
January	1	9	21	43	35	3	9	25	59	1	0	23	30
February	1	10	22	16	53	3	9	26	4	1	0	23	34
March	0	11	19	52	46	3	9	26	9	1	0	23	38
April	0	0	20	26	5	3	9	26	14	1	0	23	42
May	0	1	20	0	15	3	9	26	19	1	0	23	47
June	0	2	20	33	33	3	9	26	24	1	0	23	51
July	0	3	20	7	43	3	9	26	29	1	0	23	55
August	0	4	20	41	1	3	9	26	35	1	0	23	59
September	0	5	21	14	19	3	9	26	40	1	0	24	4
October	0	6	20	48	29	3	9	26	45	1	0	24	8
November	0	7	21	21	47	3	9	26	50	1	0	24	12
December	0	8	20	55	57	3	9	26	55	1	0	24	16

1800		Gregorian Style.											
Months.		Mean Pl. $\odot$.				M. Pl. Ap. $\odot$.				M. Pl. $1^{\circ} \gamma$.			
D ^s .		s	o	i	''	s	o	i	''	s	o	i	''
January	0	9	9	53	55	3	9	25	57	1	0	23	28
February	0	10	10	27	13	3	9	26	2	1	0	23	32
March	0	11	8	3	6	3	9	26	7	1	0	23	36
April	0	0	8	36	25	3	9	26	12	1	0	23	40
May	0	1	8	10	34	3	9	26	17	1	0	23	45
June	0	2	8	43	53	3	9	26	22	1	0	23	49
July	0	3	8	18	3	3	9	26	27	1	0	23	53
August	0	4	8	51	21	3	9	26	33	1	0	23	57
September	0	5	9	24	39	3	9	26	38	1	0	24	2
October	0	6	8	58	49	3	9	26	43	1	0	24	6
November	0	7	9	32	7	3	9	26	48	1	0	24	10
December	0	8	9	6	17	3	9	26	53	1	0	24	14

In Leap-Year take out the Month Day Motions for January and February 1 Day sooner, in both Styles.

In Leap-Year take out the Month Day Motions for January and February 1 Day sooner, in both Styles.

EXAMPLE I.

To find the Place of the Sun, Apogee, and $1^{\circ} \gamma$, for January 11, 1632, at Noon, O. S. ?

	$\odot$	Ap. $\odot$	$1^{\circ} \gamma$
O.S. Jan. 1. 1600.	9 20 12 31	3 6 0 59	0 27 35 10
Years 32	0 14 34	32 48	26 56
B. Days 10	9 51 23	2	1
O.S. Jan. 11. 1632	10 0 18 28	3 6 33 49	0 28 2 7

EXAMPLE II.

To find the Place of the Sun, Apogee, and $1^{\circ} \gamma$ for October 29, 1820, at Noon, N. S. ?

	$\odot$	Ap. $\odot$	$1^{\circ} \gamma$
N.S. Oct. 0. 1800	6 8 58 49	3 9 26 43	1 0 24 6
Years 20	0 9 6	20 30	16 50
Days 29	28 35 2	5	4
N.S. Oct. 29. 1820	7 7 42 57	3 9 47 18	1 0 41 0

Mean

Mean Motion of the SUN, and of the Fixed Stars, continued, for Months, Days, &c.

D ^s .	Mths.	M. Mot. $\odot$.				Ap. *s.	Chronological Eras before and since Christ.
		s	o	i	h		
Jan.	o	0	0	0	0	0	776 Olympiads, 4 Years each, D after Summer Solstice.
31 Feb.	o	1	0	33	18	5	4 752 Building of Rome, April 21.
59 March	o	1	28	9	11	10	8 746 Nabonassar, February 26.
90 April	o	2	28	42	30	15	12 323 Death Alex. the Great, Nov. 12.
120 May	o	3	28	16	40	20	17 45 Calendar settled by J. Cæf. Jan. 1.
151 June	o	4	28	49	58	25	21 0 Birth of Christ.
181 July	o	5	28	24	8	30	25 284 Diocletian, August 29.
212 August	o	6	28	57	26	36	29 325 Council of Nice.
243 Sept.	o	7	29	30	44	41	34 632 Hegira of Turks, July 16.
273 Oct.	o	8	29	4	54	46	38 632 Jesdegird of Persians, June 16, O.S.
304 Nov.	o	9	29	38	12	51	42 1582 Gregorian Style began October
334 Dec.	o	10	29	12	22	56	46 1752 Began in England, Sept. 3, O. S. or Sept. 14, N. S.

Take out 1 Day sooner for January and February, in Leap-year.

Days	M. Mo. $\odot$.				H. Min. Sec.	H. Min. Sec.
	o	i	h	iv		
1	0	0	59	8	0	0
2	0	1	58	17	0	0
3	0	2	57	25	0	0
4	0	3	56	33	1	0
5	0	4	55	42	1	1
6	0	5	54	50	1	1
7	0	6	53	58	1	1
8	0	7	53	7	1	1
9	0	8	52	15	1	1
10	0	9	51	23	2	1
11	0	10	50	32	2	1
12	0	11	49	40	2	2
13	0	12	48	48	2	2
14	0	13	47	57	2	2
15	0	14	47	5	2	2
16	0	15	46	13	3	2
17	0	16	45	22	3	2
18	0	17	44	30	3	2
19	0	18	43	38	3	3
20	0	19	42	47	3	3
21	0	20	41	55	3	3
22	0	21	41	3	4	3
23	0	22	40	12	4	3
24	0	23	39	20	4	3
25	0	24	38	28	4	3
26	0	25	37	37	4	4
27	0	26	36	45	4	4
28	0	27	35	53	5	4
29	0	28	35	2	5	4
30	0	29	34	10	5	4
31	1	0	33	18	5	4

EXAMPLE I.

To find the Mean Place of the Sun, his Apogee, and the first * γ , January 20, 1758, at Noon, New Style?

	M. Pla $\odot$.	Ap. $\odot$.	1* γ .
N. S. 1700	9 10 7 32	3 7 43 27	0 28 59 19
Mo. Yrs 58	— 3 10	+ 59 27	+ 48 49
Pla ^s . 1758	9 10 4 22	3 8 42 54	0 29 48 8
Mo. Jan. 20	0 19 42 47	+ + 3	+ 3
M. Pl. reqd.	9 29 47 9	3 8 42 57	0 29 48 11
		9 29 47 9	
		6 21 4 12	⊙ M. Anom.

EXAMPLE II.

To find the Place of the Sun, Apogee, and first * γ , January 1, at Midnight, 1804, New Style?

	$\odot$.	Ap. $\odot$.	1* γ .
N. S. 1800	9 9 53 55	3 9 25 57	1 0 23 28
Mo. Yrs 4	+ 1 49	+ 4 6	+ 3 22
B. Jan. 1st 1804	9 9 55 44	3 9 30 3	1 0 26 50
Hours 12	+ 29 34		
	9 10 25 18	6 0 55 15	⊙ M. Anom.

EXAMPLE III.

To find the Place of the Sun, Apogee and 1* γ , February 29, 19^h 13^m 52^s, P. M. 1848, New Style?

	$\odot$.	Ap. $\odot$.	1* γ .
N. S. 1800	9 9 53 55	3 9 25 57	1 0 23 28
Mo. Yrs 48	+ 21 51	+ 49 12	+ 40 24
February	1 0 33 18	5	4
B. Days 28	0 27 35 53	5	4
Hours 19	46 49		
Mins. 13	32	3 10 15 19	1 1 4 c
Secs 52	2		
	11 9 12 20	11 9 12 20	
		7 28 57	⊙ M. Anom.

The same may be performed by reducing the Days of the Months in the above Examples to the Old Style.

NOTE, The Gregorian is reduced to the Julian Style, (serving equally for Computation) by subtracting the Number of Days Difference. And, the contrary, by adding the Days Difference, the Julian is reduced to the Gregorian Style. — To determine universally the Number of Days Difference betwixt New and Old Style.

RULE. — From the Hundreds of CHRIST, take their Fourth, adding Two; And the Days will remain betwixt Old Style and New.

EXAMPLE. — In 27/40

$$2 + \frac{1}{4} = 8 -$$

Rem. 19 Days Difference.

EQUATION of the SUN's CENTER, or of his *Mean PLACE* or ANOMALY to the True.

Argument. SUN's Mean Anomaly.

⊙ An.	Sign	Diff.	Sig. 1	Diff.	Sig. 2	Diff.	Sig. 3	Diff.	Sig. 4	Diff.	Sig. 5	Diff.	⊙ An.
0	0	0	0	0	0	0	0	0	0	0	0	0	0
1	0	1	59	1	59	0	58	36	1	44	1	40	16
2	0	3	58	1	59	1	0	18	1	42	1	41	15
3	0	5	57	1	59	1	1	59	1	41	1	42	13
4	0	7	56	1	59	1	3	39	1	40	1	43	8
5	0	9	54	1	58	1	5	18	1	39	1	44	2
6	0	11	52	1	58	1	6	57	1	39	1	44	54
7	0	13	50	1	58	1	8	33	1	36	1	45	44
8	0	15	48	1	58	1	10	9	1	36	1	46	33
9	0	17	45	1	57	1	11	43	1	34	1	47	19
10	0	19	42	1	57	1	13	16	1	33	1	48	3
11	0	21	39	1	57	1	14	48	1	32	1	48	45
12	0	23	36	1	57	1	16	19	1	31	1	49	25
13	0	25	32	1	56	1	17	47	1	28	1	50	4
14	0	27	28	1	56	1	19	16	1	29	1	50	41
15	0	29	23	1	55	1	20	42	1	26	1	51	15
16	0	31	17	1	54	1	22	7	1	25	1	51	48
17	0	33	11	1	54	1	23	30	1	23	1	52	19
18	0	35	5	1	54	1	24	52	1	22	1	52	47
19	0	36	58	1	53	1	26	13	1	21	1	53	14
20	0	38	51	1	53	1	27	33	1	20	1	53	38
21	0	40	43	1	52	1	28	50	1	17	1	54	1
22	0	42	34	1	51	1	30	6	1	16	1	54	21
23	0	44	24	1	51	1	31	20	1	14	1	54	39
24	0	46	13	1	49	1	32	33	1	12	1	54	55
25	0	48	2	1	49	1	33	44	1	11	1	55	9
26	0	49	50	1	48	1	34	54	1	10	1	55	21
27	0	51	37	1	47	1	36	2	1	8	1	55	31
28	0	53	23	1	46	1	37	8	1	6	1	55	39
29	0	55	8	1	45	1	38	13	1	5	1	55	45
30	0	56	52	1	44	1	39	15	1	2	1	55	49
⊙ An.	+	Diff.	+	Diff.	+	Diff.	+	Diff.	+	Diff.	+	Diff.	⊙ An.
	Sig. 11.		Sig. 10		Sig. 9		Sig. 8		Sig. 7		Sig. 6		

Construction. The present Eccentricity of the Earth's Orbit is 1685 to the Semi-Transverse or mean Dist. $\odot \text{ to } \odot 100000$: Whence the Aphelion Dist. is 101685 to the Perihelion Dist. 98315—Now by this RULE. As *Aph. Dist.* : *Perihelion Dist.* :: *Tang. $\frac{1}{2}$ mean Anom.* : *Tang. $\frac{1}{2}$ true Anom.* (nearly $\frac{1}{2}$ true Anom.) which added to $\frac{1}{2}$ mean Anom. the Sum will be the correct Eccentric Anom. from Aphelion. Again, As $\sqrt{\text{Aph. Dist.}}$: $\sqrt{\text{Perib. Dist.}}$:: *Tan. $\frac{1}{2}$ cor. Eccen. An.* : *Tan. $\frac{1}{2}$ true An.*

EXAMPLE. To find the Equation to mean Anom. 30° ?

	o	i	n	Log.		Log.
$\frac{1}{2}$ Eccen. An.	14	45	43	$\frac{1}{2}$ Tan. 9.4207865	Half or 15°	Tan. 9.4280525
				Const. Log. 9.9926815 +		Const. Log. 9.9853630 +
$\frac{1}{2}$ True An.	14	31	34	.. Tan. 9.4134680	4th. $14^\circ 31' 27''$	Tan. 9.4134155
$\frac{1}{2}$ Mean Anom.	15				+ $\frac{1}{2}$ M. Anom. =	29 31 27 Cor. Eccen. An.
Rem. $\frac{1}{2}$ Equat.	0	28	26, and 56' 52''	Equation required.		

N. B. A Dot or Point after a Decimal shows the Decimal to be complete; otherwise the last Figure is understood to repeat continually.

"X"	"	"	"	"	"	"	"	"	"	"X"	"	"	"	"	"	"	"	"	"
"X"	"	"	"	"	"	"	"	"	"	"X"	"	"	"	"	"	"	"	"	"
"X"	"	"	"	"	"	"	"	"	"	"X"	"	"	"	"	"	"	"	"	"
"	"	"	"	"	"	"	"	"	"	"	"	"	"	"	"	"	"	"	"
"	10	20	30	40	50	60	70	80	90	"	10	20	30	40	50	60	70	80	90
1	16	33	5	66	83	1	16	133	15	31	516	1033	155	2066	2583	31	3616	4133	465
2	33	66	1	133	166	2	233	266	3	32	533	1066	16	2133	2666	32	3733	4266	48
3	5	1	15	2	25	3	35	4	45	33	55	11	165	22	275	33	385	44	495
4	66	133	2	266	333	4	466	533	6	34	566	1133	17	2266	2833	34	3966	4533	51
5	83	166	25	333	416	5	583	666	75	35	583	1166	175	2333	2916	35	4083	4666	525
	10	20	30	40	50	60	70	80	90		10	20	30	40	50	60	70	80	90
6	1	2	3	4	5	6	7	8	9	36	6	12	18	24	30	36	42	48	54
7	16	233	35	466	583	7	816	933	105	37	616	1233	185	2466	3083	37	4316	4933	555
8	133	266	4	533	666	8	933	1066	12	38	633	1266	19	2533	3166	38	4433	5066	57
9	15	3	45	6	75	9	105	12	135	39	65	13	195	26	325	39	455	52	585
10	166	333	5	666	833	10	1166	1333	15	40	666	1333	20	2666	3333	40	4666	5333	60
	10	20	30	40	50	60	70	80	90		10	20	30	40	50	60	70	80	90
11	183	366	55	133	916	11	1283	1466	165	41	683	1366	205	2733	3416	41	4783	5466	615
12	2	4	6	8	10	12	14	16	18	42	7	14	21	28	35	42	49	56	63
13	216	433	65	866	1083	13	1516	1733	195	43	716	1433	215	2866	3583	43	5016	5733	645
14	233	466	7	933	1166	14	1633	1866	21	44	733	1466	22	2933	3666	44	5133	5866	66
15	25	5	75	10	125	15	175	20	225	45	75	15	225	30	375	45	525	60	675
	10	20	30	40	50	60	70	80	90		10	20	30	40	50	60	70	80	90
16	266	533	8	1066	1333	16	1866	2133	24	46	766	1533	23	3066	3833	46	5366	6133	69
17	283	566	85	1133	1416	17	1983	2266	255	47	783	1566	235	3133	3916	47	5483	6266	705
18	3	6	9	12	15	18	21	24	27	48	8	16	24	32	40	48	56	64	72
19	316	633	95	1266	1583	19	2216	2533	285	49	816	1633	245	3266	4083	49	5716	6533	735
20	333	666	10	1333	1666	20	2333	2666	30	50	833</								

EXAMPLE I. *The Sun's Equation for $1^{\circ} 15^{\circ} 39' 57''$ Sun's Anomaly is required?*

18 150 | -10 20' 42" First Equation.

I 16 | -1 22 7

I : $+1\ 25=85''$ Dif. Equan : : $39'\ 57'' : 56'', 58, \&c.$
by $39\ 57$ Dif. Argument.

$+ 56$ to the first Equation.

⊙ Equation — 1 21 38 fought.

By TABLE.

$$39' \times 80'' = 52''$$

$$5 = 3,25$$

$$55,25 = 55'' 15'''$$

$$57'' \times 80'' = 76'' = 1 \text{ } 16$$

$$5 = 4''; 75 = \underline{\quad} \quad 4 \text{ } 45 \text{ iv.}$$

Proport^l Equation $\div 56 \quad 35 \quad 45$ correct.

N. B. Under *Units*, 1, 2, 3, 4, 5, 6, 7, 8, or 9, a-top, against any *N^o.* of *Min^s* or *Sec^s* on the *Side*, take out a *Decimal* more than for *tens*; and for *Tenths* 2 more; for *Hundredths* 3 more, &c.

EXAMPLES

EXAMPLES of proportioning the DIFFERENCE of EQUATIONS.

EXAMPLE II. Sun's Place at Noon, June 10, 1756, $19^{\circ} 52' 0''$ required his Right Ascension?

☉ Pl. R. A.

$$\begin{array}{r} 19^{\circ} \quad 78^{\circ} 2' 3'' \\ 20 \quad 79 \quad 7 \quad 4 \end{array}$$
As Dif. $1^{\circ} : 1' 5'' = 65'. 1''$. Dif. increasing $52'$. (Argt. above Degs.) $+ 56'. 20''. 52'''$. Prop. $\text{Æquan.} = \frac{52' \times 65'. 1''}{60'}$

By TABLE.

$$\begin{array}{r} 52' \times 60' \dots 52' \\ 5 \dots 4.33 \\ 1'' \dots 56.33 \dots 56' 20'' 0''' \\ \quad \quad \quad 52 \\ + 56 \quad 20 \quad 52 \\ \hline 78^{\circ} \quad 2 \quad 3 \quad 0 \end{array}$$

☉ R. A. required. $78^{\circ} 58' 23'' 52''$ Æquan. sought.

DECIMALLY.

$$\begin{array}{r} 65' 1'' = 65.016 \\ 52' = 52. \\ \hline 130032 \\ 325080 \\ \hline 6)3380832 \\ 56' 20'' 50''' \text{ Prop. } \text{Æq.} \end{array}$$

SEXAGESIMALLY.

Divide each $1''$ $\{ 65' \times 1'' \}$ Dif. Æq. $\times d$
 Prod. exceed $52' \{ 52' \times 0'' \}$ Dif. Arg. $\}$ togeth.
 ing 60 by 6, $52''' 130$
 (when a Fig. 325
 to the Right is cut off) taking $6)3380''$
 the Sum of the $56' 20'' 52'''$ Prop. Æquan.
 Quotients. [sought.]

SEXAGESIMALS are thus turned into DECIMALS, and the contrary, without a TABLE.

Seconds are mentally turned into Decimals of a Minute, by dividing them by 6, supposing 0 to be placed before the Remainder, continually.—The first Decimal, thence arising, will be Tenths, the second, Hundredths, &c. except when the Seconds to be turned into Decimals are under 6; then the first Decimal arising (first supposing 0 placed before the Seconds to be divided by 6) will be Hundredths, the next Thousandths, &c. where an 0 must stand in Tenths Place.

Minutes are mentally turned into Decimals of a Degree, in the same Manner; Thirds into Decimals of a Second, &c. (See an Example to the Right). And dividing the lowest Denomination by 6, then that Quotient, and the next Denomination bigger by 6, as far as Degrees, or Hours, the Decimals of Sexagesimals are thus found, generally.

Decimals (of Degrees, Hours, and of lower Order) are universally turned into Sexagesimals, by multiplying all the Decimals to the Right Hand by 6, and cutting off the Product of the first Decimal, with what is carried from the rest, for the next Denomination lower, continually.—Two or three Decimals to the Right being thus multiplied by 6, will determine the Value of the next Sexagesimal, mentally, or at Sight.

IN SEXAGESIMAL PROPORTION,

Minutes $\times$ Minutes = Seconds; Minutes $\times$ Seconds = Thirds; Seconds $\times$ Seconds = Fourths; Seconds $\times$ Thirds = Fifths. Hence, a Figure to the Right of a Sexagesimal being cut off, and all the Figures to the Left divided by 6, will give the Value of that Product, in Sexagesimals, a Denomination higher.—The Remainder, before the Figure cut off, to be placed before that Figure, for the Value of the lower Denomination.

EXAMPLE of turning SEXAGESIMALS into DECIMALS, and the contrary.

Divided by 6. $23''' \dots = 23'''$
 $34'', 383 \dots = 34'' 23$
 $29', 57305 \dots = 29' 34 23$
 $40^{\circ}, 492884259 = 40^{\circ} 29 34 23$
 Multiplied by 6. $29', 5730554 \dots$ N. B. A Dot over any Figure denotes a continual Repetition from thence.
 $34'', 383324$
 $22'' 999944$

EXAMPLE III. To find the Proportional Equation from $39' 40''$ Dif. Equation, and $53' 35''$ Difference of Argument?

By TABLE.

$$\begin{array}{r} 53' \times 30' \dots 26', 5 \\ 9 \dots 7.95 \\ \hline 34.45 \dots 34' 27'' 0''' \text{ oiv} \\ 40'' \dots 35.33 \text{ \&c.} \dots 35 20 \quad 0 \\ 39' \times 30'' \dots 19', 5 \\ 5 \dots 3.25 \\ \hline 22.75 \dots 22 45 \quad 0 \\ 35'' \times 40'' \dots 23'', 33 \dots 23 20 \end{array}$$
35 25 28 20 Prop. Æq.

DECIMALLY.

$$\begin{array}{r} 39' 40'' = 39.66 \\ 53 \quad 35 = 53.58 \\ \hline 31728 \\ 19830 \\ \hline 11898 \\ 19830 \\ \hline 6)21249828 \\ 35' 24'' \text{ fere.} \end{array}$$

SEXAGESIMALLY.

$$\begin{array}{r} 40'' \{ \text{Dif. } \text{Æq.} \} 39' \times 40'' \{ 39' \mid 40'' \\ 53'' \{ \text{Dif. Arg.} \} 53' \times 35'' \{ 53' \mid 35'' \\ \hline 120 \dots 117 \dots 195 \dots 200 \\ 200 \dots 195 \dots 117 \dots 120 \\ \hline 2120'' \dots 2067'' \dots 1365''' \dots 140 \text{ oiv} \\ 35'' 20''' \dots 34' 27'' \dots 22' 45''' \dots 23''' 20 \text{ oiv} \\ \hline 35 20''' \\ 22 45 \\ \hline 23 20 \text{ oiv} \\ \hline \text{Sum } 35 25 28 20, \text{ correctly.} \end{array}$$

By LOGISTICAL LOGARITHMS.

L. L.

Dif. Argt. $53' 35''$ $\{ 0491$ } N. B. These Logistical Logarithms proportion the easiest, but not always the quickest, as in a direct Manner, which is less disturbing to Attention, than the turning (backward and forward) over Leaves.
 Dif. Æq. $39 40$ $\{ 1797$
 Pr. Æq. $35 25$ $\{ 2288$

By Aliquot PARTS of a DEGREE

UNIVERSALLY and DIRECTLY.

[or 60'. $35''$]
 Thus $39' 40''$ of $53', 583$ 60'
 $30' \cdot \frac{1}{2} \cdot 60' \dots 26,791$ 30
 $6' \cdot \frac{1}{10} \cdot 60' \dots 5,358$ 6
 $3' \cdot \frac{1}{20} \cdot 60' \dots 2,679$ 3
 $40'' \cdot \frac{1}{9} \cdot 60' \dots 595$ 40''
 $39' 40'' \dots 35', 423$ sum
 $25'' 38$
 OR, $53' 35''$ of $39', 666$ 60'
 $48' \cdot \frac{8}{10} \times 10.31,732$ 48'
 $5' \cdot \frac{5}{10} \times 1'. 3,305$ 5
 $30'' \cdot \frac{5}{10} \times 1'. 330$ 30''
 $5' \cdot \frac{5}{10} \times 1''. 055$ 5
 $53' 35'' \dots 35', 422$ sum
 $25'' 32$
 $.6611 = 1'$
 $.01102 = 1''$
 Multy 1° and $1'$ by as many Tenths as there are 6 Min. & 6 Secs, adding for Min. & Secs. over.

Otherwise by the TABLE.

$$\begin{array}{r} 39' 40'' \mid 53', 58 \\ 39' \times 50' \dots 32', 5 \\ 3 \dots 1,95 \\ 5 \dots 3,25 \\ 08 \dots 0,52 \\ \hline 34,827 \dots 34' 49'' 37''' \\ 40'' \times 50' \dots 33'', 33 \\ 3 \dots 2 \\ \hline 35,33 \dots 35 19 \\ \hline \text{Prop. } \text{Æq.} \quad 35 \quad 24 \quad 56 \end{array}$$

NOTE. A Table of aliquot Parts, and aliquot Parts of those, will expedite the above Proportion. Universally. As $23' 12''$ to $39' 19''$ so $44' 53''$ to $76' 3'' 47'''$
 Decimally. As 23, 2 to 39, 31 so 44, 88 to $76', 044 = 76' 3''$ fere, at Sight.

The SUN's *proportional* DISTANCES from the EARTH to the Sun's Mean Distance from thence, 100000.—
The Eccentricity of the Earth's Orbit 1685 of such Parts.

Argument. SUN's Mean Anomaly.

◉M. An.	Sig. 0 ◉ à ◉	Dif.	Sig. 1 ◉ à ◉	Dif.	Sig. 2 ◉ à ◉	Dif.	Sig. 3 ◉ à ◉	Dif.	Sig. 4 ◉ à ◉	Dif.	Sig. 5 ◉ à ◉	Dif.	◉M. An.
0													0
0	101685	0	101466	14	100864	25	100028	30	99180	26	98550	15	30
1	101685	1	101452	15	100839	25	99998	30	99154	25	98535	15	29
2	101684	1	101437	15	100814	26	99968	29	99129	25	98520	14	28
3	101683	2	101422	16	100788	26	99939	29	99104	25	98506	14	27
4	101681	2	101406	17	100762	26	99910	29	99079	25	98492	14	26
5	101679	3	101389	17	100736	27	99881	29	99054	25	98479	13	25
6	101676	3	101372	17	100709	27	99852	29	99029	25	98466	12	24
7	101673	4	101355	17	100682	27	99823	29	99004	24	98454	12	23
8	101669	4	101338	18	100655	27	99794	29	98980	24	98442	11	22
9	101665	5	101320	18	100628	27	99765	29	98956	23	98431	10	21
10	101660	5	101302	18	100601	27	99736	29	98933	22	98421	10	20
11	101655	6	101284	19	100574	27	99707	29	98911	22	98411	10	19
12	101649	6	101265	19	100547	28	99678	29	98889	22	98401	9	18
13	101643	7	101246	20	100519	28	99649	29	98867	22	98392	9	17
14	101636	7	101226	20	100491	28	99620	29	98845	21	98383	8	16
15	101629	7	101206	20	100463	28	99591	29	98824	21	98375	8	15
16	101622	8	101186	21	100435	28	99562	28	98803	21	98367	7	14
17	101614	8	101165	21	100407	28	99534	28	98782	20	98360	7	13
18	101606	9	101144	22	100379	29	99506	28	98762	20	98353	6	12
19	101597	10	101122	22	100350	29	99478	28	98742	20	98347	5	11
20	101587	10	101100	23	100321	29	99450	28	98722	19	98342	5	10
21	101577	11	101077	23	100292	29	99422	28	98703	19	98337	4	9
22	101566	11	101054	23	100263	29	99394	28	98684	18	98333	4	8
23	101555	12	101031	23	100234	29	99366	27	98666	18	98329	4	7
24	101543	12	101008	23	100205	29	99339	27	98648	17	98325	3	6
25	101531	12	100985	23	100176	29	99312	27	98631	17	98322	3	5
26	101519	13	100962	24	100147	29	99285	26	98614	17	98319	2	4
27	101506	13	100938	24	100118	30	99259	27	98597	16	98317	1	3
28	101493	13	100914	25	100088	30	99232	26	98581	16	98316	1	2
29	101480	14	100889	25	100058	30	99206	26	98565	15	98315	0	1
30	101466		100864	25	100028	30	99180		98550		98315		0
◉M. An.	Sig. 11 ◉ à ◉	Dif.	Sig. 10 ◉ à ◉	Dif.	Sig. 9 ◉ à ◉	Dif.	Sig. 8 ◉ à ◉	Dif.	Sig. 7 ◉ à ◉	Dif.	Sig. 6 ◉ à ◉	Dif.	◉M. An.

CONSTRUCTION. As Sine $\frac{1}{2}$ true Anom. : Sine $\frac{1}{2}$ excent. An. :: $\sqrt{\text{Perihelion Dist.}}$: $\sqrt{\text{Dist. fr. Sun}}$; the Square of which is the Distance, or the doubled Logarithm, the Log. of the Distance from the Sun required.

EXAMPLE. To determine the Sun's proportional Distance from the Earth for 30° M. Anom.?

$\frac{1}{2}$ True Anom.	14 31 33,3 .. Sine	0,6006415 Co.
$\frac{1}{2}$ Eccen. Anom.	14 45 43 $\frac{1}{2}$.. Sine	9,4062095
$\sqrt{\text{Perih. Dist.}}$	98315 :	2,4963099

Proportional Distance from the Sun 101466 . . 5,0063218 Doubled.

N. B. In this Proportion the true and excentric Anomalies must be taken to Tenths of a Second, or to Thirds, for the proportional Distance from the Sun to come out correct : The above Table being a Work of great Labour.

LOGARITHMS of the foregoing proportional Distances of the SUN from the EARTH, to the Mean Distance 100000; and the Eccentricity of the Earth's Orbit 1685.

Argument. SUN's Mean ANOMALY.

© M. An	Sig. 0 Logar.	Dif.	Sig. 1 Logar.	Dif.	Sig. 2 Logar.	Dif.	Sig. 3 Logar.	Dif.	Sig. 4 Logar.	Dif.	Sig. 5 Logar.	Dif.	© M. An
0	5,0072568		5,0063205	599	5,0037362	1077	5,0001216	1303	4,9964241	1139	4,9936566	661	30
1	5,0072568	0	5,0062606	642	5,0036285	1077	4,9999913	1303	4,9963102	1195	4,9935905	661	29
2	5,0072526	42	5,0061964	644	5,0035208	1120	4,9998610	1260	4,9962007	1195	4,9935244	617	28
3	5,0072483	43	5,0061322	686	5,0034088	1120	4,9997350	1260	4,9960912	1096	4,9934627	617	27
4	5,0072398	85	5,0060636	728	5,0032968	1121	4,9996090	1261	4,9959816	1096	4,9934010	574	26
5	5,0072312	86	5,0059908	728	5,0031847	1164	4,9994829	1261	4,9958720	1096	4,9933436	573	25
6	5,0072184	118	5,0059180	728	5,0030683	1165	4,9993568	1262	4,9957624	1097	4,9932863	529	24
7	5,0072056	128	5,0058452	729	5,0029518	1165	4,9992306	1262	4,9956527	1053	4,9932334	530	23
8	5,0071885	171	5,0057723	771	5,0028353	1165	4,9991044	1262	4,9955474	1053	4,9931804	485	22
9	5,0071714	171	5,0056952	772	5,0027188	1165	4,9989782	1263	4,9954421	1009	4,9931319	441	21
10	5,0071501	213	5,0056180	772	5,0026023	1166	4,9988519	1263	4,9953412	966	4,9930878	442	20
11	5,0071287	214	5,0055408	814	5,0024857	1166	4,9987256	1263	4,9952446	966	4,9930436	441	19
12	5,0071031	256	5,0054594	815	5,0023691	1209	4,9985993	1264	4,9951480	966	4,9929995	397	18
13	5,0070775	256	5,0053779	858	5,0022482	1210	4,9984729	1264	4,9950514	967	4,9929598	397	17
14	5,0070475	300	5,0052921	858	5,0021272	1211	4,9983465	1264	4,9949547	923	4,9929201	354	16
15	5,0070176	299	5,0052063	859	5,0020061	1210	4,9982201	1265	4,9948624	923	4,9928847	353	15
16	5,0069877	299	5,0051204	903	5,0018851	1211	4,9980936	1221	4,9947701	923	4,9928494	309	14
17	5,0069536	341	5,0050301	900	5,0017640	1212	4,9979715	1222	4,9946778	879	4,9928185	309	13
18	5,0069193	343	5,0049401	945	5,0016428	1254	4,9978483	1223	4,9945899	880	4,9927876	265	12
19	5,0068809	384	5,0048456	944	5,0015174	1256	4,9977270	1222	4,9945019	880	4,9927611	221	11
20	5,0068381	428	5,0047512	989	5,0013918	1255	4,9976048	1223	4,9944139	835	4,9927390	220	10
21	5,0067954	427	5,0046523	988	5,0012663	1256	4,9974825	1223	4,9943304	837	4,9927170	177	9
22	5,0067483	471	5,0045535	989	5,0011407	1256	4,9973602	1224	4,9942467	792	4,9926993	177	8
23	5,0067013	470	5,0044546	989	5,0010151	1257	4,9972378	1180	4,9941675	792	4,9926816	176	7
24	5,0066500	513	5,0043557	989	5,0008894	1257	4,9971198	1181	4,9940883	749	4,9926640	133	6
25	5,0065986	514	5,0042568	989	5,0007637	1258	4,9970017	1181	4,9940134	748	4,9926507	132	5
26	5,0065473	513	5,0041579	1032	5,0006379	1257	4,9968836	1137	4,9939386	749	4,9926375	89	4
27	5,0064917	556	5,0040547	1033	5,0005122	1302	4,9967699	1182	4,9938637	705	4,9926286	44	3
28	5,0064361	556	5,0039514	1076	5,0003820	1302	4,9966517	1138	4,9937932	705	4,9926242	44	2
29	5,0063804	557	5,0038438	1076	5,0002518	1302	4,9965379	1138	4,9937227	661	4,9926198	0	1
30	5,0063205	599	5,0037362		5,0001216		4,9964241		4,9936566		4,9926198		0
© M. An	Sig. 11 Logar.	Dif.	Sig. 10 Logar.	Dif.	Sig. 9 Logar.	Dif.	Sig. 8 Logar.	Dif.	Sig. 7 Logar.	Dif.	Sig. 6 Logar.	Dif.	© M. An

EXAMPLE. To find the Logarithm of the Sun's Distance answering to the Sun's Mean Anomaly $4^{\circ} 26' 19'' 37''$?

By DECIMALS.
Anomaly $4^{\circ} 26' \dots$ 4,9939386
For $19' 37'' \dots$ — 245
Log. $\odot \rightarrow \ominus$ 4,9939141 required.

By PARTS.
 $60'$ or $1^{\circ} \dots$ 749 | $12,5=1'$
 $18' \dots, 3 \times 1^{\circ}$ 224,7 | $,208=1''$
 $1' \dots \dots \dots$ 12,5
 $36'', 6 \times 1'$ 7,5
 $1'' \dots \dots \dots$ 2
 $19' 37'' \dots$ — 244,9 | Answer.

By LOGISTICAL LOGARITHMS.
Difference 749 decreasing
749 = $12' 29''$ L. L. 0,6818
19 37 0,4855
— 245 = 4 5 | 1,1673

6)1468|7,89
— 245 fere.

Mean

EQUATION of the SUN's true ANOMALY, or PLACE, to the MEAN.

Argument. SUN's true ANOMALY.

☉ true An.	Sig. 0	Dif.	Sig. 1	Dif.	Sig. 2	Dif.	Sig. 3	Dif.	Sig. 4	Dif.	Sig. 5	Dif.	☉ true An.
0	+		+		+		+		+		+		0
0	0 0 0		0 58 34		1 40 58		1 55 51		1 39 42		0 57 17		30
1	0 2 3	2 3	1 0 19	1 45	1 41 56	0 58	1 55 49	0 6	1 38 39	1 3	0 55 31	1 46	29
2	0 4 6	2 3	1 2 3	1 44	1 42 54	0 54	1 55 43	0 6	1 37 36	1 3	0 53 45	1 46	28
3	0 6 9	2 3	1 3 45	1 42	1 43 48	0 54	1 55 37	0 9	1 36 30	1 6	0 51 59	1 46	27
4	0 8 12	2 3	1 5 27	1 42	1 44 42	0 51	1 55 28	0 11	1 35 23	1 7	0 50 11	1 48	26
5	0 10 14	2 2	1 7 9	1 42	1 45 33	0 50	1 55 17	0 12	1 34 13	1 10	0 48 23	1 48	25
6	0 12 16	2 2	1 8 47	1 38	1 46 23	0 49	1 55 5	0 17	1 33 1	1 12	0 46 33	1 50	24
7	0 14 18	2 2	1 10 26	1 39	1 47 12	0 44	1 54 48	0 18	1 31 49	1 12	0 44 43	1 50	23
8	0 16 20	2 2	1 12 2	1 36	1 47 56	0 43	1 54 30	0 16	1 30 35	1 14	0 42 52	1 51	22
9	0 18 21	2 1	1 13 37	1 35	1 48 39	0 41	1 54 14	0 23	1 29 19	1 16	0 41 0	1 52	21
10	0 20 22	2 1	1 15 11	1 34	1 49 20	0 38	1 53 51	0 24	1 28 1	1 18	0 39 7	1 53	20
11	0 22 23	2 1	1 16 44	1 33	1 49 58	0 39	1 53 27	0 25	1 26 43	1 18	0 37 15	1 52	19
12	0 24 24	1 59	1 18 14	1 30	1 50 37	0 36	1 53 2	0 28	1 25 22	1 21	0 35 21	1 54	18
13	0 26 23	1 59	1 19 45	1 31	1 51 13	0 31	1 52 34	0 29	1 24 1	1 21	0 33 27	1 54	17
14	0 28 22	1 59	1 21 13	1 28	1 51 44	0 33	1 52 5	0 34	1 22 37	1 24	0 31 32	1 55	16
15	0 30 21	1 58	1 22 40	1 27	1 52 17	0 29	1 51 31	0 32	1 21 11	1 26	0 29 36	1 56	15
16	0 32 19	1 57	1 24 4	1 24	1 52 46	0 26	1 50 59	0 35	1 19 45	1 26	0 27 41	1 55	14
17	0 34 16	1 57	1 25 27	1 23	1 53 12	0 26	1 50 24	0 38	1 18 18	1 27	0 25 44	1 57	13
18	0 36 13	1 57	1 26 49	1 22	1 53 38	0 22	1 49 46	0 39	1 16 47	1 31	0 23 47	1 57	12
19	0 38 10	1 56	1 28 11	1 22	1 54 0	0 22	1 49 7	0 45	1 15 17	1 30	0 21 50	1 57	11
20	0 40 6	1 55	1 29 28	1 17	1 54 22	0 17	1 48 22	0 43	1 13 45	1 32	0 19 50	2 0	10
21	0 42 1	1 54	1 30 45	1 15	1 54 39	0 16	1 47 39	0 44	1 12 11	1 34	0 17 52	1 58	9
22	0 43 55	1 52	1 32 0	1 12	1 54 55	0 15	1 46 55	0 47	1 10 38	1 33	0 15 53	1 59	8
23	0 45 47	1 53	1 33 12	1 13	1 55 10	0 12	1 46 8	0 49	1 9 1	1 37	0 13 54	1 59	7
24	0 47 40	1 51	1 34 25	1 11	1 55 22	0 10	1 45 19	0 53	1 7 25	1 36	0 11 56	1 58	6
25	0 49 31	1 51	1 35 36	1 8	1 55 32	0 8	1 44 26	0 53	1 5 46	1 39	0 9 57	1 59	5
26	0 51 22	1 49	1 36 44	1 6	1 55 40	0 6	1 43 33	0 56	1 4 7	1 39	0 7 58	1 59	4
27	0 53 11	1 49	1 37 50	1 5	1 55 46	0 4	1 42 37	0 56	1 2 26	1 41	0 5 59	1 59	3
28	0 55 0	1 47	1 38 55	1 1	1 55 50	0 1	1 41 41	0 58	1 0 45	1 41	0 4 0	2 0	2
29	0 56 47	1 47	1 39 56	1 2	1 55 51	0 0	1 40 43	1 1	0 59 2	1 43	0 2 0	2 0	1
30	0 58 34		1 40 58		1 55 51		1 39 42		0 57 17	1 45	0 0 0	2 0	0
☉ true An.	Sig. 11	Dif.	Sig. 10	Dif.	Sig. 9	Dif.	Sig. 8	Dif.	Sig. 7	Dif.	Sig. 6	Dif.	☉ true An.

CONSTRUCTION. From the Equation of the SUN's Center.—As the Compt. to 60' of the Dif. of any Equation, and the next Equation following, as far as the greatest, but Dif. +60' beyond the Greatest, is to that Dif. so is the former Equation of the two, to the Proportional Quantity to be added to, or subtracted from, that former Equation (according as it increases or decreases) for the correct Equation (of the true to the M. An. corresp.) to the Number of Deg. of the true, equal to the Deg. of M. An. of the Sun's Center; but with a contrary Sign. This Proportion arises from hence. As any Incr. of true An. : to corresp. Incr. or Decr. of Equation of the true to M. An. :: any other Incr. of true An. (but of the first Eq. or Compl. to whole Degrees of true An.) to what must be added to or subtracted from the first Eq. for the absolute Eq. of the increased true An. to M. Anom. correspondent.

EXAMPLE I. To find the Equation of true to M. Anom. for 1° 12' true An. Sun?

M. An.	Sun's An.	Eq. Sun Center	Log. Logar.
1° 12' } 1° 10' 43' 41" }	+1° 16' 19" = 1° 12' }	Sun's M. An.	As -58 32 9.9893 co.
1 13 } 1 11 42 13 }	+1 17 47 = 1 13 }		To + 1 28 1.6118
As Dif. Incr. 58 32 (60' - D. Eq.): 1 28 D. Eq. :: 1° 16' 19" : 1° 55" }	+1 55		So -76 19 9.8956 co.
	+ 1 18 14 Eq. reqd.		To + 1 55 1.4967
Sun's tr. An. 1° 12'	Whence M. Anom.	1° 13' 18' 14" required.	

D

EXAMPLES

EXAMPLES of the CONSTRUCTION and USE of the foregoing TABLE:

EXAMPLE II. To find the Equation for $4^{\circ} 19'$ true Anomaly?

M. An.	☉ true An.	Eq. ☉ Center
$4^{\circ} 19'$	$4^{\circ} 17' 42'' 47'''$	$+1^{\circ} 17' 13'' = 4^{\circ} 19'$
$4^{\circ} 20'$	$4^{\circ} 18' 44' 19''$	$+1^{\circ} 15' 42'' = 4^{\circ} 20'$

As Dif. Incr. 1 1 32 (D. Eq. $+60'$): 1 32 Dif. Eq. decr. $1^{\circ} 17' 13'' : 1' 56''$
 $- 1' 56''$ decr.

N. B. The foregoing Table is used with the same Facility as that of the Sun's Center.

Sun's tr. An. $4^{\circ} 19'$

$4^{\circ} 20' 15' 17''$ M. Anom. reqd. to which answers (fr. Eq. ☉ Center) $- 1^{\circ} 15' 17''$ Eq.

N. B. The foregoing Construction finds the Equation of the true from the mean Anomaly correctly enough, from the Equation of the Sun's Center, when those Equations and their Differences are correct. But near the greatest Equation, when the Differences are very small, an Error of $\frac{1}{2}''$, or $\frac{1}{4}''$, among those Equations or Differences, will create an Error of $2''$, or more, in proportioning. Wherefore the following Construction takes Place of, or exceeds the former for Correction.

To determine UNIVERSALLY and CORRECTLY, the Mean from the True Anomaly given, in all Orbits whatsoever?

RULE.—As $\sqrt{\text{Perih. Dist.}} : \sqrt{\text{Aph. Dist.}} :: \tan. \frac{1}{2} \text{ true Anom.} : \tan. \frac{1}{2} \text{ correct Excentric Anom.}$ which doubled is the Ex. Anom. Now, by the Nature of a PLANET'S or COMET'S Motion in an elliptical Orbit, and the Principles of Trigonometry. As Sem. Transf. (= Rad.) : S. Excentr. An. :: Excentricity : Length Arch L, whose Degrees D, are the Diff. of the Degrees of the Mean above those of the Excentr. Anomaly. Therefore Rad. : $59^{\circ} 29' 57'' 9'''$:: L : D, Degrees, to be added to the Excentric for the mean Anom. required.

In the Earth's Orbit, the Eccentricity is 1685 of such Parts as the Sem. Transverse or mean Dist. $\odot \text{ a } \odot = 100000$, the Perih. Dist. = 83115 and Aph. Dist. 101685, and let it be required to determine the Mean from the true Anomalies, and consequently the Equations, of 86° , 87° , 88° , 89° , 90° , 91° , 92° , 93° , 94° , and 95° . Hence, from the above Proportions,

Tr. An.	Equations.	Dif.	Tr. An.	Equations.	Dif.
$2^{\circ} 26'$	$1^{\circ} 55' 40''$	$0' 6''$	$3^{\circ} 1^{\circ}$	$1^{\circ} 55' 49''$	$0' 6''$
$2^{\circ} 27'$	$1^{\circ} 55' 46''$	$0' 4''$	$3^{\circ} 2'$	$1^{\circ} 55' 43''$	$0' 6''$
$2^{\circ} 28'$	$1^{\circ} 55' 50''$	$0' 1''$	$3^{\circ} 3'$	$1^{\circ} 55' 37''$	$0' 9''$
$2^{\circ} 29'$	$1^{\circ} 55' 51''$	$0' 0''$	$3^{\circ} 4'$	$1^{\circ} 55' 28''$	$0' 11''$
$3^{\circ} 0'$	$1^{\circ} 55' 51''$	$0' 0''$	$3^{\circ} 5'$	$1^{\circ} 55' 17''$	$0' 11''$

EXAMPLE.

Tr. An.	Equations.	Dif.	Tr. An.	Equations.	Dif.
$88^{\circ} 57' 54'' 62'''$	$88^{\circ} 57' 54'' 62'''$	$0' 0''$	$89^{\circ} 55' 49'' 62'''$	$89^{\circ} 55' 49'' 62'''$	$0' 0''$

To Conf. Log. 0.0073185 add the Log. Tan. $\frac{1}{2}$ Degrees true Anom. For Log. Tan. $\frac{1}{2}$ Degrees of Excentric Anomaly. And To Conf. Log. 9.9847226 add Log. S. Excentric An. (rej. Rad.) For Log. of a Number of Degrees to be added to the Degrees of Excentric Anom. for the Degrees of Mean Anomaly required. Whence the Equations as on the Left.

By Ward's Hypoth. As Perih. Dist. : Aph. Dist. :: Tan. $\frac{1}{2}$ tr. An. : T. $\frac{1}{2}$ m. An. nearly.

By Bullialdus's Correction } As Sem. Conj. : Sem. Transf. :: Tan. of the M. Anom. tho' } M. An. : Tan. corrected M. Anom. but a near Correction. }

Hence, const. Log. 0.0146370 + L. Tan. $\frac{1}{2}$ tr. An. = L. Tan. $\frac{1}{2}$ m. Anom. to be corrected. And const. L. 0.0000617 + L. Tan. of that m. Anom. = Log. T. corrected An. for the Earth's Orbit.

To find the Time of the Sun's Entrance into 10° ♍ in the Year 1758?

By our Tables.		Apogee ☉.	
N.S. Year 1758		$3^{\circ} 8' 42'' 54'''$	
By Palla. 1756, Mar. 30		$+ 15'$	
At the req. time	M. Place Ap. ☉	$3^{\circ} 8' 43'' 9'''$	—
	True Place ☉	$0^{\circ} 10' 0'' 0'''$	from
	True Anom. ☉	$9^{\circ} 1' 16'' 51'''$	rem.
	Corresp. Eq ⁿ .	$* - 1^{\circ} 55' 50''$	—
	M. Place then ☉	$0^{\circ} 8' 4' 10'''$	from
M. Pl. ☉ N.S. 1758		$9^{\circ} 10' 4' 12'''$	—
	March 30	$2^{\circ} 27' 59' 58'''$	rem.
		$2^{\circ} 27' 43' 21'''$	—
		$16' 37''$	rem.
Mean Time reqd.	Hours 6	$14' 47''$	—
		$1' 50''$	rem.
	Minutes 44	$1' 48' 25'''$	—
		$1' 35''$	rem.
	Seconds 39	$1' 36''$	—

RULE. Observe the Day on which the Sun entered that Sign by some Ephemeris on the Leap-Year preceding the Time required. Then from the Sun's true Place, for the Time required, take Sun's Apogee, and the Sun's true An. will remain; from whence get the Equation of true to mean An. which add to or subtract from the Sun's true Pl. (according as the Sign directs) and you will have ☉ M. Pl. for the Time required.—From which Pl. deduct M. Pl. ☉ for the Year, and out of what remains deduct the Mot. for the Month, Day, and from that Remainder the Mot. for Hours, &c. till nothing remains, and you will have the M. Time of the Sun's Entrance into any Point of the Ecliptic required.

To find the Time of the Sun's Entrance into $22^{\circ} 22'$ in 1761?

By our Tables.		Apogee ☉.	
N.S. Year 1761		$3^{\circ} 8' 45' 58'''$	
By Pal. 1756, Aug. 14		$+ 31'$	
At the req. time	M. Pl. Ap. ☉	$3^{\circ} 8' 46' 29'''$	—
	True Place ☉	$4^{\circ} 22' 22' 0'''$	From
	True Anom. ☉	$1^{\circ} 13' 35' 31'''$	rem.
	Corresp. Eq ⁿ	$* + 1^{\circ} 20' 37''$	—
	M. Place ☉	$4^{\circ} 23' 42' 37'''$	From
M. Pl. ☉ N.S. 1761		$9^{\circ} 10' 20' 21'''$	—
	Aug. 14	$7^{\circ} 13' 22' 16'''$	rem.
		$7^{\circ} 12' 45' 22'''$	—
		$36' 54''$	rem.
Mean Time reqd.	Hours 14	$34' 30''$	—
		$2' 24''$	rem.
	Minutes 58	$2' 22' 55'''$	—
		$1' 5''$	rem.
	Seconds 26	$1' 14''$	—

To find the Sun's true Places to the foregoing Times?

M. Pl. ☉		Apogee ☉.	
N.S. Yr 1758		$3^{\circ} 8' 42'' 54'''$	
March 30		$2^{\circ} 27' 43' 21'''$	
Hours 6		$14' 47''$	
Minutes 44		$1' 48' 25'''$	
Seconds 39		$1' 36''$	
M. Pl. ☉	$0^{\circ} 8' 4' 10'''$	$8^{\circ} 29' 21' 1'''$	—
Equation	$+ 1^{\circ} 55' 50''$	M. An. ☉	—
True Pl. ☉	$0^{\circ} 10' 0'' 0'''$	required.	—
N.S. Yr 1761	$9^{\circ} 10' 20' 21'''$	$3^{\circ} 8' 45' 58'''$	—
Aug. 14	$7^{\circ} 12' 45' 22'''$	$+ 31'$	—
Hours 14	$34' 30''$	$3^{\circ} 8' 46' 29'''$	—
Minutes 58	$2' 24''$	$3^{\circ} 8' 48' 53'''$	—
Seconds 26	$1' 14''$	$3^{\circ} 8' 49' 49'''$	—
M. Pl. ☉	$4^{\circ} 23' 42' 37'''$	$1^{\circ} 14' 56' 8'''$	—
Equation	$- 1^{\circ} 20' 37''$	M. An. Sun	—
True Pl. ☉	$4^{\circ} 22' 22' 0'''$	required.	—

The tr. An. every true Year, decreasing by the Increase of the Sun's Apog. in that Time. It is evident, from above *, that the Sun returns to the same Point of the Ecliptic, each true Year, later than on the former Year by the Incr. or Decr. of Eqⁿ of the Sun's true to his M. Pl. or Anom. turned into Time, and added to or subtracted from $5^h 48^m 54^s 46^t$ the Surplus of our mean Solar Yr above 365^d , in which a M. Revolution is completed.

N.B. Increase } Eqⁿ } — } The { + $365^d 5^h 48^m 54^s 46^t$, &c. = Tr. Solar Yr.
 Decrease } — } + } Time.
 Increase } + } + }
 Decrease } — } — }

Hence, the M. An. ☉ decreases in the true Year or ☉'s Return to the same Point of the Ecliptic, by Sum of the Increase of the Sun's Apog. and Dif. of Eqⁿ (of the true to M. Anom.) in the same Time; by which Decr. of the Sun's M. An. in each true Year, the Equation to the absolute Equation of Time, following, is determined.

ABSOLUTE

Argument. SUN's true PLACE in the *Ecliptic*.

tr. Pl.	Equan. 1 ^o Sun Ap. Inc. dec.	Equan. 1 ^o Sun Ap. Inc. dec.	Equan. 1 ^o Sun Ap. Inc. dec.	Equan. 1 ^o Sun Ap. Inc. dec.	Equan. 1 ^o Sun Ap. Inc. dec.	Equan. 1 ^o Sun Ap. Inc. dec.	Equan. 1 ^o Sun Ap. Inc. dec.	Equan. 1 ^o Sun Ap. Inc. dec.	Equan. 1 ^o Sun Ap. Inc. dec.	Equan. 1 ^o Sun Ap. Inc. dec.	Equan. 1 ^o Sun Ap. Inc. dec.	tr. Pl.
	- +	+ -	+ -	+ -	+ -	+ -	+ -	- +	- +	- +	- +	
9	s.	s.	s.	s.	s.	s.	s.	s.	s.	s.	s.	9
6	1	3	6	8	7	5	8 ^o 1 8 ^o	3	6	8	8	6
12	10 ^o 0 10 ^o	4	7	3	7	5	- 0 +	4	7	8	8	12
18	+ 1 -	5	7	8	6	4	1	5	7	8	7	18
24	2	5	8	8	6	3	1	5	8	8	6	24
30	2	6	8	8	5	2	2	6	8	8	5	30

I. EQUATION

I. EQUATION of TIME; of App^t to Mean and Mean to App^t.

Argument. SUN's true place.

tr. Pl.	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
Pl.	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1	0	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20
2	0	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40
3	0	59	59	59	59	59	59	59	59	59	59	59	59	59	59	59	59	59	59	59	59	59	59	59	59	59	59	59	59	59	59
4	1	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19
5	1	39	39	39	39	39	39	39	39	39	39	39	39	39	39	39	39	39	39	39	39	39	39	39	39	39	39	39	39	39	39
6	1	58	58	58	58	58	58	58	58	58	58	58	58	58	58	58	58	58	58	58	58	58	58	58	58	58	58	58	58	58	58
7	2	18	18	18	18	18	18	18	18	18	18	18	18	18	18	18	18	18	18	18	18	18	18	18	18	18	18	18	18	18	18
8	2	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37
9	2	56	56	56	56	56	56	56	56	56	56	56	56	56	56	56	56	56	56	56	56	56	56	56	56	56	56	56	56	56	56
10	3	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15
11	3	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34	34
12	3	52	52	52	52	52	52	52	52	52	52	52	52	52	52	52	52	52	52	52	52	52	52	52	52	52	52	52	52	52	52
13	4	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
14	4	28	28	28	28	28	28	28	28	28	28	28	28	28	28	28	28	28	28	28	28	28	28	28	28	28	28	28	28	28	28
15	4	46	46	46	46	46	46	46	46	46	46	46	46	46	46	46	46	46	46	46	46	46	46	46	46	46	46	46	46	46	46
16	5	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3
17	5	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20
18	5	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37	37
19	5	53	53	53	53	53	53	53	53	53	53	53	53	53	53	53	53	53	53	53	53	53	53	53	53	53	53	53	53	53	53
20	5	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
21	6	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25
22	6	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40
23	6	55	55	55	55	55	55	55	55	55	55	55	55	55	55	55	55	55	55	55	55	55	55	55	55	55	55	55	55	55	55
24	7	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
25	7	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23
26	7	36	36	36	36	36	36	36	36	36	36	36	36	36	36	36	36	36	36	36	36	36	36	36	36	36	36	36	36	36	36
27	7	48	48	48	48	48	48	48	48	48	48	48	48	48	48	48	48	48	48	48	48	48	48	48	48	48	48	48	48	48	48
28	8	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
29	8	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12
30	8	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23
tr. Pl.	+	-	+	-	+	-	+	-	+	-	+	-	+	-	+	-	+	-	+	-	+	-	+	-	+	-	+	-	+	-	+
Pl.	A.	M.	A.	M.	A.	M.	A.	M.	A.	M.	A.	M.	A.	M.	A.	M.	A.	M.	A.	M.	A.	M.	A.	M.	A.	M.	A.	M.	A.	M.	A.
	11	5	10	4	10	4	10	4	10	4	10	4	10	4	10	4	10	4	10	4	10	4	10	4	10	4	10	4	10	4	10
	8	17	11	0	11	0	11	0	11	0	11	0	11	0	11	0	11	0	11	0	11	0	11	0	11	0	11	0	11	0	11

CONSTRUCTION. The above *Table* is the *Diff.* between the Sun's true Long. and the R. A. converted into Time, which adds to Appt in the 2d and 4th *Quadratures*, and subtracts therefrom in the 1st and 3d *Quadratures*, for the *M. Time* and the contrary.

N. B. The Complement of any Diff. to 60^o or Deg. R. A. (in Tab. R. A.) converted into Time, and added to the Equat. of T. for any preceding Deg. ☉ tr. Place, as far as 8 17^o, or *great Equat. Time*, will b. the Equat. T. in the above Tab. corresp. to the succeeding Deg. of ☉ true Place— But the Diff. of R. A. above 60^o beyond 8 17^o, or the greatest Equat. converted into Time, and *subducted* fr. the Equat. of T. for any preceding Deg. ☉ true Pl. will be the Equat. T. corresponding to the succeeding Deg. thereof.

II. EQUATION of TIME; of Apparent to Mean and Mean to Apparent.

Argument. SUN's Mean Anomaly.

M. An.	Sig.	A. M.		Diff.	Sig.	A. M.		Diff.	Sig.	A. M.		Diff.	Sig.	A. M.		Diff.	Sig.	A. M.		Diff.	M. An.
		-	+			-	+			-	+			-	+			-	+		
0	0	0	0	8	3	47	6	37	7	43	6	45	3	56	30						
1	0	8	8	8	3	54	6	41	4	7	43	6	41	49	29						
2	0	16	8	8	4	1	6	45	4	7	43	6	37	42	28						
3	0	24	8	8	4	8	6	49	4	7	43	6	33	34	27						
4	0	32	8	8	4	15	6	53	4	7	43	6	29	27	26						
5	0	40	8	8	4	21	6	56	4	7	42	6	24	20	25						
6	0	47	7	7	4	28	7	0	4	7	42	6	20	12	24						
7	0	55	8	8	4	34	7	3	3	7	41	6	15	5	23						
8	1	3	8	8	4	41	7	6	3	7	40	6	10	57	22						
9	1	11	7	7	4	47	6	9	3	7	39	6	5	49	21						
10	1	19	8	8	4	53	6	12	3	7	38	6	0	42	20						
11	1	27	8	8	4	59	6	15	3	7	37	5	55	34	19						
12	1	34	7	7	5	5	6	18	3	7	35	5	49	26	18						
13	1	42	8	8	5	11	6	20	2	7	34	5	44	18	17						
14	1	50	8	8	5	17	6	23	3	7	32	5	38	10	16						
15	1	58	8	8	5	23	6	25	2	7	30	5	33	2	15						
16	2	5	7	7	5	28	5	27	2	7	28	5	27	54	14						
17	2	13	8	8	5	34	6	29	2	7	26	5	21	46	13						
18	2	20	7	7	5	39	5	31	2	7	23	5	15	38	12						
19	2	28	8	8	5	45	6	33	2	7	21	5	9	30	11						
20	2	35	7	7	5	50	5	35	2	7	19	5	3	22	10						
21	2	43	8	8	5	55	5	36	1	7	16	4	56	14	9						
22	2	50	7	7	6	0	5	37	1	7	13	4	50	6	8						
23	2	58	8	8	6	5	5	39	2	7	10	4	44	58	7						
24	3	5	7	7	6	10	5	40	1	7	7	4	37	49	6						
25	3	12	7	7	6	15	5	41	1	7	4	4	30	41	5						
26	3	19	7	7	6	20	5	41	0	7	0	4	24	33	4						
27	3	26	7	7	6	24	4	42	1	6	57	4	17	25	3						
28	3	34	8	8	6	29	5	43	1	6	53	4	10	17	2						
29	3	41	7	7	6	33	7	43	0	6	49	4	3	8	1						
30	3	47	6	6	6	37	7	43	0	6	45	3	56	0	0						
+	-				+	-		+	-		+	-		+	-						
M. An.	Sig.			Diff.	M. An.	Sig.		Diff.	M. An.	Sig.		Diff.	M. An.	Sig.		Diff.					
11					10				9				7								

CONSTRUCTION. The above *Table* is only the Equation of the Sun's Center, correspondent to the given *Mean Anomaly*, converted into Time, which subtracts from the Apparent in the *1st Six Signs*, and adds thereto in the *last Six Signs*, for the Mean Time, and the contrary.—Hence the Sum of the 1st and 11d Equations for the absolute Equation of Time, will be always equal to the Diff. of the $\odot$'s R. A. and his Mean Longitude converted into Time.

N. B. *Mean Time* is considered in *Apparent Time* correspondent to it.

EXAMPLE. To find the Mean Time, or Hour correspondent to Apparent Noon, the $\odot$'s true Place being $45^{\circ} 23'. 25''$. and $\odot$ Mean Anom. $5^{\circ} 15'. 56''$?

By Table I. $\angle 23^{\circ} 25'$. . A to M . . $+ 7^m 26^s$

By Table II. $5^{\circ} 15' 56'' \dots A \text{ to } M \dots + 1$ 54

Sum	+	9	20
Noon	12	0	0

Hours 12 9 20 required.

Mean Time correspondent to Apparent Time in the Afternoon.

REFRACTION

REFRACTION OF LIGHT (Lat. Greenwich Observatory), and **PARALLAX OF THE SUN.** The first to be subtracted from the apparent Altitude of the luminous Body, and the latter to be added to the apparent for the real Altitude of the Sun.

Argument. Apparent Altitude of the SUN.								
Ap. Alt.	Re- frac ⁿ .	Par. ○.	Ap. Alt.	Re- frac ⁿ .	Par. ○.	Ap. Alt.	Ref. ○.	Par. ○.
○.	—	+	○.	—	+	○.	—	+
°	I II	#	°	I II	#	°	#	#
0	33 45	10	21	2 28	9	56	36	6
$\frac{1}{4}$	30 24	10	22	2 11	9	57	35	6
$\frac{1}{2}$	27 35	10	23	2 5	9	58	34	5
$\frac{3}{4}$	25 11	10	24	1 59	9	59	32	5
1	23 7	10	25	1 54	9	60	31	5
$1\frac{1}{4}$	21 20	10	26	1 49	9	61	30	5
$1\frac{1}{2}$	19 46	10	27	1 44	9	62	28	5
$1\frac{3}{4}$	18 22	10	28	1 40	9	63	27	5
2	17 8	10	29	1 36	9	64	26	4
$2\frac{1}{2}$	15 2	10	30	1 32	9	65	25	4
3	13 20	10	31	1 28	9	66	24	4
$3\frac{1}{2}$	11 57	10	32	1 25	9	67	23	4
4	10 48	10	33	1 22	8	68	22	4
$4\frac{1}{2}$	9 50	10	34	1 19	8	69	21	3
5	9 2	10	35	1 16	8	70	20	3
$5\frac{1}{2}$	8 21	10	36	1 13	8	71	19	3
6	7 45	10	37	1 11	8	72	18	3
$6\frac{1}{2}$	7 14	10	38	1 8	8	73	17	2
7	6 47	10	39	1 6	8	74	16	2
$7\frac{1}{2}$	6 22	10	40	1 4	8	75	15	2
8	6 0	10	41	1 2	8	76	14	2
$8\frac{1}{2}$	5 40	10	42	1 0	8	77	13	2
9	5 22	10	43	58	7	78	12	2
$9\frac{1}{2}$	5 6	10	44	56	7	79	11	2
10	4 52	10	45	54	7	80	10	2
11	4 27	10	46	52	7	81	9	1
12	4 5	10	47	50	7	82	8	1
13	3 47	10	48	47	7	83	7	1
14	3 31	10	49	47	7	84	6	1
15	3 17	10	50	45	7	85	5	1
16	3 4	10	51	44	6	86	4	1
17	2 53	10	52	42	6	87	3	1
18	2 43	10	53	40	6	88	2	0
19	2 34	10	54	39	6	89	1	0
20	2 26	9	55	38	6	90	0	0

Hourly Motion, and Apparent Semi-Diameter of the SUN.

Arg. Sun's Mean Anomaly.							
$\odot$'s Mean Anom.	Hourly Mot. $\odot$.	Semi-Diam. $\odot$.	$\odot$'s Mean Anom.				
S	O	I	II	I	II	O	S
0	0	2	23	15	51	0	12
6	2	23	15	51	24		
12	2	23	15	52	18		
18	2	23	15	52	12		
24	2	23	15	53	6		
1	0	2	24	15	53	0	11
6	2	24	15	54	24		
12	2	24	15	55	18		
18	2	24	15	57	12		
24	2	25	15	58	6		
2	0	2	25	15	59	0	10
6	2	26	16	1	24		
12	2	26	16	2	18		
18	2	27	16	4	12		
24	2	27	16	6	6		
3	0	2	28	16	7	0	9
6	2	28	16	9	24		
12	2	29	16	11	18		
18	2	29	16	13	12		
24	2	30	16	14	6		
4	0	2	30	16	16	0	8
6	2	31	16	17	24		
12	2	31	16	18	18		
18	2	32	16	20	12		
24	2	32	16	21	6		
5	0	2	32	16	22	0	7
6	2	32	16	23	24		
12	2	33	16	23	18		
18	2	33	16	24	12		
24	2	33	16	24	6		
6	0	2	33	16	24	0	0

Acceleration of the fixed *s according to mean solar Parts of Time.

Arg. Days.			
D ^s .	Accelerat ⁿ .		
No.	h	m	s
1	0	3	56
2	0	7	52
3	0	11	48
4	0	15	44
5	0	19	39
6	0	23	35
7	0	27	31
8	0	31	27
9	0	35	23
10	0	37	19
11	0	43	15
12	0	47	11
13	0	51	7
14	0	55	3
15	0	58	58
16	1	2	54
17	1	6	50
18	1	10	46
19	1	14	42
20	1	18	38
21	1	22	34
22	1	26	30
23	1	30	26
24	1	34	22
25	1	38	17
26	1	42	13
27	1	46	9
28	1	50	5
29	1	54	1
30	1	57	57
31	2	1	53

CONSTRUCTION. To find the Sun's hourly Motion correspondent to 4^h 6^m, Sun's Mean Anomaly?

By Equation Tab. Sun's Center, p. 12
4^h 6^m M. An. + 1' 10" dif. Eq. 24^h.
+ 59 8 M. Mo. 24^h H.

Sun's tr. Mo. in 24^h 60 18 fere.

Whence 2 31 Hourly Mo. (as by Table) required.

To find $\odot$'s Apt. Sem. Diam. corresp. to 6^h M. An. the Apt. S. Diam. to 0^h M. An. being given 15' 51"?

By Tab. L. Sun's prop^d Dist. fr. $\odot$ p. 16 Reciprocally.

0^h M. An. Pr. D. $\odot$ 101685 .. L. 4,9927
Sun's ap. S. Diam. 15' 51" .. L. L. 0,5781
6^h M. An. Pr. D. $\odot$, 98316 .. L. 4,9926

Sun's ap. S. Diam. 16' 24" .. L. L. 0,5634
(as by Tab.) req^d. The ap. S. Diam. or Diam. being recip. as Sun's Dist. à $\odot$.

The fixed Stars come, in mean Time, as above, sooner to the Meridian, each Day, than on the preceding Day.

The true Time of the fixed Stars Acceleration is the Increase of the Sun's R. A. each Day converted into Time. See Tab. P. 24, 25, 26, 27.

Req. Sun's true Alt. to 15^o ap. Alt.?

15^o 0' 0"

Eq. Ref. — 3 17

Eq. Paral. + 10

$\odot$ tr. Alt. 14 56 53

DECIMALS

The Refraction of Light varies with the Latitude, Atmosphere, or Weather; especially near the Horizon. — In the Year 1695, in Latitude 65° 53', on June 14, 12 Hours P. M. the Sun's Center being depressed below the Horizon 40', at Pello, 10 Miles northward of Fornio in Western Bothnia, He was observed, by Refraction, to be two Diameters in Altitude, or above 1°. — The common Experiment of pouring Water upon a Piece of Silver, in a Basin, to bring it in Sight (when unseen before) proves the Elevation of Objects by Light passing from them through a denser to a rarer Medium, (the contrary happening from a rarer to a denser) as on the Variation of the refracting Medium the different Degrees of Refraction depend.

Rays passing through Oil of Turpentine and through Water, give the Degrees of Refraction different, proving the different Densities of those Mediums.

PARALLAX of the SUN is the Angle the Sun is seen lower from the Earth's Surface than it would be seen from its Center.

DECIMALS of a DEGREE, for proportioning SEXAGESIMAL QUANTITIES.

MINUTES.				SECONDS.				THIRDS.			
Min.	Decim.	Min.	Decim.	Sec.	Decim.	Sec.	Decim.	Th ^s .	Decim.	Th ^s .	Decim.
1	,0166	31	,5166	1	,00027	31	,00861	1	,000004629	31	,000143818
2	,0333	32	,5333	2	,00054	32	,00888	2	,000009259	32	,000148148
3	,05	33	,55	3	,00081	33	,00915	3	,000013888	33	,000152477
4	,0666	34	,5666	4	,00108	34	,00942	4	,000018518	34	,000156807
5	,0833	35	,5833	5	,00135	35	,00969	5	,000023148	35	,000161137
6	,1	36	,6	6	,00162	36	,01	6	,000027777	36	,000165466
7	,1166	37	,6166	7	,00189	37	,01027	7	,000032407	37	,000169796
8	,1333	38	,6333	8	,00216	38	,01054	8	,000037037	38	,000174125
9	,15	39	,65	9	,00243	39	,01081	9	,000041666	39	,000178455
10	,1666	40	,6666	10	,00270	40	,01108	10	,000046296	40	,000182785
11	,1833	41	,6833	11	,00297	41	,01135	11	,000050925	41	,000187114
12	,2	42	,7	12	,00324	42	,01162	12	,000055555	42	,000191444
13	,2166	43	,7166	13	,00351	43	,01189	13	,000060185	43	,000195774
14	,2333	44	,7333	14	,00378	44	,01216	14	,000064814	44	,000200103
15	,25	45	,75	15	,00405	45	,01243	15	,000069444	45	,000204433
16	,2666	46	,7666	16	,00432	46	,01270	16	,000074074	46	,000208762
17	,2833	47	,7833	17	,00459	47	,01297	17	,000078703	47	,000213092
18	,3	48	,8	18	,00486	48	,01324	18	,000083333	48	,000217422
19	,3166	49	,8166	19	,00513	49	,01351	19	,000087962	49	,000221751
20	,3333	50	,8333	20	,00540	50	,01378	20	,000092592	50	,000226081
21	,35	51	,85	21	,00567	51	,01405	21	,000097222	51	,000230411
22	,3666	52	,8666	22	,00594	52	,01432	22	,000101851	52	,000234740
23	,3833	53	,8833	23	,00621	53	,01459	23	,000106481	53	,000239070
24	,4	54	,9	24	,00648	54	,01486	24	,000111111	54	,000243400
25	,4166	55	,9166	25	,00675	55	,01513	25	,000115740	55	,000247729
26	,4333	56	,9333	26	,00702	56	,01540	26	,000120370	56	,000252059
27	,45	57	,95	27	,00729	57	,01567	27	,000125000	57	,000256388
28	,4666	58	,9666	28	,00756	58	,01594	28	,000129629	58	,000260718
29	,4833	59	,9833	29	,00783	59	,01621	29	,000134259	59	,000265048
30	,5	60	1	30	,00810	60	,01648	30	,000138888	60	,000269377

N. B. In the above Table, the Figure with a fine Stroke crosses it, denotes a continual Repetition of the Figure or Figures from that Place. And a Dot (.) or Full Point, after a Decimal, denotes its exact Value.

EXAMPLE. As $3^{\circ} 19' 23''$ is to $40^{\circ} 29' 33'' 40'''$, so is $11^{\circ} 49' 34'' 18'''$, to $144^{\circ} 6' 23'' 56'''$, &c.

By the Table

$19' = ,316666$. : As $3^{\circ} 323058$ to $40^{\circ} 4926851$ so $11^{\circ} 826194$
 $23'' = ,006388$

$,323058$

$29' = ,4833333$

$33'' = ,0091666$

$40''' = ,0001851$

$,4926851$

$49' = ,8166666$

$34'' = ,0094444$

$18''' = ,0000833$

$,8261944$

By . . .

As $3^{\circ} 323058$

To $40^{\circ} 4926851$

So $11^{\circ} 826194$

To $144^{\circ} 1066$

Logarithms

9.4784624 co. [to $144^{\circ} 10665$, &c. Decimally]

1.6073765

1.0728450

2.1586839

6',399

23'',94

56''',4

The greater Ease and Expedition of finding the Decimals to Sexagesimals, by dividing them and their Quotients, from inferior Order, by 6, than finding the same by the Table, appear from the Examples above, and on the Side.

SUN's DECLINATION from the EQUINOCTIAL, and MERIDIAN ANGLE with the ECLIPTIC. According to the Mean Ecliptic Obliquity $23^{\circ} 28' \frac{1}{2}$.

Argument. SUN's true PLACE in the ECLIPTIC.

Aries ♈ } 0 Sig.				Taurus ♉ } 1 Sig.				Gemini ♊ } 2 Sig.				Sagittarius ♐ } 8 Sig.																																			
Libra ♎ } 6				Scorpio ♏ } 7				Sagittarius ♐ } 8				Sagittarius ♐ } 8																																			
Declin.				Diff.				Declin.				Diff.				Declin.				Diff.				Declin.				Diff.																			
° ' "				° ' "				° ' "				° ' "				° ' "				° ' "				° ' "				° ' "																			
0	0	0	0	23	54	66	31	30	0	11	11	29	19	21	2	69	23	18	11	36	20	10	50	12	33	77	44	55	21	41	30																
1	0	23	54	23	54	66	31	41	0	35	11	50	21	20	50	69	34	54	11	58	20	23	23	12	10	78	6	30	21	58	20																
2	0	47	48	23	52	66	32	16	0	57	12	11	11	20	38	69	46	52	12	20	20	35	33	11	47	78	28	34	22	13	28																
3	1	11	40	23	52	66	33	13	1	21	12	31	49	20	27	69	59	12	12	44	20	47	21	11	25	78	50	47	22	28	27																
4	1	35	32	23	51	66	34	34	1	43	12	52	16	20	12	70	11	56	13	5	20	58	46	11	2	78	13	15	22	44	26																
5	1	59	23	23	48	66	36	17	2	6	13	12	28	20	0	70	25	1	13	27	21	9	48	10	38	79	35	59	22	58	25																
6	2	23	11	23	46	66	38	23	2	30	13	32	28	19	47	70	38	28	13	49	21	20	26	10	14	79	58	57	23	11	24																
7	2	46	57	23	44	66	40	53	2	52	13	52	15	19	32	70	52	17	14	11	21	30	40	9	50	80	22	8	23	26	23																
8	3	10	41	23	41	66	43	45	3	15	14	11	47	19	19	71	6	28	14	32	21	40	30	9	26	80	45	34	23	39	22																
9	3	34	22	23	37	66	47	0	3	38	14	31	6	19	4	71	21	0	14	54	21	49	56	9	2	81	9	13	23	51	21																
10	3	57	59	23	34	66	50	38	4	1	14	50	10	18	49	71	35	54	15	16	21	58	58	8	36	81	33	4	24	4	20																
11	4	21	33	23	30	66	54	39	4	24	15	8	59	18	33	71	51	10	15	37	22	7	34	8	11	81	57	8	24	15	19																
12	4	45	3	23	25	67	59	3	4	47	15	27	32	18	18	72	6	47	15	58	22	15	45	7	46	82	21	23	24	26	18																
13	5	8	28	23	20	67	3	50	5	9	15	45	50	18	2	72	25	45	16	19	22	23	31	7	21	82	45	49	24	37	17																
14	5	31	48	23	16	67	8	59	5	33	16	3	52	17	45	72	39	4	16	39	22	30	52	6	55	83	10	26	24	46	16																
15	5	55	4	23	10	67	14	32	5	55	16	21	37	17	28	72	55	43	17	1	22	37	47	6	29	83	35	12	24	56	15																
16	6	18	14	23	3	67	20	27	6	19	16	39	5	17	11	73	12	44	17	21	22	44	16	6	3	84	0	8	25	5	14																
17	6	41	17	23	58	67	26	46	6	41	16	56	16	16	54	73	30	5	17	41	22	50	19	5	37	84	25	13	25	12	13																
18	7	4	15	22	51	67	33	27	7	4	17	13	10	16	35	73	47	46	18	2	22	55	56	5	11	84	50	25	25	21	12																
19	7	27	6	22	54	67	40	31	7	29	17	29	45	16	18	74	5	48	18	22	23	1	7	4	44	84	15	46	25	28	11																
20	7	49	50	22	36	67	47	58	7	49	17	46	3	15	58	74	24	10	18	40	23	5	51	4	17	85	41	14	25	34	10																
21	8	12	26	22	29	68	55	47	8	14	18	2	1	15	39	74	42	50	19	0	23	10	8	3	51	86	6	48	25	34	9																
22	8	34	55	22	21	68	4	1	8	35	18	17	40	15	21	75	1	50	19	19	23	13	59	3	24	86	32	28	25	40	8																
23	8	57	16	22	12	68	12	36	8	58	18	33	1	15	0	75	21	9	19	39	23	17	23	2	56	86	58	13	25	45	7																
24	9	19	28	22	3	68	21	34	9	21	18	48	1	14	40	75	40	48	19	57	23	20	19	2	30	87	24	3	25	50	6																
25	9	41	31	21	54	68	30	55	9	43	19	2	41	14	20	76	0	45	20	14	23	22	49	2	30	87	49	57	25	54	5																
26	10	3	25	21	44	68	40	38	10	6	19	17	1	14	0	76	20	59	20	33	23	24	52	1	3	88	15	54	25	57	4																
27	10	25	9	21	35	69	50	44	10	29	19	31	1	13	38	76	41	32	20	51	23	26	27	1	35	88	41	53	26	59	3																
28	10	46	44	21	23	69	1	13	10	52	19	44	39	13	16	77	2	23	21	7	23	27	35	0	41	89	7	54	26	1	2																
29	11	8	7	21	12	69	12	5	11	13	19	57	55	12	55	77	23	30	21	25	23	28	16	0	14	89	33	5	26	3	1																
30	11	29	19	20	10	69	23	18			20	10	50			77	44	55			23	28	30			90	0	0			0																
Decl.				Diff.				Merid. L.				Diff.				Decl.				Diff.				Merid. L.				Diff.				Decl.				Diff.				Merid. L.				Diff.			
Virgo ♍ } 5 Sig.				Leo ♌ } 4 Sig.				Cancer ♋ } 3 Sig.				Capricorn ♐ } 9 Sig.																																			
Pisces ♓ } 11 Sig.				Aquarius ♒ } 10 Sig.																																											

SUN'S RIGHT ASCENSION, from Aries, according to the Mean Ecliptic OBLIQUITY $23^{\circ} 28' \frac{1}{2}$.

Argument. SUN'S PLACE in the ECLIPTIC.

Pl.	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1	0	55	2	55	3	27	54	15	57	23	57	48	42	62	34	90	0	0	65	25	122	11	18	62	23	152	5	45	57	14	30
2	1	50	5	55	3	28	51	38	57	31	58	51	16	62	43	91	5	25	65	24	123	13	41	62	12	153	2	59	57	6	29
3	2	45	8	55	3	29	49	9	57	41	59	53	59	62	53	92	10	49	65	24	124	15	53	62	3	154	0	5	56	58	28
4	3	40	12	55	4	30	46	50	57	50	60	56	52	63	2	93	16	13	65	22	125	17	56	61	51	154	57	3	56	50	27
5	4	35	17	55	5	31	44	40	57	59	61	59	54	63	13	94	21	35	65	20	126	19	47	61	41	155	53	53	56	42	26
6	5	30	24	55	7	32	42	39	58	9	63	3	7	63	22	95	26	55	65	18	127	21	28	61	30	156	50	35	56	34	25
7	6	25	32	55	8	33	40	48	58	18	64	6	29	63	30	96	32	13	65	15	128	22	58	61	19	157	47	9	56	28	24
8	7	20	43	55	11	34	39	6	58	29	65	9	59	63	39	97	37	28	65	12	129	24	17	60	9	158	43	37	56	21	23
9	8	15	57	55	14	35	37	35	58	38	66	13	38	63	48	98	42	40	65	9	130	25	26	60	57	159	39	58	56	14	22
10	9	11	13	55	16	36	36	13	58	49	67	17	26	63	56	99	47	49	65	5	131	26	23	60	47	160	36	12	56	7	21
11	10	6	33	55	20	37	35	2	58	58	68	21	22	63	56	100	52	54	65	0	132	27	10	60	36	161	32	19	56	1	20
12	11	1	56	55	23	38	34	0	59	10	69	25	26	64	4	101	57	54	64	56	133	27	46	60	24	162	28	21	55	55	19
13	11	57	23	55	27	39	33	10	59	20	70	29	38	64	12	103	2	50	64	51	134	28	10	60	14	163	24	16	55	50	18
14	12	52	54	55	31	40	32	30	59	30	71	33	57	64	26	104	7	41	64	44	135	28	24	60	3	164	20	6	55	46	17
15	13	48	29	55	35	41	32	0	59	41	72	38	23	64	33	105	12	25	64	39	136	28	27	59	52	165	15	52	55	39	16
16	14	44	8	55	39	42	31	41	59	52	73	42	56	64	39	106	17	4	64	33	137	28	19	59	41	166	11	31	55	35	15
17	15	39	54	55	46	43	31	33	60	3	74	47	35	64	44	107	21	37	64	26	138	28	0	59	30	167	7	6	55	31	14
18	16	35	44	55	50	44	31	36	60	14	75	52	19	64	51	108	26	3	64	19	139	27	30	59	20	168	2	37	55	27	13
19	17	31	39	55	55	45	31	50	60	24	76	57	10	64	56	109	30	22	64	12	140	26	50	59	10	168	58	4	55	23	12
20	18	27	41	56	2	46	32	14	60	36	78	2	6	65	0	110	34	34	64	4	141	26	0	58	59	169	53	27	55	20	11
21	19	23	48	56	7	47	32	50	60	47	79	7	6	65	5	111	38	38	63	56	142	24	58	58	49	170	48	47	55	16	10
22	20	20	2	56	14	48	33	37	60	57	80	12	11	65	9	112	42	34	63	48	143	23	47	58	38	171	44	3	55	14	9
23	21	16	23	56	21	49	34	34	61	9	81	17	20	65	12	113	46	22	63	39	144	22	25	58	29	172	39	17	55	11	8
24	22	12	51	56	28	50	35	43	61	19	82	22	32	65	15	114	50	1	63	30	145	20	54	58	18	173	34	28	55	8	7
25	23	9	25	56	34	51	37	2	61	30	83	27	47	65	18	115	53	31	63	22	146	19	12	58	9	174	29	36	55	7	6
26	24	6	7	56	42	52	38	32	61	41	84	33	5	65	20	116	56	53	63	13	147	17	21	57	59	175	24	43	55	5	5
27	25	2	57	56	50	53	40	13	61	51	85	38	25	65	22	118	0	6	63	2	148	15	20	57	50	176	19	48	55	4	4
28	25	59	55	56	58	54	42	4	62	3	86	43	47	65	24	119	3	8	62	53	149	13	10	57	41	177	14	52	55	4	3
29	26	57	1	57	6	55	44	7	62	12	87	49	11	65	24	120	6	1	62	43	150	10	51	57	31	178	9	55	55	4	2
30	27	54	15	57	14	56	46	19	62	23	88	54	35	65	25	121	8	44	62	34	151	8	22	57	23	179	4	58	55	3	1
Pl.	Aries	Diff.	Taurus	Diff.	Gemini	Diff.	Cancer	Diff.	Leo	Diff.	Virgo	Diff.	Pl.	Diff.	Pl.	Diff.	Pl.	Diff.	Pl.	Diff.	Pl.	Diff.	Pl.	Diff.	Pl.	Diff.	Pl.	Diff.	Pl.	Diff.	Pl.

CONSTRUCTION. As Rad. : Tang. Long. $\odot$ from next Equinoctial Point :: Cos. greatest Declination : to Tang. R. A.EXAMPLE I. To find the Sun's Right Ascension correspondent to 80° . $\odot$ Longitude from Υ ?

Com. Log.

: Rad. S.	10.0000000	Hence to the Const. Log. :	9.9624801
: : 80° T.	10.7536812	Add Tang. Long. $\odot$ from Υ .	
: : $23^{\circ} 28' \frac{1}{2}$. C. S.	9.9624801	Sum (deducting R.) = T.R.A.	
: : $79^{\circ} 7' 6''$ T.	10.7161613	[from Υ .]	

EXAMPLE II. To find the $\odot$ R. A. correspondent to $\Omega 28^{\circ}$? $\Omega 28^{\circ} = 148^{\circ}$ Suppl. = 32° .

To Const. Log. 9.9624801

Add 32° . T. . 9.7957892 $-29^{\circ} 49' 9''$ T. 9.7582693

180

Difference 150 10 51 Right Ascension required.

N. B. Having found the Right Asc. to 90° , all the rest easily follow from thence to 180° : being the Supplements of the former (taken backwards) to 180° ; so that the R. A. of any Number of Signs and Degrees of the Sun's Place being given under 90° , the Supplement thereof to 180° will be the Right Ascension to the Supplement of those Signs and Degrees to 6 Signs.

EXAMPLE. $2^{\circ} 12^{\circ}$ R. A. $70^{\circ} 29' 38''$ Supplement $3^{\circ} 18$ R. A. $109^{\circ} 30' 22$ Supplement

6 0

180 0 0 Proof.

SUN'S RIGHT ASCENSION, from Aries, according to the Mean Ecliptic OBLIQUITY, $23^{\circ} 28' \frac{1}{2}$.

Argument. SUN'S PLACE in the ECLIPTIC.

Pl.	$\triangle 6^s$	Diff.	$m 7^s$	Diff.	$\dagger 8^s$	Diff.	$\text{v} 9^s$	Diff.	$\text{w} 10^s$	Diff.	$\text{x} 11^s$	Diff.	Pl.
0	180 0 0		207 54 15		237 48 42		270 0 0		302 11 18		332 5 45		30
1	180 55 2	55 2	208 51 38	57 23	238 51 16	62 34	271 5 25	65 25	303 13 41	62 23	333 2 59	57 14	29
2	181 50 5	55 3	209 49 9	57 31	239 53 59	62 43	272 10 49	65 24	304 15 53	62 12	334 0 5	56 58	28
3	182 45 8	55 4	210 46 50	57 41	240 56 52	62 53	273 16 13	65 24	305 17 56	62 3	334 57 3	56 50	27
4	183 40 12	55 4	211 44 40	57 50	241 59 54	63 2	274 21 35	65 22	306 19 47	61 51	335 53 53	56 42	26
5	184 35 17	55 5	212 42 39	57 59	243 3 7	63 13	275 26 55	65 20	307 21 28	61 41	336 50 35	56 34	25
6	185 30 24	55 7	213 40 48	58 9	244 6 29	63 22	276 32 13	65 18	308 22 58	61 30	337 47 9	56 28	24
7	186 25 32	55 8	214 39 6	58 18	245 9 59	63 30	277 37 28	65 15	309 24 17	61 19	338 43 37	56 21	23
8	187 20 43	55 11	215 37 35	58 29	246 13 38	63 39	278 42 40	65 12	310 25 26	61 9	339 39 58	56 14	22
9	188 15 57	55 14	216 36 13	58 38	247 17 26	63 48	279 47 49	65 9	311 26 23	60 57	340 36 12	56 7	21
10	189 11 13	55 16	217 35 2	58 49	248 21 22	63 56	280 52 54	65 5	312 27 10	60 47	341 32 19	56 1	20
11	190 6 33	55 20	218 34 0	58 59	249 25 26	64 4	281 57 54	65 0	313 27 46	60 36	342 28 21	55 55	19
12	191 1 56	55 23	219 33 10	59 10	250 29 38	64 12	283 2 50	64 56	314 28 10	60 24	343 24 16	55 50	18
13	191 57 23	55 27	220 32 30	59 20	251 33 57	64 19	284 7 41	64 51	315 28 24	60 14	344 20 6	55 46	17
14	192 52 54	55 31	221 32 0	59 30	252 38 23	64 26	285 12 25	64 44	316 28 27	60 3	345 15 52	55 39	16
15	193 48 29	55 35	222 31 41	59 41	253 42 56	64 33	286 17 4	64 39	317 28 19	59 52	346 11 31	55 35	15
16	194 44 8	55 39	223 31 33	59 52	254 47 35	64 39	287 21 37	64 33	318 28 0	59 41	347 7 6	55 31	14
17	195 39 54	55 46	224 31 36	60 3	255 52 19	64 44	288 26 3	64 26	319 27 30	59 30	348 2 37	55 27	13
18	196 35 44	55 50	225 31 50	60 14	256 57 10	64 51	289 30 22	64 19	320 26 50	59 20	348 58 4	55 23	12
19	197 31 39	55 55	226 32 14	60 24	258 2 6	64 56	290 34 34	64 12	321 26 0	59 10	349 53 27	55 20	11
20	198 27 41	56 1	227 32 50	60 36	259 7 6	65 0	291 38 38	64 4	322 24 58	58 59	350 48 47	55 16	10
21	199 23 48	56 7	228 33 37	60 47	260 12 11	65 5	292 42 34	63 56	323 23 47	58 49	351 44 3	55 14	9
22	200 20 2	56 14	229 34 34	60 57	261 17 20	65 9	293 46 22	63 48	324 22 25	58 38	352 39 17	55 11	8
23	201 16 23	56 21	230 35 43	61 9	262 22 32	65 12	294 50 1	63 39	325 20 44	58 29	353 34 28	55 8	7
24	202 12 51	56 28	231 37 2	61 19	263 27 47	65 15	295 53 31	63 30	326 19 12	58 18	354 29 36	55 7	6
25	203 9 25	56 34	232 38 32	61 30	264 33 5	65 18	296 56 53	63 22	327 17 21	58 9	355 24 43	55 5	5
26	204 6 7	56 42	233 40 13	61 41	265 38 25	65 20	298 0 6	63 13	328 15 20	57 59	356 19 48	55 4	4
27	205 2 57	56 50	234 42 4	61 51	266 43 47	65 22	299 3 8	63 2	329 13 10	57 50	357 14 52	55 3	3
28	205 59 55	56 58	235 44 7	62 3	267 49 11	65 24	300 6 1	62 53	330 10 51	57 41	358 9 55	55 2	2
29	206 57 1	57 6	236 46 19	62 12	268 54 35	65 24	301 8 44	62 43	331 8 22	57 31	359 4 58	55 1	1
30	207 54 15	57 14	237 48 42	62 23	270 0 0	65 25	302 11 18	62 34	332 5 45	57 23	360 0 0	55 0	0
Pl.	Libra.	Diff.	Scorpio.	Diff.	Sagittary.	Diff.	Capricorn.	Diff.	Aquarius.	Diff.	Pisces.	Diff.	Pl.

N. B. When the Sun is in the first Quadrant of the Ecliptic, γ , δ , η , (as in the First EXAMPLE) the Fourth Term, or Arch, is the R. A. from Aries. When the Sun is in the Second Quadrant, ϵ , ζ , θ , then the Fourth Term, or Arc, must be deducted from 180° for the Sun's R. A. from γ . When the Sun is in the third Quadrant, ι , κ , λ , the Fourth Term, or Arc, must be added to 180° , for the Sun's R. A. from γ . And when the Sun is in the fourth Quadrant, ν , ω , x , the Fourth Term, or Arc, must be deducted from 360° , for the Sun's R. A. from γ .

EXAMPLE III. To find the Sun's Right Ascension correspondent to $\dagger 12^{\circ}$?
 $\dagger 12^{\circ} = 252^{\circ} = 72^{\circ}$ above 180° .

To const. Log. . . 9.9624801
 Add 72° T. . . 10.4882240

+ $70^{\circ} 29' 38''$ T. 10.4507041
 180

Sum $250^{\circ} 29' 38''$ R. A. required.
 To const. Log. . . 9.9624801
 Add 27° T. . . 9.7071659

- $25^{\circ} 2' 57''$ T. 9.6696460
 360

Diff. 334 57 3 R. A. required.

EXAMPLE IV. To find the Sun's Right Ascension correspondent to $\text{x} 3^{\circ}$?
 $\text{x} 3^{\circ} = 333^{\circ} = 153^{\circ}$ above 180° ; Supplement $= 27^{\circ}$.

N. B. The Right Ascension to the first 6 Signs of the Sun's Place being found, the Right Ascension to the last 6 Signs follow, by adding 180° to the Right Ascension of the former.

The ROYAL ASTRONOMER,

RIGHT ASCENSION of the SUN in TIME, to the present Mean OBLIQUITY of the ECLIPTIC, $23^{\circ} 28' \frac{1}{2}$.

Argument. SUN's true PLACE in the ECLIPTIC.

☉ Pl.	☿ Sig. 0	Dif.	♈ Sig. 1.	Dif.	♊ Sig. 2	Dif.	♊ Sig. 3	Dif.	♊ Sig. 4	Dif.	♊ Sig. 5	Dif.	☉ Pl.
0	h m s	m s	h m s	m s	h m s	m s	h m s	m s	h m s	m s	h m s	m s	0
1	0 3 40	3 40	1 51 37	3 50	3 51 15	4 10	6 0 0	4 22	8 8 45	4 10	10 8 23	3 49	30
2	0 7 20	3 40	1 55 27	3 50	3 55 25	4 11	6 4 22	4 21	8 12 55	4 10	10 12 12	3 49	29
3	0 11 0	3 40	1 59 17	3 50	3 59 36	4 11	6 8 43	4 22	8 17 4	4 9	10 16 1	3 49	28
4	0 14 41	3 41	2 3 7	3 52	4 3 47	4 13	6 13 5	4 21	8 21 12	4 8	10 19 49	3 48	27
5	0 18 21	3 40	2 6 59	3 52	4 8 0	4 12	6 17 26	4 22	8 25 19	4 7	10 23 36	3 47	26
6	0 22 1	3 40	2 10 51	3 52	4 12 12	4 12	6 21 43	4 21	8 29 26	4 7	10 27 23	3 47	25
7	0 25 42	3 41	2 14 43	3 52	4 16 26	4 14	6 26 9	4 21	8 33 32	4 6	10 31 9	3 46	24
8	0 29 23	3 41	2 18 36	3 53	4 20 40	4 14	6 30 30	4 21	8 37 37	4 5	10 34 54	3 45	23
9	0 33 4	3 41	2 22 30	3 54	4 24 55	4 15	6 34 51	4 21	8 41 42	4 5	10 38 40	3 46	22
10	0 36 44	3 40	2 26 24	3 54	4 29 10	4 15	6 39 11	4 20	8 45 46	4 4	10 42 25	3 45	21
11	0 40 26	3 42	2 30 20	3 56	4 33 25	4 15	6 43 32	4 21	8 49 49	4 3	10 46 9	3 44	20
12	0 44 8	3 42	2 34 16	3 56	4 37 42	4 17	6 47 52	4 20	8 53 51	4 2	10 49 53	3 44	19
13	0 47 50	3 42	2 38 13	3 57	4 41 59	4 17	6 52 11	4 19	8 57 53	4 2	10 53 37	3 44	18
14	0 51 32	3 42	2 42 10	3 57	4 46 16	4 18	6 56 31	4 20	9 1 54	4 1	10 57 20	3 43	17
15	0 55 14	3 42	2 46 8	3 58	4 50 34	4 18	7 0 50	4 19	9 5 54	4 0	11 1 3	3 43	16
16	0 58 57	3 43	2 50 7	3 59	4 54 52	4 18	7 5 9	4 19	9 9 53	3 59	11 4 46	3 43	15
17	1 2 40	3 43	2 54 6	4 0	4 59 10	4 19	7 9 26	4 18	9 13 52	3 59	11 8 28	3 42	14
18	1 6 23	3 43	2 58 6	4 1	5 3 29	4 20	7 13 44	4 18	9 17 50	3 58	11 12 10	3 42	13
19	1 10 7	3 44	3 2 7	4 2	5 7 49	4 19	7 18 1	4 17	9 21 47	3 57	11 15 52	3 42	12
20	1 13 51	3 44	3 6 9	4 2	5 12 8	4 20	7 22 18	4 17	9 25 44	3 57	11 19 34	3 42	11
21	1 17 35	3 44	3 10 11	4 2	5 16 28	4 20	7 26 35	4 17	9 29 40	3 56	11 23 16	3 42	10
22	1 21 20	3 45	3 14 14	4 3	5 20 49	4 21	7 30 50	4 15	9 33 36	3 56	11 26 56	3 40	9
23	1 25 6	3 46	3 18 18	4 4	5 25 9	4 21	7 35 5	4 15	9 37 30	3 54	11 30 37	3 41	8
24	1 28 51	3 45	3 22 23	4 5	5 29 30	4 21	7 39 20	4 15	9 41 24	3 54	11 34 18	3 41	7
25	1 32 37	3 46	3 26 28	4 5	5 33 51	4 21	7 43 34	4 14	9 45 17	3 53	11 37 59	3 41	6
26	1 36 24	3 47	3 30 34	4 6	5 38 12	4 21	7 47 48	4 14	9 49 9	3 52	11 41 39	3 40	5
27	1 40 11	3 47	3 34 41	4 7	5 42 34	4 22	7 52 0	4 12	9 53 1	3 52	11 45 19	3 40	4
28	1 43 59	3 48	3 38 48	4 7	5 46 55	4 22	7 56 13	4 13	9 56 53	3 52	11 49 0	3 41	3
29	1 47 48	3 49	3 42 56	4 8	5 51 17	4 21	8 0 24	4 11	10 0 43	3 50	11 52 40	3 40	2
30	1 51 37	3 49	3 47 5	4 9	5 55 38	4 21	8 4 35	4 11	10 4 33	3 50	11 56 20	3 40	1
☉ Pl.	Aries.	Dif.	Taurus.	Dif.	Gemini.	Dif.	Cancer.	Dif.	Leo.	Dif.	Virgo.	Dif.	☉ Pl.

N. B. The Right Ascension in Time being found, (in Tab. above) to the Degrees of Right Ascension (in the former Tables, p. 24, 25) as far as 90° , the Right Ascension from thence, in Time, to 180° , will be the Complement of the former Right Ascension (taken backwards) to 12 Hours of Time; so that the Right Ascension in Time to any Number of Signs and Degrees of the Sun's Place being given, the Complement thereof to 12 Hours will be the Right Ascension in Time to the Supplement of those Signs and Degrees of the Sun's Place to 6 Signs.

EXAMPLE $1^{\text{h}} 16^{\text{m}} \text{ R. A. } 2^{\text{h}} 54^{\text{m}} 6^{\text{s}}$

4	14	9	5	54
6	0	12	0	0

Proof. }

RIGHT

RIGHT ASCENSION of the SUN in TIME, to the present Mean OBLIQUITY of the ECLIPTIC, $23^{\circ} 28' \frac{1}{2}$.

Argument. SUN's true PLACE in the ECLIPTIC.

☉ Pl.	☉ Sig. 6	Dif.	☉ Sig. 7	Dif.	☉ Sig. 8	Dif.	☉ Sig. 9	Dif.	☉ Sig. 10	Dif.	☉ Sig. 11	Dif.	☉ Pl.
o	h m s	m s	h m s	m s	h m s	m s	h m s	m s	h m s	m s	h m s	m s	o
0	12 0 0	3 40	13 51 37	3 50	15 51 15	4 10	18 0 0	4 22	20 8 45	4 10	22 8 23	3 39	30
1	12 3 40	3 40	13 55 27	3 50	15 55 25	4 11	18 4 22	4 21	20 12 55	4 9	22 12 12	3 49	29
2	12 7 20	3 40	13 59 17	3 50	15 59 36	4 11	18 8 43	4 22	20 17 4	4 8	22 16 1	3 48	28
3	12 11 0	3 41	14 3 7	3 52	16 3 47	4 13	18 13 5	4 21	20 21 12	4 7	22 19 49	3 47	27
4	12 14 41	3 40	14 6 59	3 52	16 8 0	4 12	18 17 26	4 22	20 25 19	4 7	22 23 36	3 47	26
5	12 18 21	3 40	14 10 51	3 52	16 12 12	4 14	18 21 48	4 21	20 29 26	4 6	22 27 23	3 46	25
6	12 22 1	3 41	14 14 43	3 53	16 16 26	4 14	18 26 9	4 21	20 33 32	4 5	22 31 9	3 45	24
7	12 25 42	3 41	14 18 36	3 54	16 20 40	4 15	18 30 30	4 21	20 37 37	4 5	22 34 54	3 46	23
8	12 29 23	3 41	14 22 30	3 54	16 24 55	4 15	18 34 51	4 20	20 41 42	4 4	22 38 40	3 45	22
9	12 33 4	3 40	14 26 24	3 56	16 29 10	4 15	18 39 11	4 21	20 45 46	4 3	22 42 25	3 44	21
10	12 36 44	3 42	14 30 20	3 56	16 33 25	4 17	18 43 22	4 20	20 49 49	4 2	22 46 9	3 44	20
11	12 40 26	3 42	14 34 16	3 57	16 37 42	4 17	18 47 52	4 19	20 53 51	4 2	22 49 53	3 44	19
12	12 44 8	3 42	14 38 13	3 57	16 41 59	4 17	18 52 11	4 20	20 57 53	4 1	22 53 37	3 43	18
13	12 47 50	3 42	14 42 10	3 58	16 46 16	4 18	18 56 31	4 19	21 1 54	4 0	22 57 20	3 43	17
14	12 51 32	3 42	14 46 8	3 59	16 50 34	4 18	19 0 50	4 19	21 5 54	3 59	23 1 3	3 43	16
15	12 55 14	3 43	14 50 7	3 59	16 54 52	4 18	19 5 9	4 18	21 9 53	3 59	23 4 46	3 42	15
16	12 58 57	3 43	14 54 6	4 0	16 59 10	4 19	19 9 26	4 18	21 13 52	3 58	23 8 28	3 42	14
17	13 2 40	3 43	14 58 6	4 1	17 3 29	4 20	19 13 44	4 17	21 17 50	3 57	23 12 10	3 42	13
18	13 6 23	3 44	15 2 7	4 2	17 7 49	4 19	19 18 1	4 17	21 21 47	3 57	23 15 52	3 42	12
19	13 10 7	3 44	15 6 9	4 2	17 12 8	4 20	19 22 18	4 17	21 25 44	3 56	23 19 34	3 42	11
20	13 13 51	3 44	15 10 11	4 3	17 16 28	4 21	19 26 35	4 15	21 29 40	3 56	23 23 16	3 40	10
21	13 17 35	3 45	15 14 14	4 4	17 20 49	4 20	19 30 50	4 15	21 33 36	3 54	23 26 56	3 41	9
22	13 21 20	3 46	15 18 18	4 5	17 25 9	4 21	19 35 5	4 15	21 37 30	3 54	23 30 37	3 41	8
23	13 25 6	3 45	15 22 23	4 5	17 29 30	4 21	19 39 20	4 14	21 41 24	3 53	23 34 18	3 41	7
24	13 28 51	3 46	15 26 28	4 6	17 33 51	4 21	19 43 34	4 14	21 45 17	3 52	23 37 59	3 40	6
25	13 32 37	3 47	15 30 34	4 7	17 38 12	4 22	19 47 48	4 12	21 49 9	3 52	23 41 39	3 40	5
26	13 36 24	3 47	15 34 41	4 7	17 42 34	4 21	19 52 0	4 13	21 53 1	3 52	23 45 19	3 41	4
27	13 40 11	3 48	15 38 48	4 8	17 46 55	4 22	19 56 13	4 11	21 56 53	3 50	23 49 0	3 40	3
28	13 43 59	3 49	15 42 56	4 9	17 51 17	4 21	20 0 24	4 11	22 0 43	3 50	23 52 40	3 40	2
29	13 47 48	3 49	15 47 5	4 10	17 55 38	4 22	20 4 35	4 10	22 4 33	3 50	23 56 20	3 40	1
30	13 51 37		15 51 15		18 0 0		20 8 45		22 8 23		24 0 0		0
☉ Pl.	Libra.	Dif.	Scorpio.	Dif.	Sagittary.	Dif.	Capricorn.	Dif.	Aquarius.	Dif.	Pisces.	Dif.	☉ Pl.

N. B. The Right Ascension in Time to the first 6 Signs of the Sun's Place being found (in Table, p. 26) to the Degrees of Right Ascension (in the former Tables, p. 24, 25) as far as 180° , the Right Ascension, in Time, to the last 6 Signs, will be had by adding 12 Hours to the Right Ascension of the foregoing Table.

EXAMPLE. To find the Right Ascension in Time to the Sun's true Place $\approx 18^{\circ} 27' 47''$?

$\approx 18^{\circ}$	13^h	6^m	23^s	} Excess of the Sun's Place above Degrees $27' 47''$	} 0.3344		
	+	1	44			} Dif. Time for 1 Degree more . . . $3^m 44^s$	} 1.2061
Ho.	13	8	7	R. A. required.	$1^m 44^s$		

Logistical Logarithms.

For reducing PARTS of the EQUATOR into TIME.						For reducing TIME into PARTS of the EQUATOR.						REFRACTION and PARALLAX ☉, according to the Paris-Observatory.					
Argument. Parts of the Equator.						Argument. Hours, Minutes, Seconds, &c. of Time.						Argument. Apparent Altitude ☉					
°	Ho. Min.	°	Ho. Min.			Hours.	M.	°	Min.	°		Alt. ☉	Refract.	Alt. ☉	Refract.	Alt. ☉	Refract.
'	Min. Sec.	'	Min. Sec.				Sec.	'	"								
"	Sec. Th.	"	Sec. Th.	°	Hrs. Min.		Th.	"	"								
1	0 4	31	2 4	70	4 40	1	15	1	0 15	31	7 45	0	32 20	35	1 23	70	0 21
2	0 8	32	2 8	80	5 20	2	30	2	0 30	32	8 0	1	27 56	36	1 20	71	0 20
3	0 12	33	2 12	90	6 0	3	45	3	0 45	33	8 15	2	21 4	37	1 18	72	0 19
4	0 16	34	2 16	100	6 40	4	60	4	1 0	34	8 30	3	16 6	38	1 15	73	0 18
5	0 20	35	2 20	110	7 20	5	75	5	1 15	35	8 45	4	12 48	39	1 12	74	0 17
6	0 24	36	2 24	120	8 0	6	90	6	1 30	36	9 0	5	10 32	40	1 10	75	0 16
7	0 28	37	2 28	130	8 40	7	105	7	1 45	37	9 15	6	8 55	41	1 7	76	0 14
8	0 32	38	2 32	140	9 20	8	120	8	2 0	38	9 30	7	7 44	42	1 5	77	0 13
9	0 36	39	2 36	150	10 0	9	135	9	2 15	39	9 45	8	6 47	43	1 3	78	0 12
10	0 40	40	2 40	160	10 40	10	150	10	2 30	40	10 0	9	6 4	44	1 1	79	0 11
11	0 44	41	2 44	170	11 20	11	165	11	2 45	41	10 15	10	5 28	45	0 59	80	0 10
12	0 48	42	2 48	180	12 0	12	180	12	3 0	42	10 30	11	4 58	46	0 58	81	0 9
13	0 52	43	2 52	190	12 40	13	195	13	3 15	43	10 45	12	4 32	47	0 56	82	0 8
14	0 56	44	2 56	200	13 20	14	210	14	3 30	44	11 0	13	4 12	48	0 54	83	0 7
15	1 0	45	3 0	210	14 0	15	225	15	3 45	45	11 15	14	3 54	49	0 52	84	0 6
16	1 4	46	3 4	220	14 40	16	240	16	4 0	46	11 30	15	3 38	50	0 50	85	0 5
17	1 8	47	3 8	230	15 20	17	255	17	4 15	47	11 45	16	3 24	51	0 49	86	0 4
18	1 12	48	3 12	240	16 0	18	270	18	4 30	48	12 0	17	3 11	52	0 47	87	0 3
19	1 16	49	3 16	250	16 40	19	285	19	4 45	49	12 15	18	3 0	53	0 45	88	0 2
20	1 20	50	3 20	260	17 20	20	300	20	5 0	50	12 30	19	2 49	54	0 43	89	0 1
21	1 24	51	3 24	270	18 0	21	315	21	5 15	51	12 45	20	2 39	55	0 41	90	0 0
22	1 28	52	3 28	280	18 40	22	330	22	5 30	52	13 0	21	2 31	56	0 40	Parallax ☉.	
23	1 32	53	3 32	290	19 20	23	345	23	5 45	53	13 15	22	2 25	57	0 38	Argt. ☉ ap. Alt.	
24	1 36	54	3 36	300	20 0	24	360	24	6 0	54	13 30	23	2 18	58	0 37	Alt. +	
25	1 40	55	3 40	310	20 40	25		25	6 15	55	13 45	24	2 12	59	0 35	°	' "
26	1 44	56	3 44	320	21 20	26		26	6 30	56	14 0	25	2 6	60	0 34	0	0 10
27	1 48	57	3 48	330	22 0	27		27	6 45	57	14 15	26	2 0	61	0 33	10	0 10
28	1 52	58	3 52	340	22 40	28		28	7 0	58	14 30	27	1 55	62	0 31	20	0 9
29	1 56	59	3 56	350	23 20	29		29	7 15	59	14 45	28	1 51	63	0 30	30	0 9
30	2 0	60	4 0	360	24 0	30		30	7 30	60	15 0	29	1 46	64	0 28	40	0 8
												30	1 42	65	0 27	50	0 6
												31	1 38	66	0 26	60	0 5
												32	1 34	67	0 25	70	0 3
												33	1 30	68	0 24	80	0 2
												34	1 27	69	0 22	90	0 0
EXAMPLE.						EXAMPLE.						EXAMPLE.					
To find the Hours in 230°. 15'. 16" ?						To find the Degrees in 15 ^h . 21 ^m . 1 ^s . 4 th . of Time ?						To find the ☉ true Alt. to 7°. Alt. Apparent ?					
230°. = 15 ^h . 20 ^m .						15 ^h . = 225°.						7°. Ap. Alt. — 7'. 44". Refraction,					
15' = 1						21 ^m . = 5 15'.						+ 0 10 Parallax.					
16" = 0 1 ^s . 4 th .						1 ^s . = 0 0 15".						Sum — 7 34 deduct.					
Sum, 15 21 1 4 Answer.						4 th . = 1						From 7° 0 0 ☉ Ap. Alt.					
						Sum, 230 15 16 Anf.						☉'s true Alt. 6 52 26 required.					

For reducing PARTS of the EQUATOR, (passed over by the EARTH'S ROTATION, and PROGRESSION, or SUN's apparent Motion, from Noon to Noon) into Mean solar HOURS.

Argument. Parts of the Equator.

°	H. M. S.	°	H. M. S.	
'	M. S. Th.	'	M. S. Th.	
"	S. Th. Fo.	"	S. Th. Fo.	Ho. M. S.
1	0 3 59	31	2 3 39	70 4 39 14
2	0 7 58	32	2 7 39	80 5 19 7
3	0 11 58	33	2 11 38	90 5 59 1
4	0 15 57	34	2 15 38	100 6 38 54
5	0 19 56	35	2 19 37	110 7 18 47
6	0 23 55	36	2 23 37	120 7 58 42
7	0 27 54	37	2 27 36	130 8 38 35
8	0 31 53	38	2 31 35	140 9 18 28
9	0 35 52	39	2 35 34	150 9 58 22
10	0 39 52	40	2 39 33	160 10 38 15
11	0 43 52	41	2 43 32	170 11 18 8
12	0 47 51	42	2 47 32	180 11 58 2
13	0 51 51	43	2 51 31	190 12 37 55
14	0 55 50	44	2 55 30	200 13 17 48
15	0 59 50	45	2 59 30	210 13 57 42
16	1 3 40	46	3 3 29	220 14 37 35
17	1 7 48	47	3 7 28	230 15 17 28
18	1 11 47	48	3 11 27	240 15 57 23
19	1 15 47	49	3 15 27	250 16 37 16
20	1 19 46	50	3 19 27	260 17 17 9
21	1 23 45	51	3 23 26	270 17 57 3
22	1 27 45	52	3 27 25	280 18 36 56
23	1 31 44	53	3 31 24	290 19 16 49
24	1 35 43	54	3 35 24	300 19 56 43
25	1 39 43	55	3 39 23	310 20 36 36
26	1 43 42	56	3 43 23	320 21 16 30
27	1 47 41	57	3 47 23	330 21 56 24
28	1 51 40	58	3 51 22	340 22 36 17
29	1 55 40	59	3 55 22	350 23 16 11
30	1 59 40	60	3 59 21	360 23 56 4

EXAMPLE.

To find the mean solar Time answerable to 39°. 19'. 29". of the Earth's Rotation?

39°. = 2^h. 35^m. 34^s.
19'. = 1 15 47th.
29". = 1 55 40^{fo}.

2 36 51 42 40 reqd.

For reducing Mean solar HOURS into PARTS of the EQUATOR; passed over, from Noon to Noon, by the EARTH'S ROTATION on its AXIS and PROGRESSION in its Orbit.

Argument. Mean solar Hours, Minutes, Seconds, &c. of Time.

Hours.		M	°	'	"	M	°	'	"
		S.	'	"	'''	S.	'	"	'''
	°	'	"	'''	iv	°	'	"	iv
1	15	2	28	1	0	15	2	31	7 46 16
2	30	4	56	2	0	30	5	32	8 1 19
3	45	7	24	3	0	45	7	33	8 16 21
4	60	9	51	4	1	0	10	34	8 31 24
5	75	12	19	5	1	15	12	35	8 46 26
6	90	14	47	6	1	30	15	36	9 1 29
7	105	17	15	7	1	45	17	37	9 16 31
8	120	19	43	8	2	0	20	38	9 31 34
9	135	22	11	9	2	15	22	39	9 46 36
10	150	24	38	10	2	30	25	40	10 1 39
11	165	27	6	11	2	45	27	41	10 16 41
12	180	29	34	12	3	0	30	42	10 31 43
13	195	32	2	13	3	15	32	43	10 46 46
14	210	34	30	14	3	30	34	44	11 1 48
15	225	36	58	15	3	45	37	45	11 16 51
16	240	39	26	16	4	0	39	46	11 31 53
17	255	41	53	17	4	15	41	47	11 46 56
18	270	44	21	18	4	30	44	48	12 1 58
19	285	46	49	19	4	45	47	49	12 17 1
20	300	49	17	20	5	0	49	50	12 32 3
21	315	51	45	21	5	15	52	51	12 47 6
22	330	54	13	22	5	30	54	52	13 2 8
23	345	56	40	23	5	45	57	53	13 17 11
24	360	59	8	24	6	0	59	54	13 32 13
* 360° or 1 Revol. of the Earth's Equator is theref. performed in 23 ^h 56 ^m 4 ^s 7 th mean solar Time.									
				26	6	31	45	56	14 2 18
				27	6	46	7	57	14 17 21
				28	7	1	9	58	14 32 23
				29	7	16	11	59	14 47 26
				30	7	31	14	60	15 2 28

EXAMPLE.

To find the Deg. of the Earth's Rotation answerable to 23^h 56^m 4^s 7th of mean solar Time?

23^h = 345° 56' 40"
56^m = 14 2 18
4^s = 1 0 10"
7th = 1 45 17^{iv}

359 59 59 55 17

A Revolution nearly of the Earth's Equator, reqd.

The DIFFERENCE of Mean solar above syderial HOURS.

Argument. Mean Hours, Minutes, Sec^s, of Time.

H.	M.	S.	T.	H.	M.	S.	T.
M.	S.	T.	F.	M.	S.	T.	F.
S.	T.	F.	Fi.	S.	T.	F.	Fi.
1	0	9	51	31	5	5	33
2	0	19	43	32	5	15	24
3	0	29	34	33	5	25	15
4	0	39	25	34	5	35	7
5	0	49	17	35	5	44	58
6	0	59	8	36	5	54	50
7	1	9	0	37	6	4	41
8	1	18	51	38	6	14	32
9	1	28	42	39	6	24	24
10	1	38	34	40	6	34	15
11	1	48	25	41	6	44	6
12	1	58	17	42	6	53	58
13	2	8	8	43	7	3	49
14	2	17	59	44	7	13	41
15	2	27	51	45	7	23	32
16	2	37	42	46	7	33	23
17	2	47	33	47	7	43	15
18	2	57	25	48	7	53	6
19	3	7	16	49	8	2	58
20	3	17	8	50	8	12	49
21	3	26	59	51	8	22	40
22	3	36	50	52	8	32	32
23	3	46	42	53	8	42	23
24	3	56	33	54	8	52	14
25	4	6	24	55	9	2	6
26	4	16	16	56	9	11	57
27	4	26	7	57	9	21	49
28	4	35	59	58	9	31	40
29	4	45	50	59	9	41	31
30	4	55	41	60	9	51	23

CONSTRUCTION.

To find the Excess of 24 mean solar Hours, above 24 syderial Hours?

24^h mean = 360°. 59', 8".
10^{'''}. by Earth's Rotation.
But, 59'. = 3^m. 56^s.
Sec } 8" + 10^{'''} = 0 33th
P. }
28. } Excess 3 56 33
as by Tab. M. Time required.

N. B. A syderial Day is the Time in which a Fixed Star, or the same Point of the Equinoctial returns to the same Meridian; making a Revolution, or 360°, of the Earth's Equator.

A mean solar Day is the Time in which the SUN returns to the same Meridian; making 360°. 59'. 8". 10^{'''}. of the Earth's Equator, at a mean Rate of the Earth's Progression and Rotation: By which Means the SUN apparently retards of the Fixed Stars, or Fixed Stars accelerate of the SUN, at the mean Rate of 59'. 8". 10^{'''}, or 3^m. 56^s. 33th. of Time, each Day.

A TABLE,

A TABLE, shewing the DIFFERENCE between the Julian, Gregorian, and Solar Years, from 1600.

Years of Christ.	Solar Years from 1600.			The Lun. Cycle of 19 Years goes back in the Jul. Account. D ^s & P ^{ts} sub. fr. 3 Col.	Whole D ^s Dif. bet. 3 ^d & 5 th Col ^s , the Lun. Cycle goes for- ward in the Greg. Account	Years of Christ.	Solar Years from 1600.			The Lun. Cycle of 19 Y ^{rs} goes back in the Jul. Account. D ^s & P ^{ts} sub. fr. 9 Col.	Whole D ^s Dif. bet. 9 th and 11 th Col ^s The Lun. Cycle goes for- ward in the Greg. Account
	Gregorian Years from 1600.		Hours sub. fr. or add to Gregor ⁿ for Solar Years.				Gregorian Y ^{rs} . from 1600.		Hours add to Gregor. for Solar Years.		
	Jul. Y ^{rs} from 1600.	Days add to Jul. Y ^{rs} for Greg.					J. Y ^{rs} from 1600.	Days ad ^d to Jul. Y ^{rs} for Greg.			
	Centuries.						Cen ^s .				
No Col ^s 1.	2.	3.	4.	5.	6.	7.	8.	9.	10.	11.	12.
Add 10 Days to the Days in the 3 ^d and 9 th Col ^s , and the Sum will be the Days the Gregorian is forward of the Julian Account, or D ^s Dif. betw. N. and Old Style.											
1600		0									
1700	1	1	— 6	$\frac{1}{3}$	1	5300	37	28	+ 11	12	16
1800	2	2	— 11	$\frac{2}{3}$	1	5400	38	29	6	$12\frac{1}{4}$	17
1900	3	3	— 17	1	2	5500	39	30	0	$12\frac{2}{3}$	17
2000	4	3	+ 2	$1\frac{1}{3}$	2	5600	40	30	19	13	17
2100	5	4	— 4	$1\frac{2}{3}$	2	5700	41	31	13	$13\frac{1}{3}$	18
2200	6	5	— 2	2	3	5800	42	32	7	$13\frac{2}{3}$	18
2300	7	6	— 14	$2\frac{1}{3}$	4	5900	43	33	2	14	19
2400	8	6	+ 4	$2\frac{2}{3}$	3	6000	44	33	21	$14\frac{1}{4}$	19
2500	9	7	— 2	3	4	6100	45	34	15	$14\frac{1}{2}$	19
2600	10	8	— 7	$3\frac{1}{3}$	5	6200	46	35	9	$14\frac{3}{4}$	20
2700	11	9	— 13	$3\frac{2}{3}$	5	6300	47	36	4	15	21
2800	12	9	+ 6	4	5	6400	48	36	22	$15\frac{1}{3}$	21
2900	13	10	+ 0	$4\frac{1}{3}$	6	6500	49	37	7	$15\frac{1}{2}$	21
3000	14	11	— 5	$4\frac{2}{3}$	6	6600	50	38	11	16	22
3100	15	12	— 10	5	7	6700	51	39	6	$16\frac{1}{3}$	23
3200	16	12	+ 8	$5\frac{1}{4}$	7	6800	52	39	24	$16\frac{2}{3}$	22
3300	17	13	+ 2	$5\frac{1}{2}$	7	6900	53	40	19	17	23
3400	18	14	— 4	$5\frac{3}{4}$	8	7000	54	41	13	$17\frac{1}{3}$	24
3500	19	15	— 9	6	9	7100	55	42	7	$17\frac{2}{3}$	24
3600	20	15	+ 9	$6\frac{1}{3}$	8*	7200	56	42	26	18	24
3700	21	16	+ 4	$6\frac{2}{3}$	9	7300	57	43	21	$18\frac{1}{4}$	25
3800	22	17	— 2	7	10	7400	58	44	15	$18\frac{3}{4}$	25
3900	23	18	— 8	$7\frac{1}{3}$	10*	7500	59	45	10	19	26
4000	24	18	+ 11	$7\frac{2}{3}$	10	7600	60	45	28	$19\frac{1}{4}$	26
4100	25	19	+ 6	8	11	7700	61	46	23	$19\frac{2}{3}$	26
4200	26	20	+ 0	$8\frac{1}{3}$	12	7800	62	47	17	20	27
4300	27	21	— 5	$8\frac{2}{3}$	12	7900	63	48	12	$20\frac{1}{3}$	28
4400	28	21	+ 13	9	12	8000	64	48	30	$20\frac{2}{3}$	27
4500	29	22	+ 7	$9\frac{1}{3}$	13	8100	65	49	24	21	28
4600	30	23	+ 2	$9\frac{2}{3}$	13	8200	66	50	19	$21\frac{1}{3}$	29
4700	31	24	— 4	10	14	8300	67	51	13	$21\frac{2}{3}$	29
4800	32	24	+ 15	$10\frac{1}{3}$	14	8400	68	51	32	22	29
4900	33	25	+ 9	$10\frac{2}{3}$	14	8500	69	52	26	$22\frac{1}{3}$	30
5000	34	26	+ 4	11	15	8600	70	53	21	$22\frac{2}{3}$	0
5100	35	27	— 2	$11\frac{1}{3}$	16	&c.	&c.	&c.	&c.	&c.	&c.
5200	36	27	+ 17	$11\frac{2}{3}$	15						

EXAMPLE of the USE. In 10 Hundred Julian Years there are 10 Hundred Years Gregorian and 8 Days; and 10 Hundred Solar Years, 8 Days—7 Hours: so that the Gregorian Account will, in that Period, be 8 Days forward of the Month Days of the Julian Reckoning. And the Seasons of the Solar Years, in that Time, will be 7 Hours backward of the Gregorian Account. Also the Solar Seasons, or Years, will be 8 Days—7 Hours forward of the Month-Days in the Julian Reckoning.

CONSTRUCTION of the FOREGOING TABLE.

POPE GREGORY XIII, in the Beginning of the Month of October 1582, adjusted the Julian to the Solar Account of Time; from which Period the foregoing Table is composed; shewing, at Sight, the Difference, in Days, betwixt the Julian, Gregorian, and Solar Years. Wherein you may see how many Days the Gregorian Years, in given Centuries, advance (with Respect to Month-days) of the same Number of Years Julian. And how many Hours the Solar Years or Seasons, go forward or retreat, (with Respect to the Month-days) in the same Number of Centuries of Gregorian Years. As also the Number of Days and Hours the Solar Seasons, or Years, retreat (with Respect to Month-days) in the Julian Account. And the Number of Days the Lunar Cycle (of 19 Years) falls back in the Months of the Julian Reckoning in a given Period. By comparing which Retardation of the Lunar Cycle, in the Month days of the Julian Account, with the Month-days advanced by the Gregorian of the Julian Reckoning, in the same Period (referring to one Standard), the Difference, or Excess, will be the Days that the Lunar Cycle is carried forward in the Month-days, (in the same Period) of the Gregorian Account.

Hence, the Reason of the Golden Numbers being carried forward by certain Days, in certain Periods, in the Earl of Macclesfield's curious Table for finding Easter Full Moon; and hence the Mystry of the two last Tables in the Style-Act is unfolded.

Note, A Solar Year is here stated at $365^d 5^h 48^m 55^s$, wanting $11^m 5^s$ of a Julian Year; by which the odd Hours in the 4th and 10th Columns are computed. The Julian Anticipation of the Lunar Cycle of 19 Years is according to that of a Day in 310,7 Julian Years; whence the 5th and 11th Columns are computed near the Truth. Wherein above half a Day is taken for another whole Day; but less than half a Day more is taken for Nothing more. Our Table corrects, in some Periods, by a whole Day, the Style-Act Table. } *query*

There being 146097 Days in 400 Gregorian Years, or 3 Days less than in the same Number of Years Julian.

	d	h	m	s	th	
In 19 Gregorian Years there will be	6939	14	34	48	0	= 6939 ^d ,6075
But in 235 Synodical Months there are	6939	16	31	56	30	= 6939,688848

Greg Year
365.5.49.12

Hence the New or Full Moon goes forward }
in one Lunar Cycle, the Difference } 1 57 8 30 = ,081348

Whence, in about 233,56 Years Gregorian, if the Gregorian Year went regularly forward of the Julian, as to the Month-day of each Account, a New or Full Moon would be advanced one Month-day, by the Lunar Cycle, in the Gregorian Account. But 12 Days completely are advanced in 2800 Gregorian Years. And our Table shews that the Number of Days 3, 6, 9, 12, &c. should stand against 2200, 2900, 3700, 4400, &c. as they stand in the Style-Act Table. And that 9 and not 8 should stand against 3600, 11 and not 10 against 3900, and 21 and not 20, against 6400, correcting the Errors in the Style-Act Table.

EXAMPLE. To find the Anticipation of the Lunar Cycle in the Julian Account, for any Number of Years Julian; suppose 2300 from the Year 1600, N. S.?

As 310,7 Years : 1 Day :: 100 Years : ,322 Day, nearly; a constant Multiplier for Hundreds. Whence $23 \times ,322 = 7,4$ Days, or 7 nearest, required.

The Gregorian Month-days being then advanced 18 Days forward of the Julian Month-days, therefore $18 - 7 = 11$ Days, the Lunar Cycle is then advanced in the Month-days of the Gregorian Account, and not 10, as in the Style-Act Table.

To find the Julian, Gregorian, and Solar Years, or the different Accounts of Time, correspondent to each other?

EXAMPLE. 1400 Jul. Yrs = 1400 Greg. Yrs + 11 Days, the Gregorian Account of Time.
1400 Gr. Yrs + $1400 \times 11^m 5^s = 1400$ ————— + $10^d 18^h 36^m 40^s$, the Solar Account.

N. B. The Julian is then short of } Diff. — 5 23 20 (as by Table) the Solar Account is short
the Gregorian Account by 11 Days. } of the Gregorian.

By the above Rule the 4th and 10th Columns are constructed.

DECIMALS.

DECIMALS of a DAY.

N. B. A Figure with a fine Stroke cross it, denotes that the following Decimals repeat from thence ad infinitum. And a Dot (.) after any Figure, denotes that Decimal to be correct.

HOURS.		MINUTES.				SECONDS.			
Ho.	Decim.	Min.	Decim.	Min.	Decim.	Sec.	Decim.	Sec.	Decim.
1	0,04166	1	,000694	31	,021527	1	,0000115740	31	,0003587862
2	0,08333	2	,001388	32	,022227	2	,000023448	32	,000370370
3	0,125	3	,002083	33	,022916	3	,000034722	33	,000381944
4	0,16666	4	,002777	34	,023611	4	,000046796	34	,000393818
5	0,20833	5	,003477	35	,024308	5	,0000578703	35	,0004050925
6	0,25	6	,004166	36	,025	6	,000069444	36	,000416666
7	0,29166	7	,004861	37	,025694	7	,0000810185	37	,0004282407
8	0,33333	8	,005558	38	,026388	8	,000092892	38	,000439814
9	0,375	9	,00625	39	,027083	9	,000104166	39	,000451388
10	0,41666	10	,006944	40	,027777	10	,000115740	40	,000462962
11	0,45833	11	,007638	41	,028477	11	,0001273148	41	,0004745770
12	0,5	12	,008333	42	,029166	12	,000138888	42	,000486111
13	0,54166	13	,009027	43	,029861	13	,0001504629	43	,0004976851
14	0,58333	14	,009722	44	,030558	14	,000162037	44	,000509259
15	0,625	15	,010416	45	,03125	15	,000173611	45	,00052083
16	0,66666	16	,011111	46	,031944	16	,000185185	46	,0005324074
17	0,70833	17	,011805	47	,032638	17	,0001967592	47	,0005439814
18	0,75	18	,0125	48	,033333	18	,000208333	48	,000555558
19	0,79166	19	,013194	49	,034027	19	,0002199074	49	,0005671796
20	0,83333	20	,013888	50	,034722	20	,00023148	50	,000578704
21	0,875	21	,014583	51	,035416	21	,000243058	51	,000590277
22	0,91666	22	,015277	52	,036111	22	,000254629	52	,0006018518
23	0,95833	23	,015977	53	,036805	23	,0002662037	53	,0006134759
24	1	24	,016666	54	,0375	24	,000277777	54	,000625
25	1,04166	25	,017361	55	,038194	25	,0002893518	55	,0006365740
26	1,08333	26	,018058	56	,038888	26	,000300925	56	,000648148
27	1,125	27	,01875	57	,039583	27	,0003125	57	,000659722
28	1,16666	28	,019444	58	,040277	28	,000324074	58	,000671796
29	1,20833	29	,020138	59	,040977	29	,0003356481	59	,0006828703
30	1,25	30	,020833	60	,041666	30	,000347222	60	,000694444

A TABLE of the NUMBER of DAYS from any one Day in any MONTH, to the same Day in any other MONTH following.

Months.	Jan.	Feb.	March.	April.	May.	June.	July.	August.	Sept.	Oct.	Nov.	Dec.
January	365	31	59	90	120	151	181	212	243	273	304	334
February	334	365	28	59	89	120	150	181	212	242	273	303
March	306	337	365	31	61	92	122	153	184	214	245	275
April	275	306	334	365	30	61	91	122	153	183	214	244
May	245	276	304	335	365	31	61	92	123	153	184	214
June	214	245	273	304	334	365	30	61	92	122	153	183
July	184	215	243	274	304	335	365	31	62	92	123	153
August	153	184	212	243	273	304	334	365	31	61	92	122
September	122	153	181	212	242	273	303	334	365	30	61	91
October	92	123	151	182	212	243	273	304	335	365	31	61
November	61	92	120	151	181	212	242	273	304	334	365	30
December	31	62	90	121	151	182	212	243	274	304	335	365

N. B. One Day must be added when February of Leap-year comes between two Months.

EXAMPLE. To find the Number of Days betwixt December 19, 1759. and July 24, following?

From December 19 to July 19 following, we find, to the Right, against Dec. and under July, 213 Days, (adding 1 for Feb. of Leap-year coming between) to which 5 Days more being added, (from 19 to 24th July) the Sum is 218 Days required.

MEAN

MEAN MOTION of the SUN for MONTHS and DAYS, to Thirds of a DEGREE, for correctly constructing TABLES, pag. 5, 6, 7.

D's	Months	M. Motion $\odot$	Apog.	Stars.
		$\circ$ $'$ $''$	$\circ$ $'$ $''$	$\circ$ $'$ $''$
31	Jan. $\circ$	0 0 0	0 0 0	0 0 0
59	Feb. $\circ$	1 0 33	18 13 5	13 4 17
90	March $\circ$	1 28 9	11 26 9	56 8 9
120	April $\circ$	2 28 42	29 38 15	9 12 26
151	May $\circ$	3 28 16	39 30 20	12 16 35
181	June $\circ$	4 28 49	57 42 25	25 20 52
212	July $\circ$	5 28 24	7 44 30	28 25 1
243	August $\circ$	6 28 57	25 56 35	42 29 19
273	Sept. $\circ$	7 29 30	44 8 40	55 33 36
304	Oct. $\circ$	8 29 4	54 0 45	58 37 44
334	Nov. $\circ$	9 29 38	12 12 51	11 42 2
	Dec. $\circ$	10 29 12	22 4 56	14 46 10
Days	1	0 0 59	8 20 0	10 0 8
	2	0 1 58	16 39 0	20 0 16
	3	0 2 57	24 59 0	30 0 25
	4	0 3 56	33 19 0	40 0 33
	5	0 4 55	41 39 0	50 0 41
	6	0 5 54	49 50 1	0 49
	7	0 6 53	58 18 1	11 0 58
	8	0 7 53	6 38 1	21 1 6
	9	0 8 52	14 58 1	31 1 14
	10	0 9 51	23 17 1	41 1 23
	11	0 10 50	31 37 1	51 1 31
	12	0 11 49	39 57 2	1 1 39
	13	0 12 48	48 17 2	11 1 47
	14	0 13 47	56 36 2	21 1 56
	15	0 14 47	4 56 2	31 2 4
	16	0 15 46	13 16 2	42 2 13
	17	0 16 45	21 36 2	52 2 21
	18	0 17 44	29 55 3	2 2 29
	19	0 18 43	38 15 3	12 2 37
	20	0 19 42	46 35 3	22 2 46
	21	0 20 41	54 55 3	32 2 54
	22	0 21 41	3 14 3	42 3 2
	23	0 22 40	11 34 3	52 3 11
	24	0 23 39	19 54 4	2 3 19
	25	0 24 38	28 14 4	12 3 27
	26	0 25 37	36 33 4	23 3 35
	27	0 26 36	44 53 4	33 3 44
	28	0 27 35	53 13 4	43 3 52
	29	0 28 35	1 33 4	53 4 0
	30	0 29 34	9 52 5	3 4 9
	31	1 0 33	18 12 5	13 4 17
Correct CONSTRUCTION of Tab. p. 5, 6, 7.				
May	30	3 28 16	39 30 20	12 16 35
	30	0 29 34	9 52 5	3 4 9
May	30	4 27 50	49 22 25	15 20 44
October	19	8 29 4 54	0 45 58	37 44
	19	0 18 43	38 15 3	12 2 37
October	19	9 17 48	32 15 49	10 40 21
Sec.				

EQUATION of the POINTS of the EQUINOCTIAL, for the computed Longitudes of the SUN and MOON, from the Mean *Ecliptic* OB-
LIQUITY, $23^{\circ} 28' 30''$. Also the EQUAT. of the M. to the *tr. Ecliptic* OB-
LIQUITY. Together with the *Annual Preces.* of the EQU. from
the NUTATION of the Earth's *Axis*, during one Revol. of the D's $\odot$,
accor^d to Dr. BRADLEY's Improv^d. of the R. Observ. at Greenwich.

Arg. LONGITUDE of the MOON'S Node, from the Equinox.

Lon. D's ⊙	Eq ⁿ . to Points of Eq ^x .	Eq ⁿ . to M. Obl. of Ecl ^c .	True Obli- quity of the Ecliptic.	Eq ⁿ . to Annual M. Obl. of Ecl ^c .	Annual Prec. of Eq. P ^{ts} .	Eq ⁿ . to Points of Eq ^x .	Long. D's ⊙
°	′	″	° ′ ″	′ ″	″	′ ″	° ′
0	0	—0,	23 28 39,	+9,0	58,0	+0,	12 0
	5	—2,	39,	+9,0	57,9	+2,	25
	10	—3,9	38,9	+8,9	57,9	+3,9	20
	15	—5,8	38,7	+8,7	57,7	+5,8	15
	20	—7,7	38,5	+8,5	57,5	+7,7	10
	25	—9,6	38,2	+8,2	57,3	+9,6	5
1	0	—11,3	23 28 37,8	+7,8	57,0	+11,3	11 0
	5	—13,	37,4	+7,4	56,6	+13,	25
	10	—14,5	36,9	+6,9	56,2	+14,5	20
	15	—16,	36,4	+6,4	55,7	+16,	15
	20	—17,3	35,8	+5,8	55,2	+17,3	10
	25	—18,5	35,2	+5,2	54,7	+18,5	5
2	0	—19,6	23 28 34,5	+4,5	54,2	+19,6	10 0
	5	—20,5	33,8	+3,8	53,6	+20,5	25
	10	—21,2	33,1	+3,1	53,0	+21,2	20
	15	—21,8	32,3	+2,3	52,3	+21,8	15
	20	—22,2	31,6	+1,6	51,7	+22,2	10
	25	—22,5	30,8	+0,8	51,0	+22,5	5
3	0	—22,6	23 28 30,	+0,0	50,3	+22,6	9 0
	5	—22,5	29,2	—0,8	49,7	+22,5	25
	10	—22,2	28,4	—1,6	49,0	+22,2	20
	15	—21,8	27,7	—2,3	48,4	+21,8	15
	20	—21,2	26,9	—3,1	47,7	+21,2	10
	25	—20,5	26,2	—3,8	47,1	+20,5	5
4	0	—19,6	23 28 25,5	—4,5	46,5	+19,6	8 0
	5	—18,5	24,8	—5,2	46,0	+18,5	25
	10	—17,3	24,2	—5,8	45,5	+17,3	20
	15	—16,	23,6	—6,4	45,0	+16,	15
	20	—14,5	23,1	—6,9	44,5	+14,5	10
	25	—13,	22,6	—7,4	44,1	+13,	5
5	0	—11,3	23 28 22,2	—7,8	43,7	+11,3	7 0
	5	—9,6	21,8	—8,2	43,4	+9,6	25
	10	—7,7	21,5	—8,5	43,2	+7,7	20
	15	—5,8	21,3	—8,7	43,0	+5,8	15
	20	—3,9	21,1	—8,9	42,8	+3,9	10
	25	—2,	21,	—9,	42,8	+2,	5
6	0	—0,	23 28 21,	—9,	42,7	+0,	6 0
Lon. D's ⊙	Eq ⁿ . to Points of Eq ^x .	Eq ⁿ . to M. Obl. of Ecl ^c .	True Obli- quity of the Ecliptic.	Eq ⁿ to M. Obl. of Ecl ^c .	Annual Prec. of Eq. P ^{ts} .	Eq ⁿ . to Points of Eq ^x .	Long. D's ⊙
Ex. The Long. D's ⊙ being 4° 10′ 41″ 10″, The Sun and Moon's computed Longitudes (from the mean Ecliptic Obliquity) 10° 23′ 34″ 46″, and 0° 20′ 27″ 20″, respectively, to find the ⊙ and D's correct Places for the true ecliptic Obliquity?							
⊙ comp. Pl.				D comp. Pl.			
10° 23′ 34″ 46″				0° 20′ 27″ 20″			
By the Argument, Equation —17				Equation —17			
Correct Place ⊙ 10 23 34 29 . D . 0 20 27 3 required.							
True Ecliptic Obliquity . . . 23° 28′ 24″ about the same Position of the							
Annual Precession of the Equinox . . . 49 ′ D's ⊙.							

Radical Mean PLACES of the MOON, from 1652 to 1752, Old STYLE.

Julian Style.	M. Pla. D. s o i #	M. Pl. Ap. D. s o i #	M. Pl. Q. s o i #	Julian Style.	M. Pla. D. s o i #	M. Pl. Ap. D. s o i #	M. Pl. Q. s o i #
B. 1652	9 27 24 19	4 24 24 34	0 15 8 49	B. 1700	6 5 58 7	9 27 37 10	5 16 44 13
53	2 6 47 22	6 5 4 24	11 25 49 6	1	10 15 21 10	11 8 17 0	4 27 24 30
54	6 16 10 26	7 15 44 15	11 6 29 23	2	2 24 44 14	0 18 56 51	4 8 4 47
55	10 23 33 30	8 26 24 6	10 17 9 39	3	7 4 7 18	1 29 36 42	3 18 45 3
B. 1656	3 18 7 8	10 7 10 37	9 27 46 40	B. 1704	11 26 40 56	3 10 23 13	2 29 22 10
57	7 27 30 11	11 17 50 27	9 8 27 3	5	4 6 3 59	4 21 3 3	2 10 2 27
58	0 6 53 15	0 28 30 18	8 19 7 20	6	8 15 27 3	6 1 42 54	1 20 42 44
59	4 16 16 19	2 9 10 9	7 29 47 36	7	0 24 50 7	7 12 22 45	1 1 23 0
B. 1660	9 8 49 57	3 19 56 40	7 10 24 43	B. 1708	5 17 23 45	8 23 9 16	0 12 0 7
61	1 18 13 0	5 0 36 30	6 21 5 0	9	9 26 46 48	10 3 49 6	11 22 40 24
62	5 27 36 4	6 11 16 21	6 1 45 17	10	2 6 9 52	11 14 28 57	11 3 20 41
63	10 6 59 8	7 21 56 12	5 12 25 33	11	6 15 32 56	0 25 8 48	10 14 0 57
B. 1664	2 29 32 46	9 2 42 43	4 23 2 40	B. 1712	11 8 6 34	2 5 55 19	9 24 38 4
65	7 8 55 49	10 13 22 33	4 3 42 57	13	3 17 29 37	3 16 35 9	9 5 18 21
66	11 18 18 53	11 24 2 24	3 14 23 14	14	7 26 52 41	4 27 15 0	8 15 58 38
67	3 27 41 57	1 4 42 15	2 25 3 30	15	0 6 15 45	6 7 54 51	7 26 38 54
B. 1668	8 20 15 35	2 15 28 46	2 5 40 37	B. 1716	4 28 49 23	7 18 41 22	7 7 16 1
69	0 29 38 38	3 26 8 36	1 16 20 54	17	9 8 12 26	8 29 21 12	6 17 56 18
70	5 9 1 42	5 6 48 27	0 27 1 11	18	1 17 35 30	10 10 1 3	5 28 36 35
71	9 18 24 46	6 17 28 18	0 7 41 27	19	5 26 58 34	11 20 40 54	5 9 16 51
B. 1672	2 10 58 24	7 28 14 49	11 18 18 34	B. 1720	10 19 32 12	1 1 27 25	4 19 53 58
73	6 20 21 27	9 8 54 39	10 28 58 51	21	2 28 55 15	2 12 7 15	4 0 34 15
74	10 29 44 31	10 19 34 30	10 9 39 8	22	7 8 18 19	3 22 47 6	3 11 14 32
75	3 9 7 35	0 0 14 21	9 20 19 24	23	11 17 41 23	5 3 26 57	2 21 54 48
B. 1676	8 1 41 13	1 11 0 52	9 0 56 31	B. 1724	4 10 15 1	6 14 13 28	2 2 31 55
77	0 11 4 16	2 21 40 42	8 11 36 48	25	8 19 38 14	7 24 53 18	1 13 12 12
78	4 20 27 20	4 2 20 33	7 22 17 5	26	0 29 1 8	9 5 33 9	0 23 52 29
79	8 29 50 24	5 13 0 24	7 2 57 21	27	5 8 24 12	10 16 13 0	0 4 32 45
B. 1680	1 22 24 2	6 23 46 55	6 13 34 28	B. 1728	10 0 57 50	11 26 59 31	11 15 9 52
81	6 1 47 5	8 4 26 45	5 24 14 45	29	2 10 20 53	1 7 39 21	10 25 50 9
82	10 11 10 9	9 15 6 36	5 4 55 2	30	6 19 43 57	2 18 19 12	10 6 30 26
83	2 20 33 13	10 25 46 27	4 15 35 18	31	10 29 7 1	3 28 59 3	9 17 10 42
B. 1684	7 13 6 51	0 6 32 58	3 26 12 25	B. 1732	3 21 40 39	5 9 45 34	8 27 47 49
85	11 22 29 54	1 17 12 48	3 6 52 42	33	8 1 3 42	6 20 25 24	8 8 28 6
86	4 1 52 58	2 27 52 39	2 17 32 59	34	0 10 26 46	8 1 5 15	7 19 8 23
87	8 11 16 2	4 8 32 30	1 28 13 15	35	4 19 49 50	9 11 45 6	6 29 48 39
B. 1688	1 3 49 40	5 19 19 1	1 8 50 22	B. 1736	9 12 23 28	10 22 31 37	6 10 25 46
89	5 13 12 43	6 29 58 51	0 19 30 39	37	1 21 46 31	0 3 11 27	5 21 6 3
90	9 22 35 47	8 10 38 42	0 0 10 56	38	6 1 9 35	1 13 51 18	5 1 46 20
91	2 1 58 51	9 21 18 33	11 10 51 12	39	10 10 32 39	2 24 31 9	4 12 26 36
B. 1692	6 24 32 29	11 2 5 4	10 21 28 19	B. 1740	3 3 6 17	4 5 17 40	3 23 3 43
93	11 3 55 32	0 12 44 54	10 2 8 36	41	7 12 29 20	5 15 57 30	3 3 44 0
94	3 13 18 36	1 23 24 45	9 12 48 53	42	11 21 52 24	6 26 37 21	2 14 24 17
95	7 22 41 40	3 4 4 36	8 23 29 9	43	4 1 15 28	8 7 17 12	1 25 4 33
B. 1696	0 15 15 18	4 14 51 7	8 4 6 16	B. 1744	8 23 49 6	9 18 3 43	1 5 41 40
97	4 24 38 21	5 25 30 57	7 14 46 33	45	1 3 12 9	10 28 43 33	0 16 21 57
98	9 4 1 25	7 6 10 48	6 25 26 50	46	5 12 35 13	0 9 23 24	11 27 2 14
99	1 13 24 29	8 16 50 39	6 6 7 6	47	9 21 58 17	1 20 3 15	11 7 42 30
				B. 1748	2 14 31 55	3 0 49 46	10 18 19 37
				49	6 23 54 58	4 11 29 36	9 28 59 54
				50	11 3 18 2	5 22 9 27	9 9 40 11
				51	3 12 41 6	7 2 49 18	8 20 20 27
				B. 1752	8 5 14 44	8 13 35 49	8 0 57 34

LUNAR TABLES.

Radical

Radical Mean PLACES of the MOON, from 1752 to 1852, New STYLE.

Gregorian Style.	M. Pla. D. s o / #	M. Pl. Ap. D. s o / #	M. Pla. S. s o / #	Gregorian Style.	M. Pla. D. s o / #	M. Pl. Ap. D. s o / #	M. Pla. S. s o / #
B. 1752	3 10 18 19	8 12 22 17	8 1 32 31	1800	11 5 41 32	1 15 28 12	1 3 11 5
53	7 19 41 22	9 23 2 7	7 12 12 48	1	3 15 4 35	2 26 8 2	0 13 51 22
54	11 29 4 26	11 3 41 58	6 22 53 5	2	7 24 27 39	4 6 47 53	11 24 31 39
55	4 8 27 30	0 14 21 49	6 3 33 21	3	0 3 50 43	5 17 27 44	11 5 11 55
B. 1756	9 1 1 8	1 25 8 20	5 14 10 28	B. 1804	4 26 24 21	6 28 14 15	10 15 49 2
57	1 10 24 11	3 5 48 10	4 24 50 45	5	9 5 47 24	8 8 54 5	9 26 29 19
58	5 19 47 15	4 16 28 1	4 5 31 2	6	1 15 10 28	9 19 33 56	9 7 9 36
59	9 29 10 19	5 27 7 52	3 16 11 18	7	5 24 33 32	11 0 13 47	8 17 49 52
B. 1760	2 21 43 57	7 7 54 23	2 26 48 25	B. 1808	10 17 7 10	0 11 0 18	7 28 26 59
61	7 1 7 0	8 18 34 13	2 7 28 42	9	2 26 30 13	1 21 40 8	7 9 7 16
62	11 10 30 4	9 29 14 4	1 18 8 59	10	7 5 53 17	3 2 19 59	6 19 47 33
63	3 19 53 8	11 9 53 55	0 28 49 15	11	11 15 16 21	4 12 59 50	6 0 27 49
B. 1764	8 12 26 46	0 20 40 26	0 9 26 22	B. 1812	4 7 49 59	5 23 46 21	5 11 4 56
65	0 21 49 49	2 1 20 16	11 20 6 39	13	8 17 13 2	7 4 26 11	4 21 45 13
66	5 1 12 53	3 12 0 7	11 0 46 56	14	0 26 36 6	8 15 6 2	4 2 25 30
67	9 10 35 57	4 22 39 58	10 11 27 12	15	5 5 59 10	9 25 45 53	3 13 5 46
B. 1768	2 3 9 35	6 3 26 29	9 22 4 19	B. 1816	9 28 32 48	11 6 32 24	2 23 42 53
69	6 12 32 38	7 14 6 19	9 2 44 36	17	2 7 55 51	0 17 12 14	2 4 23 10
70	10 21 55 42	8 24 46 10	5 13 24 53	18	6 17 18 55	1 27 52 5	1 15 3 27
71	3 1 18 46	10 5 26 1	7 24 5 9	19	10 26 41 59	3 8 31 56	0 25 43 43
B. 1772	7 23 52 24	11 16 12 32	7 4 42 16	B. 1820	3 19 15 37	4 19 18 27	0 6 20 50
73	0 3 15 27	0 26 52 22	6 15 22 33	21	7 28 38 40	5 29 58 17	11 17 1 7
74	4 12 38 31	2 7 32 13	5 26 2 50	22	0 8 1 44	7 10 38 8	10 27 41 24
75	8 22 1 35	3 18 12 4	5 6 43 6	23	4 17 24 48	8 21 17 59	10 8 21 40
B. 1776	1 14 35 13	4 28 58 35	4 17 20 13	B. 1824	9 9 58 26	10 2 4 30	9 18 58 47
77	5 23 58 16	6 9 38 25	3 28 0 30	25	1 19 21 29	11 12 44 20	8 29 39 4
78	10 3 21 20	7 20 18 16	3 8 40 47	26	5 28 44 33	0 23 24 11	8 10 19 21
79	2 12 44 24	9 0 58 7	2 19 21 3	27	10 8 7 37	2 4 4 2	7 20 59 37
B. 1780	7 5 18 2	10 11 44 38	1 29 58 10	B. 1828	3 0 41 15	3 14 50 33	7 1 36 44
81	11 14 41 5	11 22 24 28	1 10 38 27	29	7 10 4 18	4 25 30 23	6 12 17 1
82	3 24 4 9	1 3 4 19	0 21 18 44	30	11 19 27 22	6 6 10 14	5 22 57 18
83	8 3 27 13	2 13 44 10	0 1 59 0	31	3 28 50 26	7 16 50 5	5 3 37 34
B. 1784	0 26 0 51	3 24 30 41	11 12 36 7	B. 1832	8 21 24 4	8 27 36 36	4 14 14 41
85	5 5 23 54	5 5 10 31	10 23 16 24	33	1 0 47 7	10 8 16 26	3 24 54 58
86	9 14 46 58	6 15 50 22	10 3 56 41	34	5 10 10 11	11 18 56 17	3 5 35 15
87	1 24 10 2	7 26 30 13	9 14 36 57	35	9 19 33 15	0 29 36 8	2 16 15 31
B. 1788	6 16 43 40	9 7 16 44	8 25 14 4	B. 1836	2 12 6 53	2 10 22 39	1 26 52 38
89	10 26 6 43	10 17 56 34	8 5 54 21	37	6 21 29 56	3 21 2 29	1 7 32 55
90	3 5 29 47	11 28 36 25	7 16 34 38	38	11 0 53 0	5 1 42 20	0 18 13 12
91	7 14 52 51	1 9 16 16	6 27 14 54	39	3 10 16 4	6 12 22 11	11 28 53 28
B. 1792	0 7 26 29	2 20 2 47	6 7 52 1	B. 1840	8 2 49 42	7 23 8 42	11 9 30 35
93	4 16 49 32	4 0 42 37	5 18 32 18	41	0 12 12 45	9 3 48 32	10 20 10 52
94	8 26 12 36	5 11 22 28	4 29 12 35	42	4 21 35 49	10 14 28 23	10 0 51 9
95	1 5 35 40	6 22 2 19	4 9 52 51	43	9 0 58 53	11 25 8 14	9 11 31 25
B. 1796	5 28 9 18	8 2 48 50	3 20 29 58	B. 1844	1 23 32 31	1 5 54 45	8 22 8 32
97	10 7 32 21	9 13 28 40	3 1 10 15	45	6 2 55 34	2 16 34 35	8 2 48 49
98	2 16 55 25	10 24 8 31	2 11 50 32	46	10 12 18 38	3 27 14 26	7 13 29 6
99	6 26 18 29	0 4 48 22	1 22 30 48	47	2 21 41 42	5 7 54 17	6 24 9 22
				B. 1848	7 14 15 20	6 18 40 48	6 4 46 29
				49	11 23 38 23	7 29 20 38	5 15 26 46
				50	4 3 1 27	9 10 0 29	4 26 7 3
				51	8 12 24 31	10 20 40 20	4 6 47 19
				B. 1852	1 4 58 9	0 1 26 51	3 17 24 26

For Advantage of taking out the Numbers after January and February, Bissextile Years Places are advanced a Day's Motion, in both Styles, correspondent to the mean Noon of January the 1st.

Mean MOTIONS of the MOON, for MONTHS and DAYS.

Days.	JANUARY.									FEBRUARY.									MARCH.											
	M. Mot. D			Ap.			Q retr.			M. Mot. D			Ap.			Q retr.			M. Mot. D			Ap.			Q retr.					
	s	o	l	l	o	l	l	o	l	l	s	o	l	l	o	l	l	s	o	l	l	o	l	l	s	o	l	l		
1	0	13	10	35	0	6	41	0	3	11	2	1	38	41	3	33	54	1	41	41	2	10	35	2	6	41	4	3	10	38
2	0	26	21	10	0	13	22	0	6	21	2	14	49	16	3	40	35	1	44	52	2	23	45	36	6	47	45	3	13	49
3	1	9	31	45	0	20	3	0	9	32	2	27	59	51	3	47	16	1	48	2	3	6	56	12	6	54	26	3	17	0
4	1	22	42	20	0	26	44	0	12	43	3	11	10	26	3	53	57	1	51	13	3	20	6	46	7	1	7	3	20	11
5	2	5	52	55	0	33	25	0	15	53	3	24	21	1	4	0	38	1	54	24	4	3	17	21	7	7	48	3	23	21
6	2	19	3	30	0	40	6	0	19	4	4	7	31	36	4	7	19	1	57	34	4	16	27	57	7	14	30	3	26	32
7	3	2	14	5	0	46	47	0	22	14	4	20	42	11	4	14	0	2	0	44	4	29	38	32	7	21	11	3	29	43
8	3	15	24	40	0	53	29	0	25	25	5	3	52	46	4	20	42	2	3	55	5	12	49	7	7	27	51	3	32	53
9	3	28	35	15	1	0	10	0	28	36	5	17	3	21	4	27	23	2	7	6	5	25	59	42	7	34	32	3	36	4
10	4	11	45	50	1	6	51	0	31	46	6	0	13	56	3	34	4	2	10	16	6	9	10	17	7	41	13	3	39	14
11	4	24	56	25	1	13	32	0	34	57	6	13	24	31	4	40	45	2	13	27	6	22	20	52	7	47	54	3	42	25
12	5	8	7	0	1	20	13	0	38	8	6	26	35	6	4	47	26	2	16	38	7	5	31	27	7	54	35	3	45	36
13	5	21	17	35	1	26	54	0	41	18	7	9	45	41	4	54	7	2	19	49	7	18	42	2	8	1	17	0	48	46
14	6	4	28	10	1	33	35	0	44	29	7	22	56	16	5	0	48	2	22	59	8	1	52	37	8	7	58	3	51	56
15	6	17	38	45	1	40	16	0	47	40	8	6	6	51	5	7	29	2	26	10	8	15	3	12	8	14	39	3	55	8
16	7	0	49	20	1	46	57	0	50	50	8	19	17	26	5	14	10	2	29	20	8	28	13	47	8	21	20	3	58	17
17	7	13	59	55	1	53	38	0	54	1	9	2	28	1	5	20	51	2	32	31	9	11	24	22	8	28	1	4	1	29
18	7	27	10	30	2	0	19	0	57	11	9	15	38	36	5	27	32	2	35	41	9	24	34	57	8	34	42	4	4	39
19	8	10	21	5	2	7	0	1	0	22	9	28	49	11	5	34	13	2	38	52	10	7	45	32	8	41	23	4	7	49
20	8	23	31	41	2	13	41	1	3	33	10	11	59	46	5	40	55	2	42	2	10	20	56	7	8	48	4	4	11	0
21	9	6	42	15	2	20	22	1	6	43	10	25	10	21	5	47	36	2	45	13	11	4	6	42	8	54	40	4	14	11
22	9	19	52	51	2	27	4	1	9	54	11	8	20	56	5	54	17	2	48	24	11	17	17	17	9	1	28	4	17	21
23	10	3	3	26	2	33	45	1	13	5	11	21	31	31	6	0	58	2	51	35	0	0	27	52	9	8	8	4	20	32
24	10	16	14	1	2	40	26	1	16	15	0	4	42	6	6	7	39	2	54	45	0	13	38	27	9	14	49	4	23	43
25	10	29	24	36	2	47	7	1	19	26	0	17	52	41	6	14	20	2	57	56	0	26	49	2	9	21	30	4	26	54
26	11	12	35	11	2	53	48	1	22	37	1	1	3	17	6	21	1	3	1	7	1	9	59	37	9	28	11	4	30	5
27	11	25	45	46	3	0	29	1	25	47	1	14	13	52	6	27	42	3	4	17	1	23	10	12	9	34	52	4	33	15
28	0	8	56	21	3	7	10	1	28	58	1	27	24	26	6	34	23	3	7	28	2	6	20	47	9	41	33	4	36	26
29	0	22	6	56	3	13	51	1	32	9	2	19	31	22	9	48	14	4	39	37	2	19	31	22	9	48	14	4	39	37
30	1	5	17	31	3	20	32	1	35	19	3	2	41	57	9	54	55	4	42	47	3	2	41	57	9	54	55	4	42	47
31	1	18	28	6	3	27	13	1	38	30	In Leap Year take out for a Day sooner in January and February.									3	15	52	32	10	1	36	4	45	57	

Mean MOTIONS of the MOON, for Months and Days, to Thirds of a DEGREE, for the correct Construction of the above Monthly Tables.

Days.	Months.	M. Mot. D.					M. Mot. Ap. D.					M. Mot. Q D. r.					Days.	M. Mot. D.					M. Mot. Ap. D.					M. Mot. Q D. r.				
		s	o	l	l	l	s	o	l	l	l	s	o	l	l	l		s	o	l	l	l	s	o	l	l	l	s	o	l	l	l
31	Jan.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	13	10	35	1	0	0	6	41	4	0	0	3	10	38
59	Feb.	0	1	18	28	5	43	0	3	27	13	9	0	1	38	29	2	0	26	21	10	3	0	0	13	22	8	0	0	6	21	16
90	Mar.	0	1	27	24	26	22	0	6	34	23	7	0	3	7	27	3	1	9	31	45	4	0	0	20	3	12	0	0	9	31	55
	Apr.	0	1	15	52	32	5	0	10	1	36	17	0	4	45	57	4	1	22	42	20	5	0	0	26	44	16	0	0	12	42	33
120	May	0	4	21	10	2	47	0	13	22	8	22	0	6	21	16	5	2	5	52	55	7	0	0	33	25	21	0	0	15	53	11
151	June	0	6	9	38	8	30	0	16	49	21	32	0	7	59	46	6	2	19	3	30	8	0	0	40	6	25	0	0	19	3	50
181	July	0	7	14	55	39	12	0	20	9	53	38	0	9	35	5	7	3	2	14	5	10	0	0	46	47	29	0	0	22	14	28
212	Aug.	0	9	3	23	44	55	0	23	37	6	47	0	11	13	35	8	1	15	24	40	11	0	0	51	28	33	0	0	25	25	6
243	Sept.	0	10	21	51	50	42	0	27	4	19	57	0	12	52	5	9	3	28	35	15	12	0	1	0	9	38	0	0	28	35	45
273	Octo.	0	11	27	9	21	20	1	0	24	52	3	0	14	27	24	10	4	11	45	50	14	0	1	6	50	42	0	0	31	46	23
304	Nov.	0	1	15	37	27	4	1	3	52	5	12	0	16	5	54	11	4	24	56	25	15	0	1	13	31	46	0	0	34	57	1
334	Dec.	0	2	20	54	57	46	1	7	12	37	18	0	17	41	13	12	5	8	7	0	17	0	1	20	12	50	0	0	38	7	40

Mean

Mean MOTIONS of the MOON, for MONTHS and DAYS.

Days.	A P R I L.									M A Y.									J U N E.												
	M. Mot. D.			Apog.			Q retr.			M. Mot. D.			Apog.			Q retr.			M. Mot. D.			Apog.			Q retr.						
	s	o	i	o	i	o	s	o	i	o	i	o	s	o	i	o	i	o	s	o	i	o	i	o	s	o	i	o	i	o	
1	3	29	3	7	10	8	17	4	49	8	5	4	20	37	13	28	49	6	24	27	6	22	48	43	16	50	3	8	2	58	
2	4	12	13	42	10	14	58	4	52	18	5	17	31	12	13	35	30	6	27	38	7	5	59	18	17	2	44	8	6	8	
3	4	25	24	17	10	21	39	4	55	30	6	0	41	48	13	42	12	6	30	49	7	19	9	53	17	9	24	8	9	19	
4	5	8	34	52	10	28	21	4	58	40	6	13	52	23	13	48	52	6	34	0	8	2	20	28	17	16	5	8	12	29	
5	5	21	45	28	10	35	1	5	1	50	6	27	2	58	13	55	33	6	37	10	8	15	31	3	17	22	46	8	15	40	
6	6	4	56	3	10	41	42	5	5	1	7	10	13	33	14	2	15	6	40	20	8	28	41	38	17	29	27	8	18	51	
7	6	18	6	38	10	48	24	5	8	12	7	23	24	8	14	8	56	6	43	31	9	11	52	13	17	36	9	8	22	1	
8	7	1	17	13	10	55	5	5	11	22	8	6	34	43	14	15	37	6	46	42	9	25	2	49	17	42	50	8	25	12	
9	7	14	27	47	11	1	46	5	14	33	8	19	45	18	14	22	18	6	49	52	10	8	13	24	17	49	31	8	28	23	
10	7	27	38	23	11	8	27	5	17	44	9	2	55	53	14	28	59	6	53	3	10	21	23	59	17	56	12	8	31	32	
11	8	10	48	58	11	15	8	5	20	54	9	16	6	28	14	35	40	6	56	14	11	4	34	34	18	2	53	8	34	44	
12	8	23	59	32	11	21	49	5	24	6	9	29	17	3	14	42	21	6	59	25	11	17	45	9	18	9	34	8	37	55	
13	9	7	10	8	11	28	30	5	27	16	10	12	27	38	14	49	2	7	2	35	0	0	55	44	18	16	15	8	41	5	
14	9	20	20	43	11	35	11	5	30	27	10	25	38	13	14	55	43	7	5	46	0	14	6	19	18	22	56	8	44	16	
15	10	3	31	17	11	41	52	5	33	37	11	8	48	48	15	2	24	7	8	56	0	27	16	54	18	29	18	8	47	27	
16	10	16	41	52	11	48	33	5	36	48	11	21	59	23	15	9	5	7	12	7	1	10	27	29	18	36	18	8	50	37	
17	10	29	52	28	11	55	14	5	39	59	0	5	9	58	15	15	46	7	15	18	1	23	38	4	18	42	59	8	53	48	
18	11	13	3	2	12	1	55	5	43	9	0	18	20	33	15	22	28	7	18	28	2	6	48	39	18	49	41	8	56	58	
19	11	26	13	38	12	8	36	5	46	20	1	1	31	8	15	29	9	7	21	39	2	19	59	14	18	56	21	9	0	9	
20	0	9	24	13	12	15	17	5	49	31	1	14	41	43	15	35	50	7	24	50	3	3	9	49	19	3	2	9	3	20	
21	0	22	34	47	12	21	59	5	52	41	1	27	52	18	15	42	31	7	28	0	3	16	20	25	19	9	44	9	0	30	
22	1	5	45	23	12	28	40	5	55	53	2	11	2	54	15	49	12	7	31	11	3	29	31	0	19	16	25	9	9	41	
23	1	18	55	57	12	35	21	5	59	3	2	24	13	28	15	55	53	7	34	22	4	12	41	35	19	23	6	9	12	61	
24	2	2	6	33	12	42	2	6	2	14	3	7	24	3	16	2	34	7	37	32	4	25	52	9	19	29	47	9	16	2	
25	2	15	17	8	12	48	43	6	5	24	3	20	34	38	16	9	15	7	40	43	5	9	2	44	19	36	28	9	19	13	
26	2	28	27	43	12	55	24	6	8	35	4	3	45	13	16	15	50	7	43	54	5	22	13	19	19	43	9	9	22	23	
27	3	11	38	17	13	2	5	6	11	45	4	16	55	48	16	22	37	7	47	4	6	5	23	54	19	49	50	9	25	34	
28	3	24	48	52	13	8	46	6	14	56	5	0	6	23	16	29	16	7	50	15	6	18	34	29	19	56	31	9	28	45	
29	4	7	59	28	13	15	27	6	18	6	5	13	16	58	16	35	59	7	53	25	7	1	45	4	20	3	12	9	31	55	
30	4	21	10	3	13	22	8	6	21	17	5	26	27	33	16	42	40	7	56	36	7	14	55	39	20	9	54	9	35	6	
31											6	9	38	8	16	49	22	7	59	46											

Mean MOTIONS of the Moon for Days, to Thirds of a DEGREE, for the correct Construction of the above Monthly Tables, continued.

Days	M. Mot. D.					M. Mot. Ap. D.					M. Mot. Q. D. r.					Days	M. Mot. D.					M. Mot. Ap. D.					M. Mot. Q. D. r.				
	s	o	i	o	o	s	o	i	o	o	s	o	i	o	o		s	o	i	o	o	s	o	i	o	o	s	o	i	o	o
13	5	21	17	35	18	0	1	26	53	54	0	0	41	18	18	23	10	3	3	25	32	0	2	33	44	36	0	1	13	4	41
14	6	4	28	10	19	0	1	33	34	58	0	0	44	28	56	24	10	16	14	0	33	0	2	40	25	40	0	1	16	15	19
15	6	17	38	45	21	0	1	40	16	3	0	0	47	39	35	25	10	29	24	35	35	0	2	47	6	44	0	1	19	25	58
16	7	0	49	20	22	0	1	46	57	7	0	0	50	50	13	26	11	12	35	10	36	0	2	53	47	49	0	1	22	36	36
17	7	13	59	55	24	0	1	53	38	11	0	0	54	0	51	27	11	25	45	45	38	0	3	0	28	53	0	1	25	47	14
18	7	27	10	30	25	0	2	0	19	15	0	0	57	11	29	28	0	8	56	20	39	0	3	7	9	57	0	1	28	57	53
19	8	10	21	5	26	0	2	7	0	19	0	1	0	22	8	29	0	22	6	5	40	0	3	13	51	1	0	1	32	8	31
20	8	23	31	40	28	0	2	13	41	24	0	1	3	32	46	30	1	5	17	30	42	0	3	20	32	5	0	1	35	19	9
21	9	6	42	15	29	0	2	20	22	28	0	1	6	43	24	31	1	18	28	5	43	0	3	27	13	9	0	1	38	29	48
22	9	19	52	50	30	0	2	27	3	32	0	1	9	54	3																

EXAMPLE of the correct Construction of the above Monthly Tables.

Nearly the same, as by the uppermost Table.

August	9	3	23	44	55	0	23	37	6	47	0	11	13	35	23
Days 20	8	23	31	40	28	0	2	13	41	24	0	1	3	32	46
August 20	5	26	55	25	23	0	25	50	48	11	0	12	17	8	9

Mean

Mean MOTIONS of the MOON, for MONTHS and DAYS.

Days.	JULY.									AUGUST.									SEPTEMBER.																	
	Mean M. D.				Ap.				Q retr.				M. Mot. D.				Ap.				Q retr.				M. Mot. D.				Ap.				Q retr.			
	s	o	i	h	s	o	i	h	s	o	i	h	s	o	i	h	s	o	i	h	s	o	i	h	s	o	i	h	s	o	i	h				
1	7	28	6	14	20	16	34		9	38	17		9	16	34	20	23	43	48	11	16	47		11	5	2	26	27	11	1	12	55	17			
2	8	11	16	49	20	23	15		9	41	27		9	29	44	55	23	50	29	11	19	57		11	18	13	1	27	17	41	12	58	27			
3	8	24	27	24	20	29	56		9	44	38		10	12	55	30	23	57	10	11	23	8		0	1	23	36	27	24	23	13	1	38			
4	9	7	37	59	20	36	37		9	47	49		10	26	6	5	24	3	50	11	26	18		0	14	34	11	27	31	3	13	4	49			
5	9	20	48	34	20	43	18		9	50	59		11	9	16	40	24	10	31	11	29	29		0	27	44	46	27	37	45	13	7	59			
6	10	3	59	9	20	49	59		9	54	10		11	22	27	15	24	17	12	11	32	40		1	10	55	21	27	44	26	13	11	10			
7	10	17	9	44	20	56	41		9	57	20		0	5	37	50	24	23	53	11	35	50		1	24	5	56	27	51	7	13	14	20			
8	11	0	20	19	21	3	22		10	0	31		0	18	48	25	24	30	35	11	39	1		2	7	16	31	27	57	48	13	17	31			
9	11	13	30	54	21	10	3		10	3	42		1	1	59	0	24	37	16	11	42	12		2	20	27	6	28	4	29	13	20	40			
10	11	26	41	30	21	16	44		10	6	52		1	15	9	35	24	43	57	11	45	22		3	3	37	41	28	11	10	13	23	52			
11	0	9	52	5	21	23	25		10	10	3		1	28	20	10	24	50	38	11	48	33		3	16	48	16	28	17	51	13	27	3			
12	0	23	2	40	21	30	6		10	13	14		2	11	30	45	24	57	19	11	51	44		3	29	58	51	28	24	32	13	30	14			
13	1	6	13	15	21	36	47		10	16	24		2	24	41	20	25	4	0	11	54	54		4	13	9	26	28	31	13	13	33	24			
14	1	19	23	50	21	43	29		10	19	35		3	7	51	55	25	10	41	11	58	5		4	26	20	1	28	37	54	13	36	35			
15	2	2	34	25	21	50	9		10	22	46		3	21	2	30	25	17	22	12	1	16		5	9	30	36	28	44	35	13	39	46			
16	2	15	45	0	21	56	50		10	25	56		4	4	13	5	25	24	3	12	4	26		5	22	41	11	28	51	16	13	42	56			
17	2	28	55	35	22	3	31		10	29	7		4	17	23	40	25	30	44	12	7	37		6	5	51	46	28	57	57	13	46	7			
18	3	12	6	10	22	10	13		10	32	17		5	0	34	15	25	37	25	12	10	47		6	19	2	21	29	4	38	13	49	17			
19	3	25	16	45	22	16	53		10	35	28		5	13	44	50	25	44	6	12	13	58		7	2	12	57	29	11	19	13	52	28			
20	4	8	27	20	22	23	35		10	38	39		5	26	55	25	25	50	47	12	17	9		7	15	23	31	29	18	0	13	55	39			
21	4	21	37	55	22	30	16		10	41	49		6	10	6	0	25	57	29	12	20	19		7	28	34	7	29	24	42	13	58	49			
22	5	4	48	30	22	36	57		10	45	0		6	23	16	35	26	4	10	12	23	30		8	11	44	41	29	31	23	14	2	0			
23	5	17	59	5	22	43	38		10	48	11		7	6	27	10	26	10	51	12	26	40		8	24	55	16	29	38	4	14	5	11			
24	6	1	9	40	22	50	19		10	51	21		7	19	37	45	26	17	32	12	29	51		9	8	5	51	29	44	45	14	8	21			
25	6	14	20	15	22	57	0		10	54	32		8	2	48	20	26	24	13	12	33	2		9	21	16	26	29	51	26	14	11	32			
26	6	27	30	50	23	3	41		10	57	42		8	15	58	55	26	30	54	12	36	13		10	17	27	1	29	58	7	14	14	42			
27	7	10	41	25	23	10	22		11	0	53		8	29	9	30	26	37	35	12	39	23		10	17	37	36	30	4	48	14	17	53			
28	7	23	52	0	23	17	4		11	4	4		9	12	20	5	26	44	16	12	42	34		11	0	48	11	30	11	29	14	21	4			
29	8	7	2	35	23	23	44		11	7	14		9	25	30	40	26	50	57	12	45	43		11	13	58	46	30	18	10	14	24	14			
30	8	20	13	10	23	30	26		11	10	25		10	8	41	15	26	57	38	12	48	55		11	27	9	21	30	24	52	14	27	24			
31	9	3	23	45	23	37	7		11	13	35		10	21	51	51	27	4	20	12	52	5		From Sept. 27, the Motion D's Apogee is expressed in Degrees for Sig. and Deg.												

EXAMPLE I.

To find the Mean Place of the D, Apogee and Node, for Octob. 12th, at Noon, 1772 N. S.

	D				Ap.				Q			
	s	o	i	h	s	o	i	h	s	o	i	h
N. S. Year 1772.	7	23	52	24	11	16	12	32	7	4	42	16
Mot. Octob. 12.	5	5	16	21	1	1	45	4	15	5	33	—
N. S. Octob. 12, 1772.	0	29	8	45	0	17	57	36	6	19	36	43

required.

EXAMPLE II.

To find the Mean Place of the D, Apogee and Node for Octob. 1st, at Noon, 1772 O. S.

O. S. Year 1700.	6	5	58	7	9	27	37	10	5	16	44	13
Mot. Years 72.	6	12	50	42	1	19	48	54	10	12	36	54
Mot. Oct. 1.	0	10	19	56	0	30	31	32	14	30	36	—
									10	27	7	30
O. S. Octob. 1, 1772.	0	29	8	45	0	17	57	36	6	19	36	43
												required.

Mean MOTIONS of the MOON for MONTHS and DAYS.

Days.	OCTOBER.			NOVEMBER.			DECEMBER.		
	M. Mot. D.	Ap.	Q retr.	M. Mot. D.	Ap.	Q retr.	M. Mot. D.	Ap.	Q retr.
1	0 10 19 56	30 31 32	14 30 36	1 28 48 1	33 58 46	16 9 6	3 4 5 32	37 19 18	17 44 25
2	0 23 30 31	30 38 13	14 33 46	2 11 58 36	34 5 27	16 12 16	3 17 16 7	37 25 59	17 47 35
3	1 6 41 6	30 44 54	14 36 57	2 25 9 12	34 12 8	16 15 27	4 0 26 42	37 32 40	17 50 46
4	1 19 51 41	30 51 35	14 40 8	3 8 19 47	34 18 49	16 18 38	4 13 37 17	37 39 21	17 53 57
5	2 3 2 16	30 58 16	14 43 18	3 21 30 22	34 25 30	16 21 48	4 26 47 52	37 46 2	17 57 7
6	2 16 12 51	31 4 57	14 46 29	4 4 40 57	34 32 11	16 24 59	5 9 38 27	37 52 43	18 0 18
7	2 29 23 26	31 11 39	14 49 39	4 17 51 32	34 38 53	16 28 9	5 23 9 2	37 59 25	18 3 28
8	3 12 34 1	31 18 20	14 52 50	5 1 2 7	34 45 34	16 31 20	6 6 19 37	38 6 6	18 6 39
9	3 25 44 36	31 25 1	14 56 1	5 14 12 42	34 52 15	16 34 31	6 19 30 12	38 12 47	18 9 50
10	4 8 55 11	31 31 42	14 59 11	5 27 23 17	34 58 56	16 37 41	7 2 40 47	38 19 28	18 13 0
11	4 22 5 46	31 38 23	15 2 22	6 10 33 52	35 5 37	16 40 52	7 15 51 22	38 26 9	18 16 11
12	5 5 16 21	31 45 4	15 5 33	6 23 44 27	35 12 18	16 44 3	7 29 1 57	38 32 50	18 19 22
13	5 18 26 56	31 51 45	15 8 43	7 6 55 2	35 18 59	16 47 13	8 12 12 32	38 39 31	18 22 32
14	6 1 37 31	31 58 27	15 11 54	7 20 5 37	35 25 40	16 50 24	8 25 23 7	38 46 12	18 25 43
15	6 14 48 6	32 5 8	15 15 5	8 3 16 12	35 32 21	16 53 35	9 8 33 42	38 52 53	18 28 54
16	6 27 58 41	32 11 49	15 18 15	8 16 26 47	35 39 2	16 56 45	9 21 44 17	38 59 34	18 32 4
17	7 11 9 16	32 18 30	15 21 26	8 29 37 22	35 45 43	16 59 56	10 4 54 52	39 6 15	18 35 15
18	7 24 19 51	32 25 11	15 24 36	9 12 47 57	35 52 24	17 3 6	10 18 5 28	39 12 56	18 38 25
19	8 7 30 26	32 31 52	15 27 47	9 25 58 32	35 59 5	17 6 17	11 1 16 3	39 19 37	18 41 36
20	8 20 41 1	32 38 33	15 30 58	10 9 9 7	36 5 46	17 9 28	11 14 26 38	39 26 18	18 44 47
21	9 3 51 36	32 45 14	15 34 8	10 22 19 42	36 12 28	17 12 38	11 27 37 13	39 33 0	18 47 57
22	9 17 2 11	32 51 55	15 37 19	11 5 30 17	36 19 9	17 15 49	0 10 47 48	39 39 41	18 51 8
23	10 0 12 46	32 58 36	15 40 29	11 18 40 52	36 25 50	17 19 0	0 23 58 23	39 46 22	18 54 19
24	10 13 23 21	33 5 17	15 43 40	0 1 51 27	36 32 31	17 22 10	1 7 8 58	39 53 3	18 57 29
25	10 26 33 56	33 11 58	15 46 51	0 15 2 2	36 39 12	17 25 21	1 20 19 33	39 59 44	19 0 40
26	11 9 44 31	33 18 39	15 50 1	0 28 12 37	36 45 53	17 28 32	2 3 30 8	40 6 25	19 3 51
27	11 22 55 6	33 25 20	15 53 12	1 11 23 12	36 52 34	17 31 42	2 16 40 43	40 13 6	19 7 1
28	0 6 5 41	33 32 2	15 56 22	1 24 33 47	36 59 15	17 34 53	2 29 51 18	40 19 47	19 10 12
29	0 19 16 16	33 38 43	15 59 33	2 7 44 22	37 5 56	17 38 3	3 13 1 53	40 26 28	19 13 22
30	1 2 26 51	33 45 24	16 2 44	2 20 54 58	37 12 37	17 41 13	3 26 12 28	40 33 9	19 16 33
31	1 15 37 27	33 52 5	16 5 54				4 9 23 3	40 39 50	19 19 43

Motion of Moon's Apogee, in this Page, for Signs and Degrees is expressed in Degrees.

RADICAL MEAN PLACES and MOTIONS of the MOON, from several AUTHORITIES.

M. Pl. D. Julian Style. Jan. O. 1701.	M. Pl. Ap. D. Julian Style. Jan. O. 1701.	M. Pl. Q D. Julian Style. Jan. O. 1701.	M. Pl. D. a Julian Style. Jan. O. 1701.	M. Mot. D in 100 Julian Years.	M. Mot. Ap. D in 100 Jul. Years.	M. Mot. Q D ret. in 100 Jul. Years. r.	M. Mot. D a C in 100 Jul. Years.	According to
10 15 21 0 GREENWICH	11 8 20 0 OBSERVATORY.	4 27 24 20	0 24 37 20	10 7 50 25 1336 Revol.	3 19 11 15 11 Revol.	4 14 11 15 5 Revol.	10 7 5 5 1236 Revol.	Sir Isaac Newton's last Edition <i>Principia</i> .
10 15 20 0	11 8 20 0	4 27 23 20	0 24 36 27	10 7 50 25 Same . .	3 19 11 15 GREENW.	4 14 11 15 OBSERV.	10 7 4 52	Dr. Ed. Halley's Tables. Gael Morris, Ast. Ast. R.
10 15 21 10	11 8 15 0	4 27 24 30	0 24 37 17	10 7 52 20	3 19 11 15 PARIS	4 14 11 15 OBSERV.	10 7 6 33	Tobias Mayer, of Gottingen, Ger- many, from his Tables.
10 15 15 42	11 8 17 52	4 27 28 32	0 24 32 7	10 7 50 25	3 19 11 15	4 14 11 15	10 7 4 25	Leonard Euler, of Berlin, Prussia, F. R. S. from his Tables.
10 15 21 30 LON- DON.	11 8 6 30	4 27 26 20	0 24 35 59	10 7 50 25	3 19 11 15	4 14 11 15	10 7 5 20	Thomas Street's Caroline Tables.
10 15 16 30	11 8 5 0	4 27 18 30	0 24 36 9	10 7 48 45	3 19 10 0	4 14 12 30	10 7 5 20	
10 15 21 10 GREENWICH	11 8 17 0	4 27 24 30	0 24 37 27	10 7 50 25	3 19 11 15	4 14 11 15	10 7 4 53	Our own Tables. Above.

Hence, From a Comparison of These, with other Observations, we give the following correct

RADICAL MEAN PLACES and MOTIONS of the MOON, for Julian and Gregorian Years.

Julian Style.	M. Places D.	M. Pl. Ap. D.	M. Pl. Q.	Greg. Style.	M. Places D.	M. Pl. Ap. D.	M. Pl. Q.
Years before & since Christ.	s o / //	s o / //	s o / //	Yrs of Christ.	s o / //	s o / //	s o / //
E. 1000	5 4 16 52	7 19 33 25	6 9 47 58	B. 1600	3 16 21 52	6 7 19 4	10 4 27 14
B. 900	3 12 7 17	11 8 44 40	1 25 36 43	1700	1 11 1 42	9 26 23 38	5 17 19 10
B. 800	1 19 57 42	2 27 55 55	9 11 25 28	1800	11 5 41 32	1 15 28 12	1 3 11 5
B. 700	11 27 48 7	6 17 7 10	4 27 14 13	1900	9 0 21 22	5 4 32 46	8 19 3 1
B. 600	10 5 38 32	10 6 18 25	0 13 2 58	B. 2000	7 8 11 47	8 23 44 1	4 4 51 46
B. 500	8 13 28 57	1 25 29 40	7 28 51 43	2100	5 2 51 37	0 12 48 35	11 20 43 42
B. 400	6 21 19 22	5 14 40 55	3 14 40 28	2200	2 27 31 27	4 1 53 9	7 6 35 37
B. 300	4 29 9 47	9 3 52 10	11 0 29 13	2300	0 22 11 17	7 20 57 43	2 22 27 33
B. 200	3 7 0 12	0 23 3 25	6 16 17 58	B. 2400	11 0 1 42	11 10 8 58	10 8 16 18
B. Jul. Per. 100	1 14 50 37	4 12 14 40	2 2 6 43	2500	8 24 41 32	2 29 13 32	5 24 8 14
B. 4713. B.C. 0	11 22 41 2	8 1 25 55	9 17 55 28	2600	6 19 21 22	6 18 18 6	1 10 0 9
B. 100	10 0 31 27	11 20 37 10	5 3 44 13	2700	4 14 1 12	10 7 22 40	8 25 52 5
B. 200	8 8 21 52	3 9 48 25	0 19 32 58	B. 2800	2 21 51 37	1 26 33 55	4 11 40 50
B. 300	6 16 12 17	6 28 59 40	8 5 21 43	2900	0 16 31 27	5 15 38 29	11 27 32 46
B. 400	4 24 2 42	10 18 10 55	3 21 10 28	3000	10 11 11 17	9 4 43 3	7 13 24 41
B. 500	3 1 53 7	2 7 22 10	11 6 59 13	3100	8 5 51 7	0 23 47 37	2 29 16 37
B. 600	1 9 43 32	5 26 33 25	6 22 47 58	B. 3200	6 13 41 32	4 12 58 52	10 15 5 22
B. 700	11 17 33 57	9 15 44 40	2 8 36 43	3300	4 8 21 22	8 2 3 26	6 0 57 18
B. 800	9 25 24 22	1 4 55 55	9 24 25 28	3400	2 3 1 12	11 21 8 0	1 16 49 13
B. 900	8 3 14 47	4 24 7 10	5 10 14 13	Biflexile Years Places are advanced a Day's Motion.			
B. 1000	6 11 5 12	8 13 18 25	0 26 2 58	From Hundreds of Years <i>Gregorian</i> deduct their <i>Fourth</i> (rejecting Fractions) and the <i>Remainder</i> will be equal to the N ^o . of Days there are <i>left</i> in any N ^o . of <i>Gregorian</i> Years than are in the same N ^o . of Years <i>Julian</i> , since 1600. Whence the Diff. of Motion between any N ^o . of <i>Julian</i> and <i>Gregorian</i> Years is determined.			
B. 1100	4 18 55 37	0 2 29 40	8 11 51 43	EXAMPLES.			
B. 1200	2 26 46 2	3 21 40 55	3 27 40 28	In the Year 2735. In the Year 2215			
B. 1300	1 4 36 27	7 10 52 10	11 13 29 13	—1600 —1600			
B. 1400	11 12 26 52	11 0 3 25	6 29 17 58	$\frac{1}{4}$ 11/35 $\frac{1}{4}$ 6/15			
B. 1500	9 20 17 17	2 19 14 40	2 15 6 43	—2 —1			
B. 1600	7 28 7 42	6 8 25 55	10 0 55 28	9 Days left 5 Days left in 6 Hundred			
B. 1700	6 5 58 7	9 27 37 10	5 16 44 13	[in 11 Hund. Years <i>Greg.</i> [Years <i>Gregorian.</i>			
B. 1800	4 13 48 32	1 16 48 25	1 2 32 58				
B. 1900	2 21 38 57	5 5 59 40	8 18 21 43				
B. 2000	0 29 29 22	8 25 10 55	4 4 10 28				
Biflexile Years Places are advanced a Day's Motion.							
Julian Years.	M. Mot. D.	Mean Mot. Ap. D.	Mean Mot. Q retr.	Greg. Years.	M. Mot. D.	Mean Mot. Ap. D.	Mean Mot. Q retr.
s o / //	s o / //	s o / //	s o / //	s o / //	s o / //	s o / //	s o / //
B. 100	10 7 50 25	3 19 11 15	4 14 11 15	100	9 24 39 50	3 19 4 34	4 14 8 4
B. 200	8 15 40 50	7 8 22 30	8 28 22 30	200	7 19 19 40	7 8 9 8	8 28 16 9
B. 300	6 23 31 15	10 27 33 45	1 12 33 45	300	5 13 59 30	10 27 13 42	1 12 24 13
B. 400	5 1 21 40	2 16 45 0	5 26 45 0	B. 400	3 21 49 55	2 16 24 57	5 26 35 28
B. 500	3 9 12 5	6 5 56 15	10 10 56 15	B. 800	7 13 39 50	5 2 49 54	11 23 10 56
B. 600	1 17 2 30	9 25 7 30	2 25 7 30	B. 1200	11 5 29 45	7 19 14 51	5 19 46 24
B. 700	11 24 52 55	1 14 18 45	7 9 18 45	B. 1600	2 27 19 40	10 5 39 48	11 16 21 52
B. 800	10 2 43 20	5 3 30 0	11 23 30 0	B. 2000	6 19 9 35	0 22 4 45	5 12 57 20
B. 900	8 10 33 45	8 22 41 15	4 7 41 15	B. 2400	10 10 59 30	3 8 29 42	11 9 32 48
B. 1000	6 18 24 10	0 11 52 30	8 21 52 30	B. 2800	2 2 49 25	5 24 54 39	5 6 8 16
B. 2000	1 6 48 20	0 23 45 0	5 13 45 0	B. 3200	5 24 39 20	8 11 19 36	11 2 43 44
B. 3000	7 25 12 30	1 5 37 30	2 5 37 30	B. 3600	9 16 29 15	10 27 44 33	4 29 19 12
B. 4000	2 13 36 40	1 17 30 0	10 27 30 0	B. 4000	1 8 19 10	1 4 9 30	10 25 54 40
B. 5000	9 2 0 50	1 29 22 30	7 19 22 30	B. 6000	7 27 28 45	1 26 14 15	4 8 52 0
B. 6000	3 20 25 0	2 11 15 0	4 11 15 0				

Mean

Mean Motion of the MOON, for 99 Julian Years.

Julian Years.	Mean Mot. D. s o / //	Apogee D. s o / //	Q ret. s o / //	Julian Years.	Mean Mot. D. s o / //	Apogee D. s o / //	Q ret. s o / //
1	4 9 23 3	1 10 39 50	0 19 19 43	49	0 17 56 51	6 13 52 26	7 17 44 19
2	8 18 46 7	2 21 19 41	1 8 39 26	50	4 27 19 55	7 24 32 17	8 7 4 2
3	0 28 9 11	4 1 59 32	1 27 59 10	51	9 6 42 59	9 5 12 8	8 26 23 46
B. 4	5 20 42 49	5 12 46 3	2 17 22 3	52	1 29 16 37	10 15 58 39	9 15 46 39
5	10 0 5 52	6 23 25 53	3 6 41 46	53	6 8 39 40	11 26 38 29	10 5 6 22
6	2 9 28 56	8 4 5 44	3 26 1 29	54	10 18 2 44	1 7 18 20	10 24 26 5
7	6 18 52 0	9 14 45 35	4 15 21 13	55	2 27 25 48	2 17 58 11	11 13 55 49
B. 8	11 11 25 38	10 25 32 6	5 4 44 6	56	7 19 59 26	3 28 44 42	0 3 8 42
9	3 20 48 41	0 6 11 56	5 24 3 49	57	11 29 22 29	5 9 24 32	0 22 28 25
10	8 0 11 45	1 16 51 47	6 13 23 32	58	4 8 45 33	6 20 4 23	1 11 48 8
11	0 9 34 49	2 27 31 38	7 2 43 16	59	8 18 8 37	8 0 44 14	2 1 7 52
B. 12	5 2 8 27	4 8 18 9	7 22 6 9	60	1 10 42 15	9 11 30 45	2 20 30 45
13	9 11 31 30	5 18 57 59	8 11 25 52	61	5 20 5 18	10 22 10 35	3 9 50 28
14	1 20 54 34	6 29 37 50	9 0 45 35	62	9 29 28 22	0 2 50 26	3 29 10 11
15	6 0 17 38	8 10 17 41	9 20 5 19	63	2 8 51 26	1 13 30 17	4 18 29 55
B. 16	10 22 51 16	9 21 4 12	10 9 28 12	64	7 1 25 4	2 24 16 48	5 7 52 48
17	3 2 14 19	11 1 44 2	10 28 47 55	65	11 10 48 7	4 4 56 38	5 27 12 31
18	7 11 37 23	0 12 23 53	11 18 7 38	66	3 20 11 11	5 15 36 29	6 16 32 14
19	11 21 0 27	1 23 3 44	0 7 27 22	67	7 29 34 15	6 26 16 20	7 5 51 58
B. 20	4 13 34 5	3 3 50 15	0 26 50 15	68	0 22 7 53	8 7 2 51	7 25 14 51
21	8 22 57 8	4 14 30 5	1 16 9 58	69	5 1 30 56	9 17 42 41	8 14 34 34
22	1 2 20 12	5 25 9 56	2 5 29 41	70	9 10 54 0	10 28 22 32	9 3 54 17
23	5 11 43 16	7 5 49 47	2 24 49 25	71	1 20 17 4	0 9 2 23	9 23 14 1
B. 24	10 4 16 54	8 16 36 18	3 14 12 18	72	6 12 50 42	1 19 48 54	10 12 36 54
25	2 13 39 57	9 27 16 8	4 3 32 1	73	10 22 13 45	3 0 28 44	11 1 56 37
26	6 23 3 1	11 7 55 59	4 22 51 44	74	3 1 36 49	4 11 8 35	11 21 16 20
27	11 2 26 5	0 18 35 50	5 12 11 28	75	7 10 59 53	5 21 48 26	0 10 36 4
B. 28	3 24 59 43	1 29 22 21	6 1 34 21	76	0 3 33 31	7 2 34 57	0 29 58 57
29	8 4 22 46	3 10 2 11	6 20 54 4	77	4 12 56 34	8 13 14 47	1 19 18 40
30	0 13 45 50	4 20 42 2	7 10 13 47	78	8 22 19 38	9 23 54 38	2 8 38 23
31	4 23 8 54	6 1 21 53	7 29 33 31	79	1 1 42 42	11 4 34 29	2 27 58 7
B. 32	9 15 42 32	7 12 8 24	8 18 56 24	80	5 24 16 20	0 15 21 0	3 17 21 0
33	1 25 5 35	8 22 48 14	9 8 16 7	81	10 3 39 23	1 26 0 50	4 6 40 43
34	6 4 28 39	10 3 28 5	9 27 35 50	82	2 13 2 27	3 6 40 41	4 26 0 26
35	10 13 51 43	11 14 7 56	10 16 53 34	83	6 22 25 31	4 17 20 32	5 15 20 10
B. 36	3 6 25 21	0 24 54 27	11 6 18 27	84	11 14 59 9	5 28 7 3	6 4 43 3
37	7 15 48 24	2 5 34 17	11 25 38 10	85	3 24 22 12	7 8 46 53	6 24 2 46
38	11 25 11 28	3 16 14 8	0 14 57 53	86	8 3 45 16	8 19 26 44	7 13 22 29
39	4 4 34 32	4 26 53 59	1 4 17 37	87	0 13 8 20	10 0 6 35	8 2 42 13
B. 40	8 27 8 10	6 7 40 30	1 23 40 30	88	5 5 41 58	11 10 53 6	8 22 5 6
41	1 6 31 13	7 18 20 20	2 13 0 13	89	9 15 5 1	0 21 32 56	9 11 24 49
42	5 15 54 17	8 29 0 11	3 2 19 56	90	1 24 28 5	2 2 12 47	10 0 44 32
43	9 25 17 21	10 9 40 2	3 21 39 40	91	6 3 51 9	3 12 52 38	10 20 4 16
B. 44	2 17 50 59	11 20 26 33	4 11 2 33	92	10 26 24 47	4 23 39 9	11 9 27 9
45	6 27 14 2	1 1 6 23	5 0 22 16	93	3 5 47 50	6 4 18 59	11 28 46 52
46	11 6 37 6	2 11 46 14	5 19 41 59	94	7 15 10 54	7 14 58 50	0 18 6 35
47	3 16 0 10	3 22 26 5	6 9 1 43	95	11 24 33 58	8 25 38 41	1 7 26 19
B. 48	8 8 33 48	5 3 12 36	6 28 24 36	96	4 17 7 36	10 6 25 12	1 26 49 12
				97	8 26 30 39	11 17 5 2	2 16 8 55
				98	1 5 53 43	0 27 44 53	3 5 28 38
				99	5 15 16 47	2 8 24 44	3 24 48 22

Connect these Motions with Places for whole Hundreds, in either Style, for the present Year's Places.

RADICAL Mean PLACES of the MOON, for Years and Months, Old and New Style.

1600				1600			
Julian Style.				Gregorian Style.			
Months.	Mean Pl. D.	M. Pl. Ap. D.	M. Pl. Q.	Months.	Mean. Pl. D.	M. Pl. Ap. D.	M. Pl. Q.
Ds.	s o / "	s o / "	s o / "	Ds.	s o / "	s o / "	s o / "
January 1	7 28 7 42	6 8 25 55	10 0 55 28	January 1	3 16 21 52	6 7 19 4	10 1 27 14
February 1	9 16 35 48	6 11 53 8	9 29 16 58	February 1	5 4 49 58	6 10 46 17	9 29 48 44
March 0	9 25 32 8	6 15 0 18	9 27 48 0	March 0	5 13 46 18	6 13 53 27	9 28 19 46
April 0	11 14 0 14	6 18 27 31	9 26 9 31	April 0	7 2 14 24	6 17 20 40	9 26 41 17
May 0	0 19 17 45	6 21 48 3	9 24 34 11	May 0	8 7 31 55	6 20 41 12	9 25 5 57
June 0	2 7 45 50	6 25 15 17	9 22 55 42	June 0	9 26 0 0	6 24 8 26	9 23 27 28
July 0	3 13 3 21	6 28 35 49	9 21 20 22	July 0	11 1 17 31	6 27 28 58	9 21 52 8
August 0	5 1 31 27	7 2 3 2	9 19 41 53	August 0	0 19 45 37	7 0 56 11	9 20 13 39
September 0	6 19 59 33	7 5 30 15	9 18 3 23	September 0	2 8 13 43	7 4 23 24	9 18 35 9
October 0	7 25 17 3	7 8 50 47	9 16 28 4	October 0	3 13 31 13	7 7 43 56	9 16 59 50
November 0	9 13 45 9	7 12 18 0	9 14 49 34	November 0	5 1 59 19	7 11 11 9	9 15 21 20
December 0	10 19 2 40	7 15 38 32	9 13 14 15	December 0	6 7 16 50	7 14 31 41	9 13 46 1
1700				1700			
Julian Style.				Gregorian Style.			
Months.	Mean Pl. D.	M. Pl. Ap. D.	M. Pl. Q.	Months.	Mean. Pl. D.	M. Pl. Ap. D.	M. Pl. Q.
Ds.	s o / "	s o / "	s o / "	Ds.	s o / "	s o / "	s o / "
January 1	6 5 58 7	9 27 37 10	5 16 44 13	January 0	1 11 1 42	9 26 23 38	5 17 19 10
February 1	7 24 26 13	10 1 4 23	5 15 5 43	February 0	2 29 29 48	9 29 50 51	5 15 40 40
March 0	8 3 22 33	10 4 11 33	5 13 36 45	March 0	3 8 26 8	10 2 58 1	5 14 11 42
April 0	9 21 50 39	10 7 38 46	5 11 58 16	April 0	4 26 54 14	10 6 25 14	5 12 33 13
May 0	10 27 8 10	10 10 59 18	5 10 22 56	May 0	6 2 11 45	10 9 45 46	5 10 57 53
June 0	0 15 36 15	10 14 26 32	5 8 44 27	June 0	7 20 39 50	10 13 13 0	5 9 19 24
July 0	1 20 53 46	10 17 47 4	5 7 9 7	July 0	8 25 57 21	10 16 33 32	5 7 44 4
August 0	3 9 21 52	10 21 14 17	5 5 30 38	August 0	10 14 25 27	10 20 0 45	5 6 5 35
September 0	4 27 49 58	10 24 41 30	5 3 52 8	September 0	0 2 53 33	10 23 27 58	5 4 27 5
October 0	6 3 7 28	10 28 2 2	5 2 16 49	October 0	1 8 11 3	10 26 48 30	5 2 51 46
November 0	7 21 35 34	11 1 29 15	5 0 38 19	November 0	2 26 39 9	11 0 15 43	5 1 13 16
December 0	8 26 53 5	11 4 49 47	4 29 3 0	December 0	4 1 56 40	11 3 36 15	4 29 37 57
1800				1800			
Julian Style.				Gregorian Style.			
Months.	Mean Pl. D.	M. Pl. Ap. D.	M. Pl. Q.	Months.	Mean. Pl. D.	M. Pl. Ap. D.	M. Pl. Q.
Ds.	s o / "	s o / "	s o / "	Ds.	s o / "	s o / "	s o / "
January 1	4 13 48 32	1 16 48 25	1 2 32 58	January 0	11 5 41 32	1 15 28 12	1 3 11 5
February 1	6 2 16 38	1 20 15 38	1 0 54 28	February 0	0 24 9 38	1 18 55 25	1 1 32 35
March 0	6 11 12 58	1 23 22 48	0 29 25 30	March 0	1 3 5 58	1 22 2 35	1 0 3 37
April 0	7 29 41 4	1 26 50 1	0 27 47 1	April 0	2 21 34 4	1 25 29 48	0 28 25 8
May 0	9 4 58 35	2 0 10 33	0 26 11 41	May 0	3 26 51 35	1 28 50 20	0 26 49 48
June 0	10 23 26 40	2 3 37 47	0 24 33 12	June 0	5 15 19 40	2 2 17 34	0 25 11 19
July 0	11 28 44 11	2 6 58 19	0 22 57 52	July 0	6 20 37 11	2 5 38 6	0 23 35 59
August 0	1 17 12 17	2 10 25 32	0 21 19 23	August 0	8 9 5 17	2 9 5 19	0 21 57 30
September 0	3 5 40 23	2 13 52 45	0 19 40 53	September 0	9 27 33 23	2 12 32 32	0 20 19 0
October 0	4 10 57 53	2 17 13 17	0 18 5 34	October 0	11 2 50 53	2 15 53 4	0 18 43 41
November 0	9 29 25 59	2 20 40 30	0 16 27 4	November 0	0 21 18 59	2 19 20 17	0 17 5 11
December 0	7 4 43 30	2 24 1 2	0 14 51 45	December 0	1 26 36 30	2 22 40 49	0 15 29 52

In Leap Year take out the Month Day Motions for January and February 1 Day sooner, in both Styles.

EXAMPLE.				EXAMPLE.			
To find the Mean Place D, Apogee, and Q, for February 19, 1752, at Noon, O. S. ?				To find the Mean Place D, Apogee, and Q, for January 15, 1760, at Noon, N. S. ?			
D Pl.	Ap. D	Q		D Pl.	Ap. D	Q	
s o / "	s o / "	s o / "		s o / "	s o / "	s o / "	
1700 O. S. Feb. 1.	7 24 26 13	10 1 4 23	5 15 5 43	1700 N. S. Jan. 0.	1 11 1 42	9 26 23 38	5 17 19 10
52 Yrs. Mot.	1 29 16 37	10 15 58 39	9 15 46 39	60 Yrs. Mot.	1 10 42 15	9 11 30 45	2 20 30 45
B. Days 18.	7 27 10 30	2 0 19	57 11	B. Days 14.	6 4 28 10	1 33 35	41 18
Leap Year.			9 16 43 50	Leap Year.			2 21 12 3
1752, O. S. Feb. 19	5 20 53 20	8 19 3 21	7 28 21 53	1760 N. S. Jan. 15	8 26 12 7	7 9 27 58	2 26 7 7

Mean

Mean MOTION of the MOON continued, for Months, Days, &c.

D ^s .	Mths.	M. Mot. D.				M. Mo. Ap. D.				M. Mot. Ω re.			
		s	o	i	''	s	o	i	''	s	o	i	''
31	Jan.	0	0	0	0	0	0	0	0	0	0	0	0
59	Feb.	1	18	28	6	0	3	27	13	0	1	38	30
90	March	1	27	24	26	0	6	34	23	0	3	7	28
120	April	3	15	52	32	0	10	1	36	0	4	45	57
151	May	4	21	10	3	0	13	22	8	0	6	21	17
181	June	6	9	38	8	0	16	49	22	0	7	59	46
212	July	7	14	55	39	0	20	9	54	0	9	35	6
243	August	9	3	23	45	0	23	37	7	0	11	13	35
273	Sept.	10	21	51	51	0	27	4	20	0	12	52	5
304	Oct.	11	27	9	21	1	0	24	52	0	14	27	24
334	Nov.	1	15	37	27	1	3	52	5	0	16	5	54
	Dec.	2	20	54	58	1	7	12	37	0	17	41	13

Take out 1 Day sooner for January and February, in Leap-year.

Days	1	0	13	10	35	0	6	41	0	3	11
2	0	26	21	10		0	13	22	0	6	21
3	1	9	31	45		0	20	3	0	9	32
4	1	22	42	20		0	26	44	0	12	43
5	2	5	52	55		0	33	25	0	15	53
6	2	19	3	30		0	40	6	0	19	4
7	3	2	14	5		0	46	47	0	22	14
8	3	15	24	40		0	53	29	0	25	25
9	3	28	35	15		1	0	10	0	28	36
10	4	11	45	50		1	6	51	0	31	46
11	4	24	56	25		1	13	32	0	34	57
12	5	8	7	0		1	20	13	0	38	8
13	5	21	17	35		1	26	54	0	41	18
14	6	4	28	10		1	33	35	0	44	29
15	6	17	38	45		1	40	16	0	47	40
16	7	0	49	20		1	46	57	0	50	50
17	7	13	59	55		1	53	38	0	54	1
18	7	27	10	30		2	0	19	0	57	11
19	8	10	21	5		2	7	0	1	0	22
20	8	23	31	40		2	13	41	1	3	33
21	9	6	42	15		2	20	22	1	6	43
22	9	19	52	50		2	27	4	1	9	54
23	10	3	3	26		2	33	45	1	13	5
24	10	16	14	0		2	40	26	1	16	15
25	10	29	24	36		2	47	7	1	19	26
26	11	12	35	11		2	53	48	1	22	37
27	11	25	45	46		3	0	29	1	25	47
28	0	8	56	21		3	7	10	1	28	58
29	0	22	6	56		3	13	51	1	32	9
30	1	5	17	31		3	20	32	1	35	19
31	1	18	28	6		3	27	13	1	38	30

Ex. 2. To find the mean Places of $\mathcal{D}$, Apogee, and Ω , for February 23, 1896, N. S. at Noon; answering to February 11, 1896, O. S.?

	M. Pl. $\mathcal{D}$.				Ap. $\mathcal{D}$.				Ω .			
	s	o	i	''	s	o	i	''	s	o	i	''
O. S. 1800.	4	13	48	32	1	16	48	25	1	2	32	58
Mo. Yrs 96	4	17	7	36	10	6	25	12	1	26	49	12
Feb.	1	18	28	6	0	3	27	13	0	1	38	30
B. Days 10	4	11	45	50	1	6	51					
									1	28	59	28
Places O. S.	3	1	10	4	11	27	47	41	11	3	33	30

The Answer.

Ex. 1. To find the mean Places of the $\mathcal{D}$, Apogee, and Ω , for May 20th, 9^h 20^m 36^s, P. M. 1759. O. S.

	M. Pl. $\mathcal{D}$.				Ap. $\mathcal{D}$.				Ω .			
	s	o	i	''	s	o	i	''	s	o	i	''
O. S. 1700	6	5	58	7	9	27	37	10	5	16	44	13
Mot. Yrs. 59	8	18	8	37	8	0	44	14	2	1	7	52
May - -	4	21	10	2	0	13	22	8	0	6	21	17
Days 20	8	23	31	40	2	13	41			1	3	33
Hours 9	4	56	28		2	30				1	11	
Minutes 20		10	59			6						3
Seconds 36			20			0			2	8	33	56
M. Places O. S.	4	13	56	13	6	13	59	49	3	8	10	17
11 Days Mo. -	4	24	56	25	-	1	13	32		+	34	57
M. Places N. S.	11	18	59	48	6	12	46	17	3	8	45	14

H.	M. Mo. $\mathcal{D}$.				Ap. $\mathcal{D}$.				H.	M. Mo. $\mathcal{D}$.				Ap. $\mathcal{D}$.			
	o	i	''	'''	o	i	''	'''		o	i	''	'''	o	i	''	'''
Min.	1	11	11	11	1	11	11	11	Min.	1	11	11	11	1	11	11	11
Sec.	1	11	iv	iv	1	11	iv	iv	Sec.	1	11	iv	iv	1	11	iv	iv
1	0	32	56	0	17	0	8		31	17	1	10		8	38	4	6
2	1	5	53	0	33	0	16		32	17	34	7		8	54	4	14
3	1	38	49	0	50	0	24		33	18	7	3		9	11	4	22
4	2	11	46	1	7	0	32		34	18	39	59		9	28	4	30
5	2	44	42	1	24	0	40		35	19	12	55		9	45	4	38
6	3	17	39	1	40	0	48		36	19	45	52		10	2	4	46
7	3	50	35	1	57	0	56		37	20	18	48		10	19	4	54
8	4	23	32	2	14	1	4		38	20	51	45		10	35	5	2
9	4	56	28	2	30	1	11		39	21	24	41		10	52	5	10
10	5	29	25	2	47	1	19		40	21	57	38		11	8	5	18
11	6	2	21	3	41	1	27		41	22	30	34		11	25	5	26
12	6	35	18	3	21	1	35		42	23	3	31		11	42	5	34
13	7	8	14	3	37	1	43		43	23	36	27		11	59	5	42
14	7	41	10	3	54	1	51		44	24	9	24		12	16	5	50
15	8	14	7	4	11	1	59		45	24	42	20		12	32	5	57
16	8	47	3	4	27	2	7		46	25	15	17		12	48	6	5
17	9	20	0	4	44	2	15		47	25	48	13		13	5	6	13
18	9	52	56	5	12	23			48	26	21	10		13	22	6	21
19	10	25	53	5	18	2	31		49	26	54	6		13	39	6	29
20	10	58	49	5	34	2	39		50	27	27	3		13	56	6	37
21	11	31	46	5	51	2	47		51	27	59	59		14	13	6	45
22	12	4	42	6	8	2	55		52	28	32	56		14	30	6	53
23	12	37	39	6	24	3	3		53	29	5	52		14	46	7	1
24	13	10	35	6	41	3	11		54	29	38	49		15	2	7	9
25	13	43	31	6	58	3	19		55	30	11	45		15	19	7	17
26	14	16	28	7	15	3	27		56	30	44	42		15	36	7	25
27	14	49	24	7	31	3	34		57	31	17	38		15	53	7	33
28	15	22	21	7	48	3	42		58	31	50	34		16	10	7	41
29	15	55	17	8	53	5	50		59	32	23	31		16	26	7	49
30	16	28	14	8	21	3	58		60	32	6	27		16	43	7	57

Ex. 3. To find the mean Places of the $\mathcal{D}$, Apogee, and Ω , for May 31st, 9^h 20^m 36^s N. S. answering to the above Time, O. S.?

	M. Pl. $\mathcal{D}$.				Ap. $\mathcal{D}$.				Ω .			
	s	o	i	''	s	o	i	''	s	o	i	''
N. S. 1700	1	11	0	42	9	26	23	38	5	17	0	10
Mot. Yrs. 59	8	18	8	37	8	0	44	14	2	1	7	52
May - -	4	21	10	2	0	13	22	8	0	6	21	17
Days 31	1	18	28	6	3	27	13			1	38	30
Hours 9	4	56	28		2	30				1	11	
Minutes 20		10	59			6						3
Seconds 36			20			0			2	9	8	53
M. Places N. S.	4	13	56	14	6	13	59	49	3	8	10	17

The same as above.

I. EQUATION of the MOON, or Annual Equation.

C.

Argument. SUN's Mean ANOMALY.

☉'s M. An.	SIGN 0.						SIGN 1.						SIGN 2.						☉'s M. An.
	D		Dif.		Apog.		D		Dif.		Apog.		D		Dif.		Apog.		
	+	-	+	-	+	-	+	-	+	-	+	-	+	-	+	-	+	-	
0	0 00				0 00		5 33				12 46		9 44				22 16		30
1	0 12	12		26	0 26	26	5 43	10		23	13 9	23	9 50	6		14	22 30	14	29
2	0 24	11		26	0 53	26	5 53	10		23	13 32	23	9 56	6		14	22 44	14	28
3	0 35	11		26	1 19	26	6 3	10		23	13 55	23	10 2	5		12	22 58	12	27
4	0 47	11		26	1 47	26	6 13	10		22	14 18	22	10 7	5		11	23 10	11	26
5	0 58	11		27	2 13	27	6 23	10		21	14 40	21	10 13	5		10	23 21	10	25
6	1 10	11		26	2 40	26	6 33	10		21	15 1	21	10 18	5		9	23 33	9	24
7	1 21	12		26	3 6	26	6 43	9		21	15 22	21	10 23	5		8	23 44	8	23
8	1 33	11		27	3 32	27	6 52	9		21	15 43	21	10 28	5		7	23 55	7	22
9	1 44	11		26	3 59	26	7 1	9		22	16 4	22	10 33	5		6	24 5	6	21
10	1 55	12		26	4 25	26	7 10	9		21	16 28	21	10 38	4		5	24 14	5	20
11	2 7	11		26	4 51	26	7 19	9		20	16 47	20	10 42	4		4	24 24	4	19
12	2 18	12		26	5 17	26	7 28	9		20	17 7	20	10 46	4		3	24 33	3	18
13	2 30	11		26	5 43	26	7 37	8		20	17 27	20	10 50	4		2	24 42	2	17
14	2 41	11		26	6 9	26	7 45	9		20	17 47	20	10 54	3		1	24 50	1	16
15	2 52	11		26	6 35	26	7 54	9		19	18 7	19	10 51	3		0	24 57	0	15
16	3 3	11		26	7 1	26	8 3	8		18	18 26	18	11 0	3		0	25 5	0	14
17	3 14	11		25	7 27	25	8 11	8		18	18 44	19	11 3	3		0	25 12	0	13
18	3 25	11		26	7 52	26	8 19	8		18	19 3	18	11 6	2		0	25 18	0	12
19	3 36	11		26	8 18	26	8 27	8		18	19 21	18	11 8	3		0	25 25	0	11
20	3 47	11		25	8 44	25	8 35	8		18	19 39	18	11 11	2		0	25 30	0	10
21	3 58	11		24	9 9	24	8 43	7		17	19 57	17	11 13	2		0	25 35	0	9
22	4 9	11		25	9 33	25	8 50	8		16	20 14	16	11 15	2		0	25 39	0	8
23	4 20	11		24	9 58	24	8 58	7		16	20 30	16	11 17	2		0	25 44	0	7
24	4 31	11		25	10 22	25	9 5	7		16	20 46	16	11 19	2		0	25 48	0	6
25	4 42	11		24	10 47	24	9 12	7		16	21 2	16	11 21	1		0	25 51	0	5
26	4 52	10		24	11 11	24	9 19	6		15	21 18	15	11 22	1		0	25 53	0	4
27	5 2	11		24	11 35	24	9 25	7		15	21 33	15	11 23	1		0	25 51	0	3
28	5 13	10		23	11 59	23	9 32	6		14	21 48	14	11 24	1		0	25 57	0	2
29	5 23	10		24	12 22	24	9 38	6		14	22 2	14	11 25	0		0	25 59	0	1
30	5 33				12 46		9 44			14	22 16		11 25	0		0	26 0		0
☉'s M. An.	D		Dif.		Apog.		D		Dif.		Apog.		D		Dif.		Apog.		☉'s M. An.
SIGN 11.																			
SIGN 10.																			
SIGN 9.																			

For the CONSTRUCTION, see the COMMENT on the MOON's THEORY.

C.

Argument. SUN's Mean ANOMALY.

☉'s M. An.	SIGN 3.					SIGN 4.					SIGN 5.					☉'s M. An.
	☽ +	☽ -	☽ +	☽ -	☽ +	☽ -	☽ +	☽ -	☽ +	☽ -	☽ +	☽ -	☽ +	☽ -		
	Dif.	Apog.	Dif.	Apog.	Dif.	Apog.	Dif.	Apog.	Dif.	Apog.	Dif.	Apog.	Dif.	Apog.		
0	11 25	26 0	9 21	0 0	10 2	22 45	8 10	4	5 52	13 14	4 45	8	30			
1	11 25	26 0	9 21	0 0	9 56	22 32	8 6	5	5 41	12 50	4 37	9	29			
2	11 25	26 0	9 21	0 0	9 50	22 18	8 1	5	5 30	12 25	4 28	9	28			
3	11 24	25 59	9 20	1 1	9 43	22 3	8 56	5	5 19	12 1	4 19	9	27			
4	11 24	25 58	9 20	2 2	9 37	21 48	7 51	5	5 8	11 37	4 10	9	26			
5	11 23	25 56	9 20	1 1	9 30	21 32	7 46	5	4 57	11 12	4 1	9	25			
6	11 23	25 54	9 19	2 2	9 23	21 17	7 41	6	4 46	10 46	3 52	9	24			
7	11 22	25 52	9 18	3 3	9 16	21 2	7 35	6	4 35	10 21	3 44	9	23			
8	11 21	25 49	9 17	4 4	9 9	20 45	7 29	7	4 24	9 55	3 35	9	22			
9	11 20	25 45	9 16	5 5	9 2	20 28	7 22	6	4 12	9 29	3 26	9	21			
10	11 18	25 41	9 15	6 6	8 55	20 11	7 16	7	4 0	9 3	3 16	10	20			
11	11 16	25 36	9 13	7 7	8 47	19 53	7 9	6	3 49	8 37	3 6	10	19			
12	11 14	25 31	9 11	8 8	8 39	19 35	7 3	6	3 37	8 11	2 56	9	18			
13	11 12	25 26	9 9	9 9	8 31	19 17	6 57	7	3 26	7 45	2 47	9	17			
14	11 10	25 21	9 7	10 10	8 23	18 59	6 50	7	3 14	7 19	2 38	10	16			
15	11 7	25 15	9 5	11 11	8 15	18 39	6 43	7	3 2	6 52	2 28	10	15			
16	11 4	25 8	9 3	12 12	8 7	18 20	6 36	7	2 51	6 25	2 18	9	14			
17	11 1	25 0	9 0	13 13	7 58	18 0	6 29	7	2 39	5 58	2 9	10	13			
18	10 57	24 52	8 57	14 14	7 49	17 40	6 22	7	2 26	5 31	1 59	10	12			
19	10 54	24 44	8 54	15 15	7 40	17 20	6 15	8	2 14	5 4	1 49	9	11			
20	10 50	24 35	8 50	16 16	7 31	16 59	6 7	8	2 2	4 37	1 40	10	10			
21	10 46	24 26	8 47	17 17	7 22	16 38	5 59	8	1 50	4 10	1 30	10	9			
22	10 41	24 17	8 44	18 18	7 13	16 16	5 51	7	1 38	3 42	1 20	10	8			
23	10 37	24 7	8 40	19 19	7 4	15 55	5 44	8	1 26	3 14	1 10	10	7			
24	10 33	23 56	8 37	20 20	6 54	15 33	5 36	8	1 14	2 46	1 0	10	6			
25	10 28	23 46	8 33	21 21	6 44	15 11	5 28	8	1 1	2 18	0 50	10	5			
26	10 23	23 34	8 29	22 22	6 33	14 48	5 19	9	0 49	1 50	0 40	10	4			
27	10 18	23 23	8 25	23 23	6 23	14 24	5 11	8	0 37	1 23	0 30	10	3			
28	10 13	23 11	8 20	24 24	6 12	14 1	5 3	9	0 25	0 56	0 20	10	2			
29	10 8	22 58	8 15	25 25	6 2	13 38	4 54	9	0 13	0 28	0 10	10	1			
30	10 2	22 45	8 10		5 52	13 14	4 45	9	0 0	0 0	0 0	10	0			
☉'s M. An.	SIGN 8.					SIGN 7.					SIGN 6.					☉'s M. An.
☽ +	☽ -	☽ +	☽ -	☽ +	☽ -	☽ +	☽ -	☽ +	☽ -	☽ +	☽ -	☽ +	☽ -	☽ +	☽ -	
Dif.	Apog.	Dif.	Apog.	Dif.	Apog.	Dif.	Apog.	Dif.	Apog.	Dif.	Apog.	Dif.	Apog.	Dif.	Apog.	

For the CONSTRUCTION, see the COMMENT on the MOON's THEORY.

II. EQUATION of the MOON, or I. *Semiannual.*
C.III. Equat. Moon.
C.*Argument.* Sun from MOON's APOGEE.*Arg.* Sun à Moon's Node.

☉ à D's Ap.	Sig. 0.	Sig. 1.	Sig. 2.	Sig. 3.	Sig. 4.	Sig. 5.	☉ à D's Ap.	Arg. prop. per.	S. 02 65	S. 12 75	S. 22 85	Arg. prop.
0	0 00	5 4	5 17	0 30	4 27	4 39	30	0	0 0	0 41	0 41	30
1	0 13	5 11	5 11	0 18	4 33	4 33	29	1	0 2	0 41	0 40	29
2	0 26	5 17	5 5	0 6	4 38	4 27	28	2	0 3	0 42	0 39	28
3	0 39	5 23	4 58	0 6	4 43	4 21	27	3	0 5	0 43	0 38	27
4	0 52	5 30	4 51	0 18	4 47	4 15	26	4	0 7	0 44	0 37	26
5	1 4	5 35	4 44	0 30	4 52	4 8	25	5	0 8	0 44	0 36	25
6	1 16	5 39	4 37	0 41	4 56	4 1	24	6	0 10	0 45	0 35	24
7	1 28	5 43	4 29	0 52	5 0	3 54	23	7	0 11	0 45	0 34	23
8	1 40	5 47	4 21	0 3	5 4	3 46	22	8	0 13	0 46	0 33	22
9	1 51	5 50	4 13	1 14	5 7	3 38	21	9	0 15	0 46	0 31	21
10	2 3	5 53	4 4	1 25	5 10	3 30	20	10	0 16	0 46	0 30	20
11	2 14	5 55	3 55	1 36	5 12	3 21	19	11	0 18	0 47	0 29	19
12	2 25	5 57	3 46	1 47	5 14	3 12	18	12	0 19	0 47	0 28	18
13	2 36	5 58	3 37	1 59	5 15	3 2	17	13	0 21	0 47	0 26	17
14	2 47	5 59	3 27	2 10	5 16	2 52	16	14	0 22	0 47	0 25	16
15	2 57	5 59	3 17	2 20	5 16	2 42	15	15	0 24	0 47	0 24	15
16	3 7	5 59	3 7	2 30	5 16	2 32	14	16	0 25	0 47	0 22	14
17	3 17	5 58	2 57	2 40	5 15	2 22	13	17	0 26	0 47	0 21	13
18	3 27	5 58	2 46	2 50	5 15	2 12	12	18	0 28	0 47	0 19	12
19	3 36	5 57	2 35	2 59	5 14	2 2	11	19	0 29	0 47	0 18	11
20	3 45	5 56	2 24	3 9	5 13	1 51	10	20	0 30	0 46	0 16	10
21	3 54	5 54	2 13	3 18	5 12	1 40	9	21	0 31	0 46	0 15	9
22	4 3	5 51	2 2	3 27	5 10	1 30	8	22	0 33	0 46	0 13	8
23	4 12	5 49	1 51	3 36	5 7	1 19	7	23	0 34	0 45	0 11	7
24	4 20	5 46	1 40	3 48	5 4	1 8	6	24	0 35	0 45	0 10	6
25	4 28	5 42	1 29	3 56	5 0	0 57	5	25	0 36	0 44	0 8	5
26	4 36	5 38	1 18	4 3	4 56	0 46	4	26	0 37	0 43	0 7	4
27	4 43	5 43	1 6	4 10	4 52	0 35	3	27	0 38	0 43	0 5	3
28	4 50	5 28	0 54	4 16	4 48	0 24	2	28	0 39	0 42	0 3	2
29	4 57	5 23	0 42	4 22	4 44	0 12	1	29	0 40	0 41	0 2	1
30	5 4	5 17	0 30	4 27	4 39	0 0	0	30	0 41	0 41	0 0	0
☉ à D's Ap.	Sig. 11.	Sig. 10.	Sig. 9.	Sig. 8.	Sig. 7.	Sig. 6.	☉ à D's Ap.	Arg. prop. per.	S. 53 11	S. 43 10	S. 33 9	Arg. prop.

For the CONSTRUCTION, see the COMMENT on the MOON's THEORY.

IV. Equat. Moon. C.				V. Eq. Moon. C.				VI. Eq. Moon. C.				VII. Equat. Moon. C.			
Arg. Mean Evection + 2 $\odot$ à D's $\odot$.				Arg. Mean Evection + 2 D M. Anom.				Arg. Mean Evection + $\odot$ M. Anom.				Arg. Mean Evection — $\odot$ M. Anom.			
Arg. pro- per.	S. 02 + 65 —	S. 12 + 75 —	S. 22 + 85 —	S. 02 + 65 —	S. 12 + 75 —	S. 22 + 85 —		S. 02 + 65 —	S. 12 + 75 —	S. 22 + 85 —		S. 02 + 65 —	S. 12 + 75 —	S. 22 + 85 —	Arg. pro- per.
0	I	II	I	II	I	II		I	II	I	II	I	II	I	0
0	0	0	0	30	0	52		0	0	1	12	2	5	0	30
1	0	1	0	31	0	52		0	2	1	15	2	7	0	29
2	0	2	0	32	0	53		0	5	1	17	2	8	0	28
3	0	3	0	32	0	53		0	7	1	19	2	9	0	27
4	0	4	0	33	0	54		0	10	1	21	2	10	0	26
5	0	5	0	34	0	54		0	12	1	23	2	11	0	25
6	0	6	0	35	0	55		0	15	1	25	2	12	0	24
7	0	7	0	36	0	55		0	17	1	27	2	13	0	23
8	0	8	0	36	0	56		0	20	1	29	2	14	0	22
9	0	9	0	37	0	56		0	22	1	31	2	15	0	21
10	0	10	0	38	0	56		0	25	1	33	2	16	0	20
11	0	11	0	39	0	57		0	27	1	35	2	17	0	19
12	0	12	0	39	0	57		0	30	1	37	2	17	0	18
13	0	13	0	40	0	57		0	32	1	39	2	18	0	17
14	0	14	0	41	0	58		0	35	1	41	2	19	0	16
15	0	15	0	42	0	58		0	37	1	43	2	20	0	15
16	0	16	0	42	0	58		0	40	1	44	2	20	0	14
17	0	17	0	43	0	58		0	42	1	46	2	21	0	13
18	0	18	0	44	0	59		0	44	1	48	2	21	0	12
19	0	19	0	45	0	59		0	46	1	49	2	22	0	11
20	0	20	0	46	0	59		0	49	1	51	2	22	0	10
21	0	28	0	47	0	59		0	51	1	52	2	23	0	9
22	0	22	0	47	0	59		0	54	1	54	2	23	0	8
22	0	23	0	48	1	0		0	56	1	55	2	24	0	7
24	0	24	0	48	1	0		0	59	1	57	2	24	0	6
25	0	25	0	49	1	0		1	1	1	58	2	24	0	5
26	0	26	0	49	1	0		1	3	2	0	2	24	0	4
27	0	27	0	50	1	0		1	5	2	1	2	25	0	3
28	0	28	0	51	1	0		1	8	2	3	2	25	0	2
29	0	29	0	51	1	0		1	10	2	4	2	25	0	1
30	0	30	0	52	1	0		1	12	2	5	2	25	0	0
Arg. pro- per.	S. 52 + 115 —	S. 42 + 105 —	S. 32 + 95 —	S. 42 + 115 —	S. 42 + 105 —	S. 32 + 95 —		S. 52 + 115 —	S. 42 + 105 —	S. 32 + 95 —		S. 52 + 115 —	S. 42 + 105 —	S. 32 + 95 —	Arg. pro- per.

N. B. Mean Evection is D M. Anom. — 2. $\odot$ à D Apogee.

For the CONSTRUCTION, see the COMMENT on the MOON'S THEORY.

VIII. Mean Elliptic EQUATION of the MOON'S CENTER, according to .05505 the Mean Eccentricity of her Orbit to Radius 1; or of 55050 Eccentricity to Sem. Transf. 1000000, or mean Distance à $\odot$.
C.

Argument. MOON'S corrected ANOMALY.

D cor. An.	Sign 0.					Sig. 1.					Sig. 2.					Sig. 3.					Sig. 4.					Sig. 5.					D cor. An.					
	—				Dif.	—				Dif.	—				Dif.	—				Dif.	—				Dif.	—				Dif.						
0	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30					
0	0	0	0	0	0	6	11	16	21	26	31	36	41	46	51	56	0	5	10	15	20	25	30	35	40	45	50	55	0	5	10					
1	0	6	11	16	21	26	31	36	41	46	51	56	0	5	10	15	20	25	30	35	40	45	50	55	0	5	10	15	20	25	30					
2	0	12	22	32	42	52	0	5	10	15	20	25	30	35	40	45	50	55	0	5	10	15	20	25	30	35	40	45	50	55	0					
3	0	18	32	46	60	74	88	0	14	28	42	56	70	84	98	112	126	140	154	168	182	196	210	224	238	252	266	280	294	308	0					
4	0	24	42	60	78	96	114	132	150	168	186	204	222	240	258	276	294	312	330	348	366	384	402	420	438	456	474	492	510	528	546					
5	0	30	52	74	96	118	140	162	184	206	228	250	272	294	316	338	360	382	404	426	448	470	492	514	536	558	580	602	624	646	668					
6	0	37	61	84	107	130	153	176	199	222	245	268	291	314	337	360	383	406	429	452	475	498	521	544	567	590	613	636	659	682	705					
7	0	43	69	95	121	147	173	199	225	251	277	303	329	355	381	407	433	459	485	511	537	563	589	615	641	667	693	719	745	771	797					
8	0	49	77	105	133	161	189	217	245	273	301	329	357	385	413	441	469	497	525	553	581	609	637	665	693	721	749	777	805	833	861					
9	0	55	84	113	142	171	199	228	257	286	315	344	373	402	431	460	489	518	547	576	605	634	663	692	721	750	779	808	837	866	895					
10	1	1	32	63	94	125	156	187	218	249	280	311	342	373	404	435	466	497	528	559	590	621	652	683	714	745	776	807	838	869	899					
11	1	7	38	69	100	131	162	193	224	255	286	317	348	379	410	441	472	503	534	565	596	627	658	689	720	751	782	813	844	875	906					
12	1	13	43	74	105	136	167	198	229	260	291	322	353	384	415	446	477	508	539	570	601	632	663	694	725	756	787	818	849	880	911					
13	1	19	49	80	111	142	173	204	235	266	297	328	359	390	421	452	483	514	545	576	607	638	669	700	731	762	793	824	855	886	917					
14	1	25	55	86	117	148	179	210	241	272	303	334	365	396	427	458	489	520	551	582	613	644	675	706	737	768	799	830	861	892	923					
15	1	31	61	92	123	154	185	216	247	278	309	340	371	402	433	464	495	526	557	588	619	650	681	712	743	774	805	836	867	898	929					
16	1	37	68	99	130	161	192	223	254	285	316	347	378	409	440	471	502	533	564	595	626	657	688	719	750	781	812	843	874	905	936					
17	1	43	74	105	136	167	198	229	260	291	322	353	384	415	446	477	508	539	570	601	632	663	694	725	756	787	818	849	880	911	942					
18	1	49	80	111	142	173	204	235	266	297	328	359	390	421	452	483	514	545	576	607	638	669	700	731	762	793	824	855	886	917	948					
19	1	55	86	117	148	179	210	241	272	303	334	365	396	427	458	489	520	551	582	613	644	675	706	737	768	799	830	861	892	923	954					
20	2	1	33	64	95	126	157	188	219	250	281	312	343	374	405	436	467	498	529	560	591	622	653	684	715	746	777	808	839	870	901					
21	2	7	39	70	101	132	163	194	225	256	287	318	349	380	411	442	473	504	535	566	597	628	659	690	721	752	783	814	845	876	907					
22	2	13	45	76	107	138	169	200	231	262	293	324	355	386	417	448	479	510	541	572	603	634	665	696	727	758	789	820	851	882	913					
23	2	19	51	82	113	144	175	206	237	268	299	330	361	392	423	454	485	516	547	578	609	640	671	702	733	764	795	826	857	888	919					
24	2	24	57	88	119	150	181	212	243	274	305	336	367	398	429	460	491	522	553	584	615	646	677	708	739	770	801	832	863	894	925					
25	2	30	63	94	125	156	187	218	249	280	311	342	373	404	435	466	497	528	559	590	621	652	683	714	745	776	807	838	869	900	931					
26	2	36	69	100	131	162	193	224	255	286	317	348	379	410	441	472	503	534	565	596	627	658	689	720	751	782	813	844	875	906	937					
27	2	41	75	106	137	168	199	230	261	292	323	354	385	416	447	478	509	540	571	602	633	664	695	726	757	788	819	850	881	912	943					
28	2	47	81	112	143	174	205	236	267	298	329	360	391	422	453	484	515	546	577	608	639	670	701	732	763	794	825	856	887	918	949					
29	2	52	87	118	149	180	211	242	273	304	335	366	397	428	459	490	521	552	583	614	645	676	707	738	769	800	831	862	893	924	955					
30	2	58	93	124	155	186	217	248	279	310	341	372	403	434	465	496	527	558	589	620	651	682	713	744	775	806	837	868	899	930	961					
D cor. An.	+ Dif.					+ Dif.					+ Dif.					+ Dif.					+ Dif.					+ Dif.					+ Dif.					D cor. An.
	Sig. 11.					Sig. 10.					Sig. 9.					Sig. 8.					Sig. 7.					Sig. 6.										

Ex. To find the Equation to $1^{\circ} 18'$ $\odot$ corrected Anomaly?

Construction, $\frac{1}{2}$ Arg. corrected An. $24^{\circ} 0' 0''$

By Tab. for red. com. Eq. $\odot$. . . C. Eq. . . . + 1 23 p. 82.

Equated $\frac{1}{2}$ Anomaly. 24 1 23 Tan.

Const. Log. + 9.6490533

Com. Log.

9.6490533

9.9521358

From the cor. $\frac{1}{2}$ Anom. deduct $\frac{1}{2}$ tr. An. 21 45 42 Tan. 9.6011891

Difference — 2 14 18 = $\frac{1}{2}$ Equat.

Doubled . . . 4 28 36 = the true Equation, req.

N. B. In the $\left\{ \begin{array}{l} \text{first} \\ \text{last} \end{array} \right\}$ 6 Signs when the $\left\{ \begin{array}{l} \text{true} \\ \text{cor.} \end{array} \right\}$ An. sub. fr: $\left\{ \begin{array}{l} \text{cor.} \\ \text{true} \end{array} \right\}$ the Sign of Equation is $\left\{ \begin{array}{l} - \\ + \end{array} \right\}$

See further on, for our universal and accurate Solution of the Keplerian Problem.

N. B. The Moon's corrected Anomaly, or above Argument, is her mean Anomaly corrected by the VII. foregoing, or First Equations.

IX.

IX. EVECTION-EQUATION of the MOON : Or Second EQUATION of the MOON's CENTER.

C.

Argument. True Evection ; Or M. Evection $\mp \frac{1}{2}$ M. Equation of the Moon's Center, according to its Title.

True E-vec.	Sign o.		Sig. 1.		Sig. 2.		Sig. 3.		Sig. 4.		Sig. 5.		True E-vec.
	—	Dif.	—	Dif.	—	Dif.	—	Dif.	—	Dif.	—	Dif.	
0	0 0 0	I 24	0 39 42	I 12	1 9 7	0 42	1 20 25	0 1	1 10 9	0 42	0 40 43	I 13	30
1	0 1 24	I 23	0 40 54	I 11	1 9 49	0 41	1 20 26	0 1	1 9 27	0 43	0 39 30	I 14	29
2	0 2 47	I 23	0 42 5	I 11	1 10 30	0 40	1 20 25	0 3	1 8 44	0 45	0 38 16	I 16	28
3	0 4 10	I 23	0 43 16	I 10	1 11 10	0 39	1 20 22	0 4	1 7 59	0 46	0 37 0	I 17	27
4	0 5 33	I 23	0 44 26	I 9	1 11 49	0 36	1 20 18	0 5	1 7 13	0 46	0 35 43	I 17	26
5	0 6 56	I 23	0 45 35	I 8	1 12 25	0 36	1 20 13	0 7	1 6 27	0 48	0 34 26	I 17	25
6	0 8 19	I 23	0 46 43	I 7	1 11 1	0 34	1 20 6	0 8	1 5 39	0 50	0 33 9	I 18	24
7	0 9 42	I 22	0 47 50	I 6	1 13 35	0 33	1 19 58	0 10	1 4 49	0 51	0 31 51	I 19	23
8	0 11 4	I 22	0 48 56	I 5	1 14 8	0 32	1 19 48	0 11	1 3 58	0 53	0 30 32	I 19	22
9	0 12 26	I 22	0 50 1	I 5	1 14 40	0 31	1 19 37	0 13	1 3 5	0 54	0 29 13	I 20	21
10	0 13 48	I 22	0 51 6	I 4	1 15 11	0 29	1 19 24	0 14	1 2 11	0 54	0 27 53	I 20	20
11	0 15 10	I 21	0 52 10	I 3	1 15 40	0 28	1 19 10	0 16	1 1 17	0 56	0 26 33	I 21	19
12	0 16 30	I 20	0 53 13	I 2	1 16 8	0 26	1 18 54	0 17	1 0 31	0 57	0 25 12	I 22	18
13	0 17 51	I 20	0 54 15	I 1	1 16 34	0 25	1 18 37	0 19	0 59 24	0 58	0 23 50	I 22	17
14	0 19 11	I 20	0 55 16	I 0	1 16 59	0 23	1 18 18	0 19	0 58 26	0 59	0 22 28	I 22	16
15	0 20 31	I 20	0 56 16	I 0	1 17 22	0 22	1 17 59	0 22	0 57 27	I 0	0 21 7	I 23	15
16	0 21 51	I 20	0 57 16	I 0	1 17 44	0 21	1 17 37	0 23	0 56 27	I 1	0 19 44	I 23	14
17	0 23 11	I 19	0 58 14	I 0	1 18 5	0 19	1 17 14	0 24	0 55 26	I 2	0 18 21	I 24	13
18	0 24 30	I 19	0 59 11	I 0	1 18 24	0 18	1 16 50	0 26	0 54 24	I 3	0 16 57	I 24	12
19	0 25 49	I 18	1 0 7	I 0	1 18 42	0 17	1 16 24	0 27	0 53 21	I 5	0 15 33	I 24	11
20	0 27 7	I 18	1 1 2	I 0	1 18 59	0 15	1 15 57	0 29	0 52 16	I 6	0 14 10	I 24	10
21	0 28 25	I 18	1 1 56	I 0	1 19 14	0 14	1 15 28	0 30	0 51 11	I 7	0 12 47	I 24	9
22	0 29 43	I 17	1 2 48	I 0	1 19 28	0 12	1 14 58	0 31	0 50 5	I 8	0 11 22	I 25	8
23	0 31 0	I 16	1 3 40	I 0	1 19 40	0 11	1 14 27	0 33	0 48 58	I 9	0 9 57	I 25	7
24	0 32 16	I 16	1 4 31	I 0	1 19 51	0 9	1 13 54	0 34	0 47 50	I 10	0 8 32	I 25	6
25	0 33 32	I 15	1 5 20	I 0	1 20 0	0 8	1 13 20	0 36	0 46 41	I 10	0 7 7	I 25	5
26	0 34 47	I 15	1 6 8	I 0	1 20 8	0 6	1 12 44	0 37	0 45 31	I 10	0 5 42	I 25	4
27	0 36 2	I 14	1 6 55	I 0	1 20 14	0 5	1 12 7	0 38	0 44 21	I 12	0 4 18	I 25	3
28	0 37 16	I 13	1 7 40	I 0	1 20 19	0 4	1 11 29	0 39	0 43 9	I 13	0 2 52	I 26	2
29	0 38 29	I 13	1 8 24	I 0	1 20 23	0 2	1 10 50	0 41	0 41 56	I 13	0 1 26	I 26	1
30	0 39 42	I 13	1 9 7	I 0	1 20 25	0 2	1 10 9	0 41	0 40 43	I 13	0 0 0	I 26	0
True E-vec.	+	Dif.	+	Dif.	+	Dif.	+	Dif.	+	Dif.	+	Dif.	True E-vec.
	Sig. 11.		Sig. 10.		Sig. 9.		Sig. 8.		Sig. 7.		Sig. 6.		

N. B. M. Anom. Δ , $\pm \frac{1}{2}$ M. Equat. Δ Center, (otherwise called $\frac{1}{2}$ the elliptic Equation) according to its Title, will be the ECCENTRIC ANOMALY of the Moon.

And the Eccentric Anomaly $\Delta - 2. \odot$ à Δ 's Apogee, will be the Argument of the EVECTION EQUATION of the Moon, as above.

For the CONSTRUCTION, see the COMMENT on the MOON's THEORY.

X. COMPOUND VARIATION-EQUATION of the MOON. C.

Argument. MOON Equated from the SUN's True PLACE.

Deq. à ☉ true.	Sig. 0.			Sig. 1.			Sig. 2.			Sig. 3.			Sig. 4.			Sig. 5.			Deq. à ☉ true.
	+		Dif.	+		Dif.	±		Dif.	—		Dif.	—		Dif.	—		Dif.	
	I	II		I	II		I	II		I	II		I	II		I	II		
0	0	0		32	19		31	12		2	0		34	26		34	3		30
1	1	19	1 19	32	56	0 37	30	29	0 43	3	19	1 19	35	4	0 38	33	21	0 42	29
2	2	38	1 19	33	30	0 31	29	43	0 48	4	38	1 19	35	40	0 33	32	36	0 47	28
3	3	57	1 18	34	1	0 28	28	55	0 49	5	57	1 18	36	13	0 30	31	49	0 50	27
4	5	15	1 18	34	29	0 26	28	6	0 51	7	15	1 17	36	43	0 28	30	59	0 51	26
5	6	33		34	55		27	15		8	32		37	11		30	8		25
6	7	50	1 17	35	19	0 24	26	22	0 53	9	48	1 16	37	37	0 26	29	14	0 54	24
7	9	6	1 16	35	40	0 21	25	27	0 55	11	4	1 16	37	59	0 22	28	17	0 57	23
8	10	22	1 16	35	58	0 18	24	30	0 57	12	19	1 15	38	19	0 20	27	19	0 58	22
9	11	38	1 16	36	14	0 16	23	31	0 59	13	34	1 15	38	36	0 17	26	19	1 0	21
10	12	53	1 15	36	27	0 13	22	30	1 1	14	48	1 14	38	51	0 15	25	17	1 2	20
11	14	7	1 14	36	37	0 10	21	27	1 3	16	12	1 14	39	3	0 12	24	13	1 4	19
12	15	20	1 13	36	44	0 7	20	23	1 4	17	15	1 13	39	12	0 9	23	7	1 6	18
13	16	31	1 11	36	49	0 4	19	17	1 6	18	27	1 12	39	18	0 6	22	0	1 7	17
14	17	40	1 9	36	50	0 1	18	10	1 7	19	37	1 10	39	22	0 4	20	51	1 9	16
15	18	48	1 8	36	49	0 1	17	1	1 9	20	46	1 9	39	21	0 1	19	41	1 10	15
16	19	55	1 7	36	45	0 4	15	50	1 11	21	53	1 7	39	20	0 1	18	29	1 12	14
17	21	0	1 5	36	38	0 7	14	38	1 12	22	58	1 5	39	15	0 5	17	16	1 13	13
18	22	4	1 4	36	29	0 9	13	26	1 14	24	1	1 3	39	8	0 7	16	1	1 15	12
19	23	7	1 3	36	17	0 12	12	13	1 13	25	3	1 2	38	58	0 10	14	44	1 17	11
20	24	8	1 1	36	3	0 14	10	59	1 14	26	4	1 1	38	45	0 13	13	27	1 17	10
21	25	7	0 59	35	46	0 17	9	43	1 16	27	4	1 0	38	29	0 16	12	9	1 18	9
22	26	4	0 57	35	26	0 20	8	27	1 16	28	2	0 58	38	10	0 19	10	51	1 18	8
23	26	59	0 55	35	3	0 23	7	10	1 17	28	58	0 56	37	49	0 21	9	32	1 19	7
24	27	51	0 52	34	38	0 25	5	52	1 18	29	52	0 54	37	25	0 24	8	12	1 20	6
25	28	41	0 50	34	10	0 28	4	34	1 18	30	43	0 51	36	58	0 27	6	51	1 21	5
26	29	29	0 48	33	39	0 31	3	16	1 18	31	32	0 49	36	28	0 30			1 21	
27	30	15	0 46	33	6	0 33	1	57	1 19	32	19	0 47	35	55	0 33	5	30	1 22	4
28	30	58	0 43	32	30	0 36	0	38	1 19	33	4	0 45	35	20	0 35	4	8	1 22	3
29	31	39	0 41	31	52	0 38	0	41	1 19	33	46	0 42	35	43	0 37	2	46	1 23	2
30	32	19	0 40	31	12	0 40	2	0	1 19	34	26	0 40	34	3	0 40	1	23	1 23	1
	—		Dif.	—		Dif.	±		Dif.	+		Dif.	+		Dif.	+		Dif.	
Deq. à ☉ true.	Sig. 11.			Sig. 10.			Sig. 9.			Sig. 8.			Sig. 7.			Sig. 6.			Deq. à ☉ true.

For the CONSTRUCTION, see the COMMENT on the MOON's THEORY.

CORRECTION of Mean VARIATION, (See Page 73.)
By Decimal Multipliers.
To be added to, or subtracted from, according to its Title,
the preceding Compound Variation.
C.

Eq. of M. Va. in Oct. or M. Dif. to Va. in Pe- rig. and Apog. $\odot \Delta \odot$.				Argument. Sun's Mean ANOMALY. Signs and Degrees.						
Arg. $\Delta \Delta \odot$. Signs and Degrees.										
Arg. pro- per.	0	1	2	0	1	2	3	4	5	Arg. pro- per.
+	+	+	+	Xrs	Xrs	Xrs	Xrs	Xrs	Xrs	
0	0	107	107	Dec.	Dec.	Dec.	Dec.	Dec.	Dec.	0
0	0	107	107	,92	,80	,48	,02	,49	,86	30
1	4	109	105	,92	,80	,47	,00	,51	,87	29
2	8	111	103	,92	,79	,45	,02	,52	,88	28
3	13	112	101	,92	,78	,43	,04	,54	,89	27
4	18	114	98	,92	,77	,41	,05	,55	,89	26
5	22	116	95	,92	,76	,40	,07	,56	,90	25
6	26	117	92	,91	,75	,38	,09	,58	,91	24
7	30	119	89	,91	,74	,36	,10	,59	,91	23
8	34	120	86	,91	,73	,35	,12	,60	,92	22
9	38	121	83	,91	,72	,34	,14	,62	,92	21
10	42	122	80	,90	,71	,33	,16	,63	,93	20
11	46	123	77	,90	,70	,31	,18	,65	,94	19
12	50	123	74	,90	,69	,30	,20	,66	,95	18
13	54	124	70	,90	,68	,28	,21	,68	,95	17
14	58	124	66	,89	,67	,26	,22	,69	,96	16
15	62	124	62	,89	,66	,25	,24	,70	,96	15
16	66	124	58	,89	,65	,24	,25	,71	,97	14
17	70	124	54	,88	,64	,22	,27	,72	,97	13
18	74	123	50	,88	,63	,20	,28	,73	,97	12
19	77	123	46	,88	,62	,18	,30	,75	,98	11
20	80	122	42	,87	,61	,17	,32	,76	,98	10
21	83	121	38	,87	,60	,15	,34	,77	,99	9
22	86	120	34	,86	,59	,14	,36	,78	,99	8
23	89	119	30	,86	,58	,12	,37	,79	,99	7
24	92	117	26	,85	,57	,10	,39	,80	,99	6
25	95	116	22	,85	,55	,09	,41	,81	1,00	5
26	98	114	18	,84	,54	,07	,42	,82	1,00	4
27	101	112	14	,83	,52	,05	,43	,83	1,00	3
28	103	111	9	,82	,51	,04	,45	,84	1,00	2
29	105	109	4	,81	,50	,03	,47	,85	1,00	1
30	107	107	0	,80	,48	,02	,49	,86	1,00	0
Arg. pro- per.	+	+	+	Xrs	Xrs	Xrs	Xrs	Xrs	Xrs	Arg. pro- per.
5	4	3		11	10	9	8	7	6	

Const. As $35' 10''$: to $124''$:: so any M. Variation : to Dif. in Seconds of a Degree, to be added to, or subtracted from, the present Mean Variation, for the Variation in Perigee or Apogee $\odot$, in the same angular Diff. $\Delta \Delta \odot$, respectively.
As Eq. $\Delta \Delta \odot$: to Cor. M. Var. to $\odot$ An. :: so 1 : to Xr, Halley. $\Delta \Delta \odot$ $0^{\circ} 25'$ Eq. $95''$, $\odot$ An. $3^{\circ} 26'$ Cor. $39''$, and $39''$ $\div 95'' = 41$, &c. Xr.

Otherwise. According to the logarithical Logarithms for correcting Halley's Mean Variation.
Given $\begin{cases} E' = \text{Minutes and Seconds in any Equation of mean Variation in Octants.} \\ S' = \text{the Sexagesimal, in Minutes and Seconds, answering to Lo. Log. Sun's mean Anomaly.} \\ e' = \text{the little Equation in Minutes and Seconds to be added to or subtracted from E, according to } \Delta \Delta \odot, \text{ for the Variation in Perigee or Apogee, respectively.} \end{cases}$
 $e'm = \text{Equation of Correction, sought, in Minutes and Seconds, to find } m?$

By Halley $E' \times \frac{S'}{60'} = E' + e'm$. Hence, by Transposition and Division $E' \times \frac{S'}{60'} - E' = m$. } Var. }
Apog. $33' 16''$
Perig. $37' 14''$

That is, $S' \text{ Dif. } 60' \times \frac{E'}{60' e'} = m = \frac{S' \text{ Dif. } 60' \text{ in Secs of a Deg.}}{213''}$ (where $E' = 35' 10''$, $e' = 2' 4''$.)

EXAMPLE. To find the Multiplier for $2^{\circ} 9'$ Sun's Mean Anomaly?
Lo. Log. Answering to
To which An. P. 73, is 0091 $58' 45''$ Dif. $60' = 1' 15'' = 75''$ } $\div 213'' = .35$, &c. nearly as above.

EXAMPLES of the USE.

THE Maximum of the foregoing Equation is the same as it is determined by Dr. Halley, the Sun being in Perigee; viz. $2' 4''$. It subtracts from the Mean Variation in the First, and Fourth Quadrature, of the Sun's Anomaly; and adds to it in the Second and Third. It is, when the Earth is in Aphelion, $1' 54''$, only.

The Readiness or Facility of working by these Multipliers is equal, at least, to that by the Logistical Logarithms.

EXAMPLE I.

$\Delta \Delta \odot 10^{\circ} 15'$. Sun's Anom. $0^{\circ} 0'$.
Gr. M. Var. $35' 10'$. Eq. of Var. $124''$ —
Correction — $1' 54''$ Dec. Xr. $,92$
Pref. Var. $33' 16''$.. C. 248
 1116 Comp. Var. $-36' 49''$
Correction — $114'' 08 = 1' 54''$
Lo. Log. 2320 Cor. Var. $-34' 55$
Gr. M. Var. $35' 10''$ For Sun's M. Anom. 0242 . See P. 73. C.
Pref. Var. $33' 16''$ 2562 .. Halley.
Correction — $1' 54''$

EXAMPLE II. See Page 72.

$\Delta \Delta \odot 0^{\circ} 25'$. Sun's Anom. $3^{\circ} 26'$.
M. Var. $26' 56''$. Eq. Var. $95''$ +
Correction + $40''$ Dec. Xr. $,42$
Pref. Variation $27' 36''$.. C. 190
 380 Comp. Var. $+28' 41''$
See P. 73, for Halley's Mean Variation. Correction + $39'' 90 = 0' 40''$ +
Lo. Log. 3479 Cor. V. + $29' 21$
M. Variation $26' 56''$ For Sun's Anom. 9896 See P. 73. C.
Pref. Variation $27' 35''$ 3375 .. Halley.
Correction + $0' 39''$

XI. Mean REDUCTION - EQUATION of the MOON to the ECLIPTIC.
C.

Arg. D in Orb. à 8 1 eq.

D à 8	0.6	1.7	2.8	D à 8
0	0	0	0	0
0	0 0	6 2	6 2	30
1	0 14	6 9	5 55	29
2	0 29	6 15	5 47	28
3	0 43	6 21	5 39	27
4	0 58	6 27	5 30	26
5	1 12	6 32	5 20	25
6	1 26	6 37	5 10	24
7	1 41	6 41	5 0	23
8	1 55	6 45	4 50	22
9	2 9	6 48	4 39	21
10	2 23	6 51	4 28	20
11	2 37	6 54	4 17	19
12	2 50	6 56	4 5	18
13	3 3	6 57	3 53	17
14	3 16	6 57	3 41	16
15	3 29	6 57	3 29	15
16	3 41	6 57	3 16	14
17	3 53	6 57	3 3	13
18	4 5	6 56	2 50	12
19	4 17	6 54	2 37	11
20	4 28	6 51	2 23	10
21	4 39	6 48	2 9	9
22	4 50	6 45	1 55	8
23	5 0	6 41	1 41	7
24	5 10	6 37	1 26	6
25	5 20	6 32	1 12	5
26	5 30	6 27	0 58	4
27	5 39	6 21	0 43	3
28	5 47	6 15	0 29	2
29	5 55	6 9	0 14	1
30	6 2	6 2	0 0	0
D à 8	+	+	+	D à 8
8	11.5	10.4	9.3	0

Note, The Seconds of Reduction, above or below the mean Reduction, are compensated for, or thrown into the Table of Variation, P. 50, together corresponding with the present Inclination of the Moon's Orb.

For, as many Seconds in the true compound Variation are omitted, viz. 24", as Dif. between the Mean, and greatest or least Excess amounts to.

MOON's Mean LATITUDE.
At the mean INCLINATION of her Orbit.
C.

Argument. D in Orb. from the Node 1 eq.

Arg. pr.	0. N	Dif.	1. N	Dif.	2. N	Dif.
0	6. S		7. S		8. S	
0	0 0 0	5 22	2 34 4	4 38	4 27 2	2 40
1	0 5 22	5 23	2 38 42	4 36	4 29 42	2 34
2	0 10 45	5 22	2 43 18	4 32	4 32 16	2 30
3	0 16 7	5 22	2 47 50	4 30	4 34 46	2 24
4	0 21 29	5 22	2 52 20	4 25	4 37 10	2 19
5	0 26 51	5 21	2 56 45	4 24	4 39 29	2 14
6	0 32 12	5 20	3 1 9	4 19	4 41 43	2 9
7	0 37 32	5 20	3 5 28	4 17	4 43 52	2 4
8	0 42 52	5 19	3 9 45	4 13	4 45 56	1 59
9	0 48 11	5 18	3 13 58	4 9	4 47 55	1 54
10	0 53 29	5 17	3 18 7	4 5	4 49 49	1 47
11	0 58 46	5 17	3 22 12	4 2	4 51 36	1 43
12	1 4 3	5 15	3 26 14	3 59	4 53 19	1 38
13	1 9 18	5 14	3 30 13	3 54	4 54 57	1 32
14	1 14 32	5 12	3 34 7	3 51	4 56 29	1 27
15	1 19 44	5 11	3 37 58	3 46	4 57 56	1 20
16	1 24 55	5 9	3 41 44	3 43	4 59 16	1 16
17	1 30 4	5 7	3 45 27	3 38	5 0 32	1 10
18	1 35 12	5 6	3 49 5	3 35	5 1 42	1 4
19	1 40 18	5 4	3 52 40	3 30	5 2 46	0 59
20	1 45 22	5 3	3 56 10	3 25	5 3 45	0 54
21	1 50 25	5 0	3 59 35	3 22	5 4 39	0 48
22	1 55 25	4 58	4 2 57	3 16	5 5 27	0 42
23	2 0 23	4 56	4 6 13	3 13	5 6 9	0 36
24	2 5 19	4 54	4 9 26	3 7	5 6 45	0 31
25	2 10 13	4 51	4 12 33	3 4	5 7 16	0 26
26	2 15 4	4 49	4 15 37	2 58	5 7 42	0 20
27	2 19 53	4 46	4 18 35	2 54	5 8 2	0 14
28	2 24 39	4 44	4 21 29	2 49	5 8 16	0 8
29	2 29 23	4 41	4 24 18	2 44	5 8 24	0 4
30	2 34 4		4 27 2		5 8 28	
Arg. pr.	11. S	Dif.	10. S	Dif.	9. S	Dif.
	5. N		4. N		3. N	

Ex. 1. Arg. Mean Lat. 9° . . M. Lat. 5° 8' 28" S.
Arg. Equation 3° . . Equat. — 8 52 N.

True Latitude D . . 4 59 36 S.

Ex. 2. Arg. M. Lat. 2° 10° . . M. Lat. 4° 49' 49" N.
Arg. Equation 5° 5° . . Equat. + 3 44 N.

True Lat. D . . 4 53 33 N.

Corresponding together with present Inclination of D's Orb.
Signs Arg.

Note, 0. 1. 2. N. A. } 6. 7. 8. S. A.
3. 4. 5. N. D. } 9. 10. 11. S. D.
N. A. North Ascending } S. A. South Ascending.
N. D. North Descending } S. D. South Descending.

EQUATION of the MOON's Mean to her True LATITUDE, for the present Inclination of her Orb. C.

Arg. 2 D à 0 — forego. Ar. Lat.

0. N	1. N	2. N	Arg. pr.
6. S	7. S	8. S	0
0 0	4 25	7 40	30
0 9	4 33	7 45	29
0 18	4 41	7 49	28
0 28	4 49	7 54	27
0 37	4 57	7 58	26
0 46	5 4	8 2	25
0 55	5 12	8 6	24
1 4	5 19	8 9	23
1 14	5 27	8 13	22
1 23	5 34	8 16	21
1 32	5 41	8 20	20
1 41	5 48	8 23	19
1 50	5 55	8 26	18
1 59	6 2	8 29	17
2 8	6 9	8 31	16
2 17	6 16	8 34	15
2 26	6 22	8 36	14
2 35	6 29	8 39	13
2 44	6 34	8 41	12
2 53	6 41	8 42	11
3 1	6 47	8 44	10
3 10	6 53	8 46	9
3 19	6 59	8 47	8
3 27	7 4	8 48	7
3 35	7 10	8 49	6
3 44	7 15	8 50	5
3 52	7 21	8 51	4
4 1	7 26	8 52	3
4 9	7 31	8 52	2
4 17	7 36	8 52	1
4 25	7 40	8 52	0
11. S	10. S	9. S	Arg. pr.
5. N	4. N	3. N	

Note, The Arg. of Mean Latitude and Argument of Equation being

Same? Name { N&N, S&S
Diff. } { N&S, S&N
the Equation will be { + }
{ - }

respectively: corresponding with the present Inclination of the Moon's Orb.

Increasing and diminishing the above Eq. in triplicate Ratio of Sun's Dist. will be nearer, viz. 0' 27" ± at a Maximum.

Mean HORIZONTAL PARALLAX of the MOON, in Syzgies.

Eccentricity 55050 to the middle State of her Orbit. Her Mean Dist fr. $\odot$ being 1000000. C.

Arg. Moon's Mean Anom. 8 eq. or eq. by the 8. or M. central Equation, only.

D's eq. An. $\circ$	Sig. 0	Sig. 1	Sig. 2	Sig. 3	Sig. 4	Sig. 5	D's eq. An. $\circ$
	+	+	+	+	+	+	
	1	1	1	1	1	1	
0	54 10	54 35	55 44	57 18	58 52	60 1	30
1	54 10	54 37	55 47	57 22	58 54	60 2	29
2	54 10	54 38	55 50	57 25	58 57	60 4	28
3	54 10	54 40	55 53	57 28	59 0	60 5	27
4	54 10	54 42	55 56	57 32	59 2	60 6	26
5	54 11	54 44	55 59	57 35	59 5	60 7	25
6	54 11	54 46	56 2	57 38	59 8	60 9	24
7	54 11	54 48	56 5	57 41	59 10	60 10	23
8	54 12	54 50	56 8	57 44	59 13	60 11	22
9	54 12	54 52	56 11	57 47	59 16	60 13	21
10	54 12	54 54	56 14	57 51	59 18	60 14	20
11	54 13	54 56	56 17	57 54	59 20	60 15	19
12	54 14	54 58	56 20	57 57	59 23	60 16	18
13	54 14	55 1	56 23	58 0	59 25	60 17	17
14	54 15	55 3	56 27	58 3	59 28	60 18	16
15	54 16	55 5	56 30	58 6	59 31	60 19	15
16	54 17	55 7	56 33	58 9	59 33	60 19	14
17	54 18	55 10	56 36	58 12	59 35	60 20	13
18	54 19	55 12	56 39	58 15	59 38	60 21	12
19	54 20	55 15	56 42	58 19	59 40	60 21	11
20	54 21	55 17	56 46	58 22	59 42	60 22	10
21	54 22	55 19	56 49	58 25	59 44	60 22	9
22	54 23	55 22	56 52	58 28	59 46	60 23	8
23	54 24	55 25	56 56	58 31	59 48	60 23	7
24	54 25	55 27	56 59	58 34	59 50	60 24	6
25	54 27	55 30	57 3	58 37	59 52	60 24	5
26	54 28	55 33	57 6	58 40	59 54	60 24	4
27	54 30	55 36	57 9	58 43	59 56	60 25	3
28	54 32	55 39	57 12	58 46	59 57	60 25	2
29	54 34	55 41	57 15	58 49	59 59	60 25	1
30	54 35	55 44	57 18	58 52	60 1	60 25	0
	+	+	+	+	+	+	
	Sig. 11	Sig. 10	Sig. 9	Sig. 8	Sig. 7	Sig. 6	

The above Mean Parallax, in Syzgies, and that in all Positions of the Moon and Sun, or in any State of the Moon's Orb, adds to the apparent for the true Altitude of the Moon; and consequently subtracts from the apparent for the true Distance from the Vertex, and the contrary.

I. EQUATION of D's hor. Parallax. C.

Arg. Mean Evection.

Arg. proper $\circ$	0	1	2
	6	7	8
	+	+	+
	1	1	1
0	42 36	21	
1	42 35	20	
2	42 35	19	
3	42 35	18	
4	42 34	18	
5	42 34	17	
6	41 34	17	
7	41 33	16	
8	41 33	15	
9	41 33	15	
10	41 32	14	
11	41 32	14	
12	40 31	13	
13	40 31	12	
14	40 31	11	
15	40 30	10	
16	40 30	9	
17	40 29	9	
18	40 29	8	
19	39 28	7	
20	39 27	6	
21	39 26	5	
22	39 25	5	
23	38 25	4	
24	38 24	4	
25	37 24	3	
26	37 23	3	
27	37 23	2	
28	36 22	1	
29	36 22	1	
30	36 21	0	
	+	+	+
	11	10	9
	5	4	3

II. EQUATION of D's hor. Parallax. C.

Arg. D's true Long. $\circ$

Arg. proper $\circ$	0	1	2
	6	7	8
	+	+	+
	1	1	1
0	15 45	30	
1	16 46	29	
2	17 46	28	
3	18 47	27	
4	19 47	26	
5	20 48	25	
6	21 49	24	
7	22 50	23	
8	23 51	22	
9	24 52	21	
10	25 52	20	
11	26 53	19	
12	27 54	18	
13	28 54	17	
14	29 55	16	
15	30 55	15	
16	31 56	14	
17	33 56	13	
18	34 57	12	
19	36 57	11	
20	37 57	10	
21	38 58	9	
22	39 58	8	
23	40 58	7	
24	41 59	6	
25	41 59	5	
26	42 59	4	
27	43 60	3	
28	43 60	2	
29	44 60	1	
30	45 60	0	
	+	+	+
	11	10	9
	5	4	3

EXAMPLES, in and out of Syzgies.

EXAMPLE I.

Equated Anom. $\circ 3^{\circ} 00'$. D $\hat{a}$ $\odot$
 $\circ 3^{\circ} 00'$ true.
 Mean H. Par. $54' 10''$ } M Ev. $\circ 3^{\circ} 00'$
 Equation $- 42$ }
 True H. Par. $53 28$ } C.
 $53 29$ } Halley.
 Eq. An. $\circ 3^{\circ} 00'$ D $\hat{a}$ $\odot$ } $3^{\circ} 60'$ -
 M. Evection 6 } $42 +$
 18 -

EXAMPLE II.

Eq. An. 6° . D $\hat{a}$ $\odot$ } 6° -
 M. Evection 6° } $42'' +$
 Mean H. Par. $60' 25''$ }
 Equation $+ 42$ }
 True H. Par. $61 7$ } C.
 $61 7$ } Halley.
 Eq. An. 6° D $\hat{a}$ $\odot$ } 9° -
 M. Evection $\circ 3'$ } $42 -$
 Equation $1' 42'' -$

EXAMPLE III.

C. Halley.
 Eq. An. $6^{\circ} 00'$. Tr. An. $5^{\circ} 18'$
 [056 Eccen.
 Mean H. Par. $60' 25''$ } $60' 17''$ } $1^{\circ} 15'$
 Evect. $3^{\circ} 10'$ } $+ 1$ } $+ 4$ } $\odot \hat{a}$ D
 D $\hat{a}$ $\odot$ } $4^{\circ} 15'$ } $- 30$ } $- 26$ } Apog.
 True H. Par. $59' 56''$ } $59' 55''$

EXAMPLE IV.

Eq. An. $3^{\circ} 10'$. D $\hat{a}$ $\odot$ } $3^{\circ} 00'$ -
 M. Evect. $3^{\circ} 7' 39''$ } $+ 0 4$
 Mean Hor. Parallax $57 22$
 True Hor. Parallax . . C. $56 26$
 By Halley . . $56 26$

We shall shew the Reduction of these Arguments of Parallax to Halley's and Halley's to these, hereafter: Or, taking + for the proper, and - for the contrary Sign,
 Our An. + Evect. Eq. + 2d Eq. Ap.
 His An. - Evect. Eq. + 2d Eq. Ap.
 [= Our An.]

- N. B. D equated Anom. — D $\hat{a}$ $\odot$ same eq. = An. Arg. where 1 Eq. Ap. D only is concerned.
 D Mean Anom. — 2. An. Argument = Mean Evection = 2. M. Pl. D $\hat{a}$ $\odot$ — D's M. Anomaly.
 D equated Anom. — An. Arg. = D $\hat{a}$ $\odot$; the D's Pl. being the same equat. with D's Anom.

EXAMPLES of Computing the True PLACES of the SUN and MOON.
C.

For the SUN's True PLACE.

Mean Time.	Sun's Longitude.	Sun's Apogee.
1733	9° 20' 58" 18"	3° 8' 17" 18"
May 27	4 24 53 25	25
Hours 20	49 17	
M. 27. S. 53	1 9	
Mean Place ☉	2 16 42 9	3 8 17 43 —
Equation ☉ Center	+ 41 50	2 16 42 9
True Place ☉	2 17 23 59	11 8 24 26 . . .
Equat. A. to M. Time } — 1 ^m 36 ^s }	. . . — 4	Or II 17 23 59 . . .
For app. Time, as above	2 17 23 55	Motion correspondent. very nearly.

To find the SUN's true PLACE.

RULE. Collect the mean Longitudes of the Sun, and his Apogee; subtracting the Latter from the Former, for the Sun's mean Anomaly; to which find the Equation of the Sun's Center, which connected with the Sun's mean Place, according to its proper Sign, gives his true Place, for Mean Time, required. With which Place connect the Mot. equal to the Equation of A. to M. Time, for ☉'s Place, apparent Time, required.

Equation of apparent to Mean Time.

For the

Sun's Mean Anomaly } + 2^m 47^s } By Tab. P. 20.
Sun's Pl. true . . . } — 4 23 }
Abfol. } A to M } Time — 1 36 + = + 3^m 57^m of Deg. M. Mo.
Equat. } M to A }
† Or proportion the Degs by the tr. diurn. Mot. 57' 14". P. 195.

For the MOON's True PLACE.

Mean Time	Moon's Longitude.	Moon's Apogee.	Moon's Node.
1733	8° 1° 3' 42"	6° 20° 25' 24"	8° 8° 28' 6"
May 27	4 16 55 48	16 22 37	7 47 4
Hours 20	10 58 49	5 34	2 39
M. 27. S. 53	15 18	8	4
Mean Place ☾	0 29 13 37	7 6 53 43	8 0 38 19
Annual Equat. I.	— 4 4	+ 9 23	— 3 22
☾ 1 Equated Apogee —	0 29 9 33	7 7 3 6	— 8 0 34 57
☾ Mean Anomaly 6 lefts Equations	5 22 6 27	5 22 6 27	0 26 19 42
	+ 0 22	2 20 41 46 . . .	4 25 44 45
☾ Anom. 7 Equated Elliptic Equation VIII.	5 22 6 49	3 1 24 41 . . .	2 ☉ à ☾'s Apogee.
	— 0 55 42	— 0 27 29 . . .	Mean Evection. 1/2 El. Eq. & 6 lefts Eqns.
☾ Anom. 8 Equated Evection Equation IX.	5 21 11 7	3 0 57 12	IX. Arg. of Evection.
	— 1 20 26		
☾ Anom. 9 Equated Comp. Var. Equation X. — 35' 54", corrected	5 19 50 41	For Corr. of ☾ Arg. ☉ M. An. M. Variat. } L. L. 0226. P. 73. Cor. — 1' 45"; Eq. 122", X 1, 86	No. Arguments of Equations.
☾ true Equated Anom. Apog. 1 Equated +	5 19 16 32	By { Mean H. Par. 60' 22" M. Evac. Eq. + 0 1 Tab. ☾ tr. à ☉ Eq. — 0 37	II. 7 10 21 5 13 . . .
☾ Orbit Longitude Reduction + 20" XI. } Equat. Precef. P. 33. }	0 26 19 38	Page 53. { True Hor. Par. 59 46 By (P. 78) Halley 59 47	III. 6 16 50 26 . . .
	+ 6 49		IV. 4 5 2 . . . 0 49
☾ in Ecliptic Observed	☿ 26 26 27	☾ Anom. 9 Equated	V. 2 16 7 . . . 1 22
	☿ 26 26 16	— ☉ à ☾'s Ap. or Ann.	VI. 2 9 48 . . . 2 16
Error	+ 0 11	☾ à ☉ . . .	VII. 3 23 0 . . . 1 34

To find the MOON's true PLACE.

RULE. Collect the mean Longitudes of the Moon, her Apogee, and Node, and then connect the Equations, from the several Arguments, as on the Left, and below.

☾ The ☾'s Ap. 1 eq. is first subtr. from ☾'s Place 1 eq. for finding the M. Anom. eq. Anomalies, and Arguments, and is lastly added to ☾'s Pl. 10 eq. for her Orbit Place.

☾ Orb Long. ☾'s Lat. XI. ☾ in Orb à ☉. I. Arg. Lat. } 2° 53' 26"
2 ☾ à ☉ — I. Ar. II. Arg. Lat. } Eq. + 8 9
☾'s true Lat. N. D. 3 1 35

Horizontal Parallax ☾ to ☾'s horizontal Diam. 60' 48" to 33' 27", Or 60' to 33'.

Equations.	Arguments.
—	☉ 2° 17° 23' 59"
+	☾'s Ap. 7 7 3 6
—	☉ à ☾ Ap. 7 10 20 53
+	or Argument Annual.
—	Double 2 20 41 46
+	2 ☾ à ☉ 8° 19° 0'
—	I. Arg. M. La. 4 25 45
+	II. Arg. Eq. Lat. 3 23 15
—	6 lefts Equations.

N. B. The ☾'s observed Longitude above is corrected by + 14", according to the Precept in the Preface to Dr. Halley's Astronomical Tables; being the Dif. of the app. above and true Semi-Diameter of the Moon, in Long. or R. A. at any Time, with Respect to ☾'s Alt. and Lat. or Declination: Dr. Halley having deduced his lunar Places, in observing either Limb of the ☾, from her app. instead of her true Diam. at those Times: 60' 48" to 33' 27" is as 60' to 33', for the horizontal Parallax to the hor. Diam. of the Moon, being as 1 to .55. Hence the constant Log. Log. 2596 being added to the Log. Log. of the hor. Par. will give the Lo. Log. of the Moon's hor. Diam. Or 1/2, + 1/10 of that Half, of the horizontal Parallax = horizontal Diameter of the Moon.

EXAMPLE. Moon's horizontal Parallax	59' 46"	Lo. Log.
	+	0017
		2596
Moon's horizontal Diameter	32' 52"	2613

Or, ☾'s hor. Par. 59' 46" = 59' 76
1/2 . . . 29 53
1/10 . . . 2 59
32' 868
52"
☾'s hor. Diam. 32' 52" as before.

☞ For common Use, where great Exactness in the Moon's Place is not required, the 6 left Equations, II, III, IV, V, VI, and VII, may be omitted, as amounting but to few Minutes; in the present Instance but to + 22". And the I, VIII, IX, X, and XI Equations (the last being the Reduction-Equation at a Mean) will be sufficient.

REMARKS

REMARKS on Mr. MAYER'S ACCELERATION of the MOON'S MOTION.

Mr. MAYER having said, that by comparing several *Periods of Eclipses* together he is ascertained of the Moon's Motion being *quicker* now than in former Ages; and that in 60 Years it is $1' 9''$ faster than *Halley's* or *our* Tables represent it. I had therefore a Mind to try, whether from solar Eclipses (whose Beginning and End are much more accurately determined than *lunar* Ones; and which, if central, or nearly so, the Error of a *Minute* in Latitude will not make any sensible Error in the *Scruples of Incidence*) I could collect any *Acceleration* beyond what the present Tables represent.

For this Purpose, I took one Period of 54 Years and another of 72: The first from 1661 to 1715; the other from 1676 to 1748.

		Equal Time at Greenwich.	
March 10, 1661,	Hevelius observed the End of the solar Eclipse at Dantzic . . .	$11^h 16^m 11''$	
	Moon's Longitude observed . . .	$\gamma 11^o 4' 9''$	
	Error of Computation . . .	$+ 5$	
April 15, 1715,	Dr. Halley observed the Beginning of the total solar Eclipse . . .	$8^h 2^m 40''$	
	Moon's Longitude observed . . .	$\delta 11^o 20' 17''$	
	Error of Computation . . .	$- 2''$	
54 Years Distance.			
June 1, 1676,	Hevelius observed the End of a solar Eclipse . . .	$10^h 23^m 30''$	
	Moon's Longitude observed . . .	$\Pi 21^o 26' 17''$	
	Error of Computation . . .	$- 4''$	
July 14, 1748,	Dr. Bradley observed, at Greenwich, the End of a solar Eclipse . . .	$12^h 15^m 48''$	
	Moon's Longitude observed . . .	$\Omega 3^o 9' 37''$	
	Error of Computation . . .	$+ 11''$	
72 Years Distance.			

According to Mayer's Hypothesis our Tables should have lost in 54 Years . . . $1' 2''$

But by the foregoing Computation the Difference is only . . . $- 7''$

According to the same Hypothesis in 72 Years, our Tables should have lost . . . $1' 23''$

But, by Comparison of the Computation, they have gained . . . $15''$

Considering the Distance of Time, between Hevelius's Days and Ours, and the great Difference of the Meridians, the near Agreement in Computation, as well as Nearness of Observation, is surprizing, after such a long Period of Time!

I am sensible that these Observations are too few to speak with Certainty upon; but they give one Reason to hope that the Mean Motions in this Age do not much differ from those set down in our Tables; though the Acceleration of the Moon's Motion, for former Ages, appears more evident, as we have shewn, and allowed for, further on.

D. C.

An Account of the foregoing TABLES.

THE Radical Mean PLACES of the SUN and MOON, (distinguished from Mean MOTIONS) are fitted to the Beginning of each Julian Year, in the Old or New Style, as expressed, answerable to December 31, at Noon: Except those Radical Mean Places for the Leap-Years, which are advanced by a Day's Motion, answerable to the 1st Day of January at Noon; and on which Account 1 Day and a Day's Motion is to be subducted in the Computation for Leap-Years, or 1 Day less taken out for, in the Months of January and February, as being the shorter Method; instead of adding a Day and Day's Motion for all the Months after February, according to the old Method in all former astronomical Tables.

The mean or middle Motions, in respect of Time (distinguished from the radical mean or middle Places) are every where proportional to the Differences of the radical mean Places, in *Revolutions, Signs, Degrees, &c.* for different Points of Time.

The radical mean Places and Motions for both Old and New Styles are inserted to suit the Purposes of Computation for Time back (before the New Style) as well as for Time forward, since the Account of Time by the New Style, was established. By this Means the Computer first reducing the Date of one Style to the other, may make a double Computation corresponding to either Style, and thereby find the Agreement in the Result of both Methods, when great Certainty or Accuracy is required: Or finding a Disagreement of both Methods, he may proceed to detect his Error, and discover Truth in the End.

The 99 Julian Years Motion, suited to both Styles, and never before contrived, serve for adding to or subtracting from the radical mean Places of the SUN or MOON; according as the Time for the mean Places sought is forward or backward, respectively, of the Time for the given Radixes; being the shortest Method of Computation that can be put in Practice, especially with the Motions for each Month-Day, as we have annexed to it; there being only one Addition or Subtraction of two Lines, and then one Addition of another Line to find the mean Place of the SUN or MOON, for any Day at Noon; whatever may be the Number of Years backward or forward of the present Year.

But we have given the Radixes from 1652 to 1752, O. S. when the Style received Alteration by Act of Parliament, and also from 1752 to 1852, N. S. for determining the radical mean Places of the Sun or Moon, for any Day at Noon, in any Year Old or New Style, of those Intervals, by one Addition of two Lines, only.

We have also given the radical mean Places of the Sun and Moon, at the Beginning of every Month, or End of the former Month, (except giving them for January 1, and February 1, at Noon, in Leap-Years) in the Years 1600, 1700, and 1800, O. and N. Style, by which the mean Places of Sun and Moon are had, for any Day at Noon, in any Year of those Centuries, by the Addition of three Lines; namely the Radix-Line, Line of the Year's Motion, and Line of the Day's Motion, in that Year; whereby the Result of one Computation may be proved by the other, corresponding with the Dates of the different Styles.

The Motions of the Sun and Moon are also given to the Beginning of each Month in the Year to be added to the Radixes, answering to the Beginning of any Year, Old or New Style, and to which the Motion for the Number of the Month-Day is to be added, for the Sake of Variety, and antient Custom.

And EXAMPLES of the USE and PRACTICE, of all these different Methods of Computation, are exhibited at the Bottom, or on the Sides of the Pages at the Beginning of our Work, to make them clear, and undoubted.

There are different Methods and Tables, for proportioning the Difference of Equations: So that the Computer may take his Choice of that Method which suits him best. This is a new Improvement we have made which was much wanted, to facilitate Computation, and give Ease to the Attention. We find, by Experience, that since Sexagesimals may be reduced, at Sight, into Decimals, (see P. 14.) that multiplying the Difference of Equation, by the Difference of Argument (reduced to Decimals) finds the proportional Difference to be added to or subtracted from the first Equation, about as soon as the Logistical Logarithm of one of them can be found (turning over Leaves) and greatly eases Attention, beyond any other Method. But, for those who are not expert or ready at Multiplication, they may use the Tables at Pages 256, 257, 258, and 259, with great Facility and Quickness.

Our Table, at Page 13, is a Novelty, first thought of; but proportions not so quick as those just mentioned.

The Tables of the solar and lunar Places and Motions are fitted to Mean Time; to which Apparent Time must be always reduced, in applying it to Use by those Tables.

PRECEPTS OF COMPUTATION.

PRECEPT I. To reduce the Apparent to equal TIME and the contrary?

By *Tab. p. 19.* Take out the Equation, at once, standing under A, correspondent to the Sun's true Place, and according to its Sign, + or —, add or subtract it, to or from the apparent for the Mean Time. Or if the Mean Time is given, take out the Equation standing under M, correspondent to the Sun's true Place, and according to its Sign, + or —, add or subtract it, to or from the Mean for the Apparent Time, required: being the *Converse* of the former Equation for reducing Apparent to Mean Time.

OTHERWISE. By *Tab. p. 20,* take out the first Equation correspondent to the Sun's Place, under A, or M, according as either the Apparent or Mean Time is given, to be reduced to the Time required, and set it down with its proper Sign, + or —; then take out the second Equation of Time answerable to the Sun's M. Anomaly, under A or M, correspondent to the Apparent or Mean Time given, in the first Equation, to be reduced to the required Time, and set this 2d Equat. down with its proper Sign, under the 1st Equation; these two Equations connected together in a Sum or Difference with the proper Sign prefix'd thereto, (adding the Equations together, if of like Signs, and putting the Sign of either to the Sum; or if of unlike Signs taking their Difference, and putting the Sign of the greater Equation to that Difference) will be the Absolute Equation of Time required, to be added to or subtracted from the Time given, according as its Sign is + or —, for reducing it to the Time required.

PREC. II. To compute the true PLACE of the SUN?

AMONG the Radixes of the Mean Places, seek the Radix to the Year in which the Sun's true Place is required, (if of modern Date) and set down the Sun's Mean Place and Apogee, in distinct Columns for the Beginning of that Year; then add the Mean Motion of the Sun and his Apogee, for the Month, Day, Hour, Minute and Second in that Year, the Sum of which will be the Sun's Mean Place and Apogee to the Mean Time required.

OR, if the Year is far back, or forward of the present Date, set down the M. Places for any of the Centuries you will find in p. 8, and afterwards set down under them the Mean Motion of the Sun and his Apogee for the Years since, and under those the Motions for the Day, Hour, &c. since those; the Sum of which (rejecting Revolutions) will be the Mean Place of the Sun, and of his Apogee for the Mean Time required.

OR, if you can find the Century, in the Month-Page 10, take out the Mean Places to the Year and Beginning of the Month, then subjoin the Motions for the Year, Day, Hour, &c. since that Century; and the Sum of which will be the Mean Place of the Sun and his Apogee, required. See *Exam. p. 10 and 11.*

NOW, from the Sun's Mean Place or Long. deduct the Place of his Apogee, and the Remainder will be the Sun's Mean Anomaly; with which (from *Table p. 12.*) take out the Equation of the Sun's Center (by making proper Proportion for the Diff. of Equation and Diff. of Argument) adding or subtracting it, to or from the Sun's Mean Place, according to the Sign of Equation + or —, and the Result will be the Sun's true Longitude, for the Mean Time, required. See *Example p. 54.* You may correct his true Place, thus computed, by *Table of Precession, p. 33.*

PREC. III. To compute the Mean and then the true PLACE of the MOON.

FROM among the Radixes of the Moon, as of the Sun, set down the Mean Place of the Moon, her Apogee and Node, in distinct Columns, drawing a Line under the radical Mean Place of the Node. Then set down in each of these respective Columns, the Mean Motions of the Moon, Apogee, and Node, for Time since the radical Time and Places. Collect the Mean Longitude or Place of the Moon, and her Apogee into two distinct Sums, in like Manner as the Mean Longitude and Apogee of the Sun is collected, by the foregoing Precept. But the Mean Motion of the Node being retrograde, you must collect the Motions of the Node into one Sum, since the first radical Time and Place, and subtract that Sum from the Node's radical Mean Place, under which a Line was drawn, and the Residue will be the present Mean Place of the Node; as the Sum of the radical Mean Places and Motions of the Moon and Apogee, before collected, will be the Mean Places of the Moon and Apogee for the Mean Time required.

PREC. IV. To apply the Moon's EQUATIONS to her Mean PLACES, to find her true PLACE.

1. FROM p. 44. (Having computed the ☉'s true Place to the Mean Time for which the ☾'s true Place is required) take out I. Equation of ☾, Apogee, and ☿, according to the Argument of ☉'s M. Anom. and set down their respective Quantities and Signs, under their like in the Mean Motions, which added and subtracted, as required, the Result will be the Moon's Place, Apogee, and Node, 1 equated.

2. Deduct the ☾'s Apogee 1 equated, from ☾'s Place 1 equated, and the ☾'s Anomaly 1 equated will remain: with which connecting the 6 left Equations, taken from their several and respective Arguments, the Result will be the Moon's Anomaly 7. equated.

3. With the Argument of the Moon's Anomaly 7. equated take out (from p. 48.) the Mean Central Equation, which (with its proper Sign) connect with the ☾'s Anom. 7. equated, and you will have the Moon's Anomaly 8 equated.

4. From the ☾'s Anom. 1 equated deduct 2 ☉ à ☾'s Apogee, or twice Ann. Arg. and the Remainder will be the Mean Evection; with which M. Evection connect (with the proper Sign) $\frac{1}{2}$ Sum of the Mean Eccentric or central Equation, and 6 left Equation, II, III, IV, V, VI, VII, and the Result will be the Arg. of Evection, for reducing the Mean to the true Central Equation. With which Arg. of Evection, take out (fr. p. 49.) the Evection Equation, and connect it (in its proper Sign) with ☾'s Anom. 8 equated, and you will have ☾'s Anom. 9 equated.

5. From the Moon's Anomaly 9 equated, deduct ☉ à ☾'s Apogee, or annual Argument, and the Remainder will be the ☾ à ☉, or Arg. of X. Equation; with which (fr. P. 50) take out the compound Variation-Equation, and correct it, according to Halley's Correction of M. Variation, P. 73, or by decimal Xrs, Page 51, connect which now correct Variation with its proper Sign, and ☾'s Anomaly 9 equated, and you will have ☾'s Anomaly 10 equated. To which adding the ☾'s Apogee 1 equated, (before deducted from ☾'s Place 1 equated) and you will have the ☾'s Orbit-Place, or Longitude in her Orbit.

6. From the Moon's Orbit-Place, deduct the Place of the Node 1 equated, and the Remainder will be the ☾ à ☿, or the Arg. for the Reduction of her Orbit-Place to the Ecciptic; and also the Argument for finding her mean Latitude. With which Argument take out the Reduction-Equation (from Page 51) and connect it (in the proper Sign) with the ☾'s Orbit-Place, (connecting also the Equation of Precession from Page 33, at the same Time, or at last, if you like it best) and you will have the Moon's true Place in the Ecciptic, required.

PREC. V. To find the MOON's true LATITUDE?

With the Argument of Reduction, or of the mean Latitude, before found, take out (from Page 52) the Moon's mean Latitude. Then deduct this Arg. from twice the Moon from the Sun, and the Remainder will be the Arg. of Equation of Latitude. With which Arg. take out the Equation of Latitude from the same Page, adding or subtracting it, to or from the mean Latitude before found, (according as it is of the same or different Name with the mean Latitude) and the Result will be the true Latitude of the Moon, required.

PREC. VI. To find the MOON's true HORIZONTAL PARALLAX?

WITH the Argument of the Moon's mean Anomaly, corrected with the Mean central Equation, or Anom. 8 equated, take out (from P. 53) the mean horizontal Parallax correspondent to the middle State of the Moon's Orbit; then, with the Argument of mean Evection, take out the first Equation of ☾'s horizontal Parallax; and lastly, with the Argument of ☾ à ☉ true, (or Residue of the annual Argument from the ☾'s Anom. 9 equated) take out the second Equation of the Moon's horizontal Parallax, connecting these two Equations, in their proper Signs, with the mean horizontal Parallax, and the Result will be the true horizontal Parallax (agreeing with Halley's prolixer Method) required. See EXAMPLES, Pages 53 and 54.

PREC. VII. To find the MOON's true horizontal DIAMETER?

ADD the const. Lo. Log. 2596 to the Lo. Log. of the true horiz. Parallax, and their Sum will be the Lo. Log. of true horizontal Diam.

Otherwise. Take the Half of the true horizontal Parallax, then take the Tenths of that Half, and the Sum of both will be the true horizontal Diameter of the Moon, required. Or multiply the horizontal Parallax by .55 for the horizontal Diameter of the Moon.

☞ The true horizontal Parallax, according to Mayer, differs from the horizontal Parallax determined by this and also Halley's Method, by almost Half a Minute of a Degree.

PREC. VIII. To find the MOON's apparent DIAMETER from her true horizontal DIAMETER?

By *Tab. 2, P. 263,* add the Seconds, answering to the Moon's Distance from her Apogee, and according to her Altitude, to the horizontal Diameter, and the Sum will be the apparent Diameter in Altitude, required.

PREC. IX. To find the MOON's true and also apparent DIAMETER in LONGITUDE and RIGHT ASCENSION?

As Cos. ☾'s Latitude or Declination : is to Rad. :: { so is ☾'s horizontal Diameter : to her true Diameter } in Long. or R.A. respectively.
 { . . . app. Diam. in Altitude : app. Diameter . . . }

See *Tab. 1, P. 263,* for Equation of true to apparent Diameter in Longitude and right Ascension, respectively, according to ☾'s Lat. or Decl. The Supplemental or Auxiliary Tables following are for different Methods of Computation.

SUPPLEMENTAL SOLAR EQUATIONS:

For determining the SUN's true Place from the mean, according to different *Radixes*, and *Hypotheses* of the Earth's Orbit.

EQUATION of the SUN's CENTER, according to *Eccentricity* of the Earth's Orbit 1692, and mean Distance à Sun, 100000.

Argument. SUN's Mean ANOMALY.

☉ M. An.	Sig. 0	Dif.	Sig. 1	Dif.	Sig. 2	Dif.	Sig. 3	Dif.	Sig. 4	Dif.	Sig. 5	Dif.	☉ M. An.
0	0 0 0	1 59	0 57 7	1 44	1 39 41	1 1	1 56 19	0 1	1 41 48	1 0	0 59 15	1 49	30
1	0 1 59	2 0	0 58 51	1 42	1 40 42	1 59	1 56 20	0 1	1 40 48	1 2	0 57 26	1 49	29
2	0 3 59	1 59	1 0 33	1 42	1 41 41	0 58	1 56 19	0 1	1 39 46	1 5	0 55 37	1 49	28
3	0 5 58	1 59	1 2 15	1 40	1 42 39	0 56	1 56 16	0 3	1 38 41	1 6	0 53 48	1 49	27
4	0 7 57	1 59	1 3 55	1 40	1 43 35	0 54	1 56 12	0 4	1 37 35	1 8	0 51 57	1 51	26
5	0 9 56	1 59	1 5 35	1 39	1 44 29	0 52	1 56 5	0 7	1 36 27	1 10	0 50 5	1 52	25
6	0 11 55	1 58	1 7 14	1 37	1 45 21	0 50	1 55 56	0 9	1 35 17	1 12	0 48 13	1 52	24
7	0 13 53	1 58	1 8 51	1 36	1 46 11	0 49	1 55 45	0 11	1 34 5	1 13	0 46 19	1 54	23
8	0 15 51	1 58	1 10 27	1 34	1 47 0	0 46	1 55 31	0 14	1 32 52	1 15	0 44 25	1 54	22
9	0 17 49	1 58	1 12 1	1 34	1 47 46	0 45	1 55 15	0 16	1 31 37	1 17	0 42 30	1 55	21
10	0 19 47	1 58	1 13 35	1 32	1 48 31	0 42	1 54 58	0 17	1 30 20	1 19	0 40 34	1 56	20
11	0 21 45	1 57	1 15 7	1 31	1 49 13	0 40	1 54 38	0 20	1 29 1	1 20	0 38 37	1 57	19
12	0 23 42	1 56	1 16 38	1 29	1 49 53	0 39	1 54 16	0 22	1 27 41	1 22	0 36 40	1 57	18
13	0 25 38	1 57	1 18 7	1 29	1 50 32	0 37	1 53 52	0 24	1 26 19	1 24	0 34 41	1 59	17
14	0 27 35	1 55	1 19 36	1 27	1 51 9	0 35	1 53 26	0 26	1 24 55	1 25	0 32 42	1 59	16
15	0 29 30	1 55	1 21 3	1 25	1 51 44	0 33	1 52 58	0 28	1 23 30	1 27	0 30 43	1 59	15
16	0 31 25	1 55	1 22 28	1 23	1 52 17	0 31	1 52 27	0 31	1 22 3	1 28	0 28 43	2 0	14
17	0 33 20	1 54	1 23 51	1 23	1 52 48	0 28	1 51 55	0 32	1 20 35	1 30	0 26 42	2 1	13
18	0 35 14	1 54	1 25 14	1 21	1 53 16	0 27	1 51 20	0 35	1 19 5	1 32	0 24 41	2 1	12
19	0 37 8	1 53	1 26 35	1 20	1 53 43	0 24	1 50 44	0 39	1 17 33	1 33	0 22 39	2 2	11
20	0 39 1	1 52	1 27 55	1 18	1 54 7	0 23	1 50 5	0 39	1 16 0	1 34	0 20 37	2 2	10
21	0 40 53	1 51	1 29 13	1 16	1 54 30	0 20	1 49 24	0 41	1 14 26	1 36	0 18 34	2 3	9
22	0 42 44	1 51	1 30 29	1 14	1 54 50	0 19	1 48 42	0 42	1 12 50	1 37	0 16 31	2 3	8
23	0 44 35	1 50	1 31 43	1 13	1 55 9	0 16	1 47 57	0 45	1 11 13	1 39	0 14 28	2 4	7
24	0 46 25	1 49	1 32 56	1 12	1 55 25	0 14	1 47 11	0 46	1 9 34	1 40	0 12 24	2 4	6
25	0 48 14	1 48	1 34 8	1 10	1 55 39	0 12	1 46 22	0 49	1 7 54	1 41	0 10 21	2 4	5
26	0 50 2	1 48	1 35 18	1 8	1 55 51	0 10	1 45 31	0 51	1 6 13	1 42	0 8 17	2 4	4
27	0 51 50	1 46	1 36 26	1 7	1 56 1	0 8	1 44 38	0 53	1 4 31	1 44	0 6 13	2 4	3
28	0 53 36	1 46	1 37 33	1 5	1 56 9	0 6	1 43 43	0 55	1 2 47	1 45	0 4 9	2 5	2
29	0 55 22	1 45	1 38 38	1 3	1 56 15	0 4	1 42 47	0 56	1 1 2	1 47	0 2 4	2 4	1
30	0 57 7		1 39 41		1 56 19		1 41 48	0 59	0 59 15		0 0 0		0
☉ M. An.	Sig. 11	Dif.	Sig. 10	Dif.	Sig. 9	Dif.	Sig. 8	Dif.	Sig. 7	Dif.	Sig. 6	Dif.	☉ M. An.

THE above Equation is according to the *Newtonian Hypothesis* of the Earth's Orbit, and the *Halleian* Computation.

To compute the Sun's Place for the Meridian of Greenwich Observatory from our Tables, according to any other Tables, correspondent to our mean Motions, and the above Hypothesis of the Earth's Orbit?

RULE. Note the Difference + — in a Table between the given *Radixes* of Sun's Place and Apogee for any common or Leap Year, and our *Radixes* (reduced to the Meridian of Greenwich) for the same Year, and connect the same Differences with the mean Places of our Tables for any Time required; and then work with the above Equation Table as with any other, for the proper Mean Anomaly of the Sun.

EXAMPLE. Connect with Mean Places

☉	Apogee	According to
— 3"	— 0' 0"	Newton,
— 10	— 5 10'	Halley,
+ 10	+ 2 0	Morris,

} Then use the Equation above.

SUPPLEMENTAL SOLAR TABLES.

LOGARITHMS of the SUN's foregoing proportional Distances from the EARTH, to the mean Distance 100000. And Eccentricity of the Earth's Orbit, 1692.

Argument. SUN's Mean ANOMALY.

☉ M. An.	Sig. 0. Logar.	Dif.	Sig. 1 Logar.	Dif.	Sig. 2 Logar.	Dif.	Sig. 3 Logar.	Dif.	Sig. 4 Logar.	Dif.	Sig. 5 Logar.	Dif.	☉ M. An.
0	5,007286		5,006347	63	5,003749	109	5,000124	129	4,996405	113	4,993620	65	30
1	5,007285	1	5,006284	64	5,003640	109	4,999995	128	4,996292	112	4,993555	64	29
2	5,007282	3	5,006220	66	5,003531	111	4,999867	128	4,996180	111	4,993491	62	28
3	5,007277	5	5,006154	67	5,003420	113	4,999739	128	4,996069	110	4,993429	60	27
4	5,007269	8	5,006087	69	5,003307	113	4,999611	128	4,995959	109	4,993369	58	26
5	5,007260	11	5,006018	72	5,003194	114	4,999483	128	4,995850	108	4,993311	56	25
6	5,007249	14	5,005946	74	5,003080	115	4,999355	128	4,995742	106	4,993255	54	24
7	5,007235	17	5,005872	75	5,002965	116	4,999227	128	4,995636	105	4,993201	51	23
8	5,007218	18	5,005797	77	5,002849	117	4,999099	128	4,995531	104	4,993150	48	22
9	5,007200	20	5,005720	78	5,002732	118	4,998971	127	4,995427	102	4,993102	47	21
10	5,007180	22	5,005642	80	5,002614	119	4,998844	127	4,995325	101	4,993055	46	20
11	5,007158	24	5,005562	82	5,002495	120	4,998717	127	4,995224	98	4,993009	43	19
12	5,007134	27	5,005480	83	5,002375	120	4,998590	127	4,995126	98	4,992966	40	18
13	5,007107	28	5,005397	85	5,002255	121	4,998463	127	4,995028	96	4,992926	38	17
14	5,007079	31	5,005312	87	5,002134	122	4,998336	126	4,994932	96	4,992888	36	16
15	5,007048	33	5,005225	89	5,002012	122	4,998210	126	4,994836	93	4,992852	34	15
16	5,007015	35	5,005136	89	5,001890	123	4,998084	124	4,994743	91	4,992818	32	14
17	5,006980	37	5,005047	91	5,001767	124	4,997960	123	4,994652	90	4,992786	29	13
18	5,006943	38	5,004956	93	5,001643	125	4,997837	123	4,994562	88	4,992757	26	12
19	5,006905	41	5,004863	95	5,001518	125	4,997714	123	4,994474	87	4,992731	25	11
20	5,006864	43	5,004768	96	5,001393	125	4,997591	123	4,994387	85	4,992706	23	10
21	5,006821	45	5,004672	97	5,001268	126	4,997468	121	4,994302	83	4,992683	20	9
22	5,006776	46	5,004575	98	5,001142	126	4,997347	121	4,994219	81	4,992663	17	8
23	5,006730	49	5,004477	100	5,001016	127	4,997226	120	4,994138	80	4,992646	15	7
24	5,006681	51	5,004377	102	5,000889	127	4,997106	119	4,994058	78	4,992631	13	6
25	5,006630	53	5,004275	102	5,000762	127	4,996987	119	4,993980	76	4,992618	11	5
26	5,006577	55	5,004173	104	5,000635	127	4,996868	118	4,993904	73	4,992607	8	4
27	5,006522	56	5,004069	106	5,000508	128	4,996750	116	4,993831	72	4,992599	6	3
28	5,006466	58	5,003963	106	5,000380	128	4,996634	115	4,993759	71	4,992593	3	2
29	5,006408	61	5,003857	108	5,000252	128	4,996519	114	4,993688	68	4,992590	1	1
30	5,006347		5,003749		5,000124		4,996405		4,993620		4,992589		0
☉ M. An.	Sig. 11 Logar.	Dif.	Sig. 10 Logar.	Dif.	Sig. 9 Logar.	Dif.	Sig. 8 Logar.	Dif.	Sig. 7 Logar.	Dif.	Sig. 6 Logar.	Dif.	☉ M. An.

The above TABLE is according to the *Newtonian Hypothesis* of the Earth's Orbit, and *Halleian Computation*. By which, and any other TABLE, for a different Eccentricity of the Earth's Orbit, the Logarithms of the *Proportional Distances*, or Distances themselves, may be determined for any third TABLE, to the same Degrees of the Sun's mean Anomaly, and other Eccentricity, different from the two former Eccentricities.

EXAMPLE. 1° 20' Sun M. An. { Log^s. 5,004768
5,004734 { Eccen^s. 1692
1680 { Prop. Dis^s. 101104
101096

$$\frac{1}{2} + \frac{1}{8} \text{ of } \frac{1}{2} = -20$$

Answer . . . 5,004748

1685

101100

To find for 1685 Eccentricity, or 7. of 12 left? Or $\frac{1}{2} + \frac{1}{8}$ of $\frac{1}{2}$ of Differences taken from first Quantity.

N. B. The Logarithms must be taken to 7 Places; to be correspondent in the 6th Place.

SUPPLEMENTAL

SUPPLEMENTAL SOLAR TABLES.

EQUATION of the SUN'S CENTER, according to Eccentricity of the Earth's Orbit, 1680, and Mean Distance of the SUN from the Earth 100000.

Argument. SUN'S Mean ANOMALY.

☉ M. An.	Sig. 0	Dif.	Sig. 1	Dif.	Sig. 2	Dif.	Sig. 3	Dif.	Sig. 4	Dif.	Sig. 5	Dif.	☉ M. An.
0	0 0 0	1 59	0 56 43	1 42	1 38 58	1 1	1 55 29	0 1	1 41 4	1 0	0 58 49	1 47	30
1	0 1 59	1 59	0 58 25	1 41	1 39 59	0 59	1 55 30	0 1	1 40 4	1 4	0 57 2	1 48	29
2	0 3 58	1 58	1 0 6	1 41	1 40 58	0 57	1 55 29	0 3	1 39 2	1 4	0 55 14	1 49	28
3	0 5 56	1 58	1 1 47	1 40	1 41 55	0 55	1 55 26	0 2	1 37 58	1 6	0 53 25	1 50	27
4	0 7 54	1 58	1 3 27	1 39	1 42 50	0 54	1 55 21	0 6	1 36 52	1 7	0 51 35	1 50	26
5	0 9 52	1 58	1 5 6	1 38	1 43 44	0 52	1 55 15	0 9	1 35 45	1 10	0 49 45	1 51	25
6	0 11 50	1 58	1 6 44	1 37	1 44 36	0 50	1 55 6	0 11	1 34 35	1 11	0 47 54	1 52	24
7	0 13 48	1 57	1 8 21	1 36	1 45 26	0 48	1 54 55	0 13	1 33 24	1 13	0 46 2	1 53	23
8	0 15 45	1 57	1 9 57	1 34	1 46 14	0 46	1 54 42	0 16	1 32 11	1 15	0 44 9	1 55	22
9	0 17 42	1 57	1 11 31	1 33	1 47 0	0 44	1 54 26	0 18	1 30 56	1 16	0 42 14	1 56	21
10	0 19 39	1 57	1 13 4	1 32	1 47 44	0 42	1 54 8	0 20	1 29 40	1 18	0 40 18	1 56	20
11	0 21 36	1 56	1 14 36	1 30	1 48 26	0 40	1 53 48	0 22	1 28 22	1 20	0 38 22	1 57	19
12	0 23 32	1 56	1 16 6	1 29	1 49 6	0 38	1 53 26	0 24	1 27 2	1 21	0 36 25	1 57	18
13	0 25 28	1 55	1 17 35	1 27	1 49 44	0 36	1 53 2	0 26	1 25 41	1 23	0 34 28	1 58	17
14	0 27 23	1 55	1 19 2	1 26	1 50 20	0 35	1 52 36	0 27	1 24 18	1 25	0 32 30	1 59	16
15	0 29 18	1 54	1 20 28	1 25	1 50 55	0 33	1 52 9	0 30	1 22 53	1 26	0 30 31	1 59	15
16	0 31 12	1 54	1 21 53	1 23	1 51 28	0 31	1 51 39	0 32	1 21 27	1 28	0 28 32	2 0	14
17	0 33 6	1 53	1 23 16	1 22	1 51 59	0 29	1 51 7	0 34	1 19 59	1 29	0 26 32	2 0	13
18	0 34 59	1 53	1 24 38	1 20	1 52 28	0 27	1 50 33	0 36	1 18 30	1 31	0 24 32	2 1	12
19	0 36 52	1 52	1 25 58	1 19	1 52 55	0 24	1 49 57	0 39	1 16 59	1 32	0 22 31	2 2	11
20	0 38 44	1 51	1 27 17	1 17	1 53 19	0 22	1 49 18	0 41	1 15 27	1 34	0 20 29	2 2	10
21	0 40 35	1 51	1 28 34	1 15	1 53 41	0 20	1 48 37	0 43	1 13 53	1 35	0 18 27	2 2	9
22	0 42 26	1 50	1 29 50	1 14	1 54 1	0 18	1 47 54	0 44	1 12 18	1 37	0 16 25	2 2	8
23	0 44 16	1 49	1 31 4	1 13	1 54 19	0 16	1 47 10	0 46	1 10 41	1 38	0 14 23	2 3	7
24	0 46 5	1 49	1 32 17	1 11	1 54 35	0 15	1 46 24	0 48	1 9 3	1 39	0 12 20	2 3	6
25	0 47 54	1 48	1 33 28	1 9	1 54 50	0 12	1 45 36	0 50	1 7 24	1 40	0 10 17	2 3	5
26	0 49 42	1 47	1 34 37	1 8	1 55 2	0 10	1 44 46	0 52	1 5 44	1 42	0 8 14	2 3	4
27	0 51 29	1 46	1 35 45	1 6	1 55 12	0 8	1 43 54	0 54	1 4 2	1 43	0 6 11	2 3	3
28	0 53 15	1 45	1 36 51	1 4	1 55 20	0 6	1 43 0	0 57	1 2 19	1 44	0 4 8	2 4	2
29	0 55 0	1 43	1 37 55	1 3	1 55 26	0 3	1 42 3	0 59	1 0 35	1 46	0 2 4	2 4	1
30	0 56 43		1 38 58		1 55 29		1 41 4		0 58 49		0 0 0		0
☉ M. An.	+	Dif.	+	Dif.	+	Dif.	+	Dif.	+	Dif.	+	Dif.	☉ M. An.
	Sig. 11		Sig. 10		Sig. 9		Sig. 8		Sig. 7		Sig. 6		

THE above Equation is according to the *Parisian Hypothesis* of the Earth's Orbit, and Computation of the Celebrated *Tobias Mayer*, of *Gottingen*, in *Germany*.

The Sun's Place for the Meridian of *Greenwich Observatory*, according to *Mr. Mayer's Tables*, is computed from our Tables, by adding 15" to Mean Place Sun, and 1' to M. Place of *Apogee* Sun, by our Tables, and then using the above Equation, correspondent to the proper M. Anomaly Sun.

N. B. Mr. Mayer's Radixes for Paris 1701,	☉ Place.	☉ Apogee.
Reduce to the Merid. of Greenwich	9° 20' 43' 35"	3° 7' 45' 30"
	W. Motion + 23	
Mayer's Correspondent Radix at Greenwich	9 20 43 58	1
Our Radixes 1701	9 20 43 43	9 7 44 30
Our Radixes reduce to Mayer's by	+ 15	+ 1 0

Difference.

SUPPLEMENTAL SOLAR TABLES.

SUN's PROPORTIONAL DISTANCES from the EARTH, according to the foregoing Eccentricity 1680, and mean Distance 100000.

Argument. SUN's mean ANOMALY.

☉ M. An.	Sig. 0 ☉ à ☿	Dif.	Sig. 1 ☉ à ☿	Dif.	Sig. 2 ☉ à ☿	Dif.	Sig. 3 ☉ à ☿	Dif.	Sig. 4 ☉ à ☿	Dif.	Sig. 5 ☉ à ☿	Dif.	☉ M. An.
0	101680	0	101462	14	100861	25	100028	30	99182	26	98553	15	30
1	101680	1	101448	15	100836	25	99998	30	99156	26	98538	15	29
2	101679	1	101433	15	100811	26	99968	29	99131	25	98524	14	28
3	101678	1	101418	16	100785	26	99939	29	99106	25	98510	14	27
4	101676	2	101402	17	100759	26	99910	29	99081	25	98496	14	26
5	101674	2	101385	17	100733	26	99881	29	99056	25	98483	13	24
6	101671	3	101368	17	100707	27	99852	29	99031	25	98470	13	25
7	101668	3	101351	17	100680	27	99823	29	99007	24	98458	12	23
8	101664	4	101334	17	100653	27	99794	29	98983	24	98446	12	22
9	101660	4	101316	18	100626	27	99765	29	98960	23	98435	11	21
10	101655	5	101298	18	100599	27	99736	29	98937	23	98425	10	20
11	101650	5	101280	18	100572	27	99707	29	98914	23	98415	10	19
12	101644	6	101261	19	100545	27	99678	29	98892	22	98405	10	18
13	101638	6	101242	19	100517	28	99649	29	98870	22	98396	9	17
14	101631	7	101222	20	100489	28	99620	29	98848	22	98387	9	16
15	101624	7	101202	20	100461	28	99591	29	98827	21	98379	8	15
16	101617	7	101182	20	100433	28	99563	28	98806	21	98371	8	14
17	101609	8	101161	21	100405	28	99535	28	98785	21	98364	7	13
18	101601	8	101140	21	100377	29	99507	28	98765	20	98358	6	12
19	101592	9	101118	22	100348	29	99479	28	98745	20	98352	6	11
20	101582	10	101096	22	100319	29	99451	28	98725	20	98347	5	10
21	101572	10	101074	22	100290	29	99423	28	98706	19	98342	5	9
22	101561	11	101051	23	100261	29	99495	27	98687	19	98338	4	8
23	101550	11	101028	23	100232	29	99468	27	98669	18	98334	4	7
24	101539	11	101005	23	100203	29	99441	27	98651	18	98330	4	6
25	101527	12	100982	23	100174	29	99414	27	98634	17	98327	3	5
26	101515	12	100959	23	100145	29	99387	27	98617	17	98324	3	4
27	101502	13	100935	24	100116	29	99360	26	98600	16	98322	2	3
28	101489	13	100911	25	100087	29	99334	26	98584	16	98321	1	2
29	101476	13	100886	25	100058	30	99308	26	98568	15	98320	1	1
30	101462	14	100861	25	100028		99182		98553		98320	0	0
☉ M. An.	Sig. 11 ☉ à ☿	Dif.	Sig. 10 ☉ à ☿	Dif.	Sig. 9 ☉ à ☿	Dif.	Sig. 8 ☉ à ☿	Dif.	Sig. 7 ☉ à ☿	Dif.	Sig. 6 ☉ à ☿	Dif.	☉ M. An.

THE above TABLE is according to the *Parisian Hypothesis*, of the Earth's Orbit, and the Celebrated *Mayer's* Computation.

$$\begin{array}{c} \text{☉ M. An.} \left\{ \begin{array}{l} \text{Equations,} \\ 1^{\circ} 21' 3'' \\ 1^{\circ} 15' 0'' \end{array} \right. \quad \begin{array}{l} \text{Ecc.} \\ 1692 \\ 1680 \end{array} \quad \begin{array}{l} \text{Halley} \\ \text{Mayer} \end{array} \quad \begin{array}{l} \text{Greatest Equ} \\ 1^{\circ} 56' 20'' \\ 1^{\circ} 55' 30'' \end{array} \quad \text{☉ M. An.} \left\{ \begin{array}{l} 3^{\circ} 1^{\circ} \end{array} \right.$$

$$\frac{1}{2} + \frac{1}{6} \text{ of } \frac{1}{2} \text{ Dif.} = \frac{35}{21} \quad \frac{12}{7} \quad \frac{50}{29} \text{ Dif. from 1st Quantities}$$

$$\text{Equation } \text{☉ Center } 1 \ 20 \ 42 \quad 1685 \text{ Our Tables. } 1 \ 55 \ 53$$

N. B. Tables are otherwise depressed or elevated in Proportion to the Differences of their respective Maxima, or greatest Quantities, similar to the Rule of different Eccentricities.

Thus are all the necessary Uses of the foregoing Supplemental Solar Tables explained.

SUPPLEMENTAL

SUPPLEMENTAL LUNAR TABLES:

For determining the True from the MOON's Mean PLACE, according to different METHODS and THEORIES.

I. EQUATION of the MOON, APOGEE and NODE.

Argument. SUN's Mean ANOMALY.

M. An.	Sig. 0						Sig. 1						Sig. 2						M. An.
	D	Dif.	Ap.	Dif.	8	Dif.	D	Dif.	Ap.	Dif.	8	Dif.	D	Dif.	Ap.	Dif.	8	Dif.	
0	0 0		0 0		0 0		5 47		9 49		4 40		10 7		17 8		8 9		30
1	0 12	12	0 20	20	0 9	9	5 58	11	10 7	17	4 48	8	10 14	7	17 19	10	8 14	5	29
2	0 24	12	0 41	20	0 19	10	6 9	10	10 24	18	4 57	9	10 20	6	17 29	10	8 19	5	28
3	0 36	12	1 1	21	0 29	10	6 19	10	10 42	17	5 5	8	10 26	6	17 39	9	8 24	5	27
4	0 48	12	1 22	20	0 39	10	6 29	10	10 59	17	5 13	8	10 31	5	17 48	10	8 28	4	26
5	1 0	12	1 42	20	0 48	9	6 39	10	11 16	17	5 21	8	10 37	6	17 58	10	8 32	4	25
6	1 12	12	2 3	20	0 58	10	6 49	10	11 33	16	5 29	8	10 42	5	18 7	9	8 36	4	24
7	1 24	12	2 23	20	1 8	10	6 59	10	11 49	17	5 37	8	10 47	5	18 16	8	8 40	4	23
8	1 36	12	2 43	21	1 18	10	7 9	10	12 6	16	5 45	8	10 52	4	18 24	8	8 44	4	22
9	1 48	11	3 4	20	1 27	9	7 19	10	12 22	17	5 53	7	10 56	5	18 32	7	8 48	4	21
10	1 59	12	3 24	20	1 37	10	7 28	9	12 39	16	6 0	7	11 1	5	18 39	7	8 52	4	20
11	2 11	12	3 44	20	1 46	10	7 38	8	12 55	15	6 8	7	11 5	4	18 46	7	8 55	3	19
12	2 23	12	4 4	20	1 56	9	7 46	9	13 10	16	6 15	7	11 9	4	18 53	7	8 58	3	18
13	2 35	11	4 24	20	2 5	10	7 55	9	13 26	15	6 22	8	11 14	5	19 0	6	9 2	4	17
14	2 46	12	4 44	20	2 15	9	8 4	9	13 41	15	6 30	8	11 19	5	19 6	6	9 5	3	16
15	2 58	12	5 4	20	2 24	9	8 13	9	13 56	15	6 37	7	11 22	3	19 12	6	9 7	2	15
16	3 10	12	5 24	20	2 34	10	8 22	9	14 10	15	6 44	7	11 25	3	19 18	5	9 10	3	14
17	3 22	12	5 44	19	2 43	10	8 31	8	14 25	14	6 50	7	11 28	2	19 23	5	9 12	3	13
18	3 34	12	6 3	20	2 53	9	8 39	8	14 39	14	6 57	7	11 30	2	19 28	5	9 15	2	12
19	3 46	11	6 23	20	3 2	9	8 47	8	14 53	14	7 4	7	11 32	3	19 33	4	9 17	2	11
20	3 57	12	6 43	19	3 11	9	8 55	8	15 7	14	7 11	6	11 35	3	19 37	4	9 19	2	10
21	4 9	11	7 2	19	3 20	9	9 3	8	15 21	12	7 17	6	11 38	2	19 41	3	9 21	1	9
22	4 20	11	7 21	19	3 29	9	9 11	8	15 33	13	7 23	6	11 40	2	19 44	4	9 22	2	8
23	4 31	11	7 40	19	3 38	9	9 19	8	15 46	12	7 29	6	11 42	1	19 48	3	9 24	3	7
24	4 42	11	7 59	19	3 47	9	9 27	7	15 58	13	7 35	6	11 43	2	19 51	2	9 25	1	6
25	4 53	11	8 18	18	3 56	9	9 34	7	16 11	12	7 41	5	11 45	1	19 53	2	9 26	1	5
26	5 4	11	8 36	18	4 5	9	9 41	6	16 23	12	7 46	5	11 46	1	19 55	2	9 27	1	4
27	5 15	11	8 54	19	4 14	9	9 47	7	16 35	11	7 52	6	11 47	1	19 57	1	9 28	1	3
28	5 26	11	9 13	18	4 23	8	9 54	7	16 46	11	7 58	6	11 48	0	19 58	1	9 29	0	2
29	5 37	10	9 31	18	4 31	9	10 1	6	16 57	11	8 4	5	11 48	1	19 59	1	9 29	0	1
30	5 47		9 49		4 40		10 7		17 8		8 9		11 49		20 0		9 30		0
M. An.	D	Dif.	Ap.	Dif.	8	Dif.	D	Dif.	Ap.	Dif.	8	Dif.	D	Dif.	Ap.	Dif.	8	Dif.	M. An.
	Sig. 11						Sig. 10						Sig. 9						

THE above Equations are according to the Newtonian Theory, and Halleian Computation. See Schol. Prop. 35. B. 3. Principia, Last Edit. p. 299.

Sir ISAAC NEWTON finds the greatest Equation $\mathcal{D}$ from the Theory of Gravity, $11' 49''$; but thinks, as the Earth's Eccentricity is something greater than he supposed ($16\frac{1}{2}$ to 1000, Rad. or Mean Dist. $\odot$ à $\ominus$) viz. $16\frac{1}{12}$ Eccentricity, the greatest Equation will be about $11' 51''$.

He found that the Apogee and Nodes moved faster in Perib. (where the Force of $\odot$'s Action is greater) than in Aphel.—That this Motion of the Ap. and Node is in the reciprocal triplicate Proportion of $\ominus$'s Dist. from $\odot$. That the greatest Equation $\odot$ Center, generated by $\odot$'s Motion, in reciprocal duplicate Proportion $\odot$ à $\ominus$, is $1^\circ 56' 20''$, correspondent to Eccentricity $\ominus$'s Orbit $16\frac{1}{12}$ aforesaid, which would be $2^\circ 54' 30''$ by a reciprocal triplicate Proportion of Distance of the Sun à Earth. And therefore the greatest Equations generated by the $\mathcal{D}$'s Ap.

SUPPLEMENTAL LUNAR TABLES.

I. EQUATION of the MOON, APOGEE and NODE.

Argument. SUN's Mean ANOMALY.

M. An.	Sig. 3				Sig. 4				Sig. 5				M. An.
	D	Dif.	Ap.	Q	D	Dif.	Ap.	Q	D	Dif.	Ap.	Q	
0	11 49	0	20 0	9 30	10 21	6	17 30	8 19	0 0	11	10 11	4 50	30
1	11 49	0	20 0	9 30	10 15	8	17 20	8 14	5 49	11	9 52	4 41	29
2	11 49	1	20 0	9 30	10 7	6	17 9	8 9	5 38	11	9 33	4 32	28
3	11 48	0	19 59	9 30	10 1	6	16 58	8 4	5 27	11	9 15	4 23	27
4	11 48	1	19 58	9 29	9 55	8	16 46	7 58	5 16	11	8 56	4 14	26
5	11 47	0	19 57	9 29	9 47	7	16 34	7 53	5 5	12	8 37	4 5	25
6	11 47	2	19 56	9 28	9 40	7	16 23	7 48	4 53	11	8 17	3 56	24
7	11 45	1	19 54	9 27	9 33	7	16 11	7 42	4 42	12	7 58	3 47	23
8	11 44	2	19 51	9 26	9 26	7	15 58	7 35	4 30	12	7 38	3 38	22
9	11 42	1	19 49	9 25	9 19	8	15 45	7 29	4 18	12	7 18	3 29	21
10	11 41	2	19 46	9 24	9 11	9	15 32	7 23	4 6	12	6 58	3 19	20
11	11 39	3	19 42	9 22	9 2	8	15 18	7 16	3 54	11	6 38	3 9	19
12	11 36	2	19 38	9 20	8 54	8	15 4	7 9	3 43	12	6 18	3 59	18
13	11 34	3	19 34	9 18	8 46	9	14 50	7 3	3 31	12	5 58	2 50	17
14	11 31	2	19 30	9 16	8 37	9	14 56	6 56	3 19	12	5 38	2 40	16
15	11 29	4	19 25	9 14	8 28	8	14 21	6 49	3 7	13	5 17	2 30	15
16	11 25	3	19 20	9 12	8 20	9	14 6	6 42	2 54	12	4 56	2 20	14
17	11 22	3	19 14	9 9	8 11	10	13 51	6 35	2 42	12	4 35	2 10	13
18	11 19	3	19 8	9 6	8 1	9	13 35	6 27	2 30	12	4 14	2 0	12
19	11 15	4	19 2	9 3	7 52	9	13 20	6 20	2 18	13	3 54	1 50	11
20	11 11	4	18 55	8 59	7 43	9	13 4	6 12	2 5	13	3 33	1 40	10
21	11 6	5	18 48	8 56	7 34	10	12 48	6 4	1 52	12	3 12	1 30	9
22	11 2	4	18 41	8 52	7 24	10	12 31	5 56	1 40	12	2 51	1 20	8
23	10 58	4	18 33	8 48	7 14	11	12 15	5 49	1 28	13	2 30	1 10	7
24	10 53	5	18 25	8 45	7 3	10	11 58	5 41	1 15	13	2 8	1 0	6
25	10 48	5	18 17	8 41	6 53	10	11 41	5 33	1 2	13	1 46	0 50	5
26	10 43	5	18 8	8 37	6 43	10	11 23	5 24	0 49	12	1 25	0 40	4
27	10 38	6	17 59	8 33	6 33	11	11 5	5 16	0 37	12	1 4	0 30	3
28	10 32	6	17 50	8 28	6 22	11	10 47	5 8	0 25	13	0 43	0 20	2
29	10 26	5	17 40	8 23	6 11	11	10 29	4 59	0 12	12	0 21	0 10	1
30	10 21	5	17 30	8 19	6 0	10	10 11	4 50	0 0	12	0 0	0 0	0
M. An.	D	Dif.	Ap.	Q	D	Dif.	Ap.	Q	D	Dif.	Ap.	Q	M. An.

D's Ap. and Q, are to $2^{\circ} 54' 30''$, as the M. diurnal Mot. D's Apog. and of her Q respectively are to M. diurnal Mot. $\odot$. — Hence gr. Equation, D's Apogee comes out $19' 43''$. And gr. Eq. of Moon's Qs $9' 24''$; which Dr. Halley raised to $20'$ and $9' 30''$. But D. Cooper fixed the gr. Equations D, Ap. and Q, at $11' 25''$; $25' 45''$; and $9' 24''$, respectively, as has been shewn, for correcting the Mean Erection-Eqn. or Mean Difference between the Mean and True Central Equation. Hence arise annual Equations of these three Motions, directly proportional to the Equations of $\odot$'s Center, as before shewn.

As $1^{\circ} 56' 20''$ } S. Sun's Present Equation } $\left\{ \begin{array}{l} 11' 49'' \\ 20' 0'' \\ 9' 30'' \end{array} \right\}$ } Decimals. Log. Log. Lo. Log. $\left\{ \begin{array}{l} Sx, 1016 = Eq. D = 0.9932 \\ Sx, 1719 = Eq. Ap. = 0.7646 \\ Sx, 0817 = Eq. Q = 1.0879 \end{array} \right\} \pm \text{Log. L. L. S. } \left\{ \begin{array}{l} \text{under} \\ \text{above} \end{array} \right\} 60'.$

SUPPLEMENTAL

and NAVIGATOR. SUPPLEMENTAL LUNAR TABLES.

63

H. EQUATION of the Moon: in Apogee and Perigee.

1. Semiannual.

Arg. Sun from Moon's Apogee.

⊙	Sig.	G.D.	Sig.	G.D.	Sig.	G.D.	⊙
⊙	abo.	1	7	abo.	2	8	⊙
⊙	L. eq.		L. eq.		L. eq.		⊙
ap.							ap.
0	0	0	3	5	19	3	5
1	0	7	1	3	9	19	3
2	0	15	1	3	12	20	2
3	0	22	2	3	15	20	2
4	0	30	3	3	18	20	2
5	0	37	4	3	21	21	2
6	0	45	4	3	24	21	2
7	0	52	5	3	26	21	2
8	0	59	6	3	28	21	2
9	1	6	7	3	29	21	2
10	1	13	8	3	31	22	2
11	1	20	8	3	32	22	2
12	1	27	9	3	33	22	2
13	1	34	10	3	33	22	2
14	1	40	10	3	34	22	1
15	1	47	11	3	34	22	1
16	1	53	12	3	34	22	1
17	2	0	12	3	33	22	1
18	2	6	13	3	33	22	1
19	2	12	14	3	32	22	1
20	2	18	14	3	31	22	1
21	2	23	15	3	29	21	1
22	2	29	15	3	28	21	0
23	2	34	16	3	26	21	0
24	2	39	16	3	24	21	0
25	2	44	17	3	21	21	0
26	2	49	17	3	18	20	0
27	2	53	18	3	15	20	0
28	2	57	18	3	12	20	0
29	3	1	19	3	9	19	0
30	3	5	19	3	5	19	0
⊙	+	G.D.	+	G.D.	+	G.D.	⊙
⊙	11	5	abo.	10	4	abo.	9
⊙	Sig.	L. eq.	Sig.	L. eq.	Sig.	L. eq.	⊙
ap.							ap.

Construction. Where Δ Ap. is without the
Ofants of Sun, as above, As Rad. : S. doub. Dif.
Sun Δ Δ Ap. (i.e. fr. nearest Quadrature or
Syzygy) : S. greatest Eq. : S. pref. Eq. i.e.
Rad. : S. $3' 34''$: S. double Sun Δ Δ Ap. : S.
Pref. Eq. — Use. As $3' 34''$, gr. Eq. in Ofants
Sun in Ap. : Least Eq. : as above, Sun in Ap.
: Gr. Eq. in Ofants correspondt Sun M. An. :
Pref. true Equation. Or, As $3' 34''$: G. Δ :
P. Δ : G. Δ : X P. Δ :
3' 34'' = True Eq. Hence,
the Construction of L. Log^s. correspondent to
Sun M. An. in Equat. of Δ in Ofants.
Ex. To find the Pref. Equat. Δ , for Sun Δ Δ
Ap. $0^\circ 27'$, and $2^\circ 5'$ Sun M. Anomaly?
L. Eq. G.D.
Sun Δ Δ Ap. $0^\circ 27'$: $2' 53''$: $18''$ ab. L. Eq.
+ $5' 31''$ X^r for $2^\circ 5'$
Present Equation $2' 58''$ red 18 [Sun M. An.]

DECIMAL MULTIPLIERS to the Gr. Dif. above the Least Equat. for Sec^s. to be added thereto, for the present Equation Δ .

Arg. Sun's Mean Anom.

⊙	0	1	2	3	4	5	⊙
⊙	X ^{rs}	X ^{rs}	X ^{rs}	X ^{rs}	X ^{rs}	X ^{rs}	⊙
⊙	A.						⊙
0	00	07	27	5	77	95	30
1	00	07	28	51	78	96	29
2	00	08	29	52	79	96	28
3	01	08	30	52	79	96	27
4	01	08	30	53	80	97	26
5	01	09	31	54	81	97	25
6	01	09	32	55	82	97	24
7	01	11	33	55	82	97	23
8	02	11	33	56	83	98	22
9	02	11	34	57	83	98	21
10	02	12	35	58	83	98	20
11	02	13	36	58	84	98	19
12	02	14	36	59	84	99	18
13	03	14	37	60	85	99	17
14	03	15	38	61	85	99	16
15	03	16	39	61	85	99	15
16	03	17	40	62	86	1.	14
17	03	17	40	63	86	1.	13
18	03	18	41	64	86	1.	12
19	04	19	42	65	87	1.	11
20	04	20	42	66	88	1.	10
21	04	20	43	67	89	1.	9
22	04	21	44	68	89	1.	8
23	04	22	45	69	90	1.	7
24	05	23	46	71	91	1.	6
25	05	24	46	72	92	1.	5
26	05	24	47	73	92	1.	4
27	06	25	48	74	93	1.	3
28	06	26	49	75	94	1.	2
29	07	27	49	76	95	1.	1
30	07	27	5	77	95	1.	0
⊙	X ^{rs}	X ^{rs}	X ^{rs}	X ^{rs}	X ^{rs}	X ^{rs}	⊙
⊙	11	10	9	8	7	6	⊙
⊙	A.						⊙

Construction. As $22''$ greatest Dif.
of Equation in Ofants, Δ in Apogee
and Perigee : present Dif. Eq. beta Δ
in Ap. and according to M. An. Sun :
greatest Dif. of Equation, Sun in Apo-
gee, according to Argument : present
Dif. to be added to the Equation of
the Argument.
Ex. Let Sun M. An. 4° where the
present Dif. Equation from Least = $17''$
Then $22'' : 17'' :: G. D. : 77 \times$
G.D. Proportional Equation required.

EXAMPLE of the foregoing
Case.
L. Log.
2 53 Equation. 1,3183
For $2^\circ 5'$ A. Sun — 120
2' 58'' Eq. reqd. 1,3063
correct, as before.

Greatest Equat. being in Ofants of Sun.

Arg. Sun's M. An.

⊙	Gr.	Lo.	Sun
⊙	An.	Equ.	M.
⊙	An.	Log.	An.
0	03	34	0 0 12
1	03	34	0 24
2	03	34	0 18
3	03	34	0 12
4	03	35	20 6
5	03	35	20 0 11
6	03	36	41 24
7	03	37	61 18
8	03	37	61 12
9	03	38	81 6
10	03	39	100 0 10
11	03	40	120 24
12	03	41	140 18
13	03	42	160 12
14	03	44	198 6
15	03	45	218 0 9
16	03	46	237 24
17	03	47	256 18
18	03	48	275 12
19	03	50	313 6
20	03	51	332 0 8
21	03	52	351 24
22	03	53	370 18
23	03	54	388 12
24	03	54	388 6
25	03	55	407 0 7
26	03	56	425 24
27	03	56	425 18
28	03	56	425 12
29	03	56	425 6
30	03	56	425 0 6
⊙	Gr.	Lo.	Sun
⊙	M.	Eq.	M.
⊙	An.	Log.	An.

Construction. The above
Equation of $3' 34''$ in Ap.
Sun and $3' 56''$ in Perig.
Sun, increases or diminishes
in the inverse Ratio of the
3d Power of the Sun Δ Δ ,
i.e. in the reciprocal tri-
plicate Proportion. See
Schol. Prop. 35. B. 3. P.
300. Princip.

III. EQUATION of the MOON. Sun at M. Diff. from Earth. 2. Semi-annual.

Arg. Sun from Node.

⊙	Sig.	Sig.	Sig.	⊙
⊙	abo.	1	7	abo.
⊙	8			8
0	0	0	41	0 41 30
1	0	2	41	0 40 29
2	0	3	42	0 39 28
3	0	5	43	0 38 27
4	0	7	44	0 37 26
5	0	8	44	0 36 25
6	0	10	45	0 35 24
7	0	11	45	0 34 23
8	0	13	46	0 33 22
9	0	15	46	0 31 21
10	0	16	46	0 30 20
11	0	18	47	0 29 19
12	0	19	47	0 28 18
13	0	21	47	0 26 17
14	0	22	47	0 25 16
15	0	24	47	0 24 15
16	0	25	47	0 22 14
17	0	26	47	0 21 13
18	0	28	47	0 19 12
19	0	29	47	0 18 11
20	0	30	46	0 16 10
21	0	31	46	0 15 9
22	0	33	46	0 13 8
23	0	34	45	0 11 7
24	0	35	45	0 10 6
25	0	36	44	0 8 5
26	0	37	44	0 7 4
27	0	38	43	0 5 3
28	0	39	42	0 3 2
29	0	40	41	0 2 1
30	0	41	41	0 0 0
⊙	+	+	+	⊙
⊙	11	5	10	4 9 3
⊙	Sig.	Sig.	Sig.	⊙

Construction. As Rad. S.
47'' a Maxim. : S. doub.
Sun Δ Δ : S. Present Equat.
In other Distances of Sun
 Δ Δ , this Equation, greatest
in the Ofants of the Nodes,
is reciprocally as Cube Sun
 Δ Δ . And in Perigee is
49'' and in Apogee 45''.

N. B. The above Equa-
tion, and the former two of
the Supplemental are accord-
to the Newtonian Theory.

SUPPLEMENTAL LUNAR TABLES.

SECOND EQUATION of the MOON'S APOGEE, from the *Newtonian THEORY*. With *Decimal* and *Sexagesimal* MULTIPLIERS into the *Greatest Diff.* under or above the *Mean Central EQUATION* of the MOON, for directly finding the *Present correct Diff.* from the *Mean*, and thence the whole *Central EQUATION* of the MOON. According to our *NEW METHOD* of Computing.

Argument. SUN from the MOON'S APOGEE, or Sun's True Place — Moon's Apogee 1 Equated.

Sig. o. 6.										Sig. 1. 7.										Sig. 2. 8.																																	
⊙	2d Equ.	⊙	Apog.	Dif.	X ^{rs} for	D.	Sex ^l X ^s	D.	⊙	2d Equ.	⊙	Apog.	Dif.	X ^{rs} for	D.	Sex ^l X ^s	D.	⊙	2d Equ.	⊙	Apog.	Dif.	X ^{rs} for	D.	Sex ^l X ^s	D.	⊙	2d Equ.	⊙	Apog.	Dif.	X ^{rs} for	D.	Sex ^l X ^s	D.	⊙																	
ap.	+				above		above			+				above		above			+				under		under		under		+				under		under		ap.																
0	0	0	0		1.		60	0	2	9	28	8	14	15	5718	273	34	18	11	40	16		4114	325	24	41	30	0	0	0	0		4114	325	24	41	30	0	0	0	0												
1	0	21	5	21	5	9994	15	52	58	9	42	23	13	46	5445	280	32	40	11	30	55	9	21	4443	325	26	39	29	1	0	21	5	21	4443	325	26	39	29	1	0	21	5	21										
2	0	42	9	21	4	9979	24	59	52	9	56	9	13	16	5165	285	30	59	11	20	29	10	26	4768	320	28	36	28	2	0	42	9	21	4768	320	28	36	28	2	0	42	9	21										
3	1	3	11	21	2	9955	36	59	44	10	9	25	13	16	4880	293	29	17	11	8	59	11	30	5088	313	30	32	27	3	1	3	11	21	5088	313	30	32	27	3	1	3	11	21										
4	1	24	11	21	0	9919	36	59	31	10	22	9	12	12	4587	297	27	31	11	56	23	12	36	5401	308	32	24	26	4	1	24	11	21	5401	308	32	24	26	4	1	24	11	21										
5	1	45	7	20	56	9874	45	59	15	10	34	21	12	12	4290	297	25	44	11	42	41	13	42	5709	301	34	15	25	5	1	45	7	20	5709	301	34	15	25	5	1	45	7	20										
6	2	5	59	20	52	9819	55	58	55	10	45	59	11	38	3986	304	23	55	11	27	54	14	47	6010	301	36	4	24	6	2	5	59	20	6010	301	36	4	24	6	2	5	59	20										
7	2	26	46	20	47	9754	65	58	31	10	57	2	10	26	3678	308	22	4	11	12	0	15	54	6304	294	37	49	23	7	2	26	46	20	6304	294	37	49	23	7	2	26	46	20										
8	2	47	28	20	42	9679	75	58	4	11	7	28	9	49	3364	314	20	11	11	55	2	16	58	6590	286	39	32	22	8	2	47	28	20	6590	286	39	32	22	8	2	47	28	20										
9	3	8	3	20	35	9595	84	57	34	11	17	17	9	10	3046	318	18	17	11	36	58	18	4	6868	278	41	12	21	9	3	8	3	20	6868	278	41	12	21	9	3	8	3	20										
10	3	28	39	20	27	9500	95	57	0	11	26	27	9	10	2723	323	16	20	11	56	50	19	8	7137	269	42	49	20	10	3	28	39	20	7137	269	42	49	20	10	3	28	39	20										
11	3	48	49	20	19	9396	104	56	23	11	34	57	8	30	2396	327	14	23	11	57	38	20	12	7397	260	44	23	19	11	3	48	49	20	7397	260	44	23	19	11	3	48	49	20										
12	4	8	59	20	10	9283	113	55	42	11	42	45	7	48	2065	331	12	23	11	57	38	21	14	7397	250	44	23	18	12	4	8	59	20	7397	250	44	23	18	12	4	8	59	20										
13	4	28	59	20	0	9159	124	54	57	11	49	51	7	6	1731	334	10	23	11	57	38	22	16	7647	240	45	53	17	13	4	28	59	19	7647	240	45	53	17	13	4	28	59	19										
14	4	48	47	19	48	9026	133	54	9	11	56	12	6	21	1394	337	8	22	11	57	38	23	15	7887	229	47	19	16	14	4	48	47	19	7887	229	47	19	16	14	4	48	47	19										
15	5	8	24	19	37	8884	142	53	18	12	1	48	5	36	1054	340	6	19	12	1	48	24	13	8334	218	50	0	15	15	5	8	24	19	8334	218	50	0	15	15	5	8	24	19										
16	5	27	48	19	24	8733	151	52	24	12	6	37	4	49	0712	342	4	16	12	6	37	25	9	8540	206	51	14	14	16	5	27	48	19	8540	206	51	14	14	16	5	27	48	19										
17	5	46	58	18	56	8573	160	51	26	12	10	38	4	1	0367	345	2	12	18	56	58	26	2	8735	195	52	24	13	17	5	46	58	18	8735	195	52	24	13	17	5	46	58	18										
18	6	5	54	18	40	8403	170	50	25	12	13	50	3	12	0020	347	0	7	18	40	54	26	54	8917	182	53	30	11	18	6	5	54	18	8917	182	53	30	11	18	6	5	54	18										
19	6	24	34	18	40	8225	178	49	21	12	16	11	2	21	0327	347	1	58	18	40	54	27	42	9085	168	54	30	10	19	6	24	34	18	9085	168	54	30	10	19	6	24	34	18										
20	6	42	58	18	6	8038	187	48	14	12	17	40	0	35	0675	348	4	3	18	6	42	28	27	9240	155	55	26	9	20	6	42	58	18	9240	155	55	26	9	20	6	42	58	18										
21	7	1	4	17	48	7842	196	47	3	12	18	15	0	19	1024	349	6	9	17	1	4	29	9	9382	142	56	18	8	21	7	1	4	17	7842	196	47	3	12	18	15	0	19	1024	349	6	9	17	1	4	17	1		
22	7	18	52	17	26	7637	205	45	49	12	17	56	1	15	1373	349	8	14	17	52	52	29	47	9510	128	57	4	7	22	7	18	52	17	7637	205	45	49	12	17	56	1	15	1373	349	8	14	17	52	29	47	9510	128	
23	7	36	20	17	26	7425	212	44	33	12	16	41	2	12	1722	349	10	20	17	20	20	30	23	9623	113	57	44	6	23	7	36	20	17	7425	212	44	33	12	16	41	2	12	1722	349	10	20	17	20	30	23	9623	113	
24	7	53	28	16	46	7204	220	43	13	12	14	29	3	11	2069	347	12	25	16	46	28	31	53	9722	99	58	28	5	24	7	53	28	16	7204	220	43	13	12	14	29	3	11	2069	347	12	25	16	46	28	31	53	9722	99
25	8	10	14	16	24	6975	229	41	51	12	11	18	3	11	2416	347	14	30	16	24	14	31	21	9806	84	58	50	4	25	8	10	14	16	6975	229	41	51	12	11	18	3	11	2416	347	14	30	16	24	14	31	21	9806	84
26	8	26	38	16	1	6739	236	40	26	12	7	8	4	10	2762	346	16	34	16	38	38	32	1	9876	70	59	15	3	26	8	26	38	16	6739	236	40	26	12	7	8	4	10	2762	346	16	34	16	38	32	1	9876	70	
27	8	42	39	15	36	6494	245	38	58	12	1	58	5	10	3104	342	18	37	16	39	39	32	15	9929	53	59	34	2	27	8	42	39	15	6494	245	38	58	12	1	58	5	10	3104	342	18	37	16	39	32	15	9929	53	
28	8	58	15	15	10	6242	252	37	27	11	55	47	7	14	3444	340	20	40	15	15	10	32	25	9968	39	59	48	1	28	8	58	15	15	6242	252	37	27	11	55	47	7	14	3444	340	20	40	15	15	10	32	25	9968	39
29	9	13	25	14	13	5984	258	35	54	11	48	33	8	17	3781	337	22	41	14	25	25	32	29	9992	24	59	57	1	29	9	13	25	14	5984	258	35	54	11	48	33	8	17	3781	337	22	41	14	25	32	29	9992	24	
30	9	28	8			5718	266	34	18	11	40	16			4114	333	24	41						1.		60	0	0	30	9	28	8		5718	266	34	18						4114	333	24	41					1.		
⊙	—				above		above			—					under		under			—					under		under						under		under		⊙																
⊙	2d Equ.	⊙	Apog.	Dif.	X ^{rs} for	D.	Sex ^l X ^s	D.	⊙	2d Equ.	⊙	Apog.	Dif.	X ^{rs} for	D.	Sex ^l X ^s	D.	⊙	2d Equ.	⊙	Apog.	D																															

CONSTRUCTION of 2d Equation Δ Apogee.

As Gr. Eccl^y 66782 : Least Eccl^y 43319 :: Tan. Arg of Δ Ap. : To T. 4th Arc, which deducted from the said Annual Arg. leaves the 2d Equat. Δ Ap. sought.

Ex. To find 2d Equation Δ Ap. for 1st 21st Δ Apogee?

CONSTRUCTION of the *Decimal* and *Sexagesimal* MULTIPLIERS into the *Gr. Dif.* under or above the *M. Central Equat.* See P. 66, 67 and 68 following. As 11732 gr. Dif. Eccl^y under or above the Mean : G. D. under or above M. Centr. Eq. :: Dif. Eccl^y correspondent to the Arg. of Eq. : Pref. Dif. under or above the M. Central Equation.

Hence $\frac{11732}{11732} = \text{Decimal Mult.}$ (= Parts of a Degree of Sex^l Mult.) which $\times$ d into G. Dif. Eq. = Pref. Dif. from M. Centr^l.

Eq. reqd. — For Construction of Eccentr. Δ Orb. See farther on.

EXAM. In 1st 7th 40th 27th Δ Ap. to find the Multipliers to the Gr. Dif. Eq. in the following Tables, p. 66, 67 and 68.

For 1^{st</}

SUPPLEMENTAL LUNAR TABLES.

SEMIANNUAL or SECOND EQUATION of the MOON'S APOGEE, from the HORRARIAN TABLE. With Decimal and Sexagesimal Multipliers into the Greatest Difference under and above the Mean ELLIPTIC EQUATION of the MOON'S CENTER. For finding, directly, the present correct Difference from the Mean, and thence the whole ELLIPTIC EQUATION of the MOON. According to our New Method of Computing.

Argument. SUN from the MOON'S APOGEE, or Sun's True Place — Moon's Apogee is Equated.

Sig. 0. 6.										Sig. 1. 7.										Sig. 2. 8.									
Equat.	Dif.	X ^{rs} for	D.	Sex X	D.	Equat.	Dif.	X ^{rs} for	D.	Equat.	Dif.	X ^{rs} for	D.	Equat.	Dif.	X ^{rs} for	D.	Equat.	Dif.	X ^{rs} for	D.	Equat.	Dif.	X ^{rs} for	D.	Equat.	Dif.	X ^{rs} for	D.
D Apo.		C. Eq.		for e		D Apo.		C. Eq.		D Apo.		C. Eq.		D Apo.		C. Eq.		D Apo.		C. Eq.		D Apo.		C. Eq.		D Apo.		C. Eq.	
+		above		above		+		above		+		above		+		above		+		above		+		above		+		above	
0 0 0 0	20 21	1,0000	0 0	0 0	0 3	9 7 14	13 38	5000	305	30 0	1 50	11 8 55	9 10	5000	298	30 0	1 47	10 59 45	10 10	5000	294	31 47	1 46	10 49 35	11 13	5000	286	33 33	1 43
1 0 20 21	20 21	9994	18 59 57	0 0	0 6	9 20 52	13 19	4695	311	28 18	1 52	10 59 45	10 10	5000	294	31 47	1 46	10 49 35	11 13	5000	286	33 33	1 43	10 38 22	12 15	5000	278	35 16	1 40
2 0 40 42	20 19	9976	30 59 51	0 0	0 11	9 34 1	12 39	4384	317	26 18	1 54	10 49 35	11 13	5000	286	33 33	1 43	10 38 22	12 15	5000	278	35 16	1 40	10 26 7	13 17	5000	272	36 56	1 38
3 1 1 1	20 16	9946	43 59 40	0 0	0 15	9 46 40	12 9	4067	321	24 24	1 56	10 38 22	12 15	5000	278	35 16	1 40	10 26 7	13 17	5000	272	36 56	1 38	10 12 50	14 19	5000	264	38 34	1 35
4 1 21 17	20 13	9903	55 59 25	0 0	0 20	9 58 49	11 35	3746	327	22 28	1 57	10 26 7	13 17	5000	272	36 56	1 38	10 12 50	14 19	5000	264	38 34	1 35	9 58 31	15 21	5000	255	40 9	1 32
5 1 41 30	20 10	9848	67 59 5	0 0	0 24	10 10 24	11 5	3419	329	20 31	1 59	9 58 31	15 21	5000	255	40 9	1 32	9 43 10	16 22	5000	247	41 41	1 29	9 43 10	16 22	5000	247	41 41	1 29
6 2 1 40	20 4	9781	78 58 41	0 0	0 28	10 21 29	10 29	3090	334	18 32	2 0	9 43 10	16 22	5000	247	41 41	1 29	9 26 48	17 24	5000	237	43 10	1 25	9 26 48	17 24	5000	237	43 10	1 25
7 2 21 44	19 58	9703	91 58 13	0 0	0 33	10 31 58	9 52	2756	337	16 32	2 1	9 26 48	17 24	5000	237	43 10	1 25	9 9 24	18 24	5000	220	44 35	1 23	9 9 24	18 24	5000	220	44 35	1 23
8 2 41 42	19 52	9612	102 57 40	0 0	0 36	10 41 50	9 18	2419	340	14 31	2 1	9 9 24	18 24	5000	220	44 35	1 23	8 51 0	19 24	5000	219	45 58	1 19	8 51 0	19 24	5000	219	45 58	1 19
9 3 1 34	19 44	9510	113 57 4	0 0	0 41	10 51 8	8 40	2079	343	12 28	2 3	8 51 0	19 24	5000	219	45 58	1 19	8 31 36	20 22	5000	210	47 17	1 15	8 31 36	20 22	5000	210	47 17	1 15
10 3 21 18	19 36	9397	125 56 23	0 0	0 45	10 59 48	7 59	1736	344	10 25	2 3	8 31 36	20 22	5000	210	47 17	1 15	8 11 14	21 19	5000	206	48 32	1 12	8 11 14	21 19	5000	206	48 32	1 12
11 3 40 54	19 28	9272	137 55 38	0 0	0 49	11 7 47	7 20	1392	347	8 21	2 4	8 11 14	21 19	5000	206	48 32	1 12	7 49 55	22 15	5000	200	49 44	1 9	7 49 55	22 15	5000	190	50 53	1 5
12 4 0 22	19 16	9135	147 54 49	0 0	0 53	11 15 7	6 38	1045	348	6 16	2 5	7 49 55	22 15	5000	200	49 44	1 9	7 27 40	23 19	5000	180	51 58	1 5	7 27 40	23 19	5000	180	51 58	1 5
13 4 19 38	19 6	8988	158 53 56	0 0	0 57	11 21 45	5 55	697	348	4 11	2 5	7 27 40	23 19	5000	180	51 58	1 5	7 4 31	24 1	5000	170	52 59	1 1	7 4 31	24 1	5000	170	52 59	1 1
14 4 38 44	18 54	8830	170 52 59	0 0	1 1	11 27 40	5 11	349	349	2 6	2 6	7 4 31	24 1	5000	170	52 59	1 1	6 40 30	24 50	5000	158	53 56	0 57	6 40 30	24 50	5000	158	53 56	0 57
15 4 57 38	18 42	8660	180 51 58	0 0	1 5	11 32 51	4 26	0	349	2 6	2 6	6 40 30	24 50	5000	158	53 56	0 57	6 15 40	25 38	5000	147	54 49	0 53	6 15 40	25 38	5000	147	54 49	0 53
16 5 16 20	18 29	8480	190 50 53	0 0	1 9	11 37 17	3 38	349	348	4 11	2 5	6 15 40	25 38	5000	147	54 49	0 53	5 50 2	26 23	5000	137	55 38	0 49	5 50 2	26 23	5000	137	55 38	0 49
17 5 34 49	18 13	8290	200 49 44	0 0	1 12	11 30 55	2 52	697	348	6 16	2 5	5 50 2	26 23	5000	137	55 38	0 49	5 23 39	27 5	5000	125	56 23	0 45	5 23 39	27 5	5000	125	56 23	0 45
18 5 53 2	18 59	8090	210 48 32	0 0	1 15	11 43 47	2 2	1045	348	8 21	2 5	5 23 39	27 5	5000	125	56 23	0 45	4 56 34	27 43	5000	113	57 4	0 41	4 56 34	27 43	5000	113	57 4	0 41
19 6 11 1	17 41	7880	219 47 17	0 0	1 19	11 45 49	1 12	1392	347	10 25	2 4	4 56 34	27 43	5000	113	57 4	0 41	4 28 51	28 19	5000	102	57 40	0 36	4 28 51	28 19	5000	102	57 40	0 36
20 6 28 42	17 26	7661	229 45 58	0 0	1 23	11 47 1	0 11	1736	344	12 28	2 3	4 0 32	28 52	5000	91	58 13	0 33	4 0 32	28 52	5000	91	58 13	0 33	4 0 32	28 52	5000	91	58 13	0 33
21 6 46 8	17 7	7431	238 44 35	0 0	1 26	11 47 22	0 32	2079	340	14 31	2 1	3 31 40	29 20	5000	80	58 41	0 28	3 31 40	29 20	5000	80	58 41	0 28	3 31 40	29 20	5000	80	58 41	0 28
22 7 3 15	16 48	7193	246 43 9	0 0	1 28	11 46 50	1 26	2419	337	16 32	2 1	3 2 20	29 44	5000	67	59 5	0 24	3 2 20	29 44	5000	67	59 5	0 24	3 2 20	29 44	5000	67	59 5	0 24
23 7 20 3	16 27	6947	255 41 41	0 0	1 32	11 45 24	2 20	2756	334	18 32	2 0	3 2 20	29 44	5000	67	59 5	0 24	2 2 30	30 6	5000	55	59 25	0 20	2 2 30	30 6	5000	55	59 25	0 20
24 7 36 39	16 6	6692	264 40 9	0 0	1 35	11 43 4	3 17	3090	330	20 31	1 59	2 2 30	30 6	5000	55	59 25	0 20	2 2 30	30 22	5000	43	59 40	0 15	2 2 30	30 22	5000	43	59 40	0 15
25 7 52 36	15 44	6428	274 38 34	0 0	1 38	11 39 47	4 13	3420	326	22 28	1 57	2 2 30	30 22	5000	43	59 40	0 15	2 2 30	30 38	5000	30	59 51	0 11	2 2 30	30 38	5000	30	59 51	0 11
26 8 8 20	15 21	6156	283 36 56	0 0	1 40	11 35 34	5 11	3746	321	24 24	1 54	2 2 30	30 22	5000	43	59 40	0 15	2 2 30	30 45	5000	18	59 57	0 6	2 2 30	30 45	5000	18	59 57	0 6
27 8 23 41	15 52	5878	287 35 16	0 0	1 43	11 30 23	6 9	4067	317	26 18	1 52	2 2 30	30 45	5000	18	59 57	0 6	2 2 30	30 45	5000	6	60 0	0 3	2 2 30	30 45	5000	6	60 0	0 3
28 8 38 38	14 30	5591	293 33 33	0 0	1 46	11 24 14	7 9	4384	311	28 10	1 50	2 2 30	30 45	5000	18	59 57	0 6	2 2 30	30 45	5000	6	60 0	0 3	2 2 30	30 45	5000	6	60 0	0 3
29 8 53 8	14 6	5298	299 31 47	0 0	1 47	11 17 5	8 10	4695	305	30 0	1 50	2 2 30	30 45	5000	18	59 57	0 6	2 2 30	30 45	5000	6	60 0	0 3	2 2 30	30 45	5000	6	60 0	0 3
30 9 7 14	14 0	5000	300 0	0 0	1 47	11 8 55	8 10	5000	305	30 0	1 50	2 2 30	30 45	5000	18	59 57	0 6	2 2 30	30 45	5000	6	60 0	0 3	2 2 30	30 45	5000	6	60 0	0 3
—	—	above	above	—	—	—	—	under	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Equat.	Dif.	X ^{rs} for	D.	Sex X	D.	Equat.	Dif.	X ^{rs} for	D.	Equat.	Dif.	X ^{rs} for	D.	Equat.	Dif.	X ^{rs} for	D.	Equat.	Dif.	X ^{rs} for	D.	Equat.	Dif.	X ^{rs} for	D.	Equat.	Dif.	X ^{rs} for	D.
D Apo.		C. Eq.		for e		D Apo.		C. Eq.		D Apo.		C. Eq.		D Apo.		C. Eq.		D Apo.		C. Eq.		D Apo.		C. Eq.		D Apo.		C. Eq.	
+		above		above		+		above		+		above		+		above		+		above		+		above		+		above	

Sig. 11. 5.

Sig. 10. 4.

Sig. 9. 3.

THE above Construction of the DECIMAL and SEXAGESIMAL MULTIPLIERS is according to different Eccentricities from the former; viz. 66854 the Greatest, 55237 the Mean, and 43619 the Least Eccentricity: but in the same Manner constructed. A Table of which different Eccentricities we have inserted farther on; that the Principles may be examined on which these Tables are founded.

SUPPLEMENTAL

SUPPLEMENTAL LUNAR TABLES.

IV. MEAN CENTRAL EQUATION of the MOON, from the *Newtonian* THEORY. With the *Greatest* Difference of EQUATION under and above the MEAN EQUATION correspondent to the *Least* and *Greatest* ECCENTRICITIES, 43319 and 66782 to Mean Dist. $\propto$ 1000000. According to our NEW METHOD of Computing.

Argument. MOON'S Equated ANOMALY; Or D. 3. Equated — D Apogee 2. Equated.

Sign 0.										Sign 1.									
D	Eq	A.	greatest	Dif.	In.	Mean	Dif.	Great.	D.	D	Eq	A.	greatest	Dif.	In.	Mean	Dif.	Great.	D.
			under.		fr.	Cent		Diff. E.					under		fr.	Cent		Diff. E.	
					M.	Eq.		+							M.	Eq.		+	
								above										above	
0	0	0	1 15	1 15	0	0 0 0	6 11	0 0	1 12	1	36	20	1 5	6	2 58	30	5 28	35	24
1	1	15	1 15	1 15	1	0 6 11	6 11	1 12	1 13	1	37	25	1 10	6	3 3 58	5 26	36	31	1 7
2	2	30	1 15	1 15	1	0 12 22	6 10	2 25	1 14	1	38	35	1 6	6	3 9 24	5 22	37	36	1 5
3	3	45	1 15	1 15	1	0 18 32	6 10	3 39	1 13	1	39	41	1 7	7	3 14 46	5 19	38	42	1 6
4	4	0	1 15	1 15	1	0 24 42	6 10	4 52	1 12	1	40	48	1 9	7	3 20 5	5 16	39	48	1 6
5	5	15	1 15	1 15	1	0 30 52	6 10	5 4	1 13	1	41	51	1 7	7	3 25 21	5 14	40	53	1 5
6	6	30	1 14	1 14	2	0 37 2	6 9	6 17	1 12	1	42	58	1 4	7	3 30 35	5 10	41	57	1 4
7	7	44	1 14	1 14	2	0 43 10	6 8	7 29	1 12	2	44	2	1 4	7	3 35 45	5 6	43	1	1 4
8	8	58	1 14	1 14	2	0 49 19	6 7	8 41	1 13	2	45	6	1 4	7	3 40 51	5 4	44	3	1 4
9	9	12	1 14	1 14	2	0 55 26	6 6	9 54	1 12	2	46	10	1 3	7	3 45 55	5 0	45	7	1 4
10	10	26	1 14	1 14	2	1 1 32	6 6	10 6	1 12	3	47	13	1 1	7	3 50 55	4 55	46	8	1 1
11	11	40	1 14	1 14	2	1 7 38	6 5	11 18	1 12	3	48	14	1 1	7	3 55 50	4 53	47	10	1 2
12	12	54	1 14	1 14	3	1 13 43	6 3	12 30	1 13	3	49	15	1 3	7	4 0 43	4 49	48	10	1 0
13	13	8	1 14	1 14	3	1 19 46	6 2	13 43	1 12	3	50	18	0 58	7	4 5 32	4 45	49	10	0 59
14	14	22	1 14	1 14	3	1 25 48	6 2	14 55	1 10	3	51	16	0 59	7	4 10 17	4 40	50	9	0 59
15	15	36	1 13	1 13	3	1 31 50	6 0	15 5	1 12	3	52	15	1 0	7	4 14 57	4 37	51	8	1 0
16	16	49	1 13	1 13	3	1 37 50	5 58	16 17	1 11	3	53	15	0 59	7	4 19 34	4 33	52	8	1 0
17	17	2	1 13	1 13	3	1 43 48	5 57	17 28	1 9	3	54	14	0 56	8	4 24 7	4 29	53	3	0 55
18	18	15	1 12	1 12	4	1 49 45	5 55	18 37	1 10	4	55	10	0 56	7	4 28 36	4 26	54	1	0 58
19	19	27	1 12	1 12	4	1 55 40	5 53	19 47	1 12	4	56	6	0 53	7	4 33 0	4 20	55	59	0 55
20	20	39	1 11	1 11	4	2 1 33	5 51	20 59	1 10	4	57	1	0 57	7	4 37 20	4 16	55	54	0 55
21	21	50	1 10	1 10	4	2 7 24	5 50	21 9	1 9	4	57	58	0 52	7	4 41 36	4 11	56	49	0 54
22	22	0	1 10	1 10	4	2 13 14	5 47	22 18	1 11	4	58	50	0 55	7	4 45 47	4 6	57	43	0 54
23	23	10	1 10	1 10	4	2 19 1	5 45	23 29	1 10	4	59	45	0 50	7	4 49 53	4 2	58	37	0 52
24	24	22	1 11	1 11	4	2 24 46	5 44	24 39	1 6	4	60	35	0 53	7	4 53 55	3 58	59	29	0 51
25	25	33	1 11	1 11	5	2 30 30	5 41	25 45	1 10	4	61	28	0 50	7	4 57 53	3 52	60	20	0 52
26	26	44	1 9	1 9	5	2 36 11	5 38	26 55	1 8	5	62	18	0 49	7	5 1 45	3 47	61	12	0 50
27	27	53	1 10	1 10	5	2 41 49	5 35	27 3	1 5	5	63	7	0 52	7	5 5 32	3 43	62	2	0 49
28	28	3	1 6	1 6	5	2 47 26	5 32	28 8	1 10	5	63	59	0 46	7	5 9 15	3 37	62	51	0 49
29	29	9	1 11	1 11	5	2 52 58	5 32	29 18	1 6	6	64	45	0 47	7	5 12 52	3 33	63	40	0 48
30	30	20			6	2 58 30	5 32	30 24	1 6	6	65	32		7	5 16 25		64	28	
			under.			+		above					under			+		above	
D	Eq	A.	greatest	Dif.	In.	Mean	Dif.	Great.	D.	D	Eq	A.	greatest	Dif.	In.	Mean	Dif.	Great.	D.
			diff. Eq.		fr.	Cent		Diff. E.					diff. Eq.		fr.	Cent		Diff. E.	
					M.	Eq.		+							M.	Eq.		+	

EXAMPLE I. To find the MOON'S Central Equation correspondent to $2^{\circ} 30' 0''$ D Apogee, and $1^{\circ} 11' 0''$ D Equated Anomaly?

By Tab. above. M. Cen. Eq. D

$1^{\circ} 11' 0''$ D Eqd. Anom. — $3^{\circ} 55' 50''$ — $48' 233$ G. D. Eq. under } Or, — $48' 14''$ 0.0948

By Tab. p. 64 — $31' 47''$ D. Multiplier } Sext: $\times 1.39$ 32 0.1812

Central Equation — $3' 24''$ 3 required. 434097 Prop. D. 31 47 0.2760

For Correction.
Against $1^{\circ} 11' 0''$ Equated Anom.
Increase from Mean, to be allowed in the two of three Parts from M. to Extreme, into which the G. Dif. is supposed divided. In 3^d Part nothing allowed.

very nearly. — $3' 24''$ 10 corrected.

But . . . 659 = $39' 32''$
 $39' 54''$ multiply 8
 $32' 45''$ by 6.

N.B. When the Arguments exceed whole Degrees, Proportion of the Equations must be made (as usual) for that Excess.

SUPPLEMENTAL LUNAR TABLES.

IV. MEAN CENTRAL EQUATION of the MOON, from the *Newtonian* THEORY. With the *Greatest* Difference of EQUATION, *under* and *above* the MEAN EQUATION, correspondent to the *Least* and *Greatest* ECCENTRICITIES, 43319 and 66782 to Mean Dist. a Sun 1000000. According to our NEW METHOD of Computing.

Argument. MOON's Equated ANOMALY, Or, D. 3 Equated — D Apogee 2. Equated.

Sig. 2.												Sig. 3.											
D	Eq	A.	greatest	Dif.	In.	Mean	Dif.	greatest	Dif.	De	greatest	Dif.	In & D.	Mean	Dif.	greatest	Dif.	D. & In.	Mean	Dif.	D	Eq	
			under		fr.	Cent		above		fr.	under		fr.	Cent		above		fr.	Cent				
					M.	Eq.				M.													
0	65	32			7	5 16 25	3 28	64 28	0 45	6	80 17	0 10	2	6 17 44	0 24	80 3	0 13	2	30				
1	66	19	0 47		7	5 19 53	3 21	65 13	0 47	6	80 27	0 8	2	6 18 8	0 17	80 16	0 9	2	29				
2	67	4	0 43		7	5 23 14	3 16	66 0	0 46	6	80 35	0 6	2	6 18 25	0 10	80 25	0 10	2	28				
3	67	47	0 42		7	5 26 30	3 12	66 46	0 42	6	80 41	0 6	2	6 18 35	0 3	80 35	0 8	1	27				
4	68	29	0 43		7	5 29 42	3 6	67 28	0 44	6	80 47	0 3	1	6 18 38	0 4	80 43	0 7	0	26				
5	69	12	0 41		6	5 32 48	3 0	68 12	0 42	6	80 50	0 2	0	6 18 34	0 11	80 50	0 4	0	25				
6	69	52	0 42		6	5 35 48	2 53	68 54	0 40	6	80 52	0 1	—	6 18 23	0 18	80 54	0 4	+	24				
7	70	34	0 38		6	5 38 43	2 39	69 34	0 41	6	80 53	0 0	1	6 18 5	0 24	80 58	0 4	1	23				
8	71	12	0 37		6	5 41 32	2 43	70 15	0 38	6	80 53	0 0	1	6 17 41	0 32	81 0	0 2	1	22				
9	71	49	0 36		6	5 44 15	2 38	70 53	0 35	6	80 51	0 2	1	6 17 9	0 39	81 0	0 0	1	21				
10	72	25	0 33		6	5 46 53	2 32	71 32	0 36	6	80 47	0 4	1	6 16 30	0 45	81 0	0 0	1	20				
11	72	58	0 38		6	5 49 25	2 26	72 8	0 32	6	80 42	0 5	1	6 15 45	0 53	80 55	0 5	1	19				
12	73	36	0 31		6	5 51 51	2 20	72 40	0 38	6	80 36	0 6	2	6 14 52	1 0	80 52	0 3	2	18				
13	74	7	0 33		5	5 54 11	2 13	73 18	0 33	6	80 28	0 8	2	6 13 52	1 8	80 46	0 6	2	17				
14	74	40	0 30		5	5 56 24	2 8	73 51	0 33	6	80 16	0 12	2	6 12 44	1 14	80 40	0 6	3	16				
15	75	10	0 30		5	5 58 32	2 2	74 24	0 32	6	80 5	0 11	2	6 11 30	1 21	80 30	0 10	3	15				
16	75	40	0 28		5	6 0 34	1 55	74 56	0 30	5	79 52	0 13	3	6 10 9	1 29	80 21	0 9	4	14				
17	76	8	0 27		5	6 2 29	1 49	75 26	0 30	5	79 37	0 15	3	6 8 40	1 35	80 8	0 13	4	13				
18	76	35	0 22		5	6 4 18	1 44	75 56	0 26	4	79 24	0 13	3	6 7 5	1 42	79 55	0 13	4	12				
19	77	1	0 25		4	6 6 2	1 36	76 22	0 27	4	79 3	0 21	3	6 5 23	1 49	79 40	0 15	4	11				
20	77	26	0 23		4	6 7 38	1 30	76 49	0 24	4	78 45	0 18	3	6 3 34	1 57	79 23	0 17	4	10				
21	77	49	0 22		4	6 9 8	1 24	77 13	0 27	4	78 24	0 21	4	6 1 37	2 3	79 2	0 21	4	9				
22	78	11	0 20		4	6 10 32	1 17	77 40	0 22	4	78 1	0 23	4	5 59 34	2 11	78 43	0 19	4	8				
23	78	31	0 21		4	6 11 49	1 11	78 2	0 21	4	77 36	0 25	4	5 57 23	2 17	78 21	0 22	5	7				
24	78	51	0 15		4	6 13 0	1 4	78 23	0 19	3	77 12	0 24	4	5 55 6	2 25	77 58	0 23	5	6				
25	79	6	0 19		4	6 14 4	0 57	78 42	0 19	3	76 44	0 28	5	5 52 41	2 31	77 34	0 24	5	5				
26	79	25	0 14		3	6 15 1	0 51	79 1	0 18	3	76 15	0 29	5	5 50 10	2 38	77 7	0 27	6	4				
27	79	39	0 14		3	6 15 52	0 44	79 19	0 17	3	75 46	0 29	5	5 47 32	2 46	76 48	0 29	6	3				
28	79	53	0 12		3	6 16 36	0 37	79 36	0 15	3	75 10	0 36	5	5 44 46	2 53	76 9	0 39	6	2				
29	80	5	0 12		3	6 17 13	0 34	79 51	0 12	3	74 38	0 32	5	5 41 53	2 59	75 38	0 31	6	1				
30	80	17			3	6 17 44		80 3		3	74 3	0 35	5	5 38 54		75 3	0 35	6	0				
			under		fr.	Cent		above		fr.	under		fr.	Cent		above		fr.	Cent				
D	Eq	A.	greatest	Dif.	In.	Mean	Dif.	greatest	Dif.	De	greatest	Dif.	D. & In.	Mean	Dif.	greatest	Dif.	In & D.	Mean	Dif.	D	Eq	
			diff. Eq.		M.	Eq.		diff. Eq.		M.	diff. Eq.		M.	Eq.		diff. Eq.		M.	Eq.				

Sig. 9.

Sig. 8.

The *Ease* and *Readiness* of this NEW METHOD of taking out the Equation of the MOON's CENTER, (where 2d Equation D's Apog. is retained) appear evident from the foregoing EXAMPLE.

ROYAL ASTRONOMER

SUPPLEMENTAL LUNAR TABLES.

IV. MEAN CENTRAL EQUATION of the MOON, from the *Newtonian* THEORY. With the *Greatest* Difference of EQUATION, *under* and *above* the MEAN EQUATION, correspondent to the *Least* and *Greatest* ECCENTRICITIES, 43319 and 66782 to Mean Dist. à Sun 1000000. According to our NEW METHOD of Computing.

Argument. MOON's Equated ANOMALY, Or, D. 3. Equated — D. Apogee 2. Equated.

Sig. 4.												Sig. 5.																			
D	greatest			De	Mean			greatest			In.	greatest			De	Mean			greatest			In.	D								
Eq	Dif. E.			fr.	Cent Eq.			Dif. E.			fr.	Dif. E.			fr.	Cent Eq.			Dif. E.			fr.	Eq								
An	under			M.	D			+			+	under			M.	D			+			+	A.								
0	I	II	I	II	I	II	I	II	I	II	I	I	II	I	II	I	II	I	II	I	II	I	0								
0	74	3	0	35	5	5	38	54	3	5	75	3	0	34	6	44	56	1	21	7	3	21	6	5	59	46	7	1	19	7	30
1	73	28	0	39	6	5	35	49	3	13	74	29	0	39	7	43	35	1	20	7	3	15	7	6	3	44	48	1	22	7	29
2	72	49	0	40	6	5	32	36	3	19	73	50	0	36	7	42	15	1	21	7	3	9	4	6	7	43	26	1	26	7	28
3	72	9	0	43	6	5	29	17	3	27	73	14	0	39	7	40	54	1	22	7	3	2	57	6	13	42	6	1	24	7	27
4	71	26	0	43	6	5	25	50	3	32	72	35	0	43	7	39	32	1	24	6	2	56	44	6	15	40	42	1	26	7	26
5	70	43	0	43	7	5	22	18	3	32	71	52	0	43	7	38	8	1	22	6	2	50	29	6	15	39	10	1	25	6	25
6	69	58	0	45	7	5	18	39	3	39	71	9	0	43	7	36	46	1	22	6	2	44	11	6	23	37	51	1	27	6	24
7	69	13	0	45	7	5	14	53	3	46	70	24	0	45	7	34	21	1	25	6	2	37	48	6	25	36	24	1	30	6	23
8	68	25	0	48	7	5	11	1	3	52	69	37	0	47	7	33	56	1	27	6	2	31	23	6	30	34	54	1	26	6	22
9	67	35	0	50	8	5	7	3	4	58	68	49	0	48	8	32	29	1	27	6	2	24	53	6	33	33	28	1	29	6	21
10	66	44	0	51	7	5	2	57	4	6	68	1	0	48	8	31	2	1	27	6	2	18	20	6	33	31	59	1	29	6	20
11	65	53	0	52	7	4	58	46	4	11	67	9	0	52	8	29	34	1	28	5	2	11	45	6	37	30	28	1	31	6	19
12	65	1	0	52	7	4	54	29	4	17	66	14	0	53	8	28	5	1	29	5	2	5	6	6	39	28	57	1	31	6	18
13	64	5	0	56	7	4	50	5	4	24	65	21	0	56	8	26	35	1	30	5	1	58	25	6	41	27	23	1	34	5	17
14	63	8	0	57	7	4	45	35	4	30	64	25	0	58	8	25	5	1	30	5	1	51	41	6	44	25	52	1	35	5	16
15	62	8	0	57	7	4	40	59	4	36	63	27	0	58	8	23	35	1	30	4	1	44	54	6	47	24	17	1	35	5	15
16	61	10	0	58	7	4	36	18	4	41	62	29	0	58	8	22	3	1	32	4	1	38	5	6	51	22	44	1	33	5	14
17	60	9	0	58	7	4	31	31	4	47	61	27	0	58	8	20	31	1	32	4	1	31	14	6	53	21	11	1	33	5	13
18	59	6	0	58	8	4	26	37	4	54	60	25	0	58	8	18	59	1	32	4	1	24	21	6	55	19	36	1	35	5	12
19	58	3	0	58	8	4	21	39	5	5	59	21	0	58	8	17	25	1	34	3	1	17	26	6	55	17	58	1	36	4	11
20	56	56	0	58	7	4	16	34	5	9	58	17	0	58	8	15	51	1	34	3	1	10	29	6	58	16	22	1	36	4	10
21	55	49	0	58	8	4	11	25	5	16	57	10	0	58	8	14	7	1	34	3	1	3	31	7	1	14	45	1	37	3	9
22	54	41	0	58	8	4	6	9	5	20	56	2	0	58	8	12	42	1	35	2	0	56	30	7	1	13	8	1	37	3	8
23	53	34	0	58	8	4	0	49	5	25	54	52	0	58	8	11	8	1	35	2	0	49	29	7	1	11	30	1	40	3	7
24	52	23	0	58	8	3	55	24	5	32	53	40	0	58	8	9	33	1	35	2	0	42	28	7	3	9	50	1	37	2	6
25	51	10	0	58	7	3	49	52	5	36	52	27	0	58	7	7	58	1	35	1	0	35	25	7	4	8	13	1	37	1	5
26	49	58	0	58	7	3	44	16	5	40	51	14	0	58	7	6	22	1	36	1	0	28	21	7	5	6	35	1	38	1	4
27	48	45	0	58	7	3	38	36	5	46	49	59	0	58	7	4	47	1	35	1	0	21	16	7	5	4	52	1	38	1	3
28	47	28	0	58	7	3	32	50	5	50	48	44	0	58	7	3	12	1	36	0	0	14	11	7	5	3	16	1	41	1	2
29	46	12	0	58	7	3	27	0	5	54	47	25	0	58	7	1	36	1	36	0	0	7	6	7	5	1	38	1	38	0	1
30	44	56	0	58	7	3	21	6	5	54	46	7	0	58	7	0	0	0	0	0	0	0	0	7	6	0	0	0	0	0	0
D	greatest			De	Mean			greatest			In.	greatest			De	Mean			greatest			In.	D								
Eq	Dif. E.			fr.	Cent Eq.			Dif. E.			fr.	Dif. E.			fr.	Cent Eq.			Dif. E.			fr.	Eq								
A.	under			M.	D			+			+	under			M.	D			+			+	A.								
0	I	II	I	II	I	II	I	II	I	II	I	I	II	I	II	I	II	I	II	I	II	I	0								
0	74	3	0	35	5	5	38	54	3	5	75	3	0	34	6	44	56	1	21	7	3	21	6	5	59	46	7	1	19	7	30
1	73	28	0	39	6	5	35	49	3	13	74	29	0	39	7	43	35	1	20	7	3	15	7	6	3	44	48	1	22	7	29
2	72	49	0	40	6	5	32	36	3	19	73	50	0	36	7	42	15	1	21	7	3	9	4	6	7	43	26	1	26	7	28
3	72	9	0	43	6	5	29	17	3	27	73	14	0	39	7	40	54	1	22	7	3	2	57	6	13	42	6	1	24	7	27
4	71	26	0	43	6	5	25	50	3	32	72	35	0	43	7	39	32	1	24	6	2	56	44	6	15	40	42	1	26	7	26
5	70	43	0	43	7	5	22	18	3	32	71	52	0	43	7	38	8	1	22	6	2	50	29	6	15	39	10	1	25	6	25
6	69	58	0	45	7	5	18	39	3	39	71	9	0	43	7	36	46	1	22	6	2	44	11	6	23	37	51	1	27	6	24
7	69	13	0	45	7	5	14	53	3	46	70	24	0	45	7	34	21	1	25	6	2	37	48	6	25	36	24	1	30	6	23
8	68	25	0	48	7	5	11	1	3	52	69	37	0	47	7	33	56	1	25	6	2	31	23	6	30	34	54	1	26	6	22
9	67	35	0	50	8	5	7	3	4	58	68	49	0	48	8	32	29	1	27	6	2	24	53	6	33	33	28	1	29	6	21
10	66	44	0	51	7	5	2	57	4	6	68	1	0	48	8	31	2	1	27	6	2	18	20	6	33	31	59	1	29	6	20
11	65	53	0	52	7	4	58	46	4	11	67	9	0	52	8	29	34	1	28	5	2	11	45	6	37	30	28	1	31	6	19
12	65	1	0	52	7	4	54	29	4	17	66	14	0	53	8	28	5	1	29	5	2	5	6	6	39	28	57	1	31	6	18
13	64	5	0	56	7	4	50	5	4																						

and NAVIGATOR. SUPPLEMENTAL LUNAR TABLES.

69

OR, IV. MEAN ELLIPTIC EQUATION of the MOON'S CENTER from the Horrobian THEORY. With the greatest Difference of EQUATION, under and above the MEAN, correspondent to the Least and Greatest ECCENTRICITIES, 43619 and 66854, to Mean Distance à 1000000. According to our NEW METHOD of Computing.

Argument. MOON'S Equated ANOMALY, Or Δ Equated — Δ M. Ap. Equated by 2d. Eq. Ap.

Sig. 0.												Sig. 1.													
D Eq. An.	G. D. Equat.		Dif.	In. fr. M.	Mean Cent Eq.		Dif.	G. D. Equat.		Dif.	De fr. M.	D	D	G. D. Equat.		Dif.	In. fr. M.	Mean Cent Eq.		Dif.	G. D. Equat.		Dif.	De fr. M.	D
	under				—									under				—							
0	0	0		0	0	0		0		0				35	56		6	2	59	4		35	4		0
1	1	14	14	1	0	6	12	1	12	1				37	1	5	6	3	4	33	5	29	35	4	30
2	2	28	14	1	0	12	24	1	12	1				38	8	1	7	3	10	0	5	27	36	10	29
3	3	42	15	1	0	18	35	1	13	1				39	14	1	6	3	15	23	5	23	37	15	28
4	4	57	13	1	0	24	47	1	12	1				40	20	1	6	3	20	44	5	21	38	20	27
5	6	10	15	1	0	30	58	1	11	1				41	23	3	7	3	26	1	5	17	39	24	26
6	7	25	15	2	0	37	9	6	11	1				42	28	1	5	3	31	16	5	15	40	29	25
7	8	40	13	2	0	43	19	6	10	1				43	34	1	6	3	36	27	5	11	41	31	24
8	9	53	13	2	0	49	27	6	8	2				44	41	1	7	3	41	35	5	8	42	34	23
9	11	6	12	2	0	55	36	6	9	2				45	41	1	0	3	46	39	5	4	43	37	22
10	12	18	12	2	1	1	43	6	7	3				46	44	1	3	3	51	39	5	0	44	39	21
11	13	32	14	2	1	7	51	6	8	3				47	43	0	59	7	56	36	4	57	46	42	20
12	14	44	12	3	1	13	57	6	6	3				48	44	1	1	4	1	30	4	54	46	42	19
13	15	58	11	3	1	20	2	6	5	3				49	45	0	1	4	6	20	4	50	47	42	18
14	17	9	13	3	1	26	5	6	3	3				50	44	0	59	7	4	11	6	4	46	41	17
15	18	22	13	3	1	32	7	6	2	3				51	42	0	58	7	4	15	47	4	41	49	16
16	19	35	13	3	1	38	8	6	1	3				52	40	0	58	7	4	20	24	4	37	50	15
17	20	48	12	3	1	44	8	5	0	3				53	38	0	58	8	4	24	58	4	34	51	14
18	22	0	12	4	1	50	6	5	58	4				54	35	0	57	8	4	29	28	4	30	52	13
19	23	12	12	4	1	56	2	5	56	4				55	31	0	56	7	4	33	53	4	25	53	12
20	24	24	12	4	2	1	56	5	54	4				56	27	0	56	7	4	38	14	4	21	54	11
21	25	36	10	4	2	7	49	5	53	4				57	22	0	55	7	4	42	31	4	17	55	10
22	26	46	8	4	2	13	39	5	50	4				58	15	0	53	7	4	46	43	4	12	56	9
23	27	54	11	4	2	19	27	5	48	4				59	7	0	52	7	4	50	50	4	7	57	8
24	29	5	8	4	2	25	14	5	47	4				59	58	0	51	7	4	54	53	4	3	58	7
25	30	13	8	5	2	30	58	5	44	4				60	52	0	54	7	4	58	55	4	2	58	6
26	31	23	10	5	2	36	41	5	43	4				61	37	0	45	7	5	2	43	3	48	59	5
27	32	32	8	5	2	42	21	5	40	5				62	27	0	48	7	5	6	32	3	49	60	4
28	33	40	8	5	2	47	58	5	37	5				63	15	0	48	7	5	10	15	3	43	61	3
29	34	48	8	5	2	53	32	5	34	6				64	3	0	44	7	5	13	54	3	39	62	2
30	35	56		5	2	59	4	5	32	6				64	47	7		5	17	27	3	33	63	4	1
	under				+			above						under			+			above					0
D Eq. An.	G. D. Equat.		Dif.	In. fr. M.	Mean Cent Eq.		Dif.	G. D. Equat.		Dif.	De fr. M.	D	D	G. D. Equat.		Dif.	In. fr. M.	Mean Cent Eq.		Dif.	G. D. Equat.		Dif.	De fr. M.	D
—					—									—				—							

EXAMPLE I. To find the MOON'S Elliptic or Central Equation correspondent to $1^{\circ} 11' 0''$ Δ Equated Anomaly, and $2^{\circ} 8' 0''$ Δ Apogee, according to the Horrobian THEORY?

By Table, p. 65, $\odot$ à Δ Ap. | Dec. X^r . | Ser. X^r . | under Mean Eccentricity and Equation.

$2^{\circ} 8' 0''$ | 57194 | $43' 10''$ |

$47' 43'' = 47' 71''$

7194

50358

50358

28776

Proportional Equation $34' 32574$

$19''$ &c.

By Tab. above $1^{\circ} 11' 0''$ | M. Cen. Eq. | G.D. & und. | L.L. |

$3^{\circ} 56' 36''$ | $47' 43''$ | 0.0995 |

$-34' 20''$ | $43' 10''$ | 0.1430 |

Δ Central Eq. . . . $3' 22' 16''$ | $-34' 20''$ | 0.2425 |

nearby.

Corr. to $1^{\circ} 11' 0''$ Eq. An. $+7''$

Δ Central Eq. required $3' 22' 23''$

SUPPLEMENTAL

SUPPLEMENTAL LUNAR TABLES.

V. MEAN ELLIPTIC EQUATION of the MOON'S CENTER from the *Horrobian* THEORY. With the greatest Difference of EQUATION, under and above the MEAN, correspondent to the *Least* and *Greatest* ECCENTRICITIES, 43619 and 66854, to Mean Distance à 1000000. According to our NEW METHOD of Computing.

Argument. MOON'S Equated ANOMALY, Or D : Equated — D M. Ap. Equated by 2d. Eq. Ap.

Sig. 2.													Sig. 3.												
Eq. An.	G. D. Equat.	Dif.	In. fr. M.	Mean Cent. Eq. D	Dif.	G. D. Equat. +	Dif.	De fr. M.	G. D. Equat. -	Dif.	In & D. fr. M.	Mean Cent. Eq. D	Dif.	G. D. Equat. +	Dif.	D. & In. fr. M.	Eq. An.								
	under		+	-		above			under		+	-		above											
0	64 47	0 49	7	5 17 27	3 27	63 51	0 47	6	79 29	0 6	2	6 18 59	0 24	79 18	0 11	2	30								
1	65 36	0 45	7	5 20 54	3 23	64 38	0 44	6	79 35	0 9	2	6 19 23	0 17	79 29	0 11	2	29								
2	66 21	0 46	7	5 24 17	3 18	65 23	0 43	6	79 44	0 7	2	6 19 40	0 10	79 40	0 10	1	28								
3	67 7	0 40	7	5 27 35	3 12	66 5	0 44	6	79 51	0 5	1	6 19 50	0 4	79 50	0 7	1	27								
4	67 47	0 42	7	5 30 47	3 6	66 40	0 43	6	79 56	0 6	0	6 19 54	0 3	79 57	0 5	0	26								
5	68 29	0 45	6	5 33 53	3 1	67 32	0 42	6	80 2	0 2	—	6 19 51	0 11	80 2	0 5	+	25								
6	69 14	0 35	6	5 36 54	2 55	68 14	0 41	6	80 4	0 2	1	6 19 40	0 17	80 7	0 3	1	24								
7	69 49	0 38	6	5 39 49	2 50	68 55	0 40	6	80 3	0 0	1	6 19 23	0 26	80 10	0 2	1	23								
8	70 27	0 38	6	5 42 39	2 45	69 35	0 37	6	80 3	0 2	1	6 18 57	0 32	80 12	0 0	1	22								
9	71 5	0 36	6	5 45 24	2 38	70 12	0 40	6	80 1	0 3	1	6 18 25	0 39	80 12	0 0	1	21								
10	71 41	0 36	6	5 48 2	2 33	70 52	0 34	6	79 58	0 4	1	6 17 46	0 46	80 12	0 3	1	20								
11	72 17	0 35	6	5 50 35	2 27	71 20	0 35	6	79 54	0 5	2	6 17 0	0 52	80 9	0 5	1	19								
12	72 52	0 32	6	5 53 2	2 20	72 11	0 34	6	79 49	0 8	2	6 16 8	1 0	80 4	0 6	2	18								
13	73 24	0 31	5	5 55 22	2 14	72 35	0 34	6	79 41	0 11	2	6 15 8	1 8	79 58	0 6	2	17								
14	73 55	0 30	5	5 57 36	2 8	73 9	0 32	6	79 30	0 11	2	6 14 0	1 14	79 52	0 9	3	16								
15	74 25	0 28	5	5 59 44	2 2	73 41	0 31	6	79 19	0 13	2	6 12 46	1 21	79 43	0 11	3	15								
16	74 53	0 27	5	6 1 46	1 56	74 12	0 30	5	79 6	0 13	3	6 11 25	1 29	79 32	0 11	4	14								
17	75 20	0 27	5	6 3 42	1 49	74 42	0 29	5	78 53	0 18	3	6 9 56	1 36	79 21	0 13	4	13								
18	75 47	0 26	5	6 5 31	1 44	75 11	0 27	4	78 35	0 19	3	6 8 20	1 43	79 8	0 15	4	12								
19	76 13	0 24	4	6 7 15	1 37	75 38	0 26	4	78 16	0 19	3	6 6 37	1 49	78 53	0 18	4	11								
20	76 37	0 24	4	6 8 52	1 31	76 4	0 25	4	77 57	0 22	3	6 4 48	1 57	78 35	0 18	4	10								
21	77 1	0 22	4	6 10 23	1 23	76 29	0 24	4	77 35	0 21	4	6 2 51	2 3	78 17	0 19	4	9								
22	77 23	0 23	4	6 11 46	1 17	76 53	0 24	4	77 14	0 21	4	6 0 48	2 11	77 56	0 21	4	8								
23	77 46	0 16	4	6 13 3	1 11	77 17	0 21	4	76 53	0 30	4	5 58 37	2 18	77 35	0 24	5	7								
24	78 2	0 19	4	6 14 14	1 5	77 38	0 19	3	76 23	0 27	4	5 56 19	2 25	77 11	0 25	5	6								
25	78 21	0 18	4	6 15 19	0 58	77 57	0 18	3	75 56	0 29	5	5 53 54	2 31	76 46	0 27	5	5								
26	78 39	0 15	3	6 16 17	0 51	78 15	0 17	3	75 27	0 31	5	5 51 23	2 40	76 19	0 27	6	4								
27	78 54	0 13	3	6 17 8	0 44	78 32	0 17	3	74 56	0 32	5	5 48 43	2 46	75 52	0 27	6	3								
28	79 7	0 12	3	6 17 52	0 37	78 49	0 16	3	74 24	0 33	5	5 45 57	2 53	75 22	0 30	6	2								
29	79 19	0 10	3	6 18 29	0 30	79 5	0 13	3	73 51	0 35	5	5 43 4	2 59	74 51	0 31	6	1								
30	79 29		3	6 18 59		79 18		3	73 16		5	5 40 5		74 18	0 33	6	0								
	under		+	+		above			under		+	+		above		+									
Eq. An.	G. D. Equat.	Dif.	In. fr. M.	Mean Cent. Eq. D	Dif.	G. D. Equat.	Dif.	De fr. M.	G. D. Equat.	Dif.	D. & In. fr. M.	Mean Cent. Eq. D	Dif.	G. D. Equat.	Dif.	In & D. fr. M.	Eq. An.								

Sig. 9.													Sig. 8.												
---------	--	--	--	--	--	--	--	--	--	--	--	--	---------	--	--	--	--	--	--	--	--	--	--	--	--

EXAMPLE II. To find the Equation of the MOON'S CENTER answerable to $1^{\circ} 15' 20''$ @ à D Apog. and $3^{\circ} 5' 44''$ D Equated Anomaly.

By Table p. 65. $1^{\circ} 15' 20''$ @ à D Ap. | Dec. X^r. | Sel. X^r. | under

$80' 42'' = 80,06$
 0696
 92800
 928696
 $55,7$

Eq. An. | M. Cen. Eq. | G. D. under | L. L.
 By Tab. above $3^{\circ} 5' 44''$ | $60' 39' 43''$ | $80' 4''$ | $0.1253 -$
 -56 | $0 42 X^r$ | $1.9331 +$
 D Central Equation $6 18 47$ | $-0 56$ | $1.8078 +$
 correctly | Prop^l. Eqⁿ.

For Correction from Tab. above = -0 .. for $3^{\circ} 5' 44''$ Eq. An.

SUPPLEMENTAL

SUPPLEMENTAL LUNAR TABLES.

IV. MEAN ELLIPTIC EQUATION of the MOON'S CENTER from the *Horroxian* THEORY. With the greatest Difference of EQUATION, under and above the MEAN, correspondent to the *Least* and *Greatest* ECCENTRICITIES, 43619 and 66854, to Mean Distance à $\odot$ 1000000. According to our NEW METHOD of Computing.

Argument. MOON'S Equated ANOMALY, Or Δ Equated — Δ M. Ap. Equated by 2d Eq. Ap.

Sig. 4.										Sig. 5.									
Δ Eq. An	G. D. Equat.	Dif.	De fr. M.	Mean Cent Eq. Δ	Dif.	G. D. Equat.	Dif.	In. fr. M.	Δ	G. D. Equat.	Dif.	De fr. M.	Mean Cent Eq. Δ	Dif.	G. D. Equat.	Dif.	In. fr. M.	Δ Eq. An	
	under					above				under					above				
0	73 16	0 35	5	5 40 5	3 5	74 18	0 36	7	44 38	1 17	6	3 21 48	6 0	45 41	1 19	7	30		
1	72 41	0 38	5	5 37 0	3 14	73 42	0 36	7	43 21	1 19	7	3 15 48	6 4	44 22	1 21	7	29		
2	72 3	0 39	6	5 33 46	3 21	73 6	0 37	7	42 2	1 20	7	3 9 44	6 8	43 1	1 23	7	28		
3	71 24	0 41	6	5 30 25	3 28	72 29	0 38	7	40 42	1 21	7	3 3 36	6 12	41 38	1 23	7	27		
4	70 43	0 43	6	5 26 57	3 33	71 51	0 41	7	39 21	1 24	6	2 57 24	6 18	40 15	1 22	7	26		
5	70 0	0 42	7	5 23 24	3 41	71 10	0 42	7	37 57	1 25	6	2 51 6	6 21	38 53	1 24	6	25		
6	69 18	0 44	7	5 19 43	3 45	70 28	0 46	7	36 32	1 24	6	2 44 45	6 23	37 29	1 27	6	24		
7	68 34	0 46	7	5 15 58	3 52	69 42	0 47	7	35 8	1 25	6	2 38 22	6 27	36 2	1 27	6	23		
8	67 48	0 49	7	5 12 6	3 59	68 55	0 48	7	33 43	1 26	6	2 31 55	6 31	34 35	1 28	6	22		
9	66 59	0 52	8	5 8 7	4 7	68 7	0 47	8	32 17	1 28	6	2 25 24	6 34	33 7	1 28	6	21		
10	66 7	0 51	7	5 4 0	4 12	67 20	0 52	8	30 49	1 28	6	2 18 50	6 37	31 39	1 28	6	20		
11	65 16	0 53	7	5 59 48	4 18	66 28	0 52	8	29 21	1 30	5	2 12 13	6 41	30 11	1 31	6	19		
12	64 23	0 55	7	5 55 30	4 24	65 36	0 53	8	27 51	1 29	5	2 5 32	6 43	28 40	1 31	6	18		
13	63 28	0 56	7	4 51 6	4 30	64 43	0 58	8	26 22	1 28	5	1 58 49	6 44	27 9	1 33	5	17		
14	62 32	0 59	7	4 46 36	4 36	63 45	0 57	8	24 54	1 31	5	1 52 5	6 48	25 36	1 32	5	16		
15	61 33	0 59	7	4 42 0	4 43	62 48	0 59	8	23 23	1 32	4	1 45 17	6 51	24 4	1 32	5	15		
16	60 34	1 2	7	4 37 17	4 50	61 49	0 59	8	21 51	1 32	4	1 38 26	6 58	22 32	1 34	5	14		
17	59 32	1 2	7	4 32 27	4 55	60 50	1 0	8	20 19	1 32	4	1 31 33	6 55	20 58	1 35	5	13		
18	58 30	1 2	8	4 27 32	4 59	59 50	1 4	8	18 47	1 32	3	1 24 38	6 56	19 23	1 36	5	12		
19	57 28	1 4	8	4 22 33	5 5	58 46	1 5	8	17 15	1 33	3	1 11 42	6 58	17 47	1 36	4	11		
20	56 24	1 6	7	4 17 28	5 11	57 41	1 6	8	15 42	1 33	3	1 10 44	7 1	16 11	1 35	4	10		
21	55 18	1 5	8	4 12 17	5 16	56 35	1 8	8	14 9	1 34	2	1 3 43	7 1	14 36	1 36	3	9		
22	54 13	1 6	8	4 7 1	5 21	55 27	1 9	8	12 35	1 32	2	0 56 42	7 2	13 0	1 37	3	8		
23	53 7	1 10	8	4 1 40	5 27	54 18	1 10	8	11 3	1 35	2	0 49 40	7 4	11 23	1 37	3	7		
24	51 57	1 10	8	3 56 13	5 33	53 8	1 10	8	9 28	1 36	1	0 42 36	7 6	9 46	1 36	2	6		
25	50 47	1 11	7	3 50 40	5 37	52 58	1 13	7	7 52	1 35	1	0 35 30	7 5	8 10	1 38	1	5		
26	49 36	1 14	7	3 45 3	5 43	50 45	1 13	7	6 17	1 35	1	0 28 25	7 6	6 32	1 38	1	4		
27	48 22	1 14	7	3 39 20	5 46	49 32	1 17	7	4 42	1 33	1	0 21 19	7 6	4 54	1 38	1	3		
28	47 8	1 15	7	3 33 34	5 51	48 15	1 17	7	3 9	1 34	0	0 14 13	7 7	3 16	1 38	1	2		
29	45 53	1 15	7	3 27 43	5 55	46 58	1 17	7	1 35	1 35	0	0 7 7	7 7	1 38	1 38	0	1		
30	44 38		7	3 21 48		45 41		7	0 0		0	0 0 0		0 0		0	0		
	under			+		above		+		under			+		above		+		
Δ Eq. An	G. D. Equat.	Dif.	De fr. M.	Mean Cent Eq. Δ	Dif.	G. D. Equat.	Dif.	In. fr. M.		G. D. Equat.	Dif.	De fr. M.	Mean Cent Eq. Δ	Dif.	G. D. Equat.	Dif.	In. fr. M.	Δ Eq. An	

Sig. 7.										Sig. 6.									
---------	--	--	--	--	--	--	--	--	--	---------	--	--	--	--	--	--	--	--	--

Sig. 7.

Sig. 6.

N. B. For other Methods of solving the *Elliptic Equation* of the MOON'S CENTER, See farther on.

The ROYAL ASTRONOMER SUPPLEMENTAL LUNAR TABLES.

V. EQUATION, or MOON'S Least and Greatest VARIATION, Sun in Apog. and Perigee, in respect of the Earth's Orbit: According to the *Newtonian Theory*.

Argument. Moon equated from the SUN.

Sig. 0. 6.				Sig. 1. 7.				Sig. 2. 8.			
L. Eq.	Dif.	G. D.	Per.	L. Eq.	Dif.	G. D.	Per.	L. Eq.	Dif.	G. D.	Per.
in Ap.	+	+	+	in Ap.	+	+	+	in Ap.	+	+	+
0	0	0	0	0	28	41	0	0	28	41	0
1	1	9	1	1	34	112	28	1	34	203	29
2	2	18	1	2	31	216	27	2	38	198	28
3	3	27	1	3	29	219	26	3	40	194	27
4	4	37	1	4	29	223	26	4	42	189	26
5	5	47	1	5	24	226	25	5	43	184	25
6	6	57	1	6	23	228	24	6	47	178	24
7	7	67	1	7	20	231	23	7	47	173	23
8	8	77	1	8	18	233	23	8	49	167	22
9	9	87	1	9	16	235	22	9	51	161	21
10	10	97	1	10	13	236	21	10	52	154	20
11	11	107	1	11	11	238	20	11	53	148	19
12	12	117	1	12	8	239	19	12	57	141	18
13	13	127	1	13	7	239	18	13	57	134	17
14	14	137	1	14	4	240	17	14	57	127	16
15	15	147	1	15	2	240	16	15	59	120	15
16	16	157	1	16	0	240	15	16	1	113	14
17	17	167	1	17	0	239	14	17	2	105	13
18	18	177	1	18	0	239	13	18	3	98	12
19	19	187	1	19	0	238	12	19	4	90	11
20	20	197	1	20	0	238	11	20	4	82	10
21	21	207	1	21	0	235	10	21	5	74	9
22	22	217	1	22	0	233	9	22	7	66	8
23	23	227	1	23	0	231	8	23	7	58	7
24	24	237	1	24	0	228	6	24	9	50	6
25	25	247	1	25	0	226	5	25	10	42	5
26	26	257	1	26	0	223	4	26	11	33	4
27	27	267	1	27	0	219	3	27	11	25	3
28	28	277	1	28	0	216	2	28	12	17	2
29	29	287	1	29	0	212	1	29	12	8	1
30	30	297	1	30	0	208	0	30	13	0	0

CONSTRUCTION. The Variation being without the D's Octants with $\odot$.

As Rad. : S. Doub. Diff. D fr. next Sisygy or Quadrature in Arg. as above :: S. Gr. Variation in Octants D à $\odot$, $\odot$ being in Apogee and Perigee, respectively : S. Pref. Equations in Apog. and Perig. as above, viz. Rad. : S. 33' 10" and S. 37' 10" :: S. 2. D à $\odot$: S. Pref. Equations in Ap. and Perig. $\odot$.

Or, 33' 10" : 240" :: any L. Eq. : G. D. as above.

i. e. $\frac{33' 10"}{240} \times \text{any L. Eq.} = \text{G. D. as above.}$

And, At 33' 10" gr. Eq. in Octants D à $\odot$, $\odot$ being in Ap. : Pref. Eq. in Octants D à $\odot$ according to $\odot$ M. An. :: Least Eq. Sun being in Ap. : Present Variation, required.

N. B. See Variation Tables following, p. 73 and 74.

DECIMAL MULTIPLIERS to the Greatest Dif. above the Least Variation. The Product to be added thereto for Present Variation.

Argument. SUN'S MEAN ANOMALY.

0	1	2	3	4	5
0.000	0.065	0.25	0.5	0.75	0.935
1.001	0.07	0.257	0.507	0.76	0.939
2.001	0.074	0.264	0.516	0.766	0.943
3.002	0.079	0.272	0.525	0.774	0.946
4.003	0.083	0.28	0.534	0.782	0.95
5.003	0.088	0.287	0.543	0.79	0.954
6.004	0.092	0.295	0.552	0.797	0.958
7.005	0.098	0.303	0.561	0.804	0.961
8.006	0.104	0.312	0.57	0.81	0.964
9.008	0.109	0.32	0.578	0.816	0.968
10.009	0.115	0.33	0.587	0.823	0.971
11.01	0.121	0.337	0.596	0.83	0.974
12.012	0.126	0.345	0.605	0.835	0.977
13.013	0.133	0.353	0.614	0.842	0.981
14.014	0.14	0.362	0.622	0.848	0.985
15.015	0.146	0.37	0.63	0.855	0.989
16.017	0.152	0.38	0.64	0.861	0.993
17.018	0.158	0.386	0.647	0.867	0.996
18.019	0.165	0.395	0.655	0.874	1.
19.023	0.171	0.404	0.664	0.88	1.
20.027	0.177	0.413	0.672	0.885	1.
21.031	0.183	0.422	0.68	0.891	1.
22.035	0.19	0.43	0.688	0.897	1.
23.038	0.195	0.44	0.687	0.902	1.
24.042	0.201	0.45	0.705	0.908	1.
25.046	0.21	0.457	0.713	0.913	1.
26.05	0.217	0.465	0.720	0.917	1.
27.054	0.225	0.473	0.728	0.922	1.
28.057	0.233	0.482	0.736	0.926	1.
29.061	0.241	0.49	0.743	0.931	1.
30.065	0.25	0.5	0.751	0.935	1.

CONSTRUCTION. As gr. Dif. Eq. D in Octants à Sun (Sun in Apog. and Perig.) viz. 240" : Pref. Dif. in Octants according to M. An. Sun :: G. D. above L. Eq. Sun in Ap. and Perig. : Pref. Dif. to be added to the L. Eq. Sun in Ap. for the Variation required.

Hence, Pref. D. D Oct. (= Mult.) $\times$ 240"

G. D. D out of Octants = Pref. Dif. to be added to L. Equation.

Ex. To find the Moon's Pref. Var. to $\odot$ 25° D à $\odot$ and 3° 26' Sun's M. An. ?

L. Eq. G. D.
 $\odot$ 25° D à $\odot$ 25' 24" 184"
 Prop^d Eq. $\times$ 2 11 $\times$ 72
 D Pref. Var. req^d. 27 35 368
 1288
 13248
 Prop^d Eq. $\times$ 2 11"

MOON'S Greatest VARIATION in Octants of D à $\odot$, from Apog. to Perigee $\odot$.

Arg. SUN'S M. ANOM.

0	1	2	3	4	5
0.033	0.4	0.012			
1.033	5	3 24			
2.033	7	7 18			
3.033	10	14 12			
4.033	14	22 6			
5.033	19	33 0 11			
6.033	27	50 24			
7.033	35	68 18			
8.033	44	87 12			
9.033	54	100 6			
10.034	4	130 0 10			
11.034	10	155 24			
12.034	28	180 18			
13.034	41	208 12			
14.034	54	235 6			
15.035	8	264 0 9			
16.035	22	292 24			
17.035	35	319 18			
18.035	49	347 12			
19.036	2	374 6			
20.036	15	400 0 8			
21.036	27	423 24			
22.036	38	445 18			
23.036	48	465 12			
24.036	58	485 6			
25.037	6	500 0 7			
26.037	13	514 24			
27.037	18	524 18			
28.037	22	531 12			
29.037	24	535 6			
30.037	25	537 0 6			

CONSTR. The above Eq^s in Oct. of D à $\odot$ incr. according to the Dif. of Cubes $\odot$ Dist. à $\odot$ fr. 33' 4" Var. in Per. $\odot$. Ex. Of the foregoing Case.

L. L.
 D à $\odot$ 25° 25' 24" Eq. 0.3733
 3° 26' $\odot$ M. An. -382

D Pref. Var. 27' 44" 3351 according to this Hypothesis.

Accord^s to Sir

I. Newton, G. Oct. D à $\odot$.

Var. in D Oct. $\odot$ Ap. 33' 14"

is in Sq. of D's $\odot$ Per. 37 11

Syn. rev. diry, Variation.

& Cub. $\odot$ à $\odot$

inversly.

Inversly, acc^s Oct. D à $\odot$.

to Cubes of $\odot$ Ap. 33' 26"

Dif. à $\odot$, M. $\odot$ Per. 37 1

Dif. 35' 10". Variation.

Accord^s to $\odot$ Ap. 33 16

Dr. Halley. $\odot$ Per. 37 14 Var.

SUPPLEMENTAL LUNAR TABLES.

OR, V. EQUATION of the MOON'S Mean VARIATION. According to the NEWTONIAN THEORY.

☉ in Octants, or at a Mean Dist. à ☉.

LOGISTICAL LOGARITHMS, to be added to the L. L. of the Mean Variation for the L. Log. of the correct Variation.

Arg. MOON Equated from the SUN.

Arg. SUN'S Mean ANOMALY.

☉ à	Sig. 0. 6.		Sig. 1. 7.		Sig. 2. 8.		☉ à	☉M.	Sig. 0.	Sig. 1.	Sig. 2.	Sig. 3.	Sig. 4.	Sig. 5.	☉M.
0.	+	Dif.	+	Dif.	+	Dif.	0.	An.	L. L.	L. L.	L. L.	L. L.	L. L.	L. L.	An.
0	1	2	1	2	1	2	0	0	+	+	+	+	+	+	0
0	0 0	1 14	30 27	0 36	30 27	0 38	30	0	0.0242	0.0211	0.0125	0.0004	9.9880	9.9787	30
1	1 14	1 13	31 3	0 33	29 49	0 40	29	1	0.0242	0.0209	0.0121	0.0000	9.9876	9.9785	29
2	2 27	1 13	31 36	0 31	29 9	0 32	28	2	0.0242	0.0207	0.0118	9.9995	9.9872	9.9783	28
3	3 40	1 14	32 7	0 29	28 27	0 44	27	3	0.0242	0.0205	0.0114	9.9991	9.9868	9.9781	27
4	4 54	1 12	32 36	0 27	27 43	0 47	26	4	0.0242	0.0203	0.0110	9.9987	9.9865	9.9779	26
5	6 6	1 13	33 3	0 24	25 56	0 48	25	5	0.0242	0.0201	0.0106	9.9982	9.9861	9.9777	25
6	7 19	1 11	33 27	0 21	26 8	0 50	24	6	0.0241	0.0198	0.0102	9.9978	9.9858	9.9775	24
7	8 30	1 11	33 48	0 19	25 18	0 52	23	7	0.0241	0.0195	0.0098	9.9973	9.9854	9.9773	23
8	9 41	1 11	34 7	0 17	24 26	0 54	22	8	0.0240	0.0193	0.0095	9.9969	9.9851	9.9772	22
9	10 52	1 10	34 24	0 14	23 32	0 56	21	9	0.0240	0.0190	0.0091	9.9965	9.9847	9.9770	21
10	12 2	1 8	34 38	0 12	22 36	0 57	20	10	0.0239	0.0188	0.0087	9.9961	9.9844	9.9769	20
11	13 10	1 8	34 50	0 9	21 39	0 59	19	11	0.0238	0.0185	0.0083	9.9956	9.9840	9.9767	19
12	14 18	1 7	34 59	0 6	20 40	1 0	18	12	0.0237	0.0182	0.0079	9.9952	9.9837	9.9766	18
13	15 25	1 5	35 5	0 4	19 40	1 2	17	13	0.0236	0.0179	0.0075	9.9948	9.9833	9.9764	17
14	16 30	1 3	35 9	0 1	18 38	1 3	16	14	0.0235	0.0177	0.0071	9.9944	9.9830	9.9763	16
15	17 35	1 2	35 10	0 1	17 35	1 5	15	15	0.0234	0.0174	0.0067	9.9940	9.9827	9.9761	15
16	18 38	1 0	35 9	0 4	16 30	1 5	14	16	0.0233	0.0171	0.0063	9.9936	9.9824	9.9760	14
17	19 40	0 59	35 5	0 6	15 25	1 7	13	17	0.0232	0.0168	0.0059	9.9932	9.9822	9.9759	13
18	20 40	0 57	34 59	0 9	14 18	1 8	12	18	0.0231	0.0165	0.0055	9.9928	9.9818	9.9758	12
19	21 39	0 56	34 50	0 12	13 10	1 8	11	19	0.0230	0.0162	0.0050	9.9924	9.9815	9.9758	11
20	22 36	0 54	34 38	0 14	12 2	1 10	10	20	0.0229	0.0159	0.0046	9.9920	9.9812	9.9757	10
21	23 32	0 52	34 24	0 17	10 52	1 11	9	21	0.0227	0.0156	0.0042	9.9916	9.9809	9.9756	9
22	24 26	0 50	34 7	0 19	9 41	1 11	8	22	0.0226	0.0153	0.0038	9.9912	9.9807	9.9755	8
23	25 18	0 48	33 48	0 21	8 30	1 11	7	23	0.0224	0.0149	0.0033	9.9908	9.9804	9.9755	7
24	26 8	0 47	33 27	0 24	7 19	1 13	6	24	0.0223	0.0146	0.0029	9.9904	9.9802	9.9754	6
25	26 56	0 44	33 3	0 27	6 6	1 12	5	25	0.0220	0.0142	0.0025	9.9900	9.9799	9.9753	5
26	27 43	0 42	32 36	0 29	4 54	1 14	4	26	0.0219	0.0139	0.0021	9.9896	9.9797	9.9753	4
27	28 27	0 40	32 7	0 31	3 40	1 13	3	27	0.0217	0.0135	0.0016	9.9892	9.9794	9.9753	3
28	29 9	0 38	31 36	0 33	2 27	1 13	2	28	0.0215	0.0132	0.0012	9.9888	9.9792	9.9753	2
29	29 49		31 3	0 36	1 14	1 14	1	29	0.0213	0.0128	0.0008	9.9884	9.9789	9.9753	1
30	30 27		30 27		0 0		0	30	0.0211	0.0125	0.0004	9.9880	9.9787	9.9753	0
☉ à	—	Dif.	—	Dif.	—	Dif.	☉ à	☉M.	+	+	+	+	+	+	☉M.
☉.	Sig. 11. 5.		Sig. 10. 4.		Sig. 9. 3.		☉.	An.	L. L.	L. L.	L. L.	L. L.	L. L.	L. L.	An.

N. B. The above, or our 5th, is Sir Isaac Newton's 6th Equation.

Construction. As Rad. S. 35' 10" Gr. Mean Variation in Octants of ☉ à ☉, : 2. ☉ à ☉ (from next Quadrature or Sisygy) Pref. Mean Variation, as above, ☉ being in Octants, or at a Mean Distance from ☉.

Ex. To find the correct Variation to 7° 24' Arg. of mean Variation, and 2° 5' of Sun's M. An.?

Eq.	L. L.
Arg. 7° 24' 33' 27"	2538
Sun's M. An. 2° 5'	0106 +
Cor. Var. reqd. 32' 39" reqd.	2644

CONSTRUCTION. As greatest M. Variation 35' 10", ☉ in Octants à ☉, and ☉ in Octants or at a Mean Dist. à ☉ : Present M. Variation, ☉ being in Octants :: great. Variation, ☉ in Octants of ☉, correspondent to the ☉ M. Anomaly : Present Variation required.

Hence. P. M. Var. x P. G. Var. ☉ in Oct. = Pr.

true Var. And Hence + L. L. Pref. G. Var. in Oct. — L. L. 35' 10" = L. L. to be added to, or connected with, the L. L. of the present M. Variation, for L. L. of present Correct Variation, required.

SUPPLEMENTAL

SUPPLEMENTAL LUNAR TABLES.

V. EQUATION, or MOON'S VARIATION.
According to the Horrobian Theory.

Arg. MOON from the SUN.									
D à ⊙	Sig. 0. 6		Sig. 1. 7		Sig. 2. 8		D à ⊙	D à ⊙	D à ⊙
	+	Dif.	+	Dif.	+	Dif.			
0	0	0	32	54	32	54	0	41	30
1	1	20	33	33	32	13	0	43	29
2	2	39	34	9	31	30	0	45	28
3	3	59	34	42	30	45	0	48	27
4	5	18	35	13	29	57	0	50	26
5	6	36	35	42	29	7	0	53	25
6	7	54	36	9	28	14	0	54	24
7	9	11	36	32	27	20	0	56	23
8	10	28	36	52	26	24	0	58	22
9	11	44	37	10	25	26	1	1	21
10	13	0	37	25	24	25	1	2	20
11	14	14	37	38	23	23	1	3	19
12	15	27	37	48	22	20	1	5	18
13	16	40	37	55	21	15	1	7	17
14	17	51	37	59	20	8	1	8	16
15	19	0	38	0	19	0	1	9	15
16	20	8	37	59	17	51	1	11	14
17	21	15	37	55	16	40	1	13	13
18	22	20	37	48	15	27	1	13	12
19	23	23	37	38	14	14	1	14	11
20	24	25	37	25	13	0	1	14	10
21	25	26	37	10	11	44	1	16	9
22	26	24	36	52	10	28	1	16	8
23	27	20	36	32	9	11	1	17	7
24	28	14	36	9	7	54	1	18	6
25	29	7	35	42	6	36	1	18	5
26	29	57	35	13	5	18	1	19	4
27	30	45	34	42	3	59	1	20	3
28	31	30	34	9	2	39	1	20	2
29	32	13	33	33	1	20	1	20	1
30	32	54	32	54	0	0	1	20	0
D à ⊙	—		—		—		D à ⊙	—	
⊙	Sig. 11. 5		Sig. 10. 4		Sig. 9. 3		⊙		

CONSTRUCTION.

As Rad. : S. Double Argument, or 2. D à ⊙ :: 38'
Equation at Greatest : Present Equation, required.

EXAMPLE. To find the Equation for 26° Argument ?

2 X 26° = 52° . . Sine ,788 to Rad. 1.

Gr. Eq. 38'

6304

2364

29,944

So for the Ref.

56",64 Eq. requd.

N. B. The above Eqn. and 7th. correctd. being connected will form a compd. Variation. See our 1st and 2d Var. foregoing : also Mayer's compound Var. following.

The 6th. and 7th. Eqns. to the Right are improved in Exactness by varying them in the inverse Ratio of D à ⊙, or directly, as the Moon's Hor. Parallax, supplied by the last Tables of Correction.

VI. EQUATION of the MOON, or
2d Equation of the Center. Ac-
cording to the Newtonian Theory.

Arg. D à ⊙ + D Ap. à ⊙ Ap.									
Arg. pro ^r	S. 6		S. 7		S. 8		Arg. pro ^r	D à ⊙	D à ⊙
	+	Dif.	+	Dif.	+	Dif.			
0	0	0	1	12	2	5	30		
1	0	2	1	15	2	7	29		
2	0	5	1	17	2	8	28		
3	0	7	1	19	2	9	27		
4	0	10	1	21	2	10	26		
5	0	12	1	23	2	11	25		
6	0	15	1	25	2	12	24		
7	0	17	1	27	2	13	23		
8	0	20	1	29	2	14	22		
9	0	22	1	31	2	15	21		
10	0	25	1	33	2	16	20		
11	0	27	1	35	2	17	19		
12	0	30	1	37	2	17	18		
13	0	32	1	39	2	18	17		
14	0	35	1	41	2	19	16		
15	0	37	1	43	2	20	15		
16	0	40	1	44	2	20	14		
17	0	42	1	46	2	21	13		
18	0	44	1	48	2	21	12		
19	0	46	1	49	2	22	11		
20	0	49	1	51	2	22	10		
21	0	51	1	52	2	23	9		
22	0	54	1	54	2	23	8		
23	0	56	1	55	2	24	7		
24	0	59	1	57	2	24	6		
25	1	1	1	58	2	24	5		
26	1	3	2	0	2	24	4		
27	1	5	2	1	2	25	3		
28	1	8	2	3	2	25	2		
29	1	10	2	4	2	25	1		
30	1	12	2	5	2	25	0		
Arg. pro ^r	S. 11		S. 10		S. 9		Arg. pro ^r		
⊙	5		4		3		⊙		

Argument above transforms to 2. D à ⊙ + ⊙

An. — D An. The above corrects the 1st Equation D's Center, and is properly Sir Isaac's 5th.

Construction. As Rad. : S. Pref. Argument from nearest Sine or Quadrature : 2' 25"

Equation at Greatest : Present Equation, required.

EXAMPLE. To find the Equation for 30° Argument ?

As 1 Rad. : 2',416 Gr. E. :: S. 30° = ,5

5

1',2080

12" Eq. required. So for the ref.

N. B. Dr. Gregory (*Astr.* p. 568) erroneously gives 2' 10" for 2' 25" gr. Eqn above, with contrary Sign. to those of Sir Is. Newton (see last Edit. Princip. Prop. 35. B. 3) and therefore,

The { + 2' 25" } = + { 4' 35" } Gr. Error.

Diff. { — 2' 10" } = — { 4' 35" } Error.

Which Error Leadbetter and Duntborne have strictly followed. Dr. Halley misplaced this E-

quation to precede the 4th or central that it should follow, occasioning Error and Trouble to mend it by a Corr. of the Arg. there misplaced.

VII. EQUATION of the MOON,
or, 2d Mean Variation. Ac-
cording to Gregory's Astrono-
my. P. 569.

Argument. MOON from the SUN.									
D à ⊙	S. 6		S. 7		S. 8		D à ⊙	D à ⊙	D à ⊙
	+	Dif.	+	Dif.	+	Dif.			
0	0	0	1	10	2	1	30		
1	0	2	1	12	2	2	29		
2	0	5	1	14	2	4	28		
3	0	7	1	16	2	5	27		
4	0	10	1	18	2	6	26		
5	0	12	1	20	2	7	25		
6	0	15	1	22	2	8	24		
7	0	17	1	24	2	9	23		
8	0	19	1	26	2	10	22		
9	0	22	1	28	2	11	21		
10	0	24	1	30	2	12	20		
11	0	27	1	32	2	12	19		
12	0	29	1	34	2	13	18		
13	0	32	1	35	2	14	17		
14	0	34	1	37	2	15	16		
15	0	36	1	39	2	15	15		
16	0	39	1	41	2	16	14		
17	0	41	1	42	2	16	13		
18	0	43	1	44	2	17	12		
19	0	46	1	46	2	17	11		
20	0	48	1	47	2	18	10		
21	0	50	1	49	2	18	9		
22	0	52	1	50	2	19	8		
23	0	55	1	52	2	19	7		
24	0	57	1	53	2	19	6		
25	0	59	1	55	2	19	5		
26	1	1	1	56	2	20	4		
27	1	4	1	57	2	20	3		
28	1	6	1	59	2	20	2		
29	1	8	2	0	2	20	1		
30	1	10	2	1	2	20	0		
D à ⊙	S. 11		S. 10		S. 9		D à ⊙		
⊙	5		4		3		⊙		

The above corrects 1st Variation.

CONSTR. As Rad. : S. Arg^t, or Pr. Diff. D à ⊙ :: 2' 20" Eq. at Greatest :

the Present Equation,

For { Gr. E. 2',33 = 2' 20"

30° Arg^t } Sine . . . 5

Eq. 1',165 = 1' 20" req.

So for the ref.

The above Eqn. varies according to the

Position of Moon and Sun's Ap. — When

they are together 54" add to, but when

opposite subtr. fr. 2' 20" (the M. gr. Eqn.

in Quad. Moon à Sun) for the pref. Gr.

Eqn. varying bet. the gr. 3' 14" and least

Eqn. 1' 26". While the Moon's Ap. is

also in Syagies with the Sun — and conly.

the proper Arg. = 0° and 6° Sun's An.

respectively.

But in Quad. Sun à Moon's Ap. the

Moon's Ap. and Sun's Ap. being also to-

gether, and conly. the proper Arg. = Sun's

An. 3° or 9°, 50" or 1'. (See Grig.)

SUPPLEMENTAL LUNAR TABLES.

CORRECTION of Gregory's VII. or 2d Variation.					VII. EQUATION of the MOON, or Second Mean Variation. According to the Newtonian Theory.					CORRECTION of the 7th EQUATION, or 2d Mean Variation. According to Sir Isaac Newton.					Farther CORRECTION; according to the Dif. of the MOON'S Distance from the EARTH and SUN.				
Arg. SUN'S true ANOMALY.					Arg. Moon from the Sun.					Arg. Moon's Ap. fr. Sun's Per.					Arg. M. An. D à M. An. O.				
Sun's true An.	Gr. Eq. in Qua. D à Sun.	Xrs of Mean Var.	L. L. +	Sun's true An.	D S.	S.	S.	D	Dap. Gr. Eq. in Qua. D à Sun.	Xrs of 2d M. Var.	L. L. +	Dap. Gr. Eq. in Qua. D à Sun.	Xrs of 2d M. Var.	L. L. +	A. D à A. O.	0	1	2	A. D à A. O.
0 0	1 26	,614	2166	0 12	0 0	1 30	2 36	30	0 0	1 0	,333	4772	0 12	0 0	0	+	+	+	0
6	1 26	,614	2166	24	1 0	3 1	32	29	6	1 0	,333	4772	24	0 0	0	+	+	+	24
12	1 27	,621	2066	18	2 0	6 1	35	28	12	1 2	,344	4629	18	0 0	0	+	+	+	18
18	1 28	,628	2016	12	3 0	9 1	38	27	18	1 6	,366	4358	12	1 0	1	0	20	0	35
24	1 31	,650	1871	6	4 0	13 1	40	26	24	1 11	,394	4040	6	2 0	2	0	21	0	36
1 0	1 32	,650	1823	0 11	5 0	15 1	43	25	1 0	1 16	,422	3745	0 11	3 0	2	0	21	0	36
6	1 36	,685	1638	24	6 0	19 1	45	24	6	1 23	,461	3362	24	4 0	3	0	22	0	36
12	1 41	,721	1418	18	7 0	22 1	48	23	12	1 31	,505	2963	18	5 0	4	0	23	0	37
18	1 43	,735	1333	12	8 0	24 1	51	22	18	1 40	,555	2553	12	6 0	4	0	23	0	37
24	1 46	,757	1208	6	9 0	28 1	53	21	24	1 49	,605	2179	6	7 0	5	0	24	0	37
2 0	1 51	,792	1008	0 10	10 0	31 1	56	20	2 0	2 0	,666	1761	0 10	8 0	6	0	24	0	37
6	1 56	,828	0816	24	11 0	35 1	58	19	6	2 11	,728	1380	24	9 0	6	0	25	0	38
12	2 1	,864	0633	18	12 0	37 2	1	18	12	2 23	,794	1000	18	10 0	7	0	26	0	38
18	2 6	,900	0457	12	13 0	41 2	2	17	18	2 35	,861	0650	12	11 0	7	0	26	0	38
24	2 13	,950	0223	6	14 0	44 2	5	16	24	2 47	,928	0326	6	12 0	8	0	27	0	38
3 0	2 20	1,000	0000	0 9	15 0	46 2	7	15	3 0	3 0	1,000	0000	0 9	13 0	9	0	27	0	38
6	2 25	1,035	0153	24	16 0	50 2	10	14	6	3 13	1,072	0303	24	14 0	9	0	28	0	38
12	2 30	1,071	0300	18	17 0	53 2	11	13	12	3 25	1,139	0565	18	15 0	10	0	28	0	39
18	2 35	1,107	0442	12	18 0	55 2	14	12	18	3 37	1,205	0812	12	16 0	11	0	29	0	39
24	2 43	1,164	0661	6	19 0	59 2	16	11	24	3 49	1,272	1045	6	17 0	11	0	29	0	39
4 0	2 49	1,207	0818	0 8	20 1	2 18	2 57	10	4 0	4 0	1,333	1249	0 8	18 0	12	0	30	0	39
6	2 54	1,243	1044	24	21 1	4 20	2 57	9	6	4 10	1,388	1426	24	19 0	13	0	30	0	39
12	2 59	1,278	1068	18	22 1	7 21	2 59	8	12	4 20	1,444	1579	18	20 0	13	0	31	0	39
18	3 4	1,314	1187	12	23 1	11 22	2 59	7	18	4 29	1,494	1744	12	21 0	14	0	31	0	39
24	3 6	1,328	1234	6	24 1	13 23	2 59	6	24	4 37	1,539	1872	6	22 0	15	0	32	0	40
5 0	3 9	1,350	1304	0 7	25 1	16 24	2 59	5	5 0	4 44	1,578	1980	0 7	23 0	15	0	32	0	40
6	3 14	1,385	1417	24	26 1	18 25	3 0	5	6	4 50	1,611	2071	25	24 0	16	0	33	0	40
12	3 14	1,385	1417	18	27 1	22 26	3 0	3	12	4 54	1,633	2130	18	25 0	17	0	33	0	40
18	3 14	1,385	1417	12	28 1	25 27	3 0	2	18	4 57	1,650	2175	12	26 0	17	0	34	0	40
24	3 14	1,385	1417	6	29 1	27 28	3 0	1	24	4 59	1,661	2204	6	27 0	18	0	34	0	40
6 0	3 14	1,385	1417	0 6	30 1	30 29	3 0	0	6 0	5 0	1,666	2218	0 6	28 0	19	0	34	0	40
Sun's true An.	Gr. Eq. in Qua. D à Sun.	Xrs of Mean Var.	L. L. +	Sun's true An.	D S.	S.	S.	D	Dap. Gr. Eq. in Qua. D à Sun.	Xrs of 2d M. Var.	L. L. +	Dap. Gr. Eq. in Qua. D à Sun.	Xrs of 2d M. Var.	L. L. +	A. D à A. O.	11	10	9	A. D à A. O.
0 0	1 26	,614	2166	0 12	0 0	1 30	2 36	30	0 0	1 0	,333	4772	0 12	0 0	0	+	+	+	0
6	1 26	,614	2166	24	1 0	3 1	32	29	6	1 0	,333	4772	24	0 0	0	+	+	+	24
12	1 27	,621	2066	18	2 0	6 1	35	28	12	1 2	,344	4629	18	0 0	0	+	+	+	18
18	1 28	,628	2016	12	3 0	9 1	38	27	18	1 6	,366	4358	12	1 0	1	0	20	0	35
24	1 31	,650	1871	6	4 0	13 1	40	26	24	1 11	,394	4040	6	2 0	2	0	21	0	36
1 0	1 32	,650	1823	0 11	5 0	15 1	43	25	1 0	1 16	,422	3745	0 11	3 0	2	0	21	0	36
6	1 36	,685	1638	24	6 0	19 1	45	24	6	1 23	,461	3362	24	4 0	3	0	22	0	36
12	1 41	,721	1418	18	7 0	22 1	48	23	12	1 31	,505	2963	18	5 0	4	0	23	0	37
18	1 43	,735	1333	12	8 0	24 1	51	22	18	1 40	,555	2553	12	6 0	4	0	23	0	37
24	1 46	,757	1208	6	9 0	28 1	53	21	24	1 49	,605	2179	6	7 0	5	0	24	0	37
2 0	1 51	,792	1008	0 10	10 0	31 1	56	20	2 0	2 0	,666	1761	0 10	8 0	6	0	24	0	37
6	1 56	,828	0816	24	11 0	35 1	58	19	6	2 11	,728	1380	24	9 0	6	0	25	0	38
12	2 1	,864	0633	18	12 0	37 2	1	18	12	2 23	,794	1000	18	10 0	7	0	26	0	38
18	2 6	,900	0457	12	13 0	41 2	2	17	18	2 35	,861	0650	12	11 0	7	0	26	0	38
24	2 13	,950	0223	6	14 0	44 2	5	16	24	2 47	,928	0326	6	12 0	8	0	27	0	38
3 0	2 20	1,000	0000	0 9	15 0	46 2	7	15	3 0	3 0	1,000	0000	0 9	13 0	9	0	27	0	38
6	2 25	1,035	0153	24	16 0	50 2	10	14	6	3 13	1,072	0303	24	14 0	9	0	28	0	38
12	2 30	1,071	0300	18	17 0	53 2	11	13	12	3 25	1,139	0565	18	15 0	10	0	28	0	39
18	2 35	1,107	0442	12	18 0	55 2	14	12	18	3 37	1,205	0812	12	16 0	11	0	29	0	39
24	2 43	1,164	0661	6	19 0	59 2	16	11	24	3 49	1,272	1045	6	17 0	11	0	29	0	39
4 0	2 49	1,207	0818	0 8	20 1	2 18	2 57	10	4 0	4 0	1,333	1249	0 8	18 0	12	0	30	0	39
6	2 54	1,243	1044	24	21 1	4 20	2 57	9	6	4 10	1,388	1426	24	19 0	13	0	30	0	39
12	2 59	1,278	1068	18	22 1	7 21	2 59	8	12	4 20	1,444	1579	18	20 0	13	0	31	0	39
18</																			

SUPPLEMENTAL LUNAR TABLES.

SECOND or Semiannual EQUATION of the MOON'S NODE. According to the Newtonian Theory, With Decimal Multipliers and Logistical Logarithms to be applied to the greatest Excesses of Reduction and Latitude of Δ , for finding the present Excesses thereof, respectively, above the REDUCTION and LATITUDE for the Least Inclination of the Lunar Orbit ($4^{\circ} 59' 35''$). Also the Present Excesses of Inclination above the Least Inclination.

Argument. SUN from the MOON'S NODE.

Sig. o. 6.										Sig. 1. 7.										Sig. 2. 8.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																														
⊙ a ⊙	Aqua. Ⓢ.		Dif.	Xrs to		Lo. Log. for Red. Lat.	G. Ex. above L. Inc.			⊙ a ⊙	Aqua. Ⓢ.		Dif.	Xrs. to		Lo. Log. for Red & Lat	G. Ex. above L. Inc.			⊙ a ⊙	Aquat. Ⓢ.		Dif.	Xrs. to		Lo. Log. for Red & Lat	G. Ex. above L. Inc.			⊙ a ⊙																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																				
	+	-		R ⁿ & L ^e	+			-	+		-	R ⁿ & L ^e		+	-			+	-		R ⁿ & L ^e	+		-	+			-	R ⁿ & L ^e		+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-	+	-	R ⁿ & L ^e	+	-

CONSTRUCTION of the above EQUATION Ⓢ. Com. Log.
As Conf. N^o. 59 (Sum of two Sides Δ) . . . Co. 8.2291480
To Conf. N^o. 56 (their Dif.) . . . 1.7481880
So Tan. $\odot$ a $\odot$. Suppose 46° ($\frac{1}{2}$ Sum Opp. $\angle$ s) 10.0151628
To Tan. Arch ($\frac{1}{2}$ Dif. $\angle$ s) $44^{\circ} 30' 10''$. . . 9.9924988
Deduct $\frac{1}{2}$ Dif. fr. $\frac{1}{2}$ Sum } 1 29 41 the eq. Ⓢ. So for the rest.
 $\angle$ s, rem. left $\angle$.

The Least Inclination Δ 's Orb with the Ecliptic being $4^{\circ} 59' 35''$ when $\odot$ s are in the $\odot$'s Quadratures, and greatest Inclination thereto $5^{\circ} 17' 20''$. When they are in Conjunction and Opposition with $\odot$; it follows that the greatest Excess of Inclination is $17' 45''$. See Prop 35. B. 3. Principia, for Construction and Solution of the different Inclinations of Δ 's Orbit with Ecliptic.—Now, as $17' 45''$. Gr. Excess Inclination: Pref. Ex. thereof :: Gr. Ex^s. of Reduction and of Latitude: Pref. Ex. thereof, respectively, to be added to those for the Least Inclination Δ Orbit, for the Red. and Lat. Δ , respectively required. Hence, Pref. Exc. Incl. = Decim. Xr. And Lo. Log. Pref. Ex^s.

Incl. — L. L. $17' 45''$ = Lo. Log. for the Red. and Lat. Δ ; as above which Xrs. and Lo. Log. are found for min^s. above Deg^s $\odot$ a $\odot$ by proportion of their Diff^s.

Or, As Sum Forces 115 (=59+56) 7.9393022. co.
To their Dif. . 3 (=59-56) 0.4771223
So Radius 10.0000000
To S. Gr. Aeq. $\odot$. $1^{\circ} 29' 41''$ 8.4164235
as before.

SUPPLEMENTAL LUNAR TABLES.

REDUCTION-EQUATION of the MOON from her Orbit to the EC-LIPTIC. For the least and greatest Inclination; with the greatest Excess of Reduction.

Arg. of Lat. or Moon fr. Node. 2. Eq.

Sig. 6. 6.				Sig. 1. 7.				Sig. 2. 8.			
Red. Eq.		gr. ex.		Red. Eq.		gr. ex.		Red. Eq.		gr. ex.	
for L. In.		for L. In.		for L. In.		for L. In.		for L. In.		for L. In.	
D.	Lat.	D.	Lat.	D.	Lat.	D.	Lat.	D.	Lat.	D.	Lat.
0	0	0	0	0	0	0	0	0	0	0	0
1	0	1	0	1	0	1	0	1	0	1	0
2	0	2	0	2	0	2	0	2	0	2	0
3	0	3	0	3	0	3	0	3	0	3	0
4	0	4	0	4	0	4	0	4	0	4	0
5	0	5	0	5	0	5	0	5	0	5	0
6	0	6	0	6	0	6	0	6	0	6	0
7	0	7	0	7	0	7	0	7	0	7	0
8	0	8	0	8	0	8	0	8	0	8	0
9	0	9	0	9	0	9	0	9	0	9	0
10	0	10	0	10	0	10	0	10	0	10	0
11	0	11	0	11	0	11	0	11	0	11	0
12	0	12	0	12	0	12	0	12	0	12	0
13	0	13	0	13	0	13	0	13	0	13	0
14	0	14	0	14	0	14	0	14	0	14	0
15	0	15	0	15	0	15	0	15	0	15	0
16	0	16	0	16	0	16	0	16	0	16	0
17	0	17	0	17	0	17	0	17	0	17	0
18	0	18	0	18	0	18	0	18	0	18	0
19	0	19	0	19	0	19	0	19	0	19	0
20	0	20	0	20	0	20	0	20	0	20	0
21	0	21	0	21	0	21	0	21	0	21	0
22	0	22	0	22	0	22	0	22	0	22	0
23	0	23	0	23	0	23	0	23	0	23	0
24	0	24	0	24	0	24	0	24	0	24	0
25	0	25	0	25	0	25	0	25	0	25	0
26	0	26	0	26	0	26	0	26	0	26	0
27	0	27	0	27	0	27	0	27	0	27	0
28	0	28	0	28	0	28	0	28	0	28	0
29	0	29	0	29	0	29	0	29	0	29	0
30	0	30	0	30	0	30	0	30	0	30	0

LATITUDE of the MOON for the least and greatest INCLINATION of her Orbit; ($4^{\circ} 59' 35''$, and $5^{\circ} 17' 20''$) with the greatest Excess of Latitude.

Argument of Latitude, or MOON from NODE. 2. Equated.

Sig. 6. 6.				Sig. 1. 7.				Sig. 2. 8.			
N. A.		S. A.		N. A.		S. A.		N. A.		S. A.	
D's Lat.		D's Lat.		D's Lat.		D's Lat.		D's Lat.		D's Lat.	
for L. In.		for L. In.		for L. In.		for L. In.		for L. In.		for L. In.	
D.	Lat.	D.	Lat.	D.	Lat.	D.	Lat.	D.	Lat.	D.	Lat.
0	0	0	0	0	0	0	0	0	0	0	0
1	0	1	0	1	0	1	0	1	0	1	0
2	0	2	0	2	0	2	0	2	0	2	0
3	0	3	0	3	0	3	0	3	0	3	0
4	0	4	0	4	0	4	0	4	0	4	0
5	0	5	0	5	0	5	0	5	0	5	0
6	0	6	0	6	0	6	0	6	0	6	0
7	0	7	0	7	0	7	0	7	0	7	0
8	0	8	0	8	0	8	0	8	0	8	0
9	0	9	0	9	0	9	0	9	0	9	0
10	0	10	0	10	0	10	0	10	0	10	0
11	0	11	0	11	0	11	0	11	0	11	0
12	0	12	0	12	0	12	0	12	0	12	0
13	0	13	0	13	0	13	0	13	0	13	0
14	0	14	0	14	0	14	0	14	0	14	0
15	0	15	0	15	0	15	0	15	0	15	0
16	0	16	0	16	0	16	0	16	0	16	0
17	0	17	0	17	0	17	0	17	0	17	0
18	0	18	0	18	0	18	0	18	0	18	0
19	0	19	0	19	0	19	0	19	0	19	0
20	0	20	0	20	0	20	0	20	0	20	0
21	0	21	0	21	0	21	0	21	0	21	0
22	0	22	0	22	0	22	0	22	0	22	0
23	0	23	0	23	0	23	0	23	0	23	0
24	0	24	0	24	0	24	0	24	0	24	0
25	0	25	0	25	0	25	0	25	0	25	0
26	0	26	0	26	0	26	0	26	0	26	0
27	0	27	0	27	0	27	0	27	0	27	0
28	0	28	0	28	0	28	0	28	0	28	0
29	0	29	0	29	0	29	0	29	0	29	0
30	0	30	0	30	0	30	0	30	0	30	0

CONSTRUCTION. As Rad. : S. twice Arg. Lat. or S. 2. Δ a Δ :: C. Red. Eq. : Present Red. Eq. i. e. in the present least and greatest Inclinations Δ Orb. S. double Arg. (taking Rad. = 1) $\times$ d into $6' 533$ or $6' 32''$, and also into $7' 333$ or $7' 20''$, respectively.

Ex. To find the Reduction to $1^{\circ} 21'$. Arg. of Lat. or Δ a Δ , and $1^{\circ} 17' \odot$ a Δ ? By Tab. foregoing $\odot$ a Δ $1^{\circ} 17' \dots \times 1,4695$ By Tab. above $\dots \Delta$ a Δ $1^{\circ} 21' \odot$ Ex. Rⁿ. 47"

Δ a Δ Redⁿ 32865
1,21, -6' 24", L. Inclⁿ. 18780

+ 22 22, 0665
Redⁿ - 6' 46 for ? $\odot$ a Δ L. L.
pref. Inⁿ. req^d. } By Lo. Log. $1^{\circ} 17'$ 3283
So for the rest. Gr. Ex. Red. 47" 1.8842

Proportional Reduction + 22" 2.2125 as above.

N. B. Two of the first Figures of the $\times$ r are sufficient to find the Reduction, which may always be taken out at Sight.

Const. As Rad. : S. pref. Inclⁿ. :: S. pref. Arg. Lat. or S. Δ a Δ : S. pref. Lat. Δ correspondent.

Com. Log.

Ex. As Rad. $\dots \dots \dots$ 10.0000000
To S. least Inc. $4^{\circ} 59' 35''$ 8.9396939
So S. 30° Arg. Δ a Δ $\dots$ 9.6989700
To S. $2^{\circ} 29' 39''$ Lat. req^d. 8.6386639. So for the rest.

Ex. To find the Latitude Δ to $4^{\circ} 25'$ Δ a Δ , the Argument of Latitude, and $\dots \dots \dots 9^{\circ} 23' \odot$ a Δ ?

Δ a Δ Lat. for least Inc.

By Tab. above $4^{\circ} 25' \dots \dots \dots 2^{\circ} 51' 41''$ $\dots \dots \dots$ Gr. Ex. Lat. $10' 15''$
Proportional Latitude + 1 33 } By Tab. foregoing

Lat. Δ for Present Inclⁿ. required $2^{\circ} 53' 14''$ $\dots \dots \dots$ 3045
Tab. foregoing $\odot$ a Δ L. L. $\dots \dots \dots$ 5075

By Lo. Log. $9^{\circ} 23'$ 8150
Tab. abo. $4^{\circ} 25'$ Δ a Δ } $10' 9''$ 7717
Gr. Excess Lat. } $1^{\circ} 33''$ 1.5867

Proportional Lat. as before $1^{\circ} 33''$ 1.5867

N. B. Three Figures of the Multiplier are generally sufficient to determine the Moon's Latitude.

SUPPLEMENTAL LUNAR TABLES.

The HORIZONTAL PARALLAX of the MOON, in *Conjunction* or *Opposition* of the SUN and MOON. Also 1st Part of Par. out of *Sizygies*. To be added to the apparent for the true Altitude of the MOON; or subtracted from the apparent for the true Dist. of the MOON from the *Vertex*, and the contrary.

Horizontal Parallax to the Middle State of the Moon's Orbit, or Mean Eccentricity 55050 to Mean Dist. $\odot$ à $\ominus$ 1000000. With the Greatest Differences of Parallax, under and above the Mean Parallax correspondent to the Greatest and Least Eccentricities, A (above) and U (under) the Mean Eccentricity.

Argument. True Anomaly of the MOON.

For Horizontal Parallax Δ , in Ecc. above or under the Mean Eccy.

Correction of the first Part of the Moon's Horizontal PARALLAX out of *Conjunction* and *Opposition*, or *Sizygies*. By Decim^l Multipliers and Logistical Logarithms.

Argument. True Anomaly of the MOON.												Ar. $\odot$ à Δ Ap. 1 eq.				Arg. MOON fr. SUN true.			
Tr	Sig. 0		Sig. 1		Sig. 2		Sig. 3		Sig. 4		Sig. 5		Tr	O. 1. 2.		Tr	D à		D à
	Mean	G.D	Mean	G.D	Mean	G.D	Mean	G.D	Mean	G.D	Mean	G.D		6.	7.		Dec.	Log.	
An	Hor.		Hor.		Hor.		Hor.		Hor.		Hor.		An	ab.	ab.	An	à	Log.	à
	Pa. D.	A. U	Pa. D.	A. U	Pa. D.	A. U	Pa. D.	A. U	Pa. D.	A. U	Pa. D.	A. U		X ^{rs} of G. D.	Parallax.		à	Log.	à
D	+	-	+	-	+	-	+	-	+	-	+	-	D	+	-	D	+	-	+
	0	1	2	3	4	5	6	7	8	9	10	11		0	1		0	1	2
0	54	4	35	54	29	31	55	38	15	57	12	4	58	47	25	59	56	40	30
1	54	4	35	54	31	30	55	41	15	57	16	5	58	49	25	59	57	40	29
2	54	4	35	54	32	30	55	44	14	57	19	5	58	52	26	59	59	40	28
3	54	4	35	54	34	29	55	47	14	57	22	6	58	55	27	60	1	40	27
4	54	4	35	54	36	29	55	50	14	57	26	6	58	57	27	60	2	40	26
5	54	5	35	54	38	28	55	53	13	57	29	7	59	0	28	60	3	41	25
6	54	5	35	54	40	28	55	56	12	57	32	8	59	3	29	60	5	41	24
7	54	5	35	54	42	28	55	59	12	57	36	8	59	5	29	60	6	41	23
8	54	6	35	54	44	27	56	2	11	57	39	9	59	8	30	60	7	42	22
9	54	6	35	54	46	27	56	5	11	57	42	10	59	11	30	60	9	42	21
10	54	6	35	54	48	27	56	8	10	57	46	10	59	13	30	60	10	42	20
11	54	7	35	54	50	26	56	11	9	57	49	11	59	15	31	60	11	43	19
12	54	8	35	54	52	26	56	14	8	57	52	12	59	18	31	60	12	43	18
13	54	8	35	54	55	26	56	17	7	57	55	13	59	20	31	60	13	43	17
14	54	9	34	54	57	25	56	20	6	57	58	14	59	23	32	60	14	43	16
15	54	10	34	54	59	25	56	23	5	58	1	15	59	26	33	60	15	43	15
16	54	11	34	55	1	24	56	26	5	58	4	15	59	28	33	60	15	43	14
17	54	12	34	55	4	23	56	30	5	58	7	15	59	30	34	60	16	44	13
18	54	13	34	55	6	22	56	33	4	58	10	16	59	33	34	60	17	44	12
19	54	14	34	55	9	22	56	36	4	58	14	17	59	35	34	60	17	44	11
20	54	15	33	55	11	21	56	40	3	58	17	18	59	37	35	60	18	44	10
21	54	16	33	55	13	20	56	43	2	58	20	19	59	39	36	60	19	44	9
22	54	17	33	55	16	20	56	46	2	58	23	19	59	41	36	60	19	44	8
23	54	18	33	55	19	19	56	50	1	58	26	20	59	43	37	60	19	45	7
24	54	20	33	55	21	19	56	53	0	58	29	21	59	45	37	60	20	46	6
25	54	21	33	55	24	19	56	57	1	58	32	21	59	47	37	60	20	46	5
26	54	22	32	55	27	18	57	0	2	58	35	22	59	49	38	60	20	46	4
27	54	24	32	55	30	18	57	3	2	58	38	23	59	51	39	60	21	46	3
28	54	26	32	55	33	17	57	6	3	58	41	23	59	52	39	60	21	46	2
29	54	28	31	55	35	16	57	9	4	58	44	24	59	54	40	60	21	46	1
30	54	29	31	55	38	15	57	12	4	58	47	25	59	56	40	60	21	46	0
	+	-	+	-	+	-	+	-	+	-	+	-							
Tr	Sig. 11		Sig. 10		Sig. 9		Sig. 8		Sig. 7		Sig. 6		Tr	O. 1. 2.		Tr	D à		D à
	Mean	A. U	Mean	A. U	Mean	A. U	Mean	A. U	Mean	A. U	Mean	A. U		ab.	un.		à	Log.	à
An	Hor.	G. D	Hor.	G. D	Hor.	G. D	Hor.	G. D	Hor.	G. D	Hor.	G. D	An	11.	10.	An	à	Log.	à
	Pa. D.		Pa. D.		Pa. D.		Pa. D.		Pa. D.		Pa. D.			5.	4.		à	Log.	à

EXAMPLE. To find the first Part of the Moon's Horizontal Parallax out of *Conjunction* and *Opposition* of the Sun and Moon, for 4^h 23^m True An. 1st 24th $\odot$ à Δ 's Apog. 1 Equated, and 3rd Arg. Moon à Sun?

By Tab. above 4^h 23^m . . 59' 43" . . + 37" above Mean Par^x for Gr. Eccy above M. Eccy.
 — 7 . . 20 X^r for 1st 24th $\odot$ à Δ 's Ap. Eccy. under M. Eccy.

1st Part of Hor. Par^x ex. Syz. req. . 59' 36" . . — 7" 40 Prop. Parallax to be sub. from the Mean Parallax.
 N. B. Because Δ — Δ An. = Δ Apogee. — Therefore $\odot$ — Δ + Δ An. correct = $\odot$ — Δ Apogee correct; and when $\odot$ à Δ , or Δ à $\odot$, (or which is the same $\odot$ — Δ , or Δ — $\odot$) = 0, or 6s, in *Conjunction* and *Opposition*, respectively, then Moon's An. true (or Δ An. $\pm$ 6s) = Annual Arg^t. correct, or $\odot$ à Δ 's Ap. true: the same of m. Dist.
 Hence, it is observable, in *Sizygies*, or *Conjunction* and *Opposition* of Sun and Moon, that the Argument of Moon's true Anom. will be the same in Signs and Degrees as the Arg^t. for the X^r in Tab. of $\odot$ à Δ 's Apog. supposing that Argument correct (the Dif. when the Arg^t. is $\odot$ à Δ 's Ap. 1 Equated, being 2^d Equation of Δ Ap. with a contrary Sign, and Equation Δ 's Center, connected with the mean Anom. corrected, for nearly the true Anom. and the contrary). The Horizontal Parallax in a former Tab. is determined, according to the Argument of Moon's mean Anomaly corrected, and Argument of Mean Evection corrected, (like the mean central and Evection Equations) and the Correction of Parallax by Argument of Moon à Sun.

N. B. The Horizontal Parallax is to the apparent Diameter of the Moon, as 1 to .55, or as 60 to 33. Hence the Conf. Lo. Log. 2596, being added to the Lo. Log. of Hor. Par. will give the Lo. Log. of the Moon's apparent Diameter. Or $\frac{1}{2} + \frac{1}{10}$ of that $\frac{1}{2}$ of Hor Par^x = Apt. Diam. Moon.

Conf. of Dec. X^{rs} from the Log. Log^s.

Log. L. 63 . 59' 8"
 Or $\frac{1}{6}$ 59, 1333, &c.
 .98555, &c.

Comp^t to 1. .0145—
 D. X^r cor^t to L. Log. 63.
 And so for the rest.

Ex. 1. Pt 59' 36" Px ex. Syzy +29
 Δ à $\odot$ 3rd . . +63

Px. out of Syzy 58' 44" +92
 Or 59' 36" = 59' 6
 — 62 . . .0145 X^r.

req^d. 58 44 2980
 [cor^t Par^x. 2384
 596
 ,86420
 52nd

SUPPLEMENTAL LUNAR TABLES.

The ECCENTRICITIES of the MOON, and Difference of those Eccentricities from the Mean Eccentricity 55237, the Greatest being 66854, and Least 43619; according to Horrox, or Flamsteed. With the Difference of Eccentricities from the Mean 55050; according to Sir Isaac Newton, Rad. or Mean Dist. Δ à $\ominus$ being 1000000.

Argument. SUN from the MOON's APOGEE; or Sun's true Pl. — m. Pl. Δ Ap. 1 Equated.

⊙ à Δ Ap.	Sig. o. 6			Sig. 1. 7			Sig. 2. 8			⊙ à Δ Ap.
	Eccentr. Δ Orb. Horrox.	Differ. from the Mean.	Diff. Ecc. fr. Mean. Newton.	Eccentr. Δ Orb. Horrox.	Diff. from the Mean.	Diff. Ecc. fr. Mean. Newton.	Eccentr. Δ Orb. Horrox.	Diff. from the Mean.	Diff. Ecc. fr. Mean. Newton.	
0	66854	11617	11732	61045	5808	6708	49429	5808	4827	30
1	66847	11610	11726	60691	5459	6388	49082	6155	5213	29
2	66826	11589	11708	60330	5093	6060	48741	6496	5594	28
3	66791	11554	11679	59962	4725	5725	48408	6829	5969	27
4	66741	11504	11638	59589	4352	5382	48085	7152	6337	26
5	66678	11441	11585	59210	3973	5033	47769	7468	6698	25
6	66600	11363	11520	58827	3590	4677	47463	7774	7051	24
7	66509	11272	11444	58439	3202	4315	47167	8070	7396	23
8	66404	11167	11356	58047	2810	3947	46880	8357	7731	22
9	66286	11049	11257	57652	2415	3574	46604	8633	8058	21
10	66154	10917	11146	57254	2017	3195	46337	8900	8373	20
11	66008	10771	11024	56854	1617	2811	46082	9155	8678	19
12	65850	10613	10891	56451	1214	2432	45838	9399	8973	18
13	65679	10442	10746	56047	810	2031	45606	9631	9253	17
14	65495	10258	10590	55642	405	1636	45385	9852	9522	16
15	65298	10061	10423	55237	± 0	1237	45176	10061	9778	15
16	65089	9852	10246	54832	405	835	44979	10258	10020	14
17	64868	9631	10058	54427	810	431	44794	10443	10248	13
18	64636	9399	9859	54023	1214	+ 24	44623	10614	10461	12
19	64392	9155	9650	53620	1617	— 384	44466	10771	10659	11
20	64137	8900	9430	53220	2017	792	44320	10917	10841	10
21	63870	8633	9200	52822	2415	1202	44188	11049	11007	9
22	63594	8357	8960	52427	2810	1611	44070	11167	11157	8
23	63307	8070	8711	52035	3202	2020	43965	11272	11290	7
24	63011	7774	8452	51647	3590	2428	43874	11363	11406	6
25	62705	7468	8183	51264	3973	2835	43796	11441	11505	5
26	62389	7152	7906	50885	4352	3240	43733	11504	11586	4
27	62066	6829	7619	50512	4725	3642	43683	11554	11649	3
28	61733	6496	7323	50144	5093	4041	43648	11589	11694	2
29	61392	6155	7020	49783	5454	4436	43627	11610	11722	1
30	61045	5808	6708	49429	5808	4827	43619	11618	11731	0
⊙ à Δ Ap.	Eccentr. Δ Orb. Horrox.	Diff. from Mean.	Diff. Ecc. fr. Mean. Newton.	Eccentr. Δ Orb. Horrox.	Diff. from Mean.	Diff. Ecc. fr. Mean. Newton.	Eccentr. Δ Orb. Horrox.	Diff. from Mean.	Diff. Ecc. fr. Mean. Newton.	⊙ à Δ Ap.
		above	above		above	above.		under.	under.	
					under.	under.				

CONSTRUCTION of the Eccentricities of the Moon's Orbit. According to Sir Isaac Newton.

As S. 2d Equation Δ Ap. : $\frac{1}{2}$ Diff. between the greatest and least Eccentricity $11731\frac{1}{2}$:: S. Double annual Argument, Δ 's Ap. Or S.

2. $\odot$ à Δ 's Ap. : Present Eccentricity.

N. B. The Eccentricity is Greatest when Apog. Δ is in Sun's Syzygies, and Least when in the Quadratures.—These and other Inequalities of the Moon's Motion are augmented and diminished in the Triplicate Proportion of the Sun's apparent Diameter. And, moreover, the Moon's VARIATION is farther augmented or diminished in the duplicate Proportion of the Time between the Quadratures.—Though in Astronomical Computations this last Inequality is commonly thrown into and confounded with the Equation of the Moon's Center. See *Principia*, Prop. 22. B. 3.

DIFFERENT METHODS

Of Computing the

ELLIPTIC EQUATION of the MOON'S CENTER.

SUPPLEMENTAL LUNAR TABLES.

CONSTANT LOGARITHMS, correspondent to the Different ECCENTRICITIES of the MOON'S Orbit, and Sem. Transverse 1000000. For taking out the EQUATION or CORRECTION to Half the Moon's Equated ANOMALY in the following Pages, and thence to find the ELLIPTIC EQUATION of the MOON'S CENTER.

Argument. SUN from the MOON'S Apogee, or ☉ true Pl. — D Ap. 1. Equated.

Sig. 0. 6.					Sig. 1. 7.					Sig. 2. 8.				
Con. Log. for Aeq. D Cent.	Dif.	Ecc. D orb parts.	Dif. pts.	Xrs of ex. of Ar	Con. Log. for Aeq. D Cent.	Dif.	Ecc. D orb parts.	Dif. pts.	Xrs of ex. of Ar	Con. Log. for Aeq. D Cent.	Dif.	Ecc. D orb parts.	Dif. pts.	Xrs of ex. of Ar
0	9,941912	5	66782	6	9,946292	279	61758	320	5,33	9,956339	336	50223	686	6,43
1	9,941917	15	66776	18	9,946571	286	61438	328	5,46	9,956675	332	49837	381	6,35
2	9,941932	26	66758	29	9,946857	292	61110	335	5,58	9,957007	326	49456	375	6,25
3	9,941958	36	66729	41	9,947149	299	60775	346	5,71	9,957333	320	49081	368	6,13
4	9,941994	46	66681	53	9,947448	304	60432	349	5,81	9,957653	315	48713	361	6,01
5	9,942040	56	66635	65	9,947752	310	60083	356	5,93	9,957968	307	48352	353	5,88
6	9,942096	67	66570	76	9,948062	315	59727	362	6,03	9,958275	300	47999	345	5,75
7	9,942163	76	66494	88	9,948377	320	59365	368	6,14	9,958575	292	47654	335	5,58
8	9,942239	87	66406	99	9,948697	326	58997	373	6,21	9,958867	284	47319	327	5,45
9	9,942326	96	66307	111	9,949023	330	58624	379	6,31	9,959151	275	46992	315	5,25
10	9,942422	107	66196	122	9,949353	334	58245	384	6,4	9,959426	265	46677	305	5,08
11	9,942529	116	66074	133	9,949687	338	57861	388	6,46	9,959691	255	46372	294	4,9
12	9,942645	126	65941	145	9,950025	342	57473	392	6,53	9,959946	245	46078	281	4,68
13	9,942771	136	65796	156	9,950367	345	57081	395	6,58	9,960191	234	45797	269	4,48
14	9,942907	146	65640	167	9,950712	347	56686	399	6,65	9,960425	223	45528	256	4,26
15	9,943053	155	65473	177	9,951059	350	56287	402	6,7	9,960648	210	45272	242	4,03
16	9,943208	164	65296	188	9,951409	352	55885	404	6,73	9,960858	198	45030	228	3,8
17	9,943372	173	65108	199	9,951761	354	55481	407	6,78	9,961056	186	44802	213	3,55
18	9,943545	183	64909	209	9,952115	355	55074	408	6,8	9,961242	172	44589	198	3,3
19	9,943728	191	64700	220	9,952470	357	54666	408	6,8	9,961414	159	44391	189	3,03
20	9,943919	201	64480	230	9,952827	356	54258	410	6,83	9,961573	144	44209	166	2,76
21	9,944120	209	64250	240	9,953183	357	53848	409	6,81	9,961717	131	44043	150	2,5
22	9,944329	217	64010	249	9,953540	356	53439	409	6,81	9,961848	115	43893	133	2,21
23	9,944546	226	63761	259	9,953896	355	53030	408	6,8	9,961963	101	43760	116	1,91
24	9,944772	234	63502	269	9,954251	354	52622	407	6,78	9,962064	86	43644	99	1,65
25	9,945006	242	63233	277	9,954605	352	52215	405	6,75	9,962150	71	43545	81	1,35
26	9,945248	250	62956	287	9,954957	350	51810	402	6,7	9,962221	55	43464	63	1,05
27	9,945498	257	62669	296	9,955307	348	51408	399	6,65	9,962276	49	43401	45	,75
28	9,945755	265	62373	303	9,955655	344	51009	395	6,58	9,962315	24	43356	28	,46
29	9,946020	272	62070	312	9,955999	340	50614	391	6,51	9,962339	8	43328	9	,15
30	9,946292		61758		9,956339		50223			9,962347		43319		
Con. Log. for Aeq. D Cent.	Dif.	Ecc. D orb.	Dif.	Xrs of ex. of Ar	Con. Log. for Aeq. D Cent.	Dif.	Ecc. D orb.	Dif.	Xrs of ex. of Ar	Con. Log. for Aeq. D Cent.	Dif.	Ecc. D orb.	Dif.	Xrs of ex. of Ar
Sig. 11. 5.					Sig. 10. 4.					Sig. 9. 3.				

SUPPLEMENTAL LUNAR TABLES.

EQUATION, or CORRECTION of half the Equated ANOMALY of the MOON, for finding, at once, the Elliptic Equation of the MOON'S CENTER.

Argument. MOON'S EQUATED ANOMALY; or Δ 3 equated — Ap. 2 equated.

D eq. A. Lo	Sig. 0.					Sig. 1.					Sig. 2.					D eq. A. Lo
	9.942	9.947	9.952	9.957	9.962	9.942	9.947	9.952	9.957	9.962	9.942	9.947	9.952	9.957	9.962	
0	0 0	0 0	0 0	0 0	0 0	1 42	1 25	1 9	0 55	0 43	1 52	1 33	1 15	1 0	0 46	30
1	0 4	0 3	0 3	0 2	0 2	1 44	1 27	1 11	0 57	0 44	1 51	1 31	1 14	0 59	0 46	29
2	0 8	0 7	0 5	0 4	0 3	1 46	1 28	1 12	0 58	0 45	1 49	1 30	1 13	0 58	0 45	28
3	0 12	0 10	0 8	0 6	0 5	1 48	1 30	1 14	0 59	0 46	1 47	1 28	1 11	0 57	0 44	27
4	0 16	0 13	0 11	0 9	0 7	1 49	1 31	1 15	1 0	0 46	1 45	1 26	1 10	0 56	0 43	26
5	0 20	0 16	0 13	0 11	0 8	1 51	1 33	1 16	1 1	0 47	1 43	1 25	1 8	0 54	0 42	25
6	0 24	0 20	0 16	0 13	0 10	1 53	1 34	1 17	1 1	0 48	1 40	1 23	1 7	0 53	0 41	24
7	0 28	0 23	0 19	0 15	0 12	1 54	1 35	1 18	1 2	0 48	1 38	1 21	1 5	0 52	0 40	23
8	0 31	0 26	0 21	0 17	0 13	1 56	1 36	1 19	1 3	0 49	1 36	1 19	1 4	0 50	0 39	22
9	0 35	0 29	0 24	0 19	0 15	1 57	1 37	1 20	1 4	0 49	1 33	1 17	1 2	0 49	0 38	21
10	0 39	0 33	0 27	0 21	0 17	1 58	1 38	1 20	1 4	0 50	1 30	1 14	1 0	0 47	0 36	20
11	0 43	0 36	0 29	0 23	0 18	1 59	1 39	1 21	1 5	0 50	1 28	1 12	0 58	0 46	0 35	19
12	0 46	0 39	0 32	0 25	0 20	2 0	1 40	1 22	1 5	0 51	1 25	1 9	0 56	0 44	0 34	18
13	0 50	0 42	0 34	0 28	0 21	2 0	1 40	1 22	1 5	0 51	1 22	1 7	0 54	0 42	0 32	17
14	0 54	0 45	0 37	0 30	0 23	2 1	1 41	1 22	1 6	0 51	1 19	1 4	0 52	0 41	0 31	16
15	0 57	0 48	0 39	0 32	0 24	2 1	1 41	1 23	1 6	0 51	1 16	1 2	0 50	0 39	0 30	15
16	1 1	0 51	0 42	0 33	0 26	2 2	1 41	1 23	1 6	0 51	1 12	0 59	0 47	0 37	0 28	14
17	1 4	0 54	0 44	0 35	0 27	2 2	1 41	1 23	1 6	0 51	1 2	0 56	0 45	0 35	0 27	13
18	1 8	0 57	0 46	0 37	0 29	2 2	1 42	1 23	1 6	0 51	1 5	0 53	0 43	0 33	0 25	12
19	1 11	0 59	0 48	0 39	0 30	2 2	1 41	1 23	1 6	0 51	1 2	0 50	0 41	0 31	0 24	11
20	1 14	1 2	0 51	0 41	0 32	2 2	1 41	1 23	1 6	0 51	1 58	0 47	0 38	0 29	0 22	10
21	1 18	1 4	0 53	0 42	0 33	2 1	1 41	1 22	1 5	0 50	0 55	0 44	0 35	0 27	0 20	9
22	1 21	1 7	0 55	0 44	0 34	2 1	1 40	1 22	1 5	0 50	0 51	0 41	0 33	0 25	0 19	8
23	1 24	1 9	0 57	0 46	0 36	2 0	1 40	1 21	1 4	0 50	0 47	0 38	0 30	0 23	0 17	7
24	1 26	1 12	0 59	0 47	0 37	1 59	1 39	1 21	1 4	0 50	0 44	0 35	0 28	0 21	0 16	6
25	1 29	1 14	1 1	0 49	0 38	1 58	1 38	1 20	1 3	0 49	0 40	0 32	0 25	0 19	0 14	5
26	1 32	1 17	1 3	0 50	0 39	1 57	1 37	1 19	1 3	0 49	0 36	0 28	0 22	0 17	0 12	4
27	1 34	1 19	1 4	0 52	0 40	1 56	1 36	1 19	1 2	0 48	0 32	0 25	0 20	0 15	0 11	3
28	1 37	1 21	1 6	0 53	0 41	1 55	1 35	1 18	1 1	0 48	0 28	0 22	0 17	0 12	0 9	2
29	1 39	1 23	1 7	0 54	0 42	1 54	1 34	1 17	1 1	0 47	0 24	0 19	0 14	0 10	0 7	1
30	1 42	1 25	1 9	0 55	0 43	1 52	1 33	1 15	1 0	0 46	0 20	0 15	0 11	0 8	0 6	0
D eq. An	Sig. 11.					Sig. 10.					Sig. 9.					D eq. An

CONSTRUCTION of the above Table and SOLUTION of the Keplerian PROBLEM. See *Larus of the Moon's Motion at the End of Vol. 2. Principia*, p. 58. Put t = Sem. Tranf. of an elliptical Orbit; n = Eccentricity, c = Sem. Conj. and $r = \angle 57^\circ 17' 44'' : 48'' = 57^\circ$, 2957795 = Radius.

$$B = \angle = \frac{n}{2t} r; E = \frac{c}{2t} B; S = \frac{4\pi}{3c} \left. \begin{array}{l} \text{Say Rad. : S. } \angle 2 A :: B : b \\ \text{Rad. : S. } \angle 2 A \pm 2b :: E : e \\ \text{Cube Rad. : Cub. S. } \angle A \pm b :: S : s. \end{array} \right\}$$

being greatest Δ Equations.

$A = \angle$ of mean Anomaly given.

N. B. The small Δ Equat. b is always of the same Sign with the Δ Equat. e , and in planetary Orbits always near the double of that Δ Equat.

Then $\angle$ of corrected Mean Anomaly will be $\left\{ \begin{array}{l} A + e + s \\ A - e + s \\ A + e - s \\ A - e - s \end{array} \right\} \begin{array}{l} 1 \\ 2 \\ 3 \\ 4 \end{array} \left\{ \begin{array}{l} \text{Quadrants} \\ \text{from} \\ \text{Apelion.} \end{array} \right. \left\{ \begin{array}{l} \frac{1}{2} \times e + s \\ \frac{1}{2} \times -e + s \\ \frac{1}{2} \times e - s \\ \frac{1}{2} \times -e - s \end{array} \right. \left\{ \begin{array}{l} \text{Correction,} \\ \text{or} \\ \Delta \text{ Equat.} \\ \text{of } \frac{1}{2} \text{ Anomaly.} \end{array} \right.$

The corrected Anomaly being known, say as *Apb. Dist.* : *Perib. Dist.* :: $T : \frac{1}{2}$ corrected An. : $T : \frac{1}{2}$ true Anomaly.

Whence the true Anomaly and the Δ Equation, in all the Planetary Orbits, are found.

Correction in very Eccentric Orbits.

The rays are reciprocally in the Square of the Velocity about the upper Focus or assumed Center of Motion. As the Rectangle of Distances fr. Foci : Rectangle Sem. Axes :: Difference between the Mean and assum'd Motion : Error of the assum'd $\angle$ at the Upper Focus.

SUPPLEMENTAL

SUPPLEMENTAL LUNAR TABLES.

EQUATION or CORRECTION of half the Equated ANOMALY of the MOON, for finding, at once, the Elliptic Equation of the MOON'S CENTER.

Argument. MOON'S Equated ANOMALY, or D 3. equated — D Ap. 2 equated.

D eq. An.	Sig. 3.					Sig. 4.					Sig. 5.					D eq. An.
Log.	9.942	9.947	9.952	9.957	9.962	9.942	9.947	9.952	9.957	9.962	9.942	9.947	9.952	9.957	9.962	Log.
0	0 20	0 15	0 11	0 8	0 6	1 26	1 13	1 0	0 49	0 39	1 36	1 21	1 6	0 53	0 42	30
1	0 16	0 12	0 9	0 6	0 4	1 29	1 15	1 2	0 50	0 40	1 35	1 19	1 5	0 52	0 41	29
2	0 12	0 9	0 6	0 4	0 2	1 30	1 16	1 3	0 51	0 41	1 33	1 18	1 4	0 51	0 40	28
3	0 8	0 5	0 3	0 2	0 1	1 32	1 18	1 5	0 52	0 42	1 31	1 16	1 2	0 50	0 39	27
4	0 4	0 2	0 0	—	—	1 34	1 20	1 6	0 53	0 42	1 28	1 14	1 1	0 49	0 38	26
5	—	—	1	2	0 2	1 36	1 21	1 7	0 54	0 43	1 26	1 12	0 59	0 47	0 37	25
6	0 4	0 4	0 5	0 5	0 4	1 38	1 23	1 8	0 55	0 44	1 24	1 10	0 57	0 46	0 36	24
7	0 8	0 8	0 7	0 7	0 6	1 39	1 24	1 9	0 56	0 44	1 21	1 8	0 56	0 44	0 35	23
8	0 12	0 11	0 10	0 9	0 8	1 41	1 25	1 10	0 57	0 45	1 18	1 6	0 54	0 43	0 34	22
9	0 16	0 14	0 13	0 11	0 10	1 42	1 26	1 11	0 57	0 45	1 16	1 3	0 52	0 41	0 33	21
10	0 20	0 18	0 16	0 13	0 11	1 44	1 27	1 12	0 58	0 46	1 13	1 1	0 50	0 40	0 31	20
11	0 24	0 21	0 18	0 16	0 13	1 45	1 28	1 13	0 58	0 46	1 10	0 58	0 48	0 38	0 30	19
12	0 27	0 24	0 21	0 18	0 14	1 45	1 29	1 13	0 59	0 46	1 7	0 56	0 46	0 37	0 29	18
13	0 31	0 27	0 24	0 20	0 16	1 46	1 29	1 13	0 59	0 46	1 3	0 53	0 43	0 35	0 27	17
14	0 35	0 31	0 26	0 22	0 18	1 47	1 30	1 14	1 0	0 47	1 0	0 50	0 41	0 33	0 26	16
15	0 39	0 34	0 29	0 24	0 19	1 47	1 30	1 14	1 0	0 47	0 57	0 48	0 39	0 31	0 24	15
16	0 43	0 37	0 31	0 26	0 21	1 47	1 30	1 14	1 0	0 47	0 53	0 45	0 37	0 29	0 23	14
17	0 46	0 40	0 34	0 28	0 22	1 48	1 30	1 15	1 0	0 47	0 50	0 42	0 34	0 27	0 21	13
18	0 50	0 43	0 36	0 30	0 24	1 48	1 30	1 15	1 0	0 47	0 46	0 39	0 32	0 25	0 20	12
19	0 53	0 46	0 38	0 32	0 25	1 48	1 30	1 15	1 0	0 47	0 43	0 36	0 29	0 23	0 18	11
20	0 57	0 48	0 41	0 33	0 27	1 47	1 30	1 14	1 0	0 47	0 39	0 33	0 27	0 21	0 17	10
21	1 0	0 51	0 43	0 35	0 28	1 47	1 30	1 14	0 59	0 47	0 35	0 29	0 24	0 19	0 15	9
22	1 3	0 54	0 45	0 37	0 29	1 46	1 29	1 13	0 59	0 46	0 31	0 26	0 21	0 17	0 13	8
23	1 6	0 56	0 47	0 39	0 31	1 45	1 29	1 13	0 58	0 46	0 28	0 23	0 19	0 15	0 12	7
24	1 9	0 59	0 49	0 40	0 32	1 44	1 28	1 12	0 58	0 45	0 24	0 20	0 16	0 13	0 10	6
25	1 12	1 1	0 51	0 42	0 33	1 43	1 27	1 12	0 57	0 45	0 20	0 16	0 13	0 11	0 8	5
26	1 15	1 4	0 53	0 43	0 34	1 42	1 26	1 11	0 57	0 44	0 16	0 13	0 11	0 9	0 7	4
27	1 18	1 6	0 55	0 45	0 36	1 41	1 25	1 10	0 56	0 44	0 12	0 10	0 8	0 7	0 5	3
28	1 21	1 9	0 57	0 46	0 37	1 40	1 24	1 9	0 55	0 43	0 8	0 7	0 5	0 4	0 3	2
29	1 23	1 11	0 59	0 48	0 38	1 38	1 22	1 8	0 54	0 43	0 4	0 3	0 3	0 2	0 2	1
30	1 26	1 13	1 0	0 49	0 39	1 36	1 21	1 6	0 53	0 42	0 0	0 0	0 0	0 0	0 0	0
D eq. An.	Sig. 8.					Sig. 7.					Sig. 6.					D eq. An.

Ex. The Eccentricity being 0.060 and Mean An. 30° to find the Correction, or Equation of $\frac{1}{2}$ M. Anom. and also the correct Equation of the Orbit?
 $B = 6' 11'' 16''' = 103132^{\circ}$; Gr. E. $= 3' 51'' 15''' = 051463^{\circ}$; $S = 29' 45''' = 008267^{\circ}$ —Now, $b = 5' 21'' 29''' = 0893^{\circ}$;
 $e = 2' 40'' 42''' = 04464^{\circ}$; $s = 3' 44''' = 00104^{\circ}$, [nearly according to Mr. Machin].

Hence, $A + e + s = 30^{\circ} 2' 44'' 26'''$ the Corrected Anomaly.

Log.

The half $15^{\circ} 1' 22'' 13'''$ Tang. 9.4287444 . . [1' 22'' 13''' Equation, or Correction.

For Eccentricity, 06 Const. Log. 9.9478220

$13^{\circ} 23' 13'' 15'''$. . Tan. 9.3765664

Double 26 46 26 30—

Correct Equation . . $3^{\circ} 13' 33'' 30'''$ required. So for the rest.

EXAMP. To find the Moon's Central Equation correspondent to $1^{\circ} 11'$ D Equated Anom. and $2^{\circ} 8'$ Sun Δ D Apogee?

By Tab. p. 67. . . Arg. $1^{\circ} 11' = 41^{\circ} 0' 0''$

Correction or Equation . . . $+ 59$

Corrected Anom. T. . . . $20 30 59$

By Tab. p. 81. . . . Const. Log.

$\frac{1}{2}$ True Anom. T. . . . $18 47 54$

Doubled . . . $37 35 48$

Equation of D's Center — $3 24 12$ required.

Equated Anomaly.

Com. Logars.

9.5731163

9.958867 for Eq. D Center.

9.5319833

—, or deduct from Eqd. Anomaly.

Eccentr ^s and Equations.				
Ecc ^s . to Rad. 1.	E.	Dif.	S.	Dif.
0.040	1 23	22	9	4
0.045	1 45	24	13	4
0.050	2 09	27	17	6
0.055	2 36	27	23	7
0.060	3 08	30	30	8
0.065	3 38	32	38	8
0.070	4 14	36	47	9

The above Table,
by Mr. Machin.

SUPPLEMENTAL LUNAR TABLES.

ELLIPTIC EQUATION of the MOON'S CENTER in all States of Her Variable Orbit: From Excentricity 43319 to 66782; the Mean Distance being 1000000. According to the Newtonian Theory.

Argument. The MOON'S Equated ANOMALY, Or Moon 3 Equated — Moon Ap. z. Equated.

SIGN o.

In. of De fr. M. for +	D. eq. An.	45000 Eq. D.	Dif.	D. Eq. for D. 5000	50000 Eq. D.	Dif.	D. Eq. for D. 5000	55000 Eq. D.	Dif.	D. Eq. for D. 5000	60000 Eq. D.	Dif.	D. Eq. for D. 5000	65000 Eq. D.	Dif.	D. Eq. for D. 5000	D. eq. An.
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1	1	5	7	0	5	39	0	6	10	0	6	42	0	7	13	0	30
0	2	10	13	1	11	17	1	12	21	1	13	23	1	14	25	1	28
1	3	15	20	1	16	56	1	18	31	1	20	5	1	21	38	1	27
1	4	20	26	2	22	34	2	24	41	2	26	46	2	28	50	2	26
2	5	25	32	2	28	12	2	30	50	2	33	27	2	36	2	2	25
3	6	30	37	3	33	50	3	37	0	3	40	7	3	43	13	3	24
2	7	35	42	3	39	26	3	43	8	3	46	47	3	50	23	3	23
3	8	40	47	4	45	3	4	49	16	4	53	26	4	57	33	4	22
4	9	45	51	4	50	39	4	55	23	4	58	4	4	61	42	4	21
3	10	50	54	5	56	13	5	1	29	5	6	42	5	11	50	5	20
3	11	55	56	5	1	47	5	7	35	5	13	18	5	18	57	7	19
5	12	1	58	6	7	21	6	13	39	6	14	53	6	26	3	6	18
6	13	5	58	6	12	53	6	19	42	6	45	26	6	33	7	4	17
6	14	10	58	7	18	24	7	25	44	6	16	33	7	40	11	7	16
7	15	15	56	7	23	54	7	31	45	7	47	39	7	47	13	7	15
6	16	20	53	8	29	22	8	37	45	8	17	46	8	54	13	8	14
6	17	25	49	9	1	34	8	43	43	8	47	52	8	1	12	8	13
7	18	30	44	9	31	15	9	49	39	9	18	58	9	8	8	9	12
7	19	35	37	10	2	45	9	55	34	9	48	5	9	15	3	9	11
6	20	40	29	10	32	1	10	26	27	10	18	2	10	21	56	10	10
7	21	45	19	11	3	56	10	56	18	10	48	18	10	28	47	10	9
7	22	50	8	11	33	1	11	26	7	11	18	24	11	35	36	11	8
7	23	54	55	12	3	6	11	56	54	11	48	30	11	42	23	11	7
8	24	59	39	12	34	12	12	26	39	12	18	36	12	49	7	12	6
9	25	4	23	13	4	17	12	55	22	12	47	43	12	55	48	12	5
9	26	9	4	13	34	22	13	25	36	13	16	49	13	3	27	13	4
8	27	13	43	14	3	27	13	55	41	13	46	55	13	37	9	13	3
9	28	18	20	14	33	32	14	24	47	14	15	3	14	53	15	13	2
9	29	22	55	15	2	37	14	53	52	14	44	3	14	34	22	14	1
10	30	27	27	15	32	42	15	22	58	15	12	3	15	3	28	14	0
+ 5000 D.	D. eq. An.	Eq. D.	Dif.	5000 D. Eq.	Eq. D.	Dif.	5000 D. Eq.	Eq. D.	Dif.	5000 D. Eq.	Eq. D.	Dif.	5000 D. Eq.	Eq. D.	Dif.	5000 D. Eq.	D. eq. An.
		45000			50000			55000			60000			65000			5000 D.

SIGN 11.

EXAMPLE I. To find the Moon's Central Equation correspondent to 23° 8' Sun, a Moon's Ap. and 13° 11' Moon Equated Anomaly, according to the Newtonian Theory?

Eccentricity of Orb. Ecc. above 45000.
 By Tab. p. 81. Arg^t Sun a Moon Ap. 1 Equated gives 47319 2319 doubled = 4638
 Equation for Eccentricity 45000.
 By Tab. p. 85. Arg^t 13° 11' D Eq. An. — 3° 14' 36" Excess of Eqⁿ. for 5000 20' 37" = 20,61
 Pr. Equation +9 33 fere 4638
 D Central Equation — 3 24 9 correctly, required. See p. 83. 27828
 L. L. 92760
 Or Excs. Eccy. 23' 19" = 38' 39" 1910
 Excess Eqⁿ 4639
 5000 = 83 20 1426 Const.
 Prop^d. Eq. req^d. 9 34 7975 req^d.
 Rule. To L. L. 1426 Const. } Cutting off 4 Fig. to the Right where the }
 Add L. L. Excess Eccy. } xing Dec^s are cut off }
 and L. L. Excess Eqⁿ. } for 10000 Divisor. }
 Sum L. L. Prop. Eq. to be }
 added to the Central Eq. for the Excess }
 of Eccentricity above the last. }

Hence, the taking out of the Eccentricity of Moon's Orb. (in p. 81.) and then the Central Equation (as above) is easily and readily performed, either by Decimal Multipliers, or Logistical Logarithms, including 2d Equation of the Moon's Apogee.

SUPPLEMENTAL LUNAR TABLES.

ELLIPTIC EQUATION of the MOON'S CENTER, according to the *Newtonian* Theory.Argument: The MOON'S Equated Anomaly, or Δ 3 Equated — Δ Ap. 2 Equated.

SIGN I.

De fr. M for +		45000		Dif.	D.Eq. for D. 5000		50000		Dif.	D.Eq. for D. 5000		55000		Dif.	D.Eq. for D. 5000		60000		Dif.	D.Eq. for D. 5000		65000		Dif.	D.Eq. for D. 5000		D eq. An	of In. fr. M for —											
		Eq. D.	—		Eq. D.	—	Eq. D.	—		Eq. D.	—	Eq. D.	—		Eq. D.	—	Eq. D.	—		Eq. D.	—	Eq. D.	—																
10	0	2	27	27	4	31	15	32	2	42	59	4	59	15	22	2	58	21	5	28	15	12	3	13	33	5	57	15	3	3	28	36	6	25	14	54	30	9	
9	1	2	31	58	4	27	16	0	2	47	58	4	57	15	51	3	3	49	5	25	15	41	3	19	30	5	53	15	31	3	35	1	6	21	15	21	29	10	
11	2	2	36	25	4	26	16	30	2	52	55	4	54	16	19	3	9	14	5	22	16	9	3	25	23	5	50	15	59	3	41	22	6	18	15	49	28	10	
11	3	2	40	51	4	22	16	58	2	57	49	4	51	16	47	3	14	36	5	19	16	37	3	31	13	5	47	16	27	3	47	40	6	15	16	17	27	10	
12	4	2	45	13	4	21	17	27	3	2	40	4	48	17	15	3	19	55	5	16	17	5	3	37	0	5	44	16	55	3	53	55	6	11	16	45	26	10	
11	5	2	49	34	4	17	17	54	3	7	28	4	45	17	43	3	25	11	5	13	17	33	3	42	44	5	40	17	22	4	0	6	8	17	11	25	11		
11	6	2	53	51	4	15	18	22	3	12	13	4	42	18	11	3	30	24	5	10	18	0	3	48	24	5	37	17	50	4	6	14	6	4	17	40	24	10	
10	7	2	58	6	4	12	18	49	3	16	55	4	40	18	39	3	35	34	5	6	18	27	3	54	1	5	34	18	17	4	12	18	6	0	18	7	23	10	
12	8	3	2	18	4	9	19	17	3	21	35	4	36	19	5	3	40	40	5	3	18	55	3	59	35	5	29	18	43	4	18	18	5	57	18	31	22	12	
12	9	3	6	27	4	6	19	44	3	26	11	4	33	19	32	3	45	43	5	0	19	21	4	5	4	5	26	19	11	4	24	15	5	52	19	1	21	10	
12	10	3	10	33	4	3	20	11	3	30	44	4	29	19	59	3	50	43	4	55	19	47	4	10	30	5	22	19	37	4	30	7	5	52	19	27	20	10	
12	11	3	14	36	4	0	20	37	3	35	13	4	26	20	25	3	55	38	4	55	20	14	4	15	52	5	19	20	3	4	35	55	5	48	19	52	19	11	
11	12	3	18	36	3	56	21	3	3	39	39	4	23	20	52	4	0	31	4	33	20	40	4	21	11	5	14	20	28	4	41	39	5	44	20	16	18	12	
13	13	3	22	32	3	54	21	30	3	44	2	4	19	21	17	4	5	19	4	48	21	6	4	26	25	5	10	20	54	4	47	19	5	40	20	42	17	12	
12	14	3	26	26	3	50	21	55	3	48	21	4	15	21	43	4	10	4	4	45	21	31	4	31	35	5	5	21	19	4	52	54	5	35	21	7	16	12	
12	15	3	30	16	3	46	22	20	3	52	36	4	12	22	8	4	14	44	4	40	21	56	4	36	40	5	5	21	45	4	58	25	5	31	21	34	15	11	
13	16	3	34	2	3	43	22	46	3	56	48	4	8	22	33	4	19	21	4	37	22	21	4	41	42	5	2	22	10	5	3	52	5	27	21	59	14	11	
14	17	3	37	45	3	40	23	11	4	0	56	4	3	22	57	4	23	53	4	32	22	47	4	46	40	4	58	22	33	5	9	13	5	21	22	19	13	14	
11	18	3	41	25	3	36	23	34	4	4	59	4	0	23	23	4	28	22	4	29	23	10	4	51	32	4	52	22	58	5	14	30	5	17	22	46	12	12	
11	19	3	45	1	3	33	23	58	4	8	59	4	57	23	47	4	32	46	4	24	23	34	4	56	20	4	48	23	23	5	19	43	5	13	23	12	11	11	
12	20	3	48	34	3	28	24	22	4	12	56	3	52	24	10	4	37	6	4	20	23	58	5	1	4	4	44	23	46	5	24	50	5	7	23	34	10	12	
13	21	3	52	2	3	25	24	46	4	16	48	3	52	24	33	4	41	21	4	15	24	21	5	5	42	4	38	24	10	5	29	52	5	2	23	59	9	11	
11	22	3	55	27	3	20	25	8	4	20	35	3	47	24	57	4	45	32	4	11	24	44	5	10	16	4	34	24	33	5	34	49	4	57	24	22	8	11	
13	23	3	58	47	3	18	25	32	4	24	19	3	44	25	19	4	49	38	4	6	25	7	5	14	45	4	29	24	56	5	39	41	4	52	24	44	7	11	
11	24	4	2	5	3	13	25	53	4	27	58	3	39	25	42	4	53	40	4	2	25	29	5	19	9	4	24	25	18	5	44	27	4	46	25	7	6	11	
11	25	4	5	18	3	9	26	15	4	31	33	3	35	26	4	4	57	37	3	57	25	51	5	23	28	4	19	25	40	5	49	8	4	41	25	29	5	11	
12	26	4	8	27	3	5	26	37	4	35	4	3	31	26	25	5	1	29	3	52	26	13	5	27	42	4	14	26	2	5	53	44	4	36	25	51	4	11	
12	27	4	11	32	3	0	26	58	4	38	30	3	22	26	46	5	5	16	3	47	26	35	5	31	51	4	9	26	23	5	58	14	4	30	26	11	3	12	
13	28	4	14	32	2	57	27	20	4	41	52	3	16	27	7	5	8	59	3	43	26	55	5	35	54	4	3	26	44	6	2	38	4	24	26	33	2	11	
11	29	4	17	29	2	52	27	39	4	45	8	3	13	27	28	5	12	36	3	37	27	16	5	39	52	3	58	27	5	6	6	57	4	19	26	54	1	11	
13	30	4	20	21	2	48	28	0	4	48	21	3	8	27	47	5	16	8	3	32	27	36	5	43	44	3	52	27	26	6	11	10	4	13	27	15	0	11	
+		+			+		+			+		+			+		+			+		+		+			+		+		+		+		+			+	
D eq. An		Eq. D.		Dif.	5000		D.Eq.		Dif.	50000		Eq. D.		Dif.	5000		D.Eq.		Dif.	5000		Eq. D.		Dif.	5000		D.Eq.		Dif.	5000		Eq. D.		Dif.	5000		D eq. An	5000	
45000		—			—		—			—		—			—		—			—		—			—		—			—		—			—			—	
SIGN 10.																																							

SIGN II.

EXAMPLE II. To find the Equation of the Moon's Center answerable to $1^h 15^m 20^s$ Sun at Δ Apog. and $3^h 5^m 44^s$ Δ equated Anomaly?

By Tab. p. 81. Sun at Δ Ap. 1. equat. S. $15^m 20^s$ gives	56153	double = 2306
By Tab. p. 87. Arg. $3^h 5^m 44^s$ Δ eqd An.	— 6° 18' 5". Excess of Eqn. for 5000	34' 28" = 34.46
Prop. Eqn.	+ 7 56	13836
Δ Central Equation correctly	— 6 26 1 reqd. See p. 96.	9224
Constant	1426+	9224
Excess Eccl. 19' 13"	4945+	6918
Excess Eccl. 34 28	2408+	+ 7,9464,76
Prop. Eqn. 7 56 as before.	3779	56"

The same Rules to be observed in other Cases.

Product cutting off 4 Decimals, besides the multiplying Decimals.

SUPPLEMENTAL

SUPPLEMENTAL LUNAR TABLES.

IV. ELLIPTIC EQUATION of the Moon's Centr. According to the Newtonian Theory.																																	
Argument, The Moon's Equated Anomaly, Or D 3 Equa. — Ap. 2 Equated.																																	
SIGN 2.																																	
D eq A.	45000			Dif.	D. Eq for D. 5000	50000			Dif.	D. Eq for D. 5000	55000			Dif.	D. Eq for D. 5000	60000			Dif.	D. Eq for D. 5000	65000			Dif.	D. Eq for D. 5000	D eq A.							
	Eq. D.	—	—			Eq. D.	—	—			Eq. D.	—	—			Eq. D.	—	—															
10	0	4	20	21	2	48	28	04	48	21	3	7	27	47	5	16	8	3	28	27	36	5	43	44	3	47	27	26	6	11	10		
11	1	4	23	9	2	43	28	19	4	51	28	3	7	28	38	5	19	36	3	28	27	55	5	47	31	3	47	27	45	6	15	16	
13	2	4	25	52	2	39	28	39	4	54	31	3	3	28	26	5	22	57	3	21	28	15	5	51	12	3	41	28	5	6	19	17	
12	3	4	28	31	2	35	28	57	4	57	28	2	57	28	45	5	26	13	3	16	28	35	5	54	48	3	36	28	24	6	23	12	
11	4	4	31	6	2	30	29	15	5	0	21	2	53	29	4	5	29	25	3	12	28	53	5	58	18	3	30	28	42	6	27	0	
12	5	4	33	36	2	24	29	33	5	3	9	2	48	29	21	5	32	30	3	5	29	12	6	1	42	3	24	29	1	6	30	43	
11	6	4	36	1	2	20	29	50	5	5	51	2	42	29	39	5	35	30	3	0	29	30	6	5	0	3	18	29	19	6	34	19	
12	7	4	38	21	2	16	30	8	5	8	25	2	38	29	56	5	38	25	2	55	29	47	6	8	12	3	12	29	36	6	37	48	
11	8	4	40	37	2	11	30	24	5	11	1	2	32	30	13	5	41	14	2	49	30	3	6	11	17	3	16	29	54	6	41	11	
11	9	4	42	48	2	7	30	40	5	13	28	2	27	30	29	5	43	57	2	43	30	20	6	14	17	3	0	30	10	6	44	27	
10	10	4	44	55	2	1	30	55	5	15	50	2	22	30	45	5	46	35	2	38	30	35	6	17	10	2	53	30	27	6	47	37	
10	11	4	46	56	1	56	31	10	5	18	6	2	16	31	0	5	49	0	2	31	30	52	6	19	58	2	48	30	42	6	50	40	
10	12	4	48	52	1	52	31	25	5	20	17	2	11	31	15	5	51	32	2	26	31	6	6	22	38	2	40	30	58	6	53	36	
8	13	4	50	44	1	46	31	38	5	22	22	2	5	31	30	5	53	52	2	20	31	6	6	22	38	2	35	30	58	6	53	36	
9	14	4	52	30	1	42	31	52	5	24	22	2	0	31	30	5	53	52	2	13	31	21	6	25	13	2	28	31	12	6	56	25	
9	15	4	54	12	1	36	32	5	26	17	1	55	31	43	5	56	5	2	8	31	36	6	27	41	2	21	31	26	6	59	7		
9	16	4	55	48	1	31	32	18	5	28	6	1	49	32	9	6	0	15	1	2	31	49	6	30	2	2	31	40	7	1	42		
9	17	4	57	19	1	26	32	30	5	29	49	1	43	32	21	6	2	10	1	55	32	16	32	16	2	8	2	14	31	54	7	4	10
8	18	4	58	45	1	21	32	41	5	31	26	1	37	32	33	6	3	59	1	49	32	26	6	36	25	1	1	32	7	7	6	31	
6	19	5	0	6	1	15	32	51	5	32	57	1	26	32	45	6	5	42	1	43	32	38	6	38	20	1	55	32	20	7	8	45	
7	20	5	1	21	1	10	33	2	5	34	23	1	20	32	55	6	7	18	1	36	32	49	6	40	7	1	47	32	31	7	10	51	
7	21	5	2	31	1	5	33	12	5	35	43	1	14	33	5	6	8	48	1	30	33	0	6	41	48	1	41	32	53	7	14	41	
6	22	5	3	36	1	0	33	21	5	36	57	1	9	33	15	6	10	12	1	24	33	9	6	43	21	1	33	33	5	7	16	26	
7	23	5	4	36	0	54	33	30	5	38	6	1	2	33	23	6	11	29	1	17	33	19	6	44	48	1	19	33	14	7	18	2	
6	24	5	5	30	0	48	33	38	5	39	8	0	56	33	32	6	12	40	1	11	33	27	6	46	7	1	13	33	23	7	19	30	
6	25	5	6	18	0	43	33	46	5	40	4	0	50	33	40	6	13	44	1	4	33	36	6	47	20	1	13	33	31	7	20	51	
4	26	5	7	1	0	38	33	52	5	40	53	0	49	33	48	6	14	41	0	57	33	44	6	48	25	1	5	33	39	7	22	4	
3	27	5	7	39	0	32	33	58	5	41	37	0	44	33	55	6	15	32	0	51	33	51	6	49	23	0	58	33	47	7	23	10	
3	28	5	8	11	0	26	34	4	5	42	15	0	38	34	1	6	16	16	0	44	33	58	6	50	14	0	51	33	54	7	24	8	
3	29	5	8	37	0	21	34	10	5	42	47	0	32	34	6	6	16	53	0	37	34	3	6	50	56	0	42	34	1	7	24	57	
4	30	5	8	58	0	15	34	14	5	43	12	0	25	34	11	6	17	23	0	30	34	10	6	51	33	0	37	34	6	7	25	39	
SIGN 7.																																	

EXAMPLE III. To find the Equation of an Orbit correspondent to 60000 Eccentricity, and 30° , or 1° Equated Mean Anomaly? See p. 60. Laws of the Moon's Mean Motion, at the End of Vol. II. Principia.

By p. 84, under Eccentricity 60000 (the same as 56 to 1 Rad.) and against 30° on the Side we find $3^\circ 13' 33''$ required. Mr. Machin has deduced it by his Method $3^\circ 13' 34''$. By Bullialdus's Correction of Ward's Hypothesis.

Com. Logarithms.

As Sem. Conj. $\sqrt{1.06 \times .94}$ 0.0007831 co.
 To Sem. Transverse 0.0000000
 So T. M. Anom. 30° 9.7614394
 To T. of corrected M. An. $30^\circ 2' 41''$: 9.7622225
 Correction of M. Anom. $+2' 41''$ Difference.
 Correction $\frac{1}{2}$ M. Anom. $+1^\circ 20\frac{1}{2}'$ the half, or Eqn
 M. Anom. corrected $15^\circ 1' 20\frac{1}{2}'$

Now, As Ap. Dist. 1.06 9.9746941 co.
 To Perih. Dist.94 9.9731279
 So T. $15^\circ 1' 20\frac{1}{2}'$ }
 $\frac{1}{2}$ M. Anom. corrected. } 9.4287300
 To T. $\frac{1}{2}$ true An. $13^\circ 23' 12''$ 9.3765520
 Double 26 46.24
 Equid M. An. 30° 0.0000000
 Equation $-3^\circ 13' 36''$ the Difference.

SUPPLEMENTAL

SUPPLEMENTAL LUNAR TABLES.

IV. ELLIPTIC EQUATION of the MOON'S Center, according to the Newtonian Theory.

Argument. The MOON'S Equated Anomaly, Or D 3 Equated — D Ap. 2 Equated.

SIGN. 3.

Argument. The Moon's Equated Anomaly, Or D 3 Equated — D Ap. 2 Equated.																					
SIGN. 3.																					
In. of De fr. M. for	45000				50000				55000				60000				65000				In. of De fr. M. for
	Eq. D.	Dif.	D.Eq. for D.	Eq. D.	Dif.	D.Eq. for D.	Eq. D.	Dif.	D.Eq. for D.	Eq. D.	Dif.	D.Eq. for D.	Eq. D.	Dif.	D.Eq. for D.	Eq. D.	Dif.	D.Eq. for D.			
+	—		5000	—		5000	—		5000	—		5000	—		5000	—		5000	—		
11	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18		
3	0	5	8	58	0	16	34	14	5	43	12	0	20	34	11	6	17	23	0	24	
3	1	5	9	14	0	10	34	18	5	43	32	0	13	34	15	6	17	47	0	17	
2	2	5	9	24	0	4	34	21	5	43	45	0	7	34	19	6	18	4	0	10	
2	3	5	9	28	0	2	34	24	5	43	52	0	0	34	22	6	18	14	0	3	
1	4	5	9	26	0	7	34	26	5	43	52	0	6	34	25	6	18	17	0	4	
0	5	5	9	19	0	13	34	27	5	43	46	0	12	34	27	6	18	13	0	11	
0	6	5	9	6	0	19	34	28	5	43	34	0	18	34	28	6	18	2	0	18	
1	7	5	8	47	0	24	34	29	5	43	16	0	25	34	28	6	17	44	0	24	
1	8	5	8	23	0	30	34	28	5	42	51	0	31	34	29	6	17	20	0	32	
0	9	5	7	53	0	36	34	27	5	42	20	0	37	34	28	6	16	48	0	39	
0	10	5	7	17	0	41	34	26	5	41	43	0	44	34	26	6	16	9	0	45	
2	11	5	6	36	0	48	34	23	5	40	59	0	51	34	25	6	15	24	0	53	
3	12	5	5	48	0	53	34	20	5	40	8	0	56	34	23	6	14	31	1	0	
2	13	5	4	55	0	58	34	17	5	39	12	1	3	34	19	6	13	31	1	7	
3	14	5	3	57	1	5	34	12	5	38	9	1	10	34	15	6	12	24	1	14	
4	15	5	2	52	1	10	34	7	5	36	59	1	16	34	11	6	11	10	1	21	
5	16	5	1	42	1	16	34	1	5	35	43	1	22	34	6	6	9	49	1	29	
4	17	5	0	26	1	22	33	55	5	34	21	1	29	33	59	6	8	20	1	35	
5	18	4	59	4	1	27	33	48	5	32	52	1	35	33	53	6	6	45	1	42	
6	19	4	57	37	1	33	33	40	5	31	17	1	41	33	46	6	5	3	1	49	
6	20	4	56	4	1	39	33	32	5	29	36	1	48	33	38	6	3	14	1	57	
6	21	4	54	25	1	44	33	23	5	27	48	1	54	33	29	6	1	7	2	3	
7	22	4	52	41	1	50	33	13	5	25	54	2	1	33	20	5	59	14	2	11	
8	23	4	50	51	1	56	33	2	5	23	53	2	6	33	10	5	57	3	2	17	
7	24	4	48	55	2	1	32	52	5	21	47	2	13	32	59	5	54	46	2	25	
7	25	4	46	54	2	7	32	40	5	19	34	2	20	32	47	5	52	21	2	31	
9	26	4	44	47	2	13	32	27	5	17	14	2	25	32	36	5	49	50	2	38	
8	27	4	42	34	2	18	32	15	5	14	49	2	32	32	23	5	47	12	2	45	
9	28	4	40	16	2	23	32	1	5	12	17	2	38	32	10	5	44	27	2	53	
9	29	4	37	53	2	29	31	46	5	9	39	2	44	31	55	5	41	34	2	59	
9	30	4	35	24	3		31	31	5	6	55			31	40	5	38	35	2		
+	D	+					5000	+						5000	+						
	Eq. D.	Dif.	D.Eq.	Eq. D.	Dif.	D.Eq.	Eq. D.	Dif.	D.Eq.	Eq. D.	Dif.	D.Eq.	Eq. D.	Dif.	D.Eq.	Eq. D.	Dif.	D.Eq.	Eq. D.		
	45000			50000			55000			60000			65000								

SIGN 8.

THE foregoing Example wrought out according to Dr. Halley's Method.

Conf. Log.

Eccentricity .06 to Rad. 1. . or 60000 to Rad. 1000000 gives 9.947824 . . 1° 5' 14' Sun à D Ap.

 $\frac{1}{2}$ Equated M. Anom. 15°

P. 82 . . Correction 0 1' 22"

15 1 22 Tang. $\frac{1}{2}$ M. An. corrected . . 9.4287374 Log. +Tang. $\frac{1}{2}$ True Anom. 13° 23' 12" 44" 9.3765614

Doubled . 26 46 25 28 True An.

Equated M. Anom. 30

By Dr. Halley the Correct Equation — 3° 13' 34" 32" required,

By Bullialdus . — 3 13 36 0

+ 1' 28" Error.

SUPPLEMENTAL

SUPPLEMENTAL LUNAR TABLES.

IV. ELLIPTIC EQUATION of the MOON'S *Center*, According to the Newtonian Theory.

Argument. The Moon's Equated Anomaly, Or $\text{D } 3.$ Equa^d. — $\text{D } \text{Ap. } 2.$ Equa^d.

SIGN 4.

fr. M for		SIGN 4.																																																In fr. M for	
D	Eq A.	45000				Dif.	D. Eq 50000				Dif.	D. Eq 55000				Dif.	D. Eq 60000				Dif.	D. Eq 65000				Dif.	D. Eq 5000				Dif.	D. Eq A.	D																		
		Eq. D.					Eq. D.					Eq. D.					Eq. D.					Eq. D.					Eq. D.																								
		—					—					—					—					—					—																								
		°	'	″	‴		°	'	″	‴		°	'	″	‴		°	'	″	‴		°	'	″	‴		°	'	″	‴				°	'	″	‴	°	'	″	‴										
9	0	4	35	24		31	31	5	6	55		31	40	5	38	35		31	52	6	10	27		32	2	6	42	29		32	12	30																			
9	1	14	32	49	2	35	31	16	5	4	5	2	50	31	25	5	35	30	3	5	31	36	6	7	6	3	21	32	2	6	42	29	3	35	32	12	30														
10	2	24	30	9	2	40	30	59	5	1	8	2	57	31	9	5	32	17	3	13	31	36	6	7	6	3	28	31	48	6	38	54	3	44	32	0	29														
10	3	34	27	24	2	45	30	42	4	58	6	3	2	31	9	5	32	17	3	19	31	21	6	3	38	3	36	31	32	6	35	10	3	44	31	43	28														
10	4	44	24	34	2	50	30	24	4	54	58	3	8	30	52	5	28	58	3	26	31	4	6	0	2	3	43	31	16	6	31	18	3	52	31	28	27														
12	5	54	21	38	2	56	30	5	4	51	43	3	15	30	34	5	25	32	3	32	30	47	5	56	19	3	50	30	59	5	27	18	4	0	8	31	11	26													
12	6	4	18	37	3	1	29	46	4	48	23	3	20	29	58	5	18	21	3	39	30	10	5	48	31	3	58	30	41	6	23	10	4	16	30	53	25														
11	7	74	15	30	3	7	29	27	4	44	57	3	26	29	58	5	14	35	3	46	30	10	5	48	31	4	5	30	23	6	18	54	4	16	30	36	24														
12	8	84	12	19	3	11	29	6	4	41	25	3	32	29	38	5	10	43	3	52	29	51	5	44	26	4	12	30	4	6	14	30	4	24	30	17	23														
14	9	94	9	3	3	16	28	44	4	37	47	3	38	29	18	5	6	45	3	58	29	31	5	40	14	4	19	29	44	6	9	58	4	32	29	51	22														
13	10	4	5	41	3	22	28	23	4	34	4	3	43	28	36	5	2	40	4	5	29	10	5	35	55	4	26	29	24	6	5	19	4	39	29	38	21														
13	11	4	2	14	3	27	28	1	4	30	15	3	47	28	14	4	58	29	4	11	28	27	5	26	56	4	33	28	41	5	55	37	4	55	28	55	19														
12	12	3	58	42	3	32	27	39	4	26	21	3	54	27	51	4	54	12	4	17	28	4	5	22	16	4	40	28	18	5	50	34	5	3	28	32	18														
11	13	3	55	6	3	36	27	15	4	22	21	4	0	27	27	4	49	48	4	24	27	41	5	17	29	4	47	27	55	5	45	24	5	10	28	9	17														
12	14	3	51	25	3	41	26	51	4	18	16	4	5	27	3	4	45	19	4	29	27	17	5	12	36	4	53	27	31	5	40	7	5	17	27	45	16														
13	15	3	47	40	3	45	26	25	4	14	5	4	11	26	38	4	40	43	4	36	26	53	5	7	36	5	0	27	6	5	34	42	5	25	27	19	15														
13	16	3	43	49	3	51	26	0	4	9	49	4	16	26	13	4	36	2	4	41	26	27	5	2	29	5	7	26	42	5	29	11	5	31	26	57	14														
13	17	3	39	54	3	55	25	34	4	5	28	4	21	25	47	4	31	15	4	47	26	1	4	57	16	5	13	26	16	5	23	32	5	39	26	31	13														
14	18	3	35	54	4	0	25	7	4	1	1	4	27	25	21	4	26	22	4	53	25	35	4	51	57	5	19	25	49	5	17	46	5	46	26	3	12														
14	19	3	31	50	4	4	24	40	3	56	30	4	31	24	54	4	21	24	4	58	25	7	4	46	31	5	26	25	22	5	11	53	5	53	25	37	11														
13	20	3	27	42	4	8	24	12	3	51	54	4	36	24	25	4	16	19	5	5	24	40	4	40	59	5	32	24	54	5	5	53	6	0	25	10	10														
15	21	3	23	29	4	13	23	43	3	47	12	4	42	23	58	4	11	10	5	9	24	11	4	35	21	5	38	24	26	4	59	47	6	6	24	41	9														
15	22	3	19	12	4	17	23	14	3	42	26	4	46	23	29	4	5	55	5	15	23	42	4	29	37	5	44	23	57	4	53	34	6	13	24	12	8														
13	23	3	14	50	4	22	22	46	3	37	36	4	50	22	59	4	0	35	5	20	23	12	4	23	47	5	50	23	27	4	47	14	6	20	23	42	7														
15	24	3	10	25	4	25	22	15	3	32	40	4	56	22	30	3	55	10	5	25	22	42	4	17	52	5	55	22	56	4	40	48	6	26	23	10	6														
13	25	3	5	56	4	29	21	45	3	27	41	4	59	21	58	3	49	39	5	31	22	12	4	11	51	6	1	22	25	4	34	16	6	31	22	38	5														
13	26	3	1	22	4	34	21	14	3	22	36	5	5	21	27	3	44	3	5	36	21	41	4	5	44	6	7	21	54	4	27	38	6	38	22	7	4														
12	27	2	56	45	4	37	20	43	3	17	28	5	8	20	55	3	38	23	5	40	21	9	3	59	32	6	12	21	22	4	20	54	6	44	21	35	3														
13	28	2	52	5	4	40	20	10	3	12	15	5	13	20	23	3	32	38	5	45	20	36	3	53	14	6	18	20	50	4	14	4	6	50	21	14	2														
12	29	2	47	20	4	45	19	38	3	6	58	5	17	19	50	3	26	48	5	50	20	4	3	46	52	6	22	20	16	4	7	8	6	56	20	28	1														
12	30	2	42	32	4	48	19	5	3	1	37	5	21	19	17	3	20	54	5	54	19	30	3	40	24	6	28	19	43	4	0	7	7	1	19	56	0														
5000 D.		+		D Eq A.		45000		Dif.		5000		D. Eq		+		D Eq A.		45000		Dif.		5000		D. Eq		+		D Eq A.		45000		Dif.		5000		D. Eq		+		D Eq A.		45000									

SIGN 7.

To determine the EQUATION of the Mean to the True Anomaly, or Mean to the true Place in all the Planetary Orbits except that of Mercury, for which the following Rule comes near the Truth.

RULE (deduced from expressions 1st and 2d p. 49, *Palladium*, 1757.)

Art. 1. As Radius is to Cofine of the given Mean Anomaly, so is the Eccentricity of the Orbit more its Fourth, (Rad. or mean Dist. being r .) to a Fourth Proportional.

2. Add this *Fourth Proportional* to the *Sem. Transverse* ($= 1$.) if the *Mean Anomaly* given be *less* 90° , or *more* than 270° ; or *Subtract* it from the *Sem. Transverse* if the *Mean Anom.* be *more* than 90° or *less* than 270° .

3. Then say, As the *Sum*, in one Case, or *Difference* in the other, is to double the said *Eccentricity*, so is the *Sine* of the given *Mean Anomaly* to the *Proportional Sine* of an Arc P.

4. The *Minutes* or *Seconds* of another Arc, answerable to *one third* of the *Cube* of this proportional *Sine* of an Arc P, being deducted from the Degrees Minutes and Seconds of the said Proportional Arc P, the Residue

N. B. This Arc of Difference to be subtracted from the Mean Anomaly found in the 1st, being

N. B. This Art of Difference to be subtracted from the Mean Anomaly for the true in the first Six Signs of the Argument, and to be added thereto in the last Six Signs.

the last six signs.

SUPPLEMENTAL

SUPPLEMENTAL LUNAR TABLES.

OR THUS. EQUATION of the MEAN to the present ELLIPTIC EQUATION of the MOON'S CENTER.

Arg. Oà D's Apogee, 1. equated.			SIGN O.															
			Argument. Equated ANOMALY of the MOON; Or, D 3. equated — Ap. D 2. equated.															
			0°	2°	4°	6°	8°	10°	12°	14°	16°	18°	20°	22°	24°	26°	28°	30°
			+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
			I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I
6	0	12	0	2 26	4 53	7 16	9 42	12 9	14 32	16 57	19 18	21 39	23 59	26 19	28 40	30 56	33 12	35 26
	2	28	0	2 25	4 52	7 15	9 40	12 7	14 30	16 54	19 15	21 35	23 55	26 14	28 34	30 50	33 6	35 20
	4	26	0	2 24	4 50	7 13	9 37	12 2	14 25	16 48	19 8	21 28	23 47	26 5	28 24	30 40	32 54	35 8
	6	24	0	2 23	4 47	7 8	9 31	11 56	14 17	16 38	18 57	21 16	23 34	25 51	28 9	30 23	32 36	35 48
	8	22	0	2 20	4 43	7 3	9 24	11 46	14 5	16 24	18 41	20 57	23 12	25 28	27 44	29 56	32 8	34 18
	10	20	0	2 18	4 38	6 55	9 13	11 32	13 49	16 6	18 21	20 35	22 46	24 59	27 14	29 23	31 32	33 40
	12	18	0	2 15	4 32	6 46	9 1	11 17	13 30	15 44	17 56	20 7	22 17	24 27	26 37	28 44	30 50	32 54
	14	16	0	2 11	4 24	6 35	8 46	10 59	13 8	15 19	17 27	19 34	21 41	23 47	25 53	27 57	30 0	32 1
	16	14	0	2 8	4 17	6 23	8 3	10 37	12 42	14 48	16 52	18 55	20 57	22 59	25 2	27 2	29 1	30 59
	18	12	0	2 2	4 7	6 10	8 11	10 14	12 15	14 17	16 16	18 15	20 14	22 11	24 9	26 4	27 57	29 48
	20	10	0	1 57	3 57	5 54	7 50	9 47	11 42	13 39	15 33	17 26	19 19	21 12	23 5	24 55	26 44	28 32
	22	8	0	1 51	3 46	5 39	7 28	9 19	11 8	12 59	14 47	16 35	18 23	20 10	21 56	23 41	25 25	27 8
	24	6	0	1 45	3 33	5 19	7 3	8 48	10 31	12 17	14 0	15 41	17 23	19 4	20 46	22 24	24 2	25 39
	26	4	0	1 39	3 20	4 59	6 36	8 14	9 49	11 28	13 3	14 38	16 14	17 49	19 24	20 56	22 27	23 58
	28	2	0	1 31	3 5	4 38	6 7	7 38	9 6	10 39	12 8	13 35	15 4	16 31	17 59	19 23	20 48	22 13
1	0	5	0	1 28	2 55	4 16	5 37	6 59	8 20	9 44	11 5	12 26	13 48	15 9	16 31	17 47	19 4	20 21
7	2	11	28	0	1 15	2 34	3 51	5 4	6 20	7 33	8 49	10 2	11 15	12 29	13 42	14 55	16 5	17 15
	4	26	0	1 6	2 16	3 25	4 1	5 38	6 43	7 51	8 56	10 1	11 6	12 8	13 8	14 15	15 21	16 24
	6	24	0	0 58	1 59	2 59	3 56	4 54	5 51	6 50	7 46	8 43	9 40	10 36	11 32	12 26	13 20	14 15
	8	22	0	0 50	1 42	2 31	3 19	4 8	4 55	5 46	6 34	7 21	8 9	8 53	9 35	10 27	11 16	12 2
	10	20	0	0 40	1 22	2 2	2 41	3 21	3 59	4 40	5 19	5 58	6 37	7 14	7 53	8 30	9 7	9 44
	12	18	0	0 30	1 3	1 33	2 2	2 33	3 1	3 33	4 3	4 32	5 1	5 30	5 59	6 28	6 50	7 24
	14	16	0	0 20	0 43	1 3	1 22	1 44	2 3	2 24	2 44	3 4	3 23	3 42	4 1	4 21	4 41	5 1
	16	14	0	0 10	0 23	0 33	0 42	0 53	1 2	1 15	1 25	1 35	1 44	1 54	2 4	2 16	2 26	2 34
	18	12	0	0 0	0 0	0 1	0 1	0 2	0 1	0 4	0 4	0 3	0 3	0 3	0 4	0 3	0 5	0 5
	20	10	0	0 10	0 18	0 29	0 40	0 49	1 0	1 9	1 19	1 30	1 39	1 48	1 57	2 6	2 15	2 25
	22	8	0	0 20	0 38	1 1	1 22	1 40	2 1	2 20	2 41	3 1	3 20	3 40	3 59	4 17	4 36	4 56
	24	6	0	0 30	0 59	1 32	2 3	2 33	3 4	3 33	4 4	4 33	5 3	5 33	6 1	6 29	6 58	7 27
	26	4	0	0 41	1 20	2 2	2 43	3 23	4 5	4 45	5 26	6 5	6 44	7 24	8 2	8 40	9 17	9 55
	28	2	0	0 51	1 40	2 33	3 24	4 14	5 5	5 56	6 46	7 34	8 23	9 13	10 1	10 49	11 36	12 24
2	0	4	0	0 1	2 0	3 3	4 4	5 4	6 5	7 6	8 6	9 4	10 3	11 1	11 59	12 56	13 52	14 48
8	2	10	28	0	1 11	2 20	3 33	4 44	5 52	7 3	8 13	9 23	10 31	11 39	12 47	13 53	15 0	16 6
	4	26	0	1 21	2 40	4 2	5 22	6 40	8 0	9 19	10 38	11 55	13 12	14 29	15 45	17 1	18 16	19 28
	6	24	0	1 30	2 59	4 30	5 59	7 26	8 55	10 22	11 50	13 17	14 43	16 9	17 33	18 57	20 20	21 42
	8	22	0	1 38	3 16	4 58	6 34	8 9	9 47	11 26	13 3	14 36	16 16	17 43	19 16	20 48	22 19	23 48
	10	20	0	1 47	3 33	5 23	7 7	8 50	10 36	12 20	14 5	15 49	17 31	19 13	20 53	22 33	24 12	25 48
	12	18	0	1 55	3 49	5 47	7 38	9 29	11 22	13 13	15 6	16 58	18 47	20 36	22 24	24 11	25 57	27 41
	14	16	0	1 2	4 4	6 9	8 6	10 4	12 5	14 0	16 0	18 1	19 57	21 53	23 47	25 41	27 34	29 27
	16	14	0	2 9	4 17	6 28	8 32	10 36	12 44	14 47	16 52	18 58	21 1	23 2	25 27	27 3	29 1	30 57
	18	12	0	2 14	4 28	6 47	8 56	11 4	13 17	15 26	17 38	19 49	21 57	24 4	26 9	28 15	30 18	32 19
	20	10	0	2 19	4 38	7 2	9 15	11 28	13 46	16 0	18 16	20 33	22 46	24 57	27 7	29 17	31 24	33 30
	22	8	0	2 23	4 46	7 14	9 31	11 48	14 11	16 28	18 48	21 8	23 25	25 40	27 54	30 8	32 19	34 29
	24	6	0	2 26	4 53	7 24	9 44	12 5	14 30	16 51	19 14	21 38	23 58	26 16	28 33	30 50	33 5	35 17
	26	4	0	2 28	4 57	7 31	9 54	12 17	14 44	17 7	19 32	21 58	24 20	26 41	28 59	31 19	33 36	35 50
	28	2	0	2 30	4 59	7 34	9 57	12 22	14 52	17 10	19 43	22 10	24 33	26 55	29 15	31 36	33 54	36 9
3	0	3	0	2 31	5 2	7 37	10 1	12 26	14 56	17 20	19 47	22 15	24 39	27 2	29 22	31 43	34 2	36 18
9	9			—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Oà D's Ap. 1. equated.			30°	28°	26°	24°	22°	20°	18°	16°	14°	12°	10°	8°	6°	4°	2°	0°
			SIGN II.															

N. B. If any Number of Minutes should chance to be misprinted, it will appear by the foregoing and following Quantities, at Sight.

SUPPLEMENTAL

EQUATION of the MEAN to the present ELLIPTIC EQUATION of the MOON'S CENTER.

SIGN 1.

Argument. Equated ANOMALY of the MOON, Or, $\text{D } 3.$ equated — Ap. $\text{D } 2.$ equated.

SIGN 10.

22

SUPPLEMENTAL

SUPPLEMENTAL LUNAR TABLES.

EQUATION of the MEAN to the present ELLIPTIC EQUATION of the MOON'S CENTER.

Arg. @ à D's Apogee, 1. equated.		SIGN 2.															
		Argument. Equated ANOMALY of the MOON; Or, D 3. Equated—Ap. D 2. equated.															
		0°	2°	4°	6°	8°	10°	12°	14°	16°	18°	20°	22°	24°	26°	28°	30°
		+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
6	0 12	64 25	66 0	67 30	68 56	70 15	71 31	72 43	73 51	74 55	75 56	76 48	77 36	78 21	79 1	79 34	80 3
	2	64 12	65 46	67 16	68 43	70 2	71 18	72 29	73 37	74 41	75 42	76 34	77 21	78 6	78 46	79 19	79 47
	4	63 31	65 25	66 54	68 18	69 38	70 54	72 6	73 12	74 16	75 17	76 8	76 55	77 39	78 19	78 52	79 20
	6	63 15	64 48	66 15	67 39	68 58	70 13	71 24	72 30	73 34	74 35	75 25	76 12	76 56	77 35	78 7	78 31
	8	62 20	63 51	65 17	66 41	67 58	69 12	70 21	71 27	72 29	73 25	74 15	75 5	75 47	76 26	76 58	77 26
	10	61 10	62 39	64 5	65 29	66 46	67 55	69 3	70 7	71 8	72 7	72 56	73 42	74 25	75 2	75 33	76 0
12	18	59 49	61 16	62 40	64 0	65 14	66 15	67 31	68 33	69 33	70 30	71 17	72 1	72 42	73 21	73 52	74 18
14	16	58 10	59 35	60 56	62 15	63 26	64 35	65 40	66 40	67 38	68 33	69 20	70 3	70 43	71 19	71 49	72 14
16	14	56 20	57 40	58 57	60 1	61 24	62 28	63 30	64 28	65 24	66 19	67 4	67 46	68 25	68 58	69 27	69 53
18	12	54 10	55 29	56 45	57 56	59 3	60 6	61 5	62 4	62 59	63 50	64 33	65 13	65 50	66 24	66 52	67 16
20	10	51 51	53 6	54 17	55 27	56 31	57 33	58 34	59 26	60 16	61 5	61 46	62 24	62 59	63 31	63 58	64 20
22	8	49 17	50 29	51 37	52 43	53 46	54 42	55 37	56 29	57 18	58 4	58 34	59 19	59 52	60 23	60 48	61 10
24	6	46 37	47 44	48 48	49 51	50 48	51 43	52 34	53 24	54 11	54 54	55 31	56 4	56 35	57 4	57 28	57 48
26	4	43 32	44 35	45 34	46 32	47 26	48 17	49 5	49 51	50 35	51 15	51 49	52 20	52 48	53 16	53 39	53 57
28	2	40 21	41 19	42 14	43 8	43 58	44 45	45 29	46 13	46 52	47 29	48 1	48 30	48 56	49 22	49 42	49 59
30	0	36 59	37 52	38 42	39 31	40 17	41 0	41 39	42 19	42 56	43 30	43 59	44 26	44 49	45 13	45 31	45 47
7	2 11	33 27	34 14	34 59	35 43	36 24	37 4	37 41	38 15	38 48	39 20	39 47	40 11	40 32	40 52	41 9	41 23
	4	29 45	30 28	31 8	31 46	32 24	33 0	33 32	34 2	34 31	35 0	35 24	35 45	36 4	36 24	36 38	36 50
	6	25 50	26 27	27 2	27 36	28 9	28 39	29 8	29 34	29 59	30 24	30 45	31 3	31 19	31 36	31 49	31 59
	8	21 47	22 20	22 50	23 18	23 45	24 11	24 35	24 58	25 19	25 40	25 57	26 12	26 25	26 39	26 50	26 58
	10	17 36	18 4	18 29	19 52	19 14	19 35	19 54	20 12	20 30	20 48	21 1	21 12	21 23	21 35	21 43	21 49
12	18	13 19	13 42	14 2	14 19	14 35	14 51	15 6	15 20	15 34	15 48	15 57	16 5	16 13	16 23	16 30	16 36
14	16	8 58	9 15	9 29	9 40	9 51	10 2	10 12	10 22	10 32	10 40	10 46	10 52	10 57	11 3	11 8	11 14
16	14	4 30	4 43	4 51	4 56	5 3	5 8	5 13	5 18	5 23	5 29	5 31	5 33	5 35	5 39	5 43	5 47
18	12	0 9	0 3	0 9	0 8	0 9	0 9	0 9	0 9	0 9	0 10	0 9	0 9	0 8	0 10	0 18	0 11
		±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±
20	10	4 22	4 30	4 37	4 41	4 46	4 52	4 57	5 1	5 4	5 7	5 12	5 17	5 20	5 20	5 22	5 23
22	8	8 56	9 8	9 20	9 32	9 43	9 53	10 1	10 12	10 22	10 29	10 36	10 43	10 50	10 54	10 58	11 1
24	6	13 29	13 47	14 6	14 24	14 40	14 55	15 10	15 24	15 37	15 47	15 58	16 9	16 19	16 23	16 30	16 36
26	4	17 59	18 24	18 48	19 12	19 33	19 54	20 13	20 33	20 50	20 54	21 19	21 32	21 45	21 52	22 0	22 8
28	2	22 27	22 58	23 27	23 56	24 21	24 49	25 13	25 37	25 59	26 16	26 34	26 51	27 6	27 15	27 27	27 37
30	0	26 50	27 26	28 2	28 37	29 9	29 35	30 8	30 38	31 4	31 25	31 46	32 5	32 24	32 35	32 48	33 0
8	2 10	31 6	31 49	32 30	33 10	33 48	34 24	34 57	35 31	36 0	36 24	36 48	37 11	37 32	37 37	38 2	38 14
	4	35 16	36 4	36 51	37 36	38 20	39 0	39 36	40 14	40 47	41 14	41 41	42 8	42 33	42 49	43 5	43 19
	6	39 14	40 9	41 2	41 52	42 39	43 24	44 7	44 47	45 24	45 55	46 25	46 53	47 20	47 38	47 57	48 13
	8	43 3	44 2	45 0	45 55	46 37	47 36	48 22	49 7	49 48	50 23	50 55	51 26	51 55	52 14	52 34	52 52
	10	46 40	47 44	48 46	49 46	50 42	51 35	52 24	53 13	53 57	54 34	55 9	55 42	56 13	56 36	56 58	57 17
12	18	50 1	51 10	52 17	53 21	54 20	55 56	56 10	57 2	57 49	58 29	59 7	59 42	60 15	60 39	61 2	61 22
14	16	53 6	54 19	55 29	56 38	57 41	58 41	59 38	60 33	61 23	62 6	62 45	63 22	63 57	64 23	64 47	65 9
16	14	55 53	57 10	58 25	59 36	60 43	61 46	62 46	63 43	64 35	65 21	66 2	66 41	67 18	67 45	68 11	68 34
18	12	58 21	59 42	61 1	62 15	63 24	64 29	65 32	66 32	67 26	68 14	67 57	69 37	70 16	70 44	71 11	71 34
20	10	60 29	61 54	63 15	64 32	65 43	66 50	67 35	68 56	69 53	70 44	71 27	72 8	72 49	73 19	73 46	74 10
22	8	62 17	63 42	65 5	66 25	67 38	68 48	69 54	70 58	71 56	72 47	73 33	74 16	74 56	75 26	75 56	76 20
24	6	63 41	65 9	66 34	67 56	69 10	70 21	71 30	72 35	73 33	74 22	75 11	75 56	76 38	77 10	77 39	78 3
26	4	64 41	66 10	67 37	69 0	70 16	71 28	72 37	73 43	74 43	75 37	76 28	77 9	77 51	78 23	78 52	79 17
28	2	65 16	66 45	68 12	69 36	70 53	72 6	73 16	74 22	75 23	76 17	77 6	77 51	78 33	79 5	79 34	79 59
30	0	65 30	67 0	68 28	69 52	71 9	72 22	73 33	74 39	75 40	76 34	77 21	78 6	78 49	79 21	79 50	80 15
9	9	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
② à D's Ap. 1. equated.		30°	28°	26°	24°	22°	20°	18°	16°	14°	12°	10°	8°	6°	4°	2°	0°
SIGN 9																	

SUPPLEMENTAL

SUPPLEMENTAL LUNAR TABLES.

EQUATION of the MEAN to the present ELLIPTIC EQUATION of the MOON'S CENTER.

Arg. $\odot$ à D's Apogee, 1. equated.		SIGN 3.															
		Argument. Equated ANOMALY of the MOON; Or, D 3. equated -- Ap. D 2. equated.															
		0°	2°	4°	6°	8°	10°	12°	14°	16°	18°	20°	22°	24°	26°	28°	30°
		+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
		I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I
		0	6	12	18	24	30	36	42	48	54	0	6	12	18	24	30
6	0	12	0	80	3	80	23	80	41	80	53	80	57	80	48	80	36
2	28	79	47	80	8	80	26	80	38	80	43	80	42	80	33	80	20
4	26	79	20	79	41	79	58	80	10	80	15	80	13	80	4	79	52
6	24	78	35	78	55	79	13	79	25	79	29	79	27	79	18	79	7
8	22	77	26	77	45	78	2	78	14	78	19	78	17	78	8	77	57
10	20	76	0	76	19	76	36	76	46	76	50	76	49	76	40	76	29
12	18	74	18	74	37	74	53	75	2	75	6	75	5	74	56	74	46
14	16	72	14	72	32	72	47	72	57	73	2	73	0	72	51	72	41
16	14	69	53	70	13	70	28	70	34	70	38	70	38	70	31	70	19
18	12	67	16	67	33	67	46	68	54	67	58	67	57	67	49	67	39
20	10	64	20	64	36	64	49	64	57	65	1	64	59	64	52	64	43
22	8	61	10	61	24	61	36	61	44	61	47	62	46	61	40	61	31
24	6	57	48	58	2	58	14	58	20	58	22	58	21	58	6	58	4
26	4	53	57	54	10	54	18	54	26	54	28	54	23	54	17	54	1
28	2	49	59	50	10	50	17	50	23	50	25	50	23	50	17	50	10
30	0	45	47	45	57	46	3	46	2	46	10	46	12	46	7	46	4
7	2	11	28	41	23	41	32	41	41	41	45	41	47	41	46	41	35
4	26	36	50	36	57	37	4	37	7	37	9	37	9	37	5	37	0
6	24	31	39	32	6	32	12	32	15	32	16	32	16	32	13	32	9
8	22	26	58	27	4	27	10	27	12	27	14	27	11	27	11	27	7
10	20	21	49	21	54	21	59	22	1	22	2	22	0	21	58	21	49
12	18	16	36	16	39	16	41	16	42	16	43	16	41	16	40	16	33
14	16	11	14	11	16	11	18	11	17	11	17	11	16	11	16	11	10
16	14	5	47	5	46	5	49	5	55	5	49	5	45	5	44	5	46
18	12	0	11	0	10	0	10	0	10	0	10	0	9	0	11	0	8
20	10	5	23	5	27	5	28	5	28	5	27	5	30	5	26	5	25
22	8	11	1	11	5	11	6	11	8	11	6	11	9	11	4	11	2
24	6	16	36	16	41	16	44	16	46	16	46	16	45	16	40	16	37
26	4	22	8	22	15	22	19	22	21	22	20	22	19	22	18	22	12
28	2	27	37	27	46	27	50	27	51	27	51	27	50	27	48	27	41
30	0	33	0	33	10	33	15	33	17	33	17	33	14	33	11	33	3
8	2	10	28	38	14	38	25	38	30	38	33	38	34	38	32	38	28
4	26	43	18	43	30	43	37	43	41	43	39	43	38	43	33	43	23
6	24	48	13	48	26	48	33	48	37	48	37	48	33	48	27	48	15
8	22	52	52	53	5	53	13	53	17	53	17	53	13	53	28	52	54
10	20	57	17	57	30	57	38	57	43	57	43	57	39	57	33	57	27
12	18	61	22	61	37	61	46	61	51	61	49	61	43	61	35	61	21
14	16	65	9	65	24	65	33	65	39	65	37	65	33	65	26	65	12
16	14	68	34	68	50	68	59	69	3	69	2	68	57	68	50	68	32
18	12	71	34	71	50	71	59	72	5	72	3	71	58	71	50	71	31
20	10	74	10	74	27	74	37	74	43	74	41	74	35	74	27	74	8
22	8	76	20	76	37	76	46	76	51	77	49	76	43	76	33	76	11
24	6	78	3	78	22	78	32	78	37	78	34	78	28	78	21	78	0
26	4	79	17	79	34	79	45	79	51	79	48	79	42	79	34	79	12
28	2	79	59	80	17	80	27	80	33	80	30	80	24	80	16	79	54
30	0	80	15	80	33	80	44	80	51	80	48	80	42	80	30	80	12
9	9	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Arg. $\odot$ à D's Ap 1. equated.		30°	28°	26°	24°	22°	20°	18°	16°	14°	12°	10°	8°	6°	4°	2°	0°

SIGN 8.

SUPPLEMENTAL.

SUPPLEMENTAL LUNAR TABLES.

EQUATION of the MEAN to the present ELLIPTIC EQUATION of the MOON'S CENTER.

Arg. ☉ & Δ's Apogee, 1. equated.		SIGN 4. Argument. Equated ANOMALY of the MOON; Or, Δ 3. equated — Ap. Δ 2. equated.															
		0°	2°	4°	6°	8°	10°	12°	14°	16°	18°	20°	22°	24°	26°	28°	30°
		+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
		1°	1°	1°	1°	1°	1°	1°	1°	1°	1°	1°	1°	1°	1°	1°	1°
6	0 12 0	75 0	73 45	72 27	71 4	69 30	67 53	66 9	64 18	62 22	60 22	58 12	55 56	53 34	51 7	48 38	46 2
	2 28	74 46	73 30	72 13	70 5	69 18	67 41	65 57	64 7	62 12	60 10	58 0	55 45	53 24	50 57	48 28	45 53
	4 26	74 21	73 6	71 48	70 25	68 53	67 17	65 34	63 53	61 48	59 49	57 40	55 26	53 8	50 40	48 12	45 37
	6 24	73 38	72 24	71 8	69 45	68 13	66 38	64 57	63 8	61 13	59 14	57 7	54 54	52 36	50 11	47 45	45 12
	8 22	72 33	71 20	70 5	68 4	67 14	65 39	63 58	62 11	60 18	58 21	56 15	54 5	51 49	49 26	47 2	44 32
	10 20	71 11	70 0	68 46	67 27	65 59	64 27	62 48	61 2	59 11	57 17	55 13	53 4	50 50	48 29	46 9	43 42
	12 18	69 34	68 25	67 13	65 54	64 28	62 58	61 23	59 38	57 49	55 58	53 57	51 51	49 41	47 24	45 6	42 41
	14 16	67 39	66 30	65 20	64 4	62 40	61 13	59 39	57 58	56 12	54 24	52 27	50 25	48 19	46 5	43 51	41 30
	16 14	65 28	64 22	63 15	62 0	60 39	59 15	57 44	56 7	54 24	52 39	50 46	48 48	46 48	44 37	42 26	40 7
	18 12	62 57	61 54	60 48	59 37	58 19	56 58	55 29	53 57	52 19	50 36	48 46	46 53	44 57	42 51	40 47	38 39
	20 10	60 11	59 10	58 8	57 0	55 45	54 27	53 3	51 34	50 0	48 23	46 39	44 52	43 0	40 59	38 59	36 54
	22 8	57 12	56 14	55 15	54 11	52 59	51 44	50 25	49 0	47 30	45 58	44 19	42 37	40 52	38 57	37 2	35 3
	24 6	54 3	53 8	52 12	51 11	50 34	48 53	47 38	46 17	44 52	43 24	41 52	40 16	38 37	36 47	34 59	33 7
	26 4	50 26	49 35	48 42	47 45	46 41	45 36	44 25	43 10	41 51	40 29	39 4	37 35	36 2	34 19	32 59	30 54
	28 2	46 42	45 54	45 6	44 14	43 14	42 14	41 9	40 1	38 45	37 28	36 0	34 47	33 23	31 46	30 13	28 36
1	0 5 0	42 44	42 0	41 17	40 28	39 38	38 43	37 40	36 35	35 28	34 18	33 5	31 51	31 35	29 5	27 39	26 10
7	2 11 28	38 38	37 58	37 18	36 33	35 45	34 55	34 1	33 3	32 2	30 59	29 53	28 43	27 28	26 13	24 58	23 38
	4 26	34 21	33 46	33 11	32 32	31 48	31 3	30 14	29 23	28 28	27 32	26 34	25 31	24 25	23 18	22 11	21 0
	6 24	29 44	29 17	28 48	28 14	27 36	26 57	26 14	25 29	24 42	23 54	23 4	22 9	21 11	20 12	19 15	18 13
	8 22	25 10	24 43	24 17	23 48	23 16	22 44	22 8	21 30	20 50	20 9	19 27	18 41	17 51	17 2	16 15	15 22
	10 20	20 21	19 59	19 38	19 15	19 49	18 24	17 55	17 24	16 51	16 19	15 45	15 7	14 27	13 47	13 8	12 25
	12 18	14 26	15 9	14 54	14 37	14 17	13 57	13 34	13 11	12 46	12 23	11 57	11 29	10 57	10 27	9 58	9 25
	14 16	10 24	10 13	10 3	9 51	9 37	9 24	9 10	8 54	8 37	8 21	8 4	7 45	7 23	7 3	6 44	6 21
	16 14	5 18	5 13	5 9	5 2	4 54	4 48	4 41	4 32	4 24	4 17	4 9	3 58	3 45	3 35	3 27	3 14
	18 12	0 9	0 8	0 9	0 8	0 8	0 8	0 7	0 7	0 7	0 9	0 9	0 7	0 5	0 4	0 6	0 4
	20 10	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±
	22 8	5 3	4 58	4 52	4 47	4 42	4 35	4 29	4 21	4 13	4 3	3 57	3 51	3 43	3 33	3 19	3 6
	24 6	10 16	10 7	9 54	9 43	9 31	9 17	9 5	8 46	8 29	8 12	7 58	7 42	7 23	7 4	6 39	6 16
	26 4	15 27	15 12	14 54	14 37	14 18	13 57	13 37	13 13	12 48	12 20	11 56	11 29	11 11	10 32	9 59	9 26
	28 2	20 35	20 15	19 51	19 28	19 2	18 35	18 8	17 36	17 3	16 27	15 53	15 17	14 40	14 1	13 16	12 33
	2 0 4 0	29 38	25 14	24 44	24 14	23 43	23 8	22 33	21 53	21 13	20 30	19 46	19 1	18 15	17 25	16 30	15 37
2	0 4 0	30 37	30 8	29 33	28 50	28 18	27 37	26 56	26 8	25 19	24 27	23 36	22 42	21 45	20 46	19 42	18 39
8	2 10 28	35 26	34 52	34 13	33 32	32 47	31 59	31 10	30 15	29 19	28 21	27 19	26 17	25 10	24 1	22 48	21 35
	4 26	40 8	39 28	38 43	37 57	37 6	36 12	35 17	34 15	33 12	32 6	30 55	29 42	28 28	27 11	25 48	24 26
	6 24	44 38	43 54	43 4	42 12	41 15	40 15	39 14	38 4	36 54	35 41	34 21	33 1	31 39	30 12	28 40	27 8
	8 22	48 54	48 5	47 11	46 15	45 13	43 6	42 59	41 43	40 26	39 5	37 31	36 10	34 40	33 5	31 24	29 42
	10 20	52 57	51 4	51 50	50 34	48 55	47 44	46 32	45 11	43 49	42 25	40 47	39 9	37 32	35 49	33 59	32 9
	12 18	56 43	55 45	54 42	53 37	52 23	51 6	49 49	48 22	46 52	45 18	43 38	41 56	40 12	38 21	36 25	34 25
	14 16	60 10	59 8	58 1	56 53	55 36	54 14	52 51	51 18	49 43	48 3	46 12	44 28	42 38	40 39	38 34	36 31
	16 14	63 17	62 12	61 2	59 50	58 30	57 3	55 36	53 58	52 18	50 33	48 40	46 46	44 59	42 46	40 35	38 22
	18 12	66 3	64 55	63 42	62 27	61 2	59 32	58 0	56 18	54 34	52 45	50 47	48 39	46 47	44 37	42 29	40 3
	20 10	68 26	67 15	65 59	64 41	63 13	61 41	60 6	58 20	56 32	54 39	52 37	50 54	48 28	46 13	43 51	42 29
	22 8	70 26	69 14	67 56	66 35	65 5	63 29	61 51	60 2	58 11	56 16	54 10	52 3	49 54	47 36	45 9	42 40
	24 6	71 59	70 45	69 25	68 3	66 31	64 54	63 14	61 23	59 29	57 32	55 21	53 10	50 58	48 36	46 8	43 37
	26 4	73 6	71 51	70 30	69 6	67 32	65 54	64 13	62 19	60 23	58 22	56 11	53 59	51 46	49 22	46 51	44 18
	28 2	73 45	72 30	71 7	69 42	68 8	66 28	64 45	62 50	60 54	58 53	56 40	54 27	52 13	49 48	47 15	44 40
3	0 3 0	74 0	72 44	71 22	69 57	68 22	66 42	64 59	63 4	61 7	59 5	56 53	54 40	52 24	49 58	47 15	44 50
9	9	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
☉ & Δ's Ap. 1. equated.		30°	28°	26°	24°	22°	20°	18°	16°	14°	12°	10°	8°	6°	4°	2°	0°
SIGN 7.																	

SUPPLEMENTAL LUNAR TABLES.

EQUATION of the MEAN to the present ELLIPTIC EQUATION of the MOON'S CENTER.

SIGN 5.

Arg. $\odot$ à D's Apogee, 1. equated.		Argument. Equated ANOMALY of the MOON; Or, D 3. equated — Ap. D 2. equated.															
		0°	2°	4°	6°	8°	10°	12°	14°	16°	18°	20°	22°	24°	26°	28°	30°
		+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
		I	II	I	II	I	II	I	II	I	II	I	II	I	II	I	II
6	0 12 0	46 2	43 21	40 38	37 48	34 53	31 55	28 55	25 48	22 41	19 31	16 19	13 6	9 51	6 35	3 17	0
	2 28	45 53	43 13	40 31	37 41	34 46	31 50	28 50	25 43	22 36	19 27	16 16	13 4	9 49	6 32	3 17	0
	4 26	45 37	42 58	40 16	37 27	34 33	31 37	28 39	25 33	22 28	19 21	16 10	12 59	9 46	6 30	3 15	0
	6 24	45 12	42 34	39 54	37 6	34 14	31 20	28 23	25 19	22 16	19 9	16 1	12 51	9 40	6 27	3 14	0
	8 22	44 32	41 56	39 18	36 33	33 44	30 52	27 57	24 57	21 57	18 52	15 57	12 40	9 31	6 21	3 11	0
	10 20	43 42	41 8	38 33	35 53	33 6	30 17	27 26	24 27	21 31	18 31	15 29	12 26	9 21	6 13	3 7	0
12	18	42 41	40 12	37 40	35 2	32 20	29 35	26 49	23 55	21 2	18 5	15 8	12 9	9 8	6 5	3 3	0
14	16	41 30	39 4	36 37	34 3	31 25	28 46	26 4	23 15	20 26	17 35	14 42	11 49	8 52	5 54	2 57	0
16	14	40 7	37 45	35 23	32 57	30 23	27 49	25 11	22 26	19 44	17 0	14 12	11 24	8 34	5 42	2 51	0
18	12	38 39	36 24	34 7	31 43	29 15	26 46	24 16	21 38	19 2	16 24	13 42	11 0	8 16	5 31	2 46	0
20	10	36 54	34 45	32 34	30 18	27 57	25 35	23 11	20 40	18 10	15 38	13 4	10 30	7 54	5 15	2 38	0
22	8	35 3	33 0	30 56	28 45	26 32	24 17	22 0	19 38	17 16	14 51	12 25	9 59	7 31	4 59	2 29	0
24	6	33 7	31 10	29 12	27 10	25 3	22 56	20 47	18 32	16 18	14 2	11 44	9 26	7 5	4 42	2 21	0
26	4	30 54	29 5	27 15	25 20	23 22	21 24	19 24	17 18	15 13	13 6	10 57	8 47	6 36	4 22	2 12	0
28	2	28 36	26 56	25 15	23 28	21 39	19 49	17 57	16 1	14 5	12 8	10 46	8 8	6 7	4 4	2 2	0
1 0	5 0	26 10	24 37	23 5	21 28	19 47	18 6	16 25	14 39	12 53	11 6	9 16	7 27	5 36	3 43	1 51	0
7	2 11 28	23 38	22 15	20 51	19 23	17 52	16 21	14 49	13 13	11 39	10 1	8 22	6 43	5 3	3 21	1 41	0
	4 26	21 10	19 47	18 33	17 14	15 53	14 32	13 10	11 46	10 22	8 55	7 26	5 58	4 29	2 59	1 30	0
	6 24	18 13	17 9	16 6	14 59	13 47	12 36	12 26	10 12	8 59	7 44	6 27	5 10	3 53	2 35	1 18	0
	8 22	15 22	14 28	13 35	12 37	11 38	10 39	9 39	8 36	7 34	6 31	5 26	4 22	3 17	2 11	1 6	0
	10 20	12 25	11 42	10 59	10 11	9 24	8 36	7 48	6 57	6 8	5 17	4 24	3 32	2 39	1 46	0 54	0
12	18	9 25	8 52	8 19	7 43	7 8	6 33	5 55	5 15	4 39	4 0	3 10	2 41	2 0	1 21	0 41	0
14	16	6 21	5 58	5 37	5 13	4 49	4 25	4 0	3 33	3 8	2 42	2 15	1 49	1 20	0 54	0 28	0
16	14	3 14	3 3	2 53	2 40	2 28	2 16	2 4	1 49	1 36	1 23	1 9	0 56	0 41	0 28	0 15	0
18	12	0 4	0 5	0 6	0 4	0 5	0 5	0 4	0 3	0 4	0 3	0 2	0 0	0 0	0 0	0 1	0
		±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	
20	10	3 6	2 55	2 41	2 30	2 20	2 8	1 56	1 46	1 32	1 18	1 5	0 53	0 42	0 28	0 13	0
22	8	6 16	5 54	5 29	5 6	4 43	4 19	3 55	3 32	3 5	2 39	2 14	1 48	1 23	0 56	0 27	0
24	6	9 25	8 51	8 15	7 42	7 7	6 31	5 53	5 18	4 39	3 59	3 21	2 41	2 3	1 22	0 41	0
26	4	12 33	11 48	11 0	10 16	9 30	8 41	7 51	7 2	6 11	5 20	4 27	3 33	2 42	1 49	0 54	0
28	2	15 37	14 42	13 44	12 48	11 49	10 48	9 46	8 45	7 42	6 39	5 32	4 26	3 21	2 14	1 7	0
2 0	4 0	18 39	17 32	16 23	15 17	14 6	12 53	11 39	10 25	9 11	7 58	6 37	5 17	4 0	2 40	1 19	0
8	2 10 28	21 55	20 18	18 58	17 42	16 19	14 55	13 29	12 4	10 38	9 13	7 40	6 7	4 37	3 5	1 32	0
	4 26	24 26	22 58	21 28	20 1	18 28	16 52	15 15	13 39	12 1	10 25	8 40	6 56	5 13	3 29	1 44	0
	6 24	27 8	25 31	23 52	22 15	20 31	18 45	16 58	15 10	13 21	11 32	9 37	7 41	5 47	3 53	1 57	0
	8 22	29 42	27 57	26 8	24 21	22 28	20 32	18 36	16 38	14 37	12 37	10 30	8 24	6 20	4 15	2 7	0
	10 20	32 9	30 15	28 17	26 21	24 18	22 14	20 8	18 0	15 49	13 39	11 22	9 6	6 52	4 35	2 17	0
12	18	34 25	32 23	30 18	28 14	26 1	23 47	21 33	19 16	16 56	14 36	12 11	9 45	7 21	4 55	2 27	0
14	16	36 31	34 22	32 8	29 55	27 36	25 14	22 51	20 25	17 57	15 28	12 55	10 21	7 48	5 13	2 36	0
16	14	38 22	36 7	33 47	31 28	29 1	26 32	24 2	21 29	18 53	16 16	13 34	10 52	8 11	5 28	2 44	0
18	12	40 3	37 42	35 15	32 49	30 16	27 40	25 4	22 24	19 41	16 57	14 9	11 20	8 33	5 42	2 51	0
20	10	42 29	39 3	36 32	34 0	31 21	28 40	25 28	23 13	20 24	17 33	14 39	11 44	8 51	5 54	2 57	0
22	8	42 40	40 11	37 36	35 0	32 17	29 32	26 45	23 55	21 1	18 6	15 7	12 6	9 5	6 4	3 1	0
24	6	43 37	41 4	38 25	35 46	32 59	30 10	27 19	24 25	21 26	18 26	15 23	12 19	9 19	6 13	3 7	0
26	4	44 18	41 41	39 0	36 18	33 29	30 37	27 43	24 47	21 45	18 42	15 36	12 30	9 27	6 19	3 9	0
28	2	44 40	42 3	39 21	36 38	33 47	30 53	27 58	25 0	21 57	18 53	15 46	12 38	9 32	6 23	3 11	0
3 0	3 0	44 50	42 12	39 29	36 45	33 53	30 59	28 4	25 5	22 2	18 57	15 50	12 41	9 34	6 23	3 11	0
9	9	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
$\odot$ à D's Ap. 1. equated.		30°	28°	26°	24°	22°	20°	18°	16°	14°	12°	10°	8°	6°	4°	2°	0°

SIGN 6.

N. B. Whether you solve the Central Equation by this Method, or former one of Eccentricities, you must go back for the other Equations, according to the Newtonian Theory, and its Improvements.

SUPPLEMENTAL

SUPPLEMENTAL LUNAR TABLES.

THE foregoing EQUATION of the MEAN to the present ELLIPTIC EQUATION is the *Difference* between the mean and present ELLIPTIC EQUATION, constructed from the different Eccentricities of the Lunar Orbit, correspondent to the respective Distances of the Sun & Moon's Apogee and equated Anomalies of the Moon.

USE of the foregoing ELLIPTIC EQUATION.

EXAMPLE I. To find the Moon's Central or Elliptic Equation, answerable to $2^{\circ} 8'$ Sun & Moon's Apogee, and $1^{\circ} 11'$ Moon's equated Anomaly?

Eq. An. $0^{\circ} 1'$
By Tab. p. 66. Agt. $1^{\circ} 11'$. . M. Eq. Moon's Center — $3^{\circ} 55' 50''$
 $\odot \Delta \text{ Ap.}$ Eq. An.
By Tab. p. 91. Agt. $2^{\circ} 8'$ under $1^{\circ} 11'$. . Equation — $31^{\circ} 40'$

See p. 83, 84. Equation of Moon's Center required . . — $3^{\circ} 24' 10''$

$\odot \Delta \text{ Ap.}$	Undr. An.	Equation.	Dif. Incr.
Agt. $2^{\circ} 8'$	$1^{\circ} 10'$	— $30' 58''$	
	12	— $32^{\circ} 41'$	+ $1^{\circ} 23''$ for 2° Incr. An.

+ $42'$ for 1° Incr. An.
— $31^{\circ} 40''$ Equation, as above.

EXAMPLE II. To find the Equation of the Moon's Center, correspondent to $1^{\circ} 15' 20''$ Sun & Moon's Apogee, and $3^{\circ} 5' 44''$ equated Anomaly?

Eq. An. $0^{\circ} 1'$
By Tab. p. 67. Agt. $3^{\circ} 5' 44''$. . M. Eq. Moon's Center — $6^{\circ} 18' 26''$
 $\odot \Delta \text{ Ap.}$ Eq. An.
By Tab. p. 93. Agt. $1^{\circ} 15' 20''$, under $3^{\circ} 5' 44''$. . Equⁿ. + $7^{\circ} 43'$

See p. 85. . . Present Equation of Moon's Center required — $6^{\circ} 26' 9''$

$\odot \Delta \text{ Ap.}$	Undr. An.	Equation.	Undr. An.	Equation.
For $1^{\circ} 14'$		+ $11' 18''$	$3^{\circ} 6'$	+ $11' 16''$
Agt. 16	$3^{\circ} 4'$	+ $5^{\circ} 49'$	$3^{\circ} 6'$	+ $5^{\circ} 55'$

+ 2° Dif. — $5^{\circ} 29'$ De. Dif. — $5^{\circ} 21'$ Decr.
Half . . . 1° — $2^{\circ} 44'$ — $3^{\circ} 38''$ — $2^{\circ} 40'$ — $3^{\circ} 33''$
 $20' = \frac{1}{2}$ of 1° . . . $20'$ — $54'$ fr. 1st — $53'$ fr. 1st.

$\odot \Delta \text{ Ap.}$ $1^{\circ} 15' 20''$ + $7^{\circ} 40'$ + $7^{\circ} 43'$
given 7 43

An. + 2° + 3 Increase Eq.
connect $\frac{1}{2}$ of 1° $1\frac{1}{2}$ } + $2^{\circ} \frac{1}{2}$ or 3 connect with $7^{\circ} 40''$
with above $\frac{1}{2}$ of 1° 1 } [above.]

Given An. $3^{\circ} 5' 44''$ + $7^{\circ} 43''$, Equation as above.

EXAMPLE III. To find the Central Equation for $11^{\circ} 22' 18'' 38''$ Sun & Moon's Ap. and $4^{\circ} 21' 27' 32''$ Moon's An.?

By Tab. $\odot \Delta \text{ Ap.}$ Undr. An. Equation. Undr. An. Equation.
p. 94. $11^{\circ} 22'$ $4^{\circ} 20'$ + $56' 15''$ $4^{\circ} 22'$ + $54' 5''$
Agt. 24 + $57^{\circ} 7'$ + $54^{\circ} 54'$

	Dif. + $52'$ Incr.	Dif. + $49'$ Incr.
+ 2°		
$\frac{1}{2}$. . . 1°	26	24
	+ 8	+ 7
$\odot \Delta \text{ Ap.}$ 11 22 18 38	56 23	54 12
given.	54 12	

An. + 2° — $2^{\circ} 11'$ Decrease Eq.
connect $\frac{1}{2}$ of 1° — $1^{\circ} 5'$ — $1^{\circ} 35''$ connect with $56' 23''$
with above $\frac{1}{2}$ of 1° — $30'$ } [above.]

Given An. $4^{\circ} 21' 27' 32''$ + $54' 48''$ Equation required.

$\Delta \text{ An.}$ $0^{\circ} 1'$
By Tab. p. 68. $4^{\circ} 21' 27' 32''$ give the M. Eq. Δ 's Center — $4^{\circ} 9' 0''$
Equation from above + $54^{\circ} 48'$

Present elliptic or central Equation required — $5^{\circ} 3' 48''$

N. B. In the above EASY METHOD, the taking out the elliptic Equation in exact Proportion for 2° either above or under the present Deg. of $\odot \Delta \text{ Ap.}$ will cause an Error sometimes of a few Seconds (not exceeding 8 or 10) because the Equat. for Anom. is not proportional to the Change of Sun & Moon's Apogee; but to the Change of Eccentricity, now rejected by some Astronomers, for easier Computation. The Remedy for which Error (of a few Seconds) is by a short Interpolation.

To find the true from the mean Anomaly in any elliptic Orbit, universally and correctly?

Let E = Arch of Excentric Anomaly, x = its natural Sine (Rad. = 1) e = Eccentricity of the Orbit (Transverse = 1) and M = Arch of mean Anomaly.

Then, by the Property of the Ellipsis and a Planet's Motion in its Orbit, $E + x = M$. And Radius being = Arch of $57^{\circ}, 295779$ Deg. $E^{\circ} + x \times 57^{\circ}, 295779 = M^{\circ}$ (where $x \times 57^{\circ}, 295779 =$ Dif. Degrees between the excentric and mean Anomalies.)

To find the Excentric Anomaly?

You may readily determine it by the Pen and a Table of natural Sines from the above Equation; assuming two different excentric Anomalies adding the respective Differences, and proportioning the Errors above or under the Degrees of mean Anomaly, so as to assign a near excentric Anomaly, which may be correctly determined by a second Operation.

Or, to solve the above Equation, you may proceed thus.

1^o. Say, As Aph. Dist. : Perib. Dist. :: Tang. $\frac{1}{2}$ M. Anom. : Tang. of an Arch (nearly $\frac{1}{2}$ true Anom.) which added to $\frac{1}{2}$ M. Anomaly will be a first excentric Anomaly; which, in Orbits not very excentric, or where the mean and excentric Anomalies differ not above 3 Degrees, will be sufficiently correct.

2^o. In Orbits much excentric, add the Log. of Eccentricity to the Log. 1.7581227 (of $57^{\circ}, 295779$ the Degrees of Rad.) for a constant Log. for that Orbit; to which add the Log. Sine of the 1st approximated excentric Anomaly, and take the Number of Degrees correspondent, and subtract them from the mean Anomaly for a second approximated excentric Anomaly, whose Log. Sine is to be added to the const. Log. as before, and a new Number of Degrees taken, which are to be subtracted from the mean Anomaly, and so proceeding till an excentric Anomaly and the following Difference being added, shall equal the mean Anomaly, when the excentric Anomaly will be correct.

3^o. Then say, as Sq. Root Aph. Dist. : Sq. Root Perib. Dist. :: Tan. $\frac{1}{2}$ correct Excent. An. : Tan. $\frac{1}{2}$ true Anomaly, which doubled is the true required.

Example. The Anomaly of Mercury being 70° , and Eccentricity of that Planet's Orbit 0.20589 (of 1 the Transverse) to find the true Anomaly of Mercury, at that Time?

	Log.
As Aph. Dist. 1.20589	9.9186923 co.
To Per. Dist. 0.79411	9.8998807
So Tan. $35^{\circ} \frac{1}{2}$ M. An.	9.8452268
To Tan. $24^{\circ} 45' 17''$. .	9.6637998

	Log.
Rad. $57^{\circ}, 295779$	1.7581227
Ecc. . . . 0.20589	9.3136353
M. An. 70 00 00	Conf. Log. . . 1.0717580
1. Ex. An. 59 45 17	Log. Sine . . . 9.9364519 +
Dif. — 10 11 27	$10^{\circ}, 19094$ 1.0082099
2. Ex. An. 59 48 33	Log. Sine . . . 9.9366923 +
Dif. — 10 11 47	$10^{\circ}, 19648$ 1.0084503
3. Ex. An. 59 48 13	Log. Sine . . . 9.9366678 +
Dif. — 10 11 45	$10^{\circ}, 19591$ 1.0084258
4. Ex. An. 59 48 15	Log. Sine . . . 9.9366702 +
Dif. — 10 11 45 $\frac{1}{2}$	$10^{\circ}, 19596$ 1.0084280
Cor. Ex. An. 59 48 14 $\frac{1}{2}$	

And the same exactly by Propor. of Inc. of Ex. An. and Error, fr. 1 and 2 Operations.

Now, As $\sqrt{\text{Aph. Dist. 1.20589}}$ 9.9593461 co. } N. B. The above Method exceeds all others for its Simplicity, Universality, and Exactness.
To $\sqrt{\text{Per. Dist. 0.79411}}$ 9.9499404
So $T. \frac{1}{2} \text{ Ex. An. } 29^{\circ} 54' 7'' 15''$ 9.7597224
To $T. \frac{1}{2} \text{ Tr. An. } 25. 1.1. 12$ 9.6690089
Doubled . . . 50. 2.2. 24 True An. req.

Whence $19^{\circ} 57' 58''$ here, Equation, agreeing with Dr. Halley's Tables.

Compare the above universal Method with Mr. Simpson's intricate and limited Correction of Ward's Hypothesis, (See his Misc. Tracts fr. p. 48 to p. 57.) which we shall exhibit hereafter.

N. B. If any Error should happen by the above Method, the next Operation will correct it, proving the Truth at last.

SUPPLEMENTAL LUNAR EQUATIONS.

According to the Construction of Mr. MAYER.

ACCELERATION of the MOON's Mean MOTION: Or Equation of the D's Mean Long.			EQUATION of the NODE and Mean ANOMALY of the MOON.											
			Argument. Mean Anomaly of the SUN.											
			The single for the S, and double of which Equation for Mean Anom. D.											
Yrs b. & fin. Ch.	+	Dif.	0 M. An.	1 Sig.	2 Sig.	3 Sig.	4 Sig.	5 Sig.	6 Sig.	7 Sig.	8 Sig.	9 Sig.	10 Sig.	11 Sig.
B. 800	1 9 48	5 29	0	0 0	11	5 1	9	8 47	6	10 18	0	9 4	5	5 17
Ch. 700	1 4 19	5 15	1	0 11	11	5 10	9	8 53	5	10 18	0	8 59	5	5 8
600	0 59 4	5 1	2	0 22	10	5 19	9	8 58	5	10 18	0	8 53	6	4 58
500	0 54 3	4 48	3	0 32	10	5 28	9	9 3	6	10 18	1	8 47	6	4 48
400	0 49 15	4 35	4	0 43	10	5 37	9	9 9	5	10 17	0	8 41	6	4 38
300	0 44 40	4 21	5	0 53	11	5 46	9	9 14	5	10 17	1	8 35	6	4 28
200	0 40 19	4 8	6	1 4	10	5 55	8	9 19	5	10 16	1	8 29	7	4 18
100	0 36 11	3 55	7	1 14	11	6 3	9	9 23	4	10 15	1	8 22	7	4 8
0	0 32 16	3 41	8	1 25	10	6 12	8	9 27	4	10 14	1	8 16	6	3 58
S. 100	0 28 15	3 28	9	1 35	10	6 20	8	9 31	4	10 13	1	8 10	7	3 48
Ch. 200	0 25 7	3 14	10	1 45	10	6 28	8	9 35	4	10 12	1	8 3	7	3 38
300	0 21 53	3 1	11	1 55	10	6 36	8	9 39	4	10 10	2	7 56	7	3 27
400	0 18 52	2 47	12	2 5	10	6 44	8	9 42	3	10 8	2	7 48	7	3 17
500	0 16 5	2 34	13	2 15	10	6 52	8	9 46	4	10 6	2	7 41	7	3 6
600	0 13 31	2 21	14	2 25	10	7 0	8	9 50	4	10 4	2	7 34	7	2 56
700	0 11 10	2 7	15	2 35	10	7 8	8	9 53	3	10 2	2	7 26	7	2 45
800	0 9 3	1 54	16	2 45	10	7 16	7	9 56	3	9 59	3	7 18	8	2 35
900	0 7 9	1 41	17	2 55	10	7 23	8	9 58	2	9 56	3	7 10	8	2 24
1000	0 5 28	1 27	18	3 5	10	7 31	8	10 1	3	9 53	3	7 2	8	2 13
1100	0 4 1	1 13	19	3 15	10	7 38	7	10 3	2	9 50	3	6 54	8	2 2
1200	0 2 48	1 1	20	3 25	10	7 45	7	10 5	2	9 46	4	6 46	8	1 51
1300	0 1 47	0 47	21	3 35	9	7 52	6	10 7	2	9 43	3	6 37	9	1 40
1400	0 1 0	0 33	22	3 44	10	7 58	7	10 9	2	9 39	4	6 29	9	1 29
1500	0 0 27	0 20	23	3 54	10	8 5	7	10 11	2	9 35	4	6 20	9	1 18
1600	0 0 7	0 5	24	4 4	10	8 12	6	10 13	1	9 31	4	6 12	9	1 7
1650	0 0 2	0 2	25	4 14	9	8 18	6	10 14	1	9 27	4	6 3	9	0 56
1700	0 0 0	0 0	26	4 23	10	8 24	6	10 15	1	9 23	4	5 54	9	0 45
1750	0 0 0	0 0	27	4 33	9	8 30	6	10 16	1	9 18	5	5 45	9	0 34
1800	0 0 7	0 8	28	4 42	10	8 36	6	10 17	1	9 14	4	5 36	9	0 23
1850	0 0 15	0 12	29	4 52	9	8 42	5	10 18	0	9 9	5	5 27	10	0 12
1900	0 0 27	0 15	30	5 1	9	8 47	5	10 18	0	9 4	5	5 17	10	0 0
1950	0 0 42	0 18	OM.	—	D.	—	D.	—	D.	—	D.	—	D.	—
2000	0 1 0	0 18	An.	11 Sig.	10 Sig.	9 Sig.	8 Sig.	7 Sig.	6 Sig.	5 Sig.	4 Sig.	3 Sig.	2 Sig.	1 Sig.

REMARKS on Mr. MAYER's LUNAR TABLES: To which our OWN LUNAR TABLES are reducible in the Mean PLACES and MOTIONS: See Examples farther on.

HE did not presume (he says) to send his Tables of the Lunar Motions into the World, without being thoroughly convinced of their very near, and scarce to be hoped-for nearer Agreement with the Heavens. That briefly to tell us, he says, of more than two hundred Observations of the Moon made not only in this but in the preceding Age, scarce ten were found, from which his Tables differ one Minute and a half; a great many of them come nearer than a single Minute, and not one of them all discovered an Error in his Tables of two Minutes. Which as no Tables heretofore published perform the like, the best of them often erring four or five Minutes, he was the more unwilling his Tables should lie any longer concealed; especially, as the most celebrated Astronomers of almost every Age have ardently wished for a perfect Theory of the Moon (to which his Tables are said nearly to approach) on Account of its singular Use in Navigation.

Nor, indeed, (he says) do they seem worthy of Publication for the Sake of this particular Service, in completing the Theory only, but also on Account of that easy and expeditious Method, by which that particular Service from them may be had, in computing the Longitude and Latitude. For, if you look into other Tables, you will find, on Comparison, these larger Equations so few in Number, and the Rest of them so small, that there is no Need of having Recourse to the troublesome Method of interpolating, nor of making the Computation of the Arguments to Seconds. He has constructed these Tables, he says, with Respect to the Inequalities of Motions, from that famous Theory of the GREAT NEWTON, which that eminent Mathematician Eulerus first elegantly reduced to general analytical Equations. And, in the resolving of these Equations, after some fruitless Essays by other Methods, he has hit (as he says) upon a particular Method, pretty elegant and concise, which it would be too tedious for him to explain at large in his Preface. For which Reason he has concluded only to give such Hints as may conduce to lay open the Origin and Causes of the Inequalities exhibited in the Equation Tables, so far as it can be done without making Use of Calculations.

SUPPLEMENTAL LUNAR EQUATIONS.

According to the Construction of Mr. MAYER.

Argument proper.	I. EQUATION D.						II. EQUAT. D.			III. EQUAT. D.			IV. EQUAT. D.			V. EQUAT. D.			Argument proper.															
	Argument 1.						Argument 2.			Argument 3.			Argument 4.			Argument 5.																		
	Mean Anomaly $\odot$.						2. $\Delta a \odot + A$ g. 1.			2. $\Delta a \odot - \text{Arg. 1.}$			Arg. 2. — M. An. D.			Arg. 3. — M. An. D.																		
	0	1	2	3	4	5	0	1	2	0	1	2	0	1	2	0	1	2																
	+	+	+	+	+	+	—	—	—	—	—	—	+	+	+	+	+	+																
0	0	5	31	9	40	11	20	9	58	5	49	0	0	0	31	0	54	0	0	0	36	1	2	30										
1	0	12	5	41	9	46	11	20	9	52	5	39	0	10	28	0	47	0	10	32	0	54	0	20	56	1	35	0	10	37	1	3	29	
2	0	24	5	51	9	52	11	20	9	46	5	28	0	20	29	0	48	0	20	33	0	55	0	40	57	1	36	0	20	38	1	3	28	
3	0	35	6	1	9	58	11	20	9	39	5	17	0	30	29	0	48	0	30	34	0	55	0	60	59	1	37	0	40	39	1	4	27	
4	0	47	6	11	10	3	11	19	9	33	5	6	0	40	30	0	49	0	40	35	0	56	0	81	0	1	38	0	50	40	1	4	26	
5	0	58	6	21	10	9	11	19	9	26	4	55	0	50	31	0	49	0	60	36	0	56	0	101	1	2	39	0	60	41	1	5	25	
6	1	10	6	30	10	14	11	18	9	19	4	44	0	60	31	0	50	0	70	37	0	56	0	121	1	3	39	0	80	42	1	5	24	
7	1	21	6	40	10	19	11	17	9	12	4	33	0	70	32	0	50	0	80	38	0	57	0	131	1	5	40	0	90	43	1	6	23	
8	1	33	6	49	0	24	11	16	9	5	4	22	0	80	33	0	50	0	90	38	0	57	0	151	1	6	41	0	100	44	1	6	22	
9	1	44	6	58	10	29	11	15	8	5	4	11	0	90	34	0	51	0	100	39	0	58	0	171	1	8	41	0	120	45	1	7	21	
10	1	55	7	7	10	33	11	13	8	5	3	59	0	100	34	0	51	0	110	40	0	58	0	191	1	9	42	0	130	46	1	7	20	
11	2	7	7	16	10	37	11	11	8	4	3	48	0	110	35	0	51	0	120	41	0	58	0	211	1	10	42	0	140	47	1	8	19	
12	2	18	7	25	10	41	11	9	8	3	3	36	0	120	36	0	52	0	130	42	0	59	0	231	1	12	43	0	160	48	1	8	18	
13	2	29	7	34	10	45	11	7	8	2	3	25	0	130	37	0	52	0	140	42	0	59	0	241	1	13	44	0	170	49	1	9	17	
14	2	40	7	42	10	49	11	5	8	1	3	13	0	140	37	0	52	0	150	43	1	0	0	261	1	15	44	0	180	50	1	9	16	
15	2	51	7	51	10	52	11	2	8	1	3	1	0	150	38	0	52	0	160	44	1	0	0	281	1	16	45	0	190	51	1	10	15	
16	3	2	7	59	10	55	10	59	8	3	2	50	0	160	39	0	52	0	170	45	1	0	0	301	1	18	45	0	200	52	1	10	14	
17	3	13	8	7	10	58	10	56	7	5	2	38	0	170	39	0	53	0	180	45	1	0	0	321	1	19	46	0	220	52	1	10	13	
18	3	24	8	15	11	1	10	52	7	4	2	26	0	180	40	0	53	0	190	46	1	1	1	331	1	20	46	0	230	53	1	11	12	
19	3	35	8	23	11	3	10	49	7	3	2	14	0	190	41	0	53	0	200	47	1	1	1	351	1	22	46	0	240	54	1	11	11	
20	3	46	8	31	11	6	10	45	7	2	2	2	0	200	41	0	53	0	210	48	1	1	1	371	1	23	47	0	250	55	1	11	10	
21	3	57	8	39	11	8	10	41	7	1	1	50	0	210	42	0	53	0	220	48	1	1	1	391	1	24	47	0	260	56	1	11	9	
22	4	8	8	46	11	10	10	37	7	9	1	38	0	220	42	0	53	0	230	49	1	1	1	411	1	26	47	0	280	56	1	11	8	
23	4	18	8	54	11	12	10	33	7	0	1	26	0	230	43	0	53	0	240	50	1	1	1	421	1	27	47	0	290	57	1	11	7	
24	4	29	9	1	11	14	10	29	6	5	0	14	0	240	44	0	54	0	250	50	1	2	2	441	1	28	47	0	300	58	1	12	6	
25	4	40	9	8	11	16	10	24	6	4	0	1	0	250	44	0	54	0	260	51	1	2	2	461	1	29	48	0	310	59	1	12	5	
26	4	50	9	15	11	17	10	19	6	3	0	49	0	260	45	0	54	0	270	51	1	2	2	481	1	30	48	0	320	59	1	12	4	
27	5	0	9	21	11	18	10	14	6	2	0	37	0	270	45	0	54	0	280	52	1	2	2	491	1	31	48	0	330	59	1	12	3	
28	5	11	9	28	11	19	10	9	6	9	0	25	0	280	46	0	54	0	290	52	1	2	2	511	1	32	48	0	340	59	1	12	2	
29	5	21	9	34	11	20	10	4	5	5	0	13	0	290	46	0	54	0	300	53	1	2	2	521	1	33	48	0	350	59	1	12	1	
30	5	31	9	40	11	20	9	5	5	4	0	0	0	300	47	0	54	0	310	54	1	2	2	541	1	34	48	0	360	59	1	12	0	
Argument proper.	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Argument proper.	11	10	9	8	7	6	11	10	9	11	10	9	11	10	9	11	10	9	11	10	9	11	10	9	11	10	9	11	10	9	11	10	9	11
Argument proper.	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Argument proper.	5	4	3	2	1	0	5	4	3	5	4	3	5	4	3	5	4	3	5	4	3	5	4	3	5	4	3	5	4	3	5	4	3	5
Argument proper.	Argument as above.						2. $\Delta a \odot + \odot$						2. $\Delta a \odot - \odot$						2. $\Delta a \odot + \odot$ An.						2. $\Delta a \odot - \odot$ An.									
Argument proper.	I. EQUATION D.						II. EQUAT. D.						III. EQUAT. D.						IV. EQUAT. D.						V. EQUAT. D.									

Of Mr. MAYER'S LUNAR EQUATIONS and MOTIONS.

According to Mr. Mayer, the Moon's Motion in Longitude ought to be corrected by XIII Equations; the most considerable of which are the XI, XII, and XIII; on which almost all the Rest depend. The XI, otherwise called the Equation of the Moon's Center, proceeds from the Eccentricity of the Moon's Orbit, and follows the Keplerian Law. Its Quantity is not ascertained by the Theory itself, but must be determined by immediate Observations; and is the only one, he thinks, that may not be affected by the Sun's Action.

The XII. Equation is derived from the Power of the Sun upon the Moon, and the Eccentricity of the Lunar Orbit, jointly, and would vanish, was either the Sun's Action to cease, or the Moon's Orbit to become perfectly circular. This Equation, or rather one like it, Bullialdus, and some others, called the Evection. It strictly follows the Keplerian Law, except that it is variable from a twofold Cause. The first is the unequal Distances of the Sun from the Earth; for when the Sun is in Apogee, the Evection will be less, and when in Perigee, almost as much greater. The second Cause is the mutable Distance of the Moon from the Earth, which at the Moon's Perigee makes the Equation greater, and in its Apogee less. The mean Quantity thereof is indeed that set down in the following Equation Tables; but then that Part which is, for the aforementioned Cause, variable, is restored by the IV, V, X, and also some Part of the XIII. Equation, from whence the Reason of those Equations may be known.

By these two Equations, the XI and XII, the same Inequalities are regulated, which NEWTON, and those who have closely followed him, have explained by the variable Eccentricity of the Moon's Orbit, and the unequal Motion of the Apfides. And though that Method, Mr. Mayer asserts,

SUPPLEMENTAL LUNAR EQUATIONS.

According to the Construction of Mr. MAYER.

VI. EQUAT. D.				VII. EQUAT. D.				VIII. EQ. D.				IX. EQUAT. D.				X. EQUATION D.						Argument proper.											
Argument 6.				Argument 7.				Argument 8.				Argument 9.				Argument 10.																	
2. D à O + M.				2. D à O - M.				M. An. D -				Pl. - Pl. true				D à O - M. Anom. D.																	
An. D.				An. D.				Arg. 1.																									
0	1	2		0	1	2		0	1	2		0	1	2		0	1	2	3	4	5												
+	+	+		+	+	+		+	+	+		+	+	+		+	+	+	±	-	-												
6	7	8		6	7	8		6	7	8		6	7	8																			
-	-	-		-	-	-		-	-	-		+	+	+		+	+	+	±	-	-												
/	//	/	//	/	//	/	//	/	//	/	//	/	//	/	//	/	//	/	//	/	//												
0	0	0	45	1	18	0	0	0	29	0	50	0	0	0	41	0	41	0	3	11	3	22	0	29	2	32	2	42	30				
1	0	20	46	1	19	0	10	30	0	50	0	10	20	0	41	0	40	0	8	3	15	3	19	0	22	2	35	2	39	29			
2	0	30	48	1	19	0	20	31	0	51	0	20	21	0	42	0	39	0	15	3	19	3	15	0	15	2	39	2	36	28			
3	0	50	49	1	20	0	30	31	0	51	0	20	21	0	43	0	38	0	23	3	22	3	11	0	8	2	42	2	32	27			
4	0	60	51	1	20	0	40	32	0	52	0	30	22	0	43	0	37	0	30	3	25	3	7	0	12	45	2	28	26				
5	0	80	52	1	21	0	50	33	0	52	0	40	23	0	44	0	36	0	38	3	28	3	3	0	6	2	48	2	24	25			
6	0	90	53	1	22	0	60	34	0	53	0	40	23	0	44	0	35	0	45	3	31	2	58	0	13	2	51	2	20	24			
7	0	110	55	1	22	0	70	35	0	53	0	50	24	0	45	0	34	0	53	3	33	2	53	0	20	2	53	2	15	23			
8	0	130	56	1	23	0	80	35	0	54	0	60	24	0	45	0	32	0	1	0	36	2	48	0	27	2	55	2	11	22			
9	0	140	57	1	23	0	90	36	0	54	0	60	25	0	46	0	31	0	1	8	38	2	43	0	34	2	57	2	6	21			
10	0	160	58	1	24	0	100	37	0	54	0	70	26	0	46	0	30	0	1	15	3	40	2	38	0	41	2	59	2	1	20		
11	0	170	59	1	25	0	110	38	0	55	0	70	26	0	46	0	29	0	1	22	3	42	2	32	0	48	3	0	1	56	19		
12	0	190	1	1	25	0	120	38	0	55	0	80	27	0	47	0	27	0	1	29	3	43	2	27	0	54	3	1	1	51	18		
13	0	200	1	2	26	0	130	39	0	55	0	90	27	0	47	0	26	0	1	36	3	44	2	21	0	1	13	2	1	46	17		
14	0	220	1	3	26	0	140	40	0	56	0	90	28	0	47	0	25	0	1	43	3	44	2	16	0	1	8	3	1	40	16		
15	0	230	1	4	27	0	150	41	0	56	0	100	28	0	47	0	23	0	1	50	3	45	2	10	0	1	14	3	4	1	35	15	
16	0	250	1	5	27	0	160	41	0	56	0	110	29	0	47	0	22	0	1	56	3	45	2	4	0	1	20	3	4	1	29	14	
17	0	260	1	6	28	0	170	42	0	56	0	110	29	0	47	0	20	0	2	3	3	44	1	58	0	1	26	3	3	1	23	13	
18	0	280	1	7	28	0	180	43	0	57	0	120	30	0	47	0	19	0	2	9	3	44	1	51	0	1	32	3	3	1	17	12	
19	0	290	1	8	28	0	190	43	0	57	0	130	30	0	46	0	17	0	2	15	3	43	1	45	0	1	38	3	2	1	11	11	
20	0	310	1	9	29	0	200	44	0	57	0	130	31	0	46	0	16	0	2	21	3	43	1	38	0	1	44	3	2	1	5	10	
21	0	320	1	10	29	0	210	45	0	57	0	140	31	0	46	0	14	0	2	26	3	42	1	32	0	1	49	3	1	0	59	9	
22	0	340	1	11	29	0	220	45	0	57	0	150	32	0	45	0	13	0	2	32	3	41	1	25	0	1	55	3	0	0	53	8	
23	0	350	1	12	29	0	230	46	0	58	0	150	32	0	45	0	11	0	2	37	3	39	1	18	0	2	0	2	59	0	46	7	
24	0	370	1	13	29	0	240	46	0	58	0	160	33	0	44	0	10	0	2	43	3	37	1	11	0	2	5	2	57	0	40	6	
25	0	380	1	14	30	0	250	47	0	58	0	170	33	0	44	0	8	0	2	48	3	35	1	4	0	2	10	2	55	0	33	5	
26	0	390	1	15	30	0	260	47	0	58	0	170	34	0	43	0	6	0	2	53	3	33	0	57	0	2	15	2	53	0	27	4	
27	0	410	1	16	30	0	270	48	0	58	0	180	34	0	43	0	5	0	2	58	3	30	0	50	0	2	19	2	50	0	20	3	
28	0	420	1	16	30	0	270	49	0	58	0	190	34	0	42	0	3	0	3	2	3	28	0	43	0	2	24	2	48	0	14	2	
29	0	440	1	17	30	0	280	49	0	58	0	190	35	0	41	0	2	0	3	7	3	25	0	36	0	2	28	2	45	0	7	1	
30	0	450	1	18	30	0	290	50	0	58	0	200	35	0	41	0	0	0	3	11	3	22	0	29	0	2	32	2	42	0	0	0	
Argument proper.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	11	10	9	11	10	9	11	10	9	11	10	9	10	9	9	11	10	9	11	10	9	8	7	6	5	4	3	2	1	0	0	0	
	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	5	4	3	5	4	3	5	4	3	5	4	3	5	4	3	5	4	3	5	4	3	2	1	0	0	0	0	0	0	0	0	0	0
	Arg. As above.			As above.			M. An. D - M.			As above.			As above.						As above.														
	VI. EQUAT. D.			VII. EQUAT. D.			VIII. EQ. D.			IX. EQUAT. D.			X. EQUATION D.																				

Of Mr. MAYER'S LUNAR EQUATIONS and MOTIONS.

afferts, exactly corresponds with the Theory, yet he thinks it is a little the more difficult, and not so well adapted to a tabular Computation. For which Reason, he says, the Lunar Astronomy is highly indebted to the celebrated Eulerus, who first elegantly substituted the constant Equation of the Center together with that which is called the Evection, in the Room of the variable Eccentricity, and so adorned the Lunar Theory, which was otherwise extremely intricate, with a neat and elegant Contraction: omitting 2d large Equation D's Apogee.

The XIII. EQUATION was first discovered by Tycho, who gave it the Name of the VARIATION; Bullialdus, and others, call it the REFLECTION. The Theory itself plainly enough determines its Quantity, arising from the Action of the Sun singly, so that it would remain, if the Orbits both of the Moon and Earth had no Eccentricity. Some Part of it depends on the Quantity of the Sun's Parallax, which Mr. Mayer assumed = 11" 1/2, which he thinks not wide of Truth. It is here to be noted, that the mean Quantity of this Equation (which by Reason of the different Distances of the Sun and Moon from the Earth making it variable like the foregoing) is exactly equal to half the Evection; which, though not absolutely necessary, so far as may be discovered from the Theory, yet he thinks should not be wholly ascribed to Chance. But the Changes it undergoes, from the different Distances of the Sun, are contained in the II. and III. Equations; and those which happen on Account of the Inequality of the Moon's Distance from the Earth, are accounted for in the VI. Equation, and in some Part of the XII.

The first Equation of the Moon's Longitude is derived from the Sun's Action and also from the Eccentricity of the Earth's Orbit. For if these were both destroyed, the Equation would vanish with them. The Nature of this Equation is such, that it accelerates the Moon's Motion

SUPPLEMENTAL LUNAR EQUATIONS.

According to the Construction of Mr. MAYER.

XI.

MEAN EQUATION of the MOON'S CENTER.

Argument II. MOON'S Mean ANOMALY, corrected by the 11 preceding EQUATIONS.

Decor. An.	Sig. 0.	Sig. 1.	Sig. 2.	Sig. 3.	Sig. 4.	Sig. 5.	Decor. An.
—	Dif.	—	Dif.	—	Dif.	—	Dif.
0	0 0 0	2 58 33	5 16 28	6 17 50	5 39 0	3 21 4	30
1	0 6 11	3 4 1	5 19 55	6 18 14	5 35 53	3 15 4	29
2	0 12 22	3 9 26	5 23 17	6 18 31	5 32 40	3 9 1	28
3	0 18 32	3 14 48	5 26 34	6 18 41	5 29 20	3 2 54	27
4	0 24 42	3 20 7	5 29 46	6 18 44	5 25 53	2 56 43	26
5	0 30 52	3 25 24	5 32 53	6 18 40	5 22 21	2 50 28	25
6	0 37 2	3 30 38	5 35 54	6 18 29	5 18 42	2 44 9	24
7	0 43 11	3 35 48	5 38 49	6 18 11	5 14 56	2 37 47	23
8	0 49 19	3 40 55	5 41 38	6 17 46	5 11 4	2 31 21	22
9	0 55 27	3 45 58	5 44 21	6 17 14	5 7 5	2 24 51	21
10	1 1 54	3 50 57	5 46 58	6 16 36	5 2 59	2 18 18	20
11	1 7 40	3 55 53	5 49 29	6 15 50	4 58 47	2 11 42	19
12	1 13 45	4 0 45	5 51 55	6 14 57	4 54 29	2 5 4	18
13	1 19 48	4 5 34	5 54 15	6 13 57	4 50 5	1 58 23	17
14	1 25 50	4 10 19	5 56 29	6 12 50	4 45 36	1 51 39	16
15	1 31 52	4 15 0	5 58 37	6 11 36	4 41 1	1 44 52	15
16	1 37 52	4 19 37	6 0 39	6 10 15	4 36 20	1 38 3	14
17	1 43 50	4 24 10	6 2 34	6 8 47	4 31 33	1 31 12	13
18	1 49 47	4 28 38	6 4 23	6 7 12	4 26 39	1 24 18	12
19	1 55 42	4 33 2	6 6 6	6 5 29	4 21 40	1 17 23	11
20	2 1 35	4 37 22	6 7 43	6 3 39	4 16 24	1 10 27	10
21	2 7 26	4 41 38	6 9 13	6 1 42	4 11 23	1 3 29	9
22	2 13 15	4 45 49	6 10 37	5 59 38	4 6 7	0 56 30	8
23	2 19 2	4 49 55	6 11 54	5 57 28	4 0 46	0 49 29	7
24	2 24 48	4 53 57	6 13 5	5 55 11	3 55 21	0 42 27	6
25	2 30 32	4 57 55	6 14 10	5 52 47	3 49 51	0 35 23	5
26	2 36 14	5 1 48	6 15 8	5 50 16	3 44 16	0 28 19	4
27	2 41 53	5 5 36	6 15 59	5 47 38	3 38 36	0 21 15	3
28	2 47 29	5 9 19	6 16 43	5 44 53	3 32 50	0 14 10	2
29	2 53 2	5 12 56	6 17 20	5 42 0	3 27 0	0 7 5	1
30	2 58 33	5 16 28	6 17 50	5 39 0	3 21 4	0 0 0	0
Decor. An.	+ Dif.	+ Dif.	+ Dif.	+ Dif.	+ Dif.	+ Dif.	Decor. An.
	Sig. 11.	Sig. 10.	Sig. 9.	Sig. 8.	Sig. 7.	Sig. 6.	

Of Mr. MAYER'S LUNAR EQUATIONS and MOTIONS.

Motion when the Sun is in *Apogee*, and retards it, when the Sun is in *Perigee*. It has passed under various Names both in this and the last Age; but its true Quantity, which the *Theory* leaves in some Measure doubtful, Mr. Mayer has restored from a great Number of Observations. And as this Equation is variable (which no one, before him, seems to have taken due Notice of) by Reason of the variable Distances of the $\mathcal{D}$ from the Earth, he therefore found it necessary to add the VIII. EQUATION, though that Equation does not wholly account for the Change. The other Part is excellently restored by that Equation, which is to be applied to the *mean Anomaly*, and which also contains the Inequality of the Motion of the *Apogee*, so far as that varies on Account of the greater or less Distance of the Sun from the Earth.

The VII. and IX. EQUATIONS cannot so easily be explained in Words without a *Calculus*: But it may be sufficient to observe, he says, that they arise from the *Inclination* of the Moon's Orbit to the *Ecliptic*, which is the Cause, that the Sun sometimes acts obliquely on the Moon and so affects its Motion in *Longitude* a little: Add to this, the *Eccentricity* of the Moon's Orbit, which makes the Sun's Action greater when the Moon is near its *Apogee*, and less, when in its *Perigee*. There are, 'tis true, some other *Inequalities* affecting the Moon's Motion in *Longitude*, which he has not mentioned. But then some of them are united with the Equations above mentioned; others are preserved by the Equation he applied to the *mean Anomaly*; and others there are, which he has entirely neglected. The Reduction Equation to the *Ecliptic*, or the XIV. EQUATION of *Longitude*, he has fixed as invariable, though the *Inclination* of the Orbit, whence it arises, is not always the same. But if the VARIATION of XIII. Equation be somewhat diminished, as has been done, and requires Notice should be taken thereof, it amounts to the same, and is more commodious than if this Reduction Equation had been made variable. This he particularly mentions, that no Body may in this Place absurdly seek out Errors, of which he may think the Tables themselves not yet clear. He has formed two Tables to determine the Moon's Latitude, one containing its *mean Quantity*, independent of the Sun's Action; the other containing the *Variations* of Latitude arising therefrom.

If this second Latitude Table, as it may be named, be made variable in the reciprocal triplicate Ratio of the Sun's Distance from the Earth, it will, perhaps, come somewhat nearer the Truth. But still there are other Irregularities in the $\mathcal{D}$'s Latitude, not yet discovered, that Correction of them, as well as of all those yet unknown in the *Theory*, is useless, till we gain more accurate Observations of the Latitude, which may be now more carefully performed, he thinks, by Means of the more exact *Parallax* of the Moon ascertained at the End of these Equation Tables.

SUPPLEMENTAL

SUPPLEMENTAL LUNAR EQUATIONS.

According to the Construction of Mr. MAYER.

XII. EQUATION of the MOON'S EVECTION, or Second Equation of the MOON'S CENTER.

Arg. 12. Or 2. Δ a cor^d. by D's Pl. at 10 Eq. (not at 11th, of which this Eq. is 2d. Part) — M. An. cor^d. or Arg. 11th.

evection.	Sig. 0.		Sig. 1.		Sig. 2.		Sig. 3.		Sig. 4.		Sig. 5.		evection.
	—	Dif.	—	Dif.	—	Dif.	—	Dif.	—	Dif.	—	Dif.	
0	0 0 0		0 39 50	1 12	1 9 22	0 42	1 20 42	0 1	1 10 24	0 42	0 40 52	1 14	30
1	0 1 24	1 23	0 41 2	1 12	1 10 4	0 41	1 20 43	0 1	1 9 42	0 44	0 39 38	1 15	29
2	0 2 47	1 23	0 42 14	1 11	1 10 45	0 40	1 20 42	0 3	1 8 58	0 45	0 38 23	1 16	28
3	0 4 10	1 23	0 43 25	1 10	1 11 25	0 39	1 20 39	0 4	1 8 13	0 46	0 37 7	1 17	27
4	0 5 33	1 23	0 44 35	1 9	1 12 4	0 37	1 20 35	0 5	1 7 27	0 47	0 35 50	1 17	26
5	0 6 56	1 23	0 45 44	1 8	1 12 41	0 36	1 20 30	0 7	1 6 40	0 48	0 34 33	1 18	25
6	0 8 19	1 23	0 46 52	1 7	1 13 17	0 34	1 20 23	0 8	1 5 52	0 50	0 33 15	1 18	24
7	0 9 42	1 23	0 47 59	1 7	1 13 51	0 33	1 20 15	0 10	1 5 2	0 51	0 31 57	1 19	23
8	0 11 5	1 22	0 49 6	1 6	1 14 24	0 32	1 20 5	0 11	1 4 11	0 53	0 30 38	1 19	22
9	0 12 27	1 22	0 50 12	1 5	1 14 56	0 31	1 19 54	0 13	1 3 18	0 54	0 29 19	1 20	21
10	0 13 49	1 22	0 51 17	1 4	1 15 27	0 29	1 19 41	0 14	1 2 24	0 55	0 27 59	1 21	20
11	0 15 11	1 22	0 52 21	1 3	1 15 50	0 28	1 19 27	0 16	1 1 29	0 56	0 26 38	1 21	19
12	0 16 33	1 21	0 53 24	1 2	1 16 24	0 26	1 19 11	0 17	1 0 33	0 57	0 25 17	1 22	18
13	0 17 54	1 21	0 54 26	1 1	1 16 50	0 25	1 18 54	0 19	0 59 30	0 58	0 23 55	1 22	17
14	0 19 15	1 21	0 55 27	1 1	1 17 15	0 24	1 18 35	0 20	0 58 38	0 59	0 22 33	1 22	16
15	0 20 36	1 21	0 56 28	1 0	1 17 39	0 22	1 18 15	0 22	0 57 39	1 0	0 21 11	1 22	15
16	0 21 56	1 20	0 57 28	0 58	1 18 1	0 21	1 17 53	0 23	0 56 39	1 1	0 19 48	1 23	14
17	0 23 16	1 19	0 58 26	0 57	1 18 22	0 19	1 17 30	0 24	0 55 38	1 2	0 18 25	1 23	13
18	0 24 35	1 19	0 59 23	0 56	1 18 41	0 18	1 17 6	0 26	0 54 36	1 4	0 17 1	1 24	12
19	0 25 54	1 19	1 0 19	0 55	1 18 59	0 17	1 16 40	0 27	0 53 32	1 5	0 15 37	1 24	11
20	0 27 13	1 18	1 1 14	0 54	1 19 16	0 15	1 16 13	0 29	0 52 27	1 6	0 14 13	1 24	10
21	0 28 31	1 18	1 2 8	0 53	1 19 31	0 14	1 15 44	0 30	0 51 21	1 7	0 12 48	1 25	9
22	0 29 49	1 17	1 3 1	0 52	1 19 45	0 12	1 15 14	0 31	0 50 15	1 8	0 11 23	1 25	8
23	0 31 6	1 17	1 3 53	0 51	1 19 57	0 11	1 14 43	0 33	0 49 8	1 9	0 9 58	1 25	7
24	0 32 23	1 16	1 4 41	0 49	1 20 8	0 9	1 14 10	0 34	0 48 0	1 10	0 8 33	1 25	6
25	0 33 39	1 15	1 5 33	0 48	1 20 17	0 8	1 13 35	0 36	0 46 51	1 11	0 7 8	1 25	5
26	0 34 54	1 15	1 6 21	0 47	1 20 25	0 6	1 13 0	0 37	0 45 41	1 12	0 5 43	1 25	4
27	0 36 9	1 14	1 7 8	0 46	1 20 31	0 5	1 12 23	0 38	0 44 30	1 13	0 4 18	1 26	3
28	0 37 23	1 14	1 7 54	0 45	1 20 36	0 4	1 11 45	0 40	0 43 18	1 13	0 2 52	1 26	2
29	0 38 37	1 13	1 8 39	0 43	1 20 40	0 2	1 11 5	0 41	0 42 5	1 13	0 1 26	1 26	1
30	0 39 50		1 9 22		1 20 42		1 10 24		0 40 52		0 0 0	1 26	0
evection.	+	Dif.	+	Dif.	+	Dif.	+	Dif.	+	Dif.	+	Dif.	evection.
	Sig. 11.		Sig. 10.		Sig. 9.		Sig. 8.		Sig. 7.		Sig. 6.		

Of Mr. MAYER'S LUNAR EQUATIONS and MOTIONS.

He, not being furnished with such accurate Observations, could not promise that the Latitude found by his Tables will come nearer than one Minute: but, in Eclipses, the Error, he asserts, will not amount to 20". And how plain and easy the Method of computing the Latitude is by Tables thus contrived, (like our own for Facility) and how different from all Methods heretofore laid down, is too evident to require making Mention.

In settling the mean Motion of the Moon he has endeavoured to arrive at some Degree of Certainty, as well as Agreement with the Observations of ancient Times. He has examined the earliest Observations of the Lunar Eclipses made by the Babylonians, as well as Hipparchus, and Ptolemy; though they are so gross and incorrect, that he in vain attempted to bring them tolerably near their Times, by the Tables; but this will not appear strange to any one, who considers, that the Ancients, in observing the Times of these Phenomena, did not much regard a Quarter or Half an Hour. Besides which there is great Reason to suspect, that Ptolemy, from whom we have the Accounts of these Eclipses, has too boldly altered the Times of some, and adapted them to the Numbers of his own Hypotheses. Instances of which are produced by Ismael Bullialdus, in the Astron. Philol. B. III. C. 7. For this Reason no Body will impute it as a Fault in Mr. Mayer, if his Tables should be found in the Calculation of one or two of those Eclipses, not to come within half an Hour. But, notwithstanding this Difference, arising either from the Negligence of the Antients, or the Unfaithfulness of PTOLEMY, these Observations have concurrently shewn, that the Moon's Motion of old was sensibly slower than it is discovered to be in our Age. Halley, and some other Astronomers, have taken Notice of this Acceleration in the Moon's Motion; but the Quantity thereof has scarcely been well ascertained by any one. In Order there ore to determine it the more accurately, Mr. Mayer has, with great Pains and Assiduity, examined the Observations made between those of Ptolemy and our Time; viz. those of Albategnius and other Arabian Astronomers. Among which he found two Observations of solar Eclipses, that because of some singular Circumstances attending them, the Sun's Altitude taken at their Beginning and End, ought, in his Judgment, to be accounted more valuable than Gold and Silver. He does not remember that any of these Gentlemen who compiled Tables of the Moon's Motion, having made any Use of these Observations; though probably more Advantage might be reaped from those alone than from all the Observations of Ptolemy. For which Reason, and as they are particularly useful to demonstrate the Acceleration of the Moon's Motion, he thought them worthy of a Place, and so copied them, as follows, from the Prolegomena of Tycho's Historia Caelestis, where they have hitherto lain hid among others of less Account.

"In the Year of Mahomet's Flight 367, on Thursday the 28th of the latter Month Rabia, was observed at Grand Cairo, the Metropolis of Egypt, the Beginning of an Eclipse of the Sun, which, at that Time, was 15°. 43'. high, the Quantity of the Obscuration 8 Digits; at the End the Sun's Altitude was 33°. The same Year, on Sunday the 29th of the Month Sywal the Sun was eclipsed 7 ½ Digits; at the Beginning

SUPPLEMENTAL

SUPPLEMENTAL LUNAR EQUATIONS.

According to the Construction of Mr. MAYER.

XIII. EQUATION of the MOON'S VARIATION.

Argument 13. D equated from the SUN; or D's Pl. equated — O's Pl. true.

D eqd. à O.	Sign 0.		Sign 1.		Sign 2.		Sign 3.		Sign 4.		Sign 5.		D eqd. à O.
	+	Dif.	+	Dif.	±	Dif.	+	Dif.	+	Dif.	+	Dif.	
0	0 0		34 17		33 2		1 57		36 21		36 8		30
1	1 24	1 24	34 56	0 39	32 17	0 45	3 21	1 24	37 1	0 40	35 23	0 45	29
2	2 48	1 24	35 32	0 36	31 30	0 47	4 44	1 23	37 39	0 38	34 36	0 47	28
3	4 11	1 23	36 5	0 33	30 40	0 50	6 7	1 23	38 14	0 35	33 46	0 50	27
4	5 34	1 23	36 35	0 30	29 47	0 53	7 29	1 22	38 47	0 33	32 54	0 52	26
5	6 57	1 23	37 2	0 27	28 52	0 55	8 51	1 22	39 17	0 30	31 59	0 55	25
6	8 19	1 22	37 26	0 24	27 55	0 57	10 12	1 21	39 44	0 27	31 2	0 57	24
7	9 40	1 21	37 48	0 22	26 56	0 59	11 33	1 21	40 8	0 24	30 3	0 59	23
8	11 1	1 21	38 7	0 19	25 56	1 0	12 53	1 20	40 30	0 22	29 1	1 2	22
9	12 21	1 20	38 24	0 17	24 54	1 2	14 13	1 20	40 49	0 19	27 57	1 4	21
10	13 40	1 19	38 38	0 14	23 50	1 4	15 32	1 19	41 5	0 16	26 51	1 6	20
11	14 58	1 18	38 49	0 11	22 44	1 6	16 50	1 18	41 18	0 13	25 43	1 8	19
12	16 15	1 17	38 57	0 8	21 36	1 8	18 6	1 16	41 28	0 10	24 33	1 10	18
13	17 31	1 16	39 2	0 5	20 27	1 9	19 21	1 15	41 35	0 7	23 22	1 11	17
14	18 45	1 14	39 3	0 1	19 16	1 11	20 35	1 14	41 39	0 4	22 9	1 13	16
15	19 58	1 13	39 1	0 2	18 3	1 13	21 48	1 13	41 41	0 2	20 54	1 15	15
16	21 9	1 11	38 56	0 5	16 49	1 14	22 59	1 11	41 40	0 1	19 38	1 16	14
17	22 18	1 9	38 49	0 7	15 34	1 15	24 9	1 10	41 35	0 5	18 20	1 18	13
18	23 25	1 7	38 39	0 10	14 18	1 16	25 17	1 8	41 27	0 8	17 1	1 19	12
19	24 31	1 6	38 26	0 13	13 1	1 17	26 23	1 6	41 16	0 11	15 40	1 21	11
20	25 35	1 4	38 11	0 15	11 42	1 19	27 28	1 5	41 2	0 14	14 18	1 22	10
21	26 37	1 2	37 53	0 18	10 22	1 20	28 31	1 3	40 54	0 17	12 55	1 23	9
22	27 37	1 0	37 32	0 21	9 2	1 20	29 32	1 1	40 25	0 20	11 31	1 24	8
23	28 35	0 58	37 8	0 24	7 41	1 21	30 31	0 59	40 3	0 22	10 7	1 24	7
24	29 31	0 56	36 41	0 27	6 20	1 21	31 23	0 57	39 38	0 25	8 42	1 25	6
25	30 25	0 54	36 10	0 31	4 58	1 22	32 22	0 54	39 10	0 28	7 16	1 26	5
26	31 16	0 51	35 37	0 33	3 30	1 22	33 14	0 52	38 39	0 31	5 50	1 26	4
27	32 5	0 49	35 2	0 35	2 13	1 23	34 4	0 50	38 5	0 34	4 23	1 27	3
28	32 51	0 46	34 24	0 38	0 50	1 23	34 52	0 48	37 29	0 36	2 56	1 27	2
29	33 35	0 44	33 44	0 40	0 33	1 23	35 38	0 46	36 50	0 39	1 28	1 28	1
30	34 17	0 42	33 2	0 42	1 57	1 24	36 21	0 43	36 8	0 42	0 0	1 28	0
Deq. à O.	—	Dif.	—	Dif.	—	Dif.	—	Dif.	—	Dif.	—	Dif.	Deq. à O.
	Sig. 11.		Sig. 10.		Sig. 9.		Sig. 8.		Sig. 7.		Sig. 6.		

Of Mr. MAYER'S LUNAR EQUATIONS and MOTIONS.

"ning the Sun's Altitude was almost 56°. at the End 26°." Another Eclipse is mentioned in the same Place, but it being of the Moon, is therefore omitted. Schickard, who preserved them from Oblivion, adds the following REMARK; "These three Observations were made by Ibn-Junis, who by Order of Abu-Haly-Almansor, a wise Man, at that Time King of Egypt, devoted himself to the Study of Astronomy. This Author's Tables are in the Hands of James Golius, Professor at Lyons, who communicated these Things to me out of them; in which there are several other Observations both of his own and the preceding Age; the Place of Observation was near Grand Cairo." After this Account was transcribed, ready for the Press, Mr. Mayer met with the same Account of these Eclipses, in the Philosophical Transactions, No. 492, given by the celebrated R. Dunkorne, where they are applied to determine the Acceleration of the Moon's Motion; yet he was loth to envy his Readers the Passage he has quoted, as the Observations cannot be too well known among the Astronomers. The Time of the former Eclipse, reduced to Julian Years, is December 13, Ann. Dom. 977. The Beginning, according to the Altitudes taken, is found to be 8h. 24m. 24s. the End 10h. 43m. 44s. Morning; supposing the Latitude of Grand Cairo 30°. 2'. 30". as the Observations of the Moderns make it to be. The other Eclipse, according to our Way of reckoning, happened in the Year of Christ 978, June 8: the Beginning, from the observed Altitude of the Sun, 2h. 30m. 16s. the End 4h. 50m. 24s. Afternoon: With these Times, particularly those of the Ends, Mr. Mayer's Lunar Tables are said to agree within one or two Minutes, as any one (he says) will find by Trial. Whereas all the Tables before published by other Authors make the Times of these Eclipses sooner by almost half an Hour; an infallible Proof, that the Moon's Motion is now swifter than it was many Ages ago; and that the Quantity of this Acceleration has been rightly ascertained in the Equation Tables, according to Mr. Mayer, exhibited at the Beginning.

Nor, indeed, (he says) is this Acceleration of the D's Motion so small, but that it may be plainly shewn from the Observations of this and the last Age only. He has found the Moon's mean Motion in 60 of our Years (besides 802 Revolutions) to be 1° 10' 43" 24" which yet other Tables (the Authors whereof having collected the Mean Motions from the earliest Observations only compared with the Modern) make to be 1°. 10'. 41". 10". or at most 1°. 10'. 42". 15". Now, we have a great many Observations of Eclipses made above 60 Years ago, and with Care and Diligence as at this Day. Most of these Mr. Mayer has reduced to Numbers, and asserts that he has not found one, that proves his Tables, constructed upon the Moon's accelerated Motion, to err a Minute in Longitude; whereas the Error would sometimes have been equal to 3 Minutes, had he retained the Quantity, he says, commonly assumed for the Moon's Mean Motion. He has also pursued, as he asserts, a particular Method to determine, most accurately, the Quantity of this Motion, notwithstanding the Imperfection of his Tables, should there be any yet in them. He chose such Eclipses, as were distant, in Time, one Chaldean Period, that is, 223 Lunations, or, which is better, several Periods Distance from each other, and as all the Irregularities of the Moon are in a Manner restored again after such an Interval of Time, the same Errors in his Tables must needs return again, nearly. If therefore it be known how much his Tables differ from Observations, at the End of one Period, We shall also know how much they ought to disagree at the Expiration of two or more such Periods; and if the Errors are found to be unequal, it is a Sign that the Mean Motion ought to be corrected by such a Quantity as will reduce them to an Equality.

SUPPLEMENTAL LUNAR EQUATIONS.

According to the Construction of Mr MAYER.

XIV. Reduction - Equation
to the Ecliptic.

D à 8	Arg. D in Orb. à 8 cor.			D à 8
	0.6	1.7	2.8	
0	0	6	2	30
1	0	14	6	29
2	0	29	15	28
3	0	43	21	27
4	0	58	27	26
5	1	12	32	25
6	1	26	37	24
7	1	41	41	23
8	1	55	45	22
9	2	9	48	21
10	2	23	51	20
11	2	37	54	19
12	2	50	56	18
13	3	3	57	17
14	3	16	57	16
15	3	29	57	15
16	3	41	57	14
17	3	53	57	13
18	4	5	56	12
19	4	17	54	11
20	4	28	51	10
21	4	39	48	9
22	4	50	45	8
23	5	0	41	7
24	5	10	37	6
25	5	20	32	5
26	5	30	27	4
27	5	39	21	3
28	5	47	15	2
29	5	55	9	1
30	6	2	2	0
D à 8	+	+	+	D à 8
8	11.5	10.4	9.3	8

I. For the Latitude of the Moon.

Argument. D in Orb. à Long 8 corrected.

Arg. proper	0. N.		Dif.	1. N.		Dif.	2. N.		Dif.
	6. S.	7. S.		7. S.	8. S.		8. S.	9. S.	
0	0	0	0	2	34	25	4	27	38
1	0	5	23	2	39	4	4	30	18
2	0	10	46	2	43	39	4	32	53
3	0	16	9	2	48	12	4	35	22
4	0	21	32	2	52	42	4	37	47
5	0	26	54	2	57	9	4	40	7
6	0	32	16	3	1	33	4	42	21
7	0	37	37	3	5	53	4	44	30
8	0	42	58	3	10	10	4	46	34
9	0	48	18	3	14	23	4	48	33
10	0	53	37	3	18	33	4	50	27
11	0	58	55	3	22	39	4	52	15
12	1	4	11	3	26	42	4	53	58
13	1	9	27	3	30	41	4	55	36
14	1	14	42	3	34	36	4	57	8
15	1	19	55	3	38	27	4	58	35
16	1	25	6	3	42	14	4	59	56
17	1	30	16	3	45	57	4	1	12
18	1	35	24	3	49	36	4	2	22
19	1	40	31	3	53	11	4	3	26
20	1	45	36	3	56	41	4	4	25
21	1	50	39	4	0	7	4	5	19
22	1	55	40	4	3	29	4	6	7
23	2	0	39	4	6	47	4	6	50
24	2	5	36	4	9	59	4	7	26
25	2	10	30	4	13	7	4	7	57
26	2	15	22	4	16	11	4	8	23
27	2	20	12	4	19	10	4	8	43
28	2	24	59	4	22	4	4	8	57
29	2	29	43	4	24	53	4	9	5
30	2	34	25	4	27	38	4	9	8
Arg. proper	11. S.	10. S.	Dif.	10. S.	9. S.	Dif.	9. S.	8. S.	Dif.
8	5. N.	4. N.	3. N.	3. N.	3. N.	3. N.	3. N.	3. N.	3. N.

II. For the Latitude of
the Moon.

Arg. 2 D à 8 — forgoing Arg. Lat.

Arg. 2 D à 8	0. N.		1. N.	2. N.	Arg. prop
	6. S.	7. S.			
0	0	4	25	7	39
1	0	9	4	33	7
2	0	19	4	41	7
3	0	28	4	49	7
4	0	37	4	56	7
5	0	46	5	4	8
6	0	55	5	11	8
7	1	5	5	19	8
8	1	14	5	26	8
9	1	23	5	34	8
10	1	32	5	41	8
11	1	41	5	48	8
12	1	50	5	55	8
13	1	59	6	2	8
14	1	8	6	9	8
15	2	17	6	15	8
16	2	26	6	22	8
17	2	35	6	28	8
18	2	44	6	34	8
19	2	52	6	40	8
20	3	1	6	46	8
21	3	10	6	52	8
22	3	18	6	57	8
23	3	27	7	3	8
24	3	35	7	8	8
25	3	44	7	14	8
26	3	52	7	19	8
27	4	1	7	24	8
28	4	9	7	29	8
29	4	17	7	34	8
30	4	25	7	39	8
Arg. 2 D à 8	11. S.	10. S.	9. S.	8. S.	Arg. prop
8	5. N.	4. N.	3. N.	3. N.	3. N.

Of Mr. MAYER'S LUNAR EQUATIONS and SOLAR NUMBERS.

Equality. In this Manner he has examined, he says, not one or two Cycles only, but a great many thereof; so that he further asserts there are very few Eclipses, of which Observations have been made in this and the last Age, but what have undergone this Examination.

With these Tables he has also compared many Observations of the Moon that were made a little earlier, by Tycho, Walther, Regiomontanus, as well in the Syzygies as without them, and have all along found as near an Agreement as could well be expected from Observations not made to the greatest Nicety and Exactness.

With Respect to HIS SOLAR NUMBERS, of the MEAN PLACES and MOTIONS, deducible from our own Examples, and which we have exhibited, with their Equation, (p. 59.) though he had not so much left for him to do as those going before him, yet he assumed no Principle, he says, but what has stood the Test of repeated Observations. Thus he retained the Motion of the Sun's Apogee, exhibited in the Tables of Dr. Flamsteed, as agreeing nearest with Observations, especially those of Hipparchus and Albategnius. Moreover, he made the Quantity of the Solar Tropical Year invariable, agreeable to the Observations of Hipparchus, Albategnius, Walther, and, in short, of all the Astronomers of any Credit and Authority, except Ptolemy, whose Equinoxes, 'tis almost indisputably certain, were not ascertained by Observations of the Heavens; but determined by his own Tables, and the Quantity of the Solar Year, which he had received from Hipparchus. Mr. Mayer is therefore firmly of Opinion, from very substantial Reasons, that the Chronological Order of Time from Ptolemy down to us, has not been interrupted by the Omission of a Day or two. Lastly, Mr. Mayer has deduced the mean apparent Motion of the Fixed Stars, or the Precession of the Equinoxes, chiefly from Timocharis's Observations of the Moon's Appulses to the Fixed Stars, which Method seemed to him more certain than if he had taken the Longitude of those Stars from Hipparchus, and made use of them to this purpose.

Who therefore thinks he has strictly examined and closely considered every Thing, that may be justly required of any one who takes the Subject in Hand, and has left nothing untried, he thinks, that might conduce to the perfecting of the Moon's Theory; a Point far the most intricate of all in the celestial Science. And, if he is not greatly deceived, his Tables are now brought to almost as great a Degree of Perfection, as the Lunar Observations themselves can be brought by the Methods hitherto made use of by the Astronomers; who further alledges, that

SUPPLEMENTAL

SUPPLEMENTAL LUNAR EQUATIONS.

According to the Construction of Mr. MAYER.

EQUATORIAL PARALLAX, and APPARENT DIAMETER of the MOON.

MEAN HORIZONTAL PARALLAX of the MOON, for the Middle State of her ORBIT, in, or out of, Syzygies.

1. Equat. Par. for tr. State of Orb. in or out of Syzygies.

2. Equation of the D's Parallax out of Syzygies, only.

For Diameter D from Equatorial Parallax

Argument II. Or D's Corrected M. ANOMALY.

D	Sig. 0.	Sig. 1.	Sig. 2.	Sig. 3.	Sig. 4.	Sig. 5.	D
Cor	HorPar	HorPar	HorPar	HorPar	HorPar	HorPar	Cor
An.	+	+	+	+	+	+	An.
0	1	2	3	4	5	6	7
0	54	10	54	30	55	29	58
1	54	10	54	32	55	32	58
2	54	10	54	33	55	34	57
3	54	10	54	35	55	37	57
4	54	11	54	36	55	39	57
5	54	11	54	38	55	42	57
6	54	11	54	39	55	45	57
7	54	11	54	41	55	47	57
8	54	11	54	42	55	50	57
9	54	12	54	44	55	53	57
10	54	12	54	46	55	56	57
11	54	12	54	48	55	59	57
12	54	13	54	49	56	2	57
13	54	13	54	51	56	5	57
14	54	14	54	53	56	8	57
15	54	15	54	55	56	11	57
16	54	15	54	57	56	14	57
17	54	16	54	59	56	17	57
18	54	17	55	1	56	20	57
19	54	18	55	3	56	23	58
20	54	19	55	5	56	26	58
21	54	20	55	7	56	29	58
22	54	21	55	9	56	32	58
23	54	22	55	12	56	36	58
24	54	23	55	14	56	39	58
25	54	24	55	17	56	42	58
26	54	25	55	19	56	45	58
27	54	26	55	22	56	49	58
28	54	28	55	24	56	52	58
29	54	29	55	27	56	55	58
30	54	30	55	29	56	58	58
D	+	+	+	+	+	+	D
Cor	HorPar	HorPar	HorPar	HorPar	HorPar	HorPar	Cor
An.	Sig. 11.	Sig. 10.	Sig. 9.	Sig. 8.	Sig. 7.	Sig. 6.	An.

Ar. 2. D² - D cor. An. or Arg. 12.

Arg. prop.	0	1	2
0	38	33	19
1	38	32	18
2	38	32	18
3	38	32	17
4	38	31	17
5	38	31	16
6	38	31	16
7	37	30	15
8	37	30	14
9	37	30	14
10	37	29	13
11	37	29	13
12	37	28	12
13	37	28	11
14	36	28	11
15	36	27	10
16	36	27	9
17	36	26	9
18	36	26	8
19	35	25	7
20	35	25	6
21	35	24	6
22	35	24	5
23	34	23	4
24	34	23	4
25	34	22	3
26	34	21	3
27	33	21	2
28	33	20	1
29	33	20	1
30	33	19	0
Arg. prop.	11	10	9
	+	+	+
	5	4	3

Arg. D eq. a true.	0	1	2	3	4	5	Arg. pro.
0	25	12	14	27	13	14	30
1	25	12	14	27	12	14	29
2	25	11	15	27	12	15	28
3	25	10	16	27	11	16	27
4	25	9	16	27	10	17	26
5	25	8	17	27	9	18	25
6	25	8	18	27	8	18	24
7	24	7	18	26	7	19	23
8	24	6	19	26	6	20	22
9	24	5	20	26	5	20	21
10	24	4	21	25	4	21	20
11	24	3	21	25	3	22	19
12	23	2	22	24	2	22	18
13	23	1	23	24	1	23	17
14	23	0	23	23	0	23	16
15	23	1	24	23	1	24	15
16	22	1	24	22	2	24	14
17	22	2	25	22	3	25	13
18	21	3	25	21	4	25	12
19	20	4	25	21	5	25	11
20	19	5	26	20	6	26	10
21	19	6	26	19	7	26	9
22	18	7	26	19	7	26	8
23	17	8	26	18	8	26	7
24	17	9	26	17	9	27	6
25	16	10	27	16	10	27	5
26	15	10	27	16	11	27	4
27	15	11	27	15	11	28	3
28	14	12	27	14	12	28	2
29	13	13	27	13	13	28	1
30	12	14	27	13	14	28	0
Arg. pro	+	+	-	-	+	+	Arg. pro
	11	10	9	8	7	6	

Par.	Diam.
54 5	29 20
54 16	29 36
54 27	29 42
54 38	29 48
55 49	29 54
55 6	30 0
55 11	30 6
55 22	30 12
55 33	30 18
55 44	30 24
55 55	30 30
56 6	30 36
56 17	30 42
56 28	30 48
56 39	30 54
56 50	31 0
57 1	31 6
57 12	31 12
57 23	31 18
57 34	31 24
57 45	31 30
57 56	31 36
58 7	31 42
58 18	31 48
58 29	31 54
58 40	32 0
58 51	32 6
59 2	32 12
59 13	32 18
59 24	32 24
59 35	32 30
59 46	32 36
59 57	32 42
60 8	32 48
60 19	32 54
60 30	33 0
60 41	33 6
60 52	33 12
61 3	33 18
61 14	33 24
61 25	33 30

Of Mr. MAYER'S LUNAR EQUATIONS, and SOLAR and LUNAR MOTIONS.

all who have been accustomed to the handling of Instruments, and making Observations therewith, must readily allow, how liable a Person is to err a single Minute. That unless the Astronomers should hereafter discover a Method of ascertaining the Moon's Place in the Heavens by Observations to a greater Degree of Exactness, he thinks a more perfect Theory of the Moon than that by his Tables, can hardly be expected or required. See *Cassini de Thury in Memoirs de l'Acad. Roy. des Sciences, et Belles Lettres de Berlin, Année 1749. p. 364.*

According to his Tables the mean tropical solar Year contains 365^d. 5^h. 48^m. 51^s. 12th. and is, for ought appears from Observations, invariably the same; though perhaps the Quantity of the mean natural Days, or the Earth's diurnal Revolution, be not, according to the Opinion commonly received, equable.

Moreover, the mean Periodical Lunar Month he makes to be variable, and gradually, though slowly, diminishes: Its Decrease in a Century being 1st. 5th. 15th. At the Beginning of the Century 1700, the Periodical Month, was 27^d. 7^h. 43^m. 4^s. 45th. 14th. and at the Time, between Hipparchus and Ptolemy, or at the Beginning of the Christian Era, it was 27^d. 7^h. 43^m. 5^s. 31th. 30th. Hence the mean Synodical Month is continually decreasing. In the Beginning of the present Century 1700, its Quantity was 29^d. 12^h. 44^m. 2^s. 53th. 23rd. and at the Commencement of the Christian Era 29^d. 12^h. 44^m. 3^s. 15th. so that it loses in every hundred Years, if Observations can be relied on, 16th. 16th.

In ECLIPSES OF THE MOON, the Semidiameter of the Earth's Shadow, found from the common Tables, must, on Account of the Earth's Atmosphere, be augmented by the 60th Part of itself, or so many Seconds added to the said Semidiameter as there are Minutes contained in it.

Vide Commentarii Societatis Regiæ Scientiarum Gottingensis. Tomus II. ad Annum 1752. Being The Acts of the Gottingen Society. Thus have we given an impartial Account of the celebrated Mr. Mayer's Lunar and Solar Tables in general, in order that they may be judged and compared with our own; whatever Correction; or Improvements of his Tables Mr. Mayer shall think fit to make hereafter.

According to the original Construction of Mr. EULER, and late Improvement and Correction of his PLAN.

Argument. Mean Anomaly of the SUN and MOON, respectively.

LUNAR EQUATION of the SUN, according to Mr. EULER.

Argument. MOON from the SUN.

REMARKS on the SUN and MOON'S MEAN MOTIONS, according to
Mr. Leonard Euler, of the Royal Academy of Sciences at Berlin, and
F. R. S. at London.

FIRST, *Of the SUN's mean MOTION and ACCELERATION.*

Intervall

SUPPLEMENTAL LUNAR EQUATIONS.

According to the IMPROVEMENT and Correction of the celebrated Mr. EULER's PLAN.

EULER's I. EQUATION of MEAN LONGITUDE for the middle State of the MOON's Orbit. . . Maximum $6^{\circ} 12' 2''$
 Mayer's Maximum $6^{\circ} 18' 44''$

DIFFERENCE, Euler's Maximum too little . . . $6' 42''$

Hence, Mayer's XI. EQUATION divided by 56,53, and the Quotient subtracted from that Equation will give Euler's I. EQUATION, if you chuse to try it.

N. B. You may perform the DIVISION by Logistical Logarithms; as shall be shewn further on.

For EULER's I. use MAYER's XI. EQUATION.

EULER's II. EQUATION (Argument, Sun's excentric Anomaly) Maximum . . . $9' 44''$
 MAYER's EQUATION MOON's mean Anomaly, p. 97. (Argument, Sun's mean Anomaly) Maximum . . . $10' 18''$

DIFFERENCE, Euler's Maximum too little . . . $0' 34''$

Hence, MAYER's EQUATION of Moon's Anom. (p. 97) divided by 18,18 and the Quotient taken from that Equation will give Euler's II. EQUATION.

For EULER's II. use MAYER's Equation Moon's Anom. for Longitude (p. 97) with Arg. Sun's excentric Anom. of EULER, or Sun's mean Anomaly. Or, use our I. EQUATION (Arg. Sun's An.) p. 61.

EULER's III. EQUATION (Arg. Moon twice equated à Sun) . . . Maxima . . .	1st Part. $33' 27''$	2d Part. $36' 10''$
MAYER's XIII. (Argument, Moon equated à Sun) . . . Maxima . . .	$39' 31''$	$41' 41''$

DIFFERENCE, EULER's Maxima too little . . . $5' 36''$ | $5' 31''$

Hence, Mayer's first Part XIII. Equation divided by 6,97, and the Quotient taken from that Equation will give Euler's III. Equation.

Also, Mayer's latter Part XIII. Equation divided by 7,55, and the Quotient taken from that Equation will give Euler's III. Equation.

For EULER's III. use MAYER's XIII. EQUATION.

Intervals of Time forward, in Imitation of that same Correctness of Judgment shewn by Mr. Euler before him, in respect of the Sun or Earth's Acceleration, or Increase of Place at equal Intervals of Time. And therefore the further we go into Antiquity or Futurity, the greater will be the Correction of mean Motion or Place, from the stated Period whence the Correction begins; as at equal Distance of Time from that Period into Antiquity and Futurity, the Correction of the limited mean Motion will be evidently the same, according to the Hypothesis of an Acceleration of Motion, or Increase of mean Places at equal Intervals of Time forward.

N. B. Mr. Euler's limited mean Motion of the Sun in 20 Julian Years, $= 20^{\circ} 08' 09'' 12''' = 20^{\circ},0004259259$, &c. by which dividing 7305 Days in the 20 Julian Years, the Quotient will be $365^{\circ} 24' 22'' 16'''$, &c. $= 365^{\circ} 5^{\circ} 48' 47'' 57'''$, &c. his limited mean Year; but according to his Hypothesis of Acceleration of the Sun's mean Motion, still diminishes from that Quantity.

Nobody, it is presumed, (says our Author) will deny the Earth's Situation is varied, a small Matter, by the Moon's Attraction, and therefore knowing the Cause to which this Variation may be ascribed, the celebrated Euler has, in a Table, shewn how much the Sun's Place, and his Distance from the Earth may be changed with Regard to each lunar Aspect.

He is, nevertheless, doubtful whether his Corrections, which he has taken from the horizontal Parallax $12''$, will not be something too little; because, by comparing his several Corrections with former Observations, he has discovered that the Sun's horizontal Parallax is but $10''$, or probably does not exceed it above $\frac{1}{2}$ a Second. He therefore chuses to abide by his Correction of the Sun's limited mean Motion and Acceleration according to the Hypothesis he has laid down, till the Quantity of Variation can be more accurately discovered from future Observation.

He collected the Place of the Sun's Apogee, and the Excentricity of the Orbit from the Observations of the celebrated Monnierus, which he with great Study and Pains had instituted for the same Purpose, and published before that remarkable Correction from the Moon's Rising. He has made the Motion of the Apogee in Nothing different from the Precession of the Equinox, because (he says) the Arguments used by Astronomers to prove the particular Motion of the Apogee forward of the fixed Stars, appear (to him) to be insufficient. For though he allows the Resistance of the Ether, diminishing the Earth's Orbit, from whence the Acceleration of the Earth's Motion, or Decrease of the Year, is supposed to proceed, yet he has found, from this Resistance, the Position of the Line of Apides of the Sun, or Earth in her Orbit, not to be affected therewith; whence, he would rather (if any Variation of the Apogee, with Regard to the fixed Stars can be truly demonstrated) attribute it to the Action of a Comet sometimes passing very near the Earth.

Of the LUNAR NUMBERS and EQUATIONS of Mr. EULER.

The Design of which (he says) to set forth at large would take up too much Room in the Compass of his small Essay: He therefore observes, that they are taken from the same Theory of Attraction which the celebrated Sir ISAAC NEWTON introduced into the Science of Astronomy with so happy Success. That though there are several lunar Tables, published before his, founded on the same Hypothesis, yet he asserts that the Calculation to which the Principle of Attraction is relative, is exceedingly intricate, and that all those Tables widely differ from the Theory; nor does he pretend to have shewn all the unequal Motions in these Tables. Nevertheless, he has set forth all the Equations (he says) perceptible to Observation, and which are to be found after the first half Minute.

Thus far he offers his Calculation to the Public; shewing the Title and Argument of every individual Inequality or Equation, determining most of the certain Quantities of each, according to the aforesaid celebrated Theory. Some of which Quantities and Equations he was obliged to ascertain by Observation; others (he says) depend upon the Sun's Distance from the Earth, or his horizontal Parallax, which, agreeable to the latest Observations, he finds to be $12'' \frac{1}{2}$; but, afterwards, on comparing these several lunar Equations with former Observations, he finds the Sun's Parallax cannot probably be reckoned above $10''$.

He farther observes, that whoever compares his Tables with others already published, may presently discern a vast Difference both in the Titles and Arguments of the Equations. That as he makes Use of the eccentric Anomalies of the SUN and MOON, he places them by themselves, in both solar and lunar Tables, that those and the other Anomalies may be the more readily found.

That, though the Number of his Equations is larger than in any other Tables preceding his, yet he has so contrived them, that the Computation may be made much sooner, and with less Trouble from them than by former Tables.

And lastly, that, since all Causes affecting the Moon's Place and her Distance from the Earth, are so very variable, he has added the more Tables, by the Help of which, (he says) the Distance of the Moon from the Earth, and from thence her horizontal Parallax and apparent Diameter may be the more correctly and readily known.

SEE Latin Preface to the Berlin astronomical Tables, from which Tables Mr. Mayer probably derived many of his later Improvements, particularly the Acceleration of the Moon's mean Motion, by an Eye bad to Mr. Euler's Acceleration first supposed of the Sun's mean Motion, and Solution of the Moon's central Equation, afterwards imitated by Mr. Mayer, (who likewise solved the Moon's Parallax for the true State of her Orbit in like Manner) as he partly acknowledges, whence his Improvements, by confessing that the same celebrated Eulerus (Author of the abovementioned Tables) was the first who solved the Moon's Central Equation, in two Parts; without Eccentricity, (see p. 99 of our Work) and "so adorned the lunar Theory with a neat, and elegant Contraction." And who admits (see Remarks at the Bottom of Mr. Mayer's Equation Tables) that the above-mentioned Mr. Euler was the first who reduced the Newtonian Theory to general analytical Equations, which Mr. Mayer afterwards "bit upon an elegant and concise Method of resolving."

SUPPLEMENTAL LUNAR EQUATIONS.

According to the Construction of Mr. EULER.

IV. EQUATION of the MOON, or EVECTION.

Argument. Twice Dist. MOON from SUN — Excentric Anom. MOON.

Arg. pro.	Sig. 0.				Sig. 1.				Sig. 2.				Sig. 3.				Sig. 4.				Sig. 5.				Arg. pro.								
	—			Dif.	—			Dif.	—			Dif.	—			Dif.	—			Dif.	—			Dif.									
0	0	1	2	1	0	0	1	2	1	0	0	1	2	1	0	0	1	2	1	0	0	1	2	1	0	0	1	2	1	0			
0	0	0	0	1	0	36	59	1	6	1	4	5	0	38	1	14	0	1	4	5	0	36	59	1	6	0	36	59	1	6	30		
1	0	1	18	1	0	38	5	1	7	1	4	43	0	37	1	13	59	1	3	26	0	35	53	1	8	0	35	53	1	8	29		
2	0	2	34	1	0	39	12	1	6	1	5	20	0	35	1	13	57	1	2	45	0	34	45	1	9	0	34	45	1	9	28		
3	0	3	52	1	0	40	18	1	5	1	5	55	0	36	1	13	53	1	2	4	0	33	36	1	10	0	33	36	1	10	27		
4	0	5	9	1	0	41	23	1	5	1	6	31	0	32	1	13	49	1	1	20	0	32	26	1	10	0	32	26	1	10	26		
5	0	6	27	1	0	42	27	1	4	1	7	3	0	33	1	13	43	1	0	36	0	31	16	1	10	0	31	16	1	10	25		
6	0	7	44	1	0	43	30	1	3	1	7	36	0	30	1	13	35	1	0	59	51	0	30	6	1	10	0	30	6	1	10	24	
7	0	9	1	1	0	44	32	1	2	1	8	6	0	30	1	13	26	1	9	5	0	28	55	1	11	0	28	55	1	11	23		
8	0	10	18	1	0	45	34	1	2	1	8	36	0	28	1	13	16	1	10	18	0	27	43	1	12	0	27	43	1	12	22		
9	0	11	35	1	0	46	34	1	0	1	9	4	0	28	1	13	5	1	11	30	0	26	31	1	12	0	26	31	1	12	21		
10	0	12	51	1	0	47	34	1	0	1	9	32	0	28	1	12	52	1	13	41	0	25	18	1	13	0	25	18	1	13	20		
11	0	14	8	1	0	48	33	0	59	1	9	57	0	25	1	12	38	0	14	51	0	24	5	1	13	0	24	5	1	13	19		
12	0	15	23	1	0	49	31	0	58	1	10	22	0	25	1	12	22	0	16	50	0	22	51	1	14	0	22	51	1	14	18		
13	0	16	39	1	0	50	27	0	56	1	10	46	0	24	1	12	5	0	17	54	0	21	37	1	14	0	21	37	1	14	17		
14	0	17	54	1	0	51	24	0	57	1	11	7	0	21	1	11	48	0	17	53	0	20	24	1	13	0	20	24	1	13	16		
15	0	19	9	1	0	52	19	0	55	1	11	28	0	21	1	11	28	0	20	52	19	0	19	9	1	15	0	19	9	1	15	15	
16	0	20	24	1	0	53	14	0	55	1	11	48	0	20	1	11	7	0	21	51	24	0	17	54	1	15	0	17	54	1	15	14	
17	0	21	37	1	0	54	7	0	53	1	12	5	0	17	1	10	46	0	21	50	27	0	16	39	1	15	0	16	39	1	15	13	
18	0	22	51	1	0	54	58	0	51	1	12	22	0	17	1	10	22	0	24	49	31	0	15	23	1	16	0	15	23	1	16	12	
19	0	24	5	1	0	55	50	0	52	1	12	38	0	16	1	9	57	0	25	48	33	0	14	8	1	15	0	14	8	1	15	11	
20	0	25	18	1	0	56	41	0	51	1	12	52	0	14	1	9	32	0	25	47	34	0	12	51	1	17	0	12	51	1	17	10	
21	0	26	31	1	0	57	30	0	49	1	13	5	0	13	1	9	4	0	28	46	34	0	11	35	1	16	0	11	35	1	16	9	
22	0	27	43	1	0	58	18	0	48	1	13	16	0	11	1	8	36	0	28	45	34	0	10	18	1	17	0	10	18	1	17	8	
23	0	28	55	1	0	59	5	0	47	1	13	26	0	10	1	8	6	0	30	44	32	0	9	1	1	17	0	9	1	1	17	7	
24	0	30	6	1	0	59	51	0	46	1	13	35	0	9	1	7	36	0	30	43	30	0	7	44	1	17	0	7	44	1	17	6	
25	0	31	16	1	0	0	36	0	45	1	13	43	0	8	1	7	3	0	33	42	27	0	6	27	1	17	0	6	27	1	17	5	
26	0	32	26	1	0	1	20	0	44	1	13	49	0	6	1	6	31	0	32	41	23	0	5	9	1	18	0	5	9	1	18	4	
27	0	33	36	1	0	1	2	4	44	1	13	53	0	4	1	5	55	0	36	40	18	0	3	52	1	17	0	3	52	1	17	3	
28	0	34	45	1	0	1	2	45	41	1	13	57	0	4	1	5	20	0	35	39	12	0	2	34	1	18	0	2	34	1	18	2	
29	0	35	53	1	0	1	3	26	41	1	13	59	0	2	1	4	43	0	37	38	5	0	1	18	1	16	0	1	18	1	16	1	
30	0	36	59	1	0	1	4	5	39	1	14	0	0	1	1	4	5	0	38	36	59	0	0	0	1	18	0	0	0	1	18	0	
Arg. pro.	+			Dif.	+			Dif.	+			Dif.	+			Dif.	+			Dif.	+			Dif.	+			Dif.	+			Dif.	Arg. pro.
	Sig. 11.				Sig. 10.				Sig. 9.				Sig. 8.				Sig. 7.				Sig. 6.				Sig. 5.				Arg. pro.				

MAYER's Argument to the above EVECTION-EQUATION is twice the Moon from Sun — Mean Anom. Moon.

The above Maximum . . . 1° 14' 0"

MAYER's Maximum of Evection or XII. Equation . . . 1 20 43

Difference, EULER's Maximum too little . . . 6 43

Hence, MAYER's Evection-Equation divided by 12,02, and the Quotient taken from that Equation, will give the above Equation of EULER, correspondent.

For the above Equation use MAYER's XII. Equation, if you neglect the following V. Equation of EULER. But, if you retain the Vth. with the above EVECTION of EULER, they will, conjunctly used, come near MAYER's EVECTION or XII. EQUATION; and compensate for some Errors, or Inequalities, of the Lunar Orbit.

N. B. The Evection-Equation according to STREET's intricate Method of computing it (from the Chord Evection, and Argument of half the Synodical Anomaly) is supplied by the easier Computation of the Evection-Equation above. Which Equation is applied to the central Equation for the mean Eccentricity, and is therefore equal to about half of STREET's Evection, applied to the central (or correspondent eccentric) Equation, for the least Eccentricity of the Lunar Orbit.

SUPPLEMENTAL LUNAR EQUATIONS.

According to the Construction of Mr. EULER.

V. EQUATION of the MOON.

Arg. Twice Δ à $\odot$ — Twice Excent. Anom. Δ .

Arg. pro.	Sig. 0.	Sig. 1.	Sig. 2.	Sig. 3.	Sig. 4.	Sig. 5.	Arg. pro.
0	+	+	+	+	+	+	0
0	0 0	4 12	7 16	8 24	7 16	4 12	30
1	0 9	4 20	7 21	8 24	7 12	4 4	29
2	0 18	4 27	7 25	8 24	7 7	3 57	28
3	0 26	4 34	7 29	8 23	7 3	3 49	27
4	0 35	4 42	7 33	8 23	6 58	3 41	26
5	0 44	4 49	7 37	8 22	6 53	3 33	25
6	0 53	4 56	7 40	8 21	6 48	3 25	24
7	1 1	5 3	7 44	8 20	6 43	3 17	23
8	1 10	5 10	7 47	8 19	6 37	3 9	22
9	1 19	5 17	7 51	8 18	6 32	3 1	21
10	1 28	5 24	7 54	8 15	6 26	2 52	20
11	1 36	5 31	7 57	8 15	6 20	2 44	19
12	1 45	5 37	7 59	8 13	6 15	2 36	18
13	1 53	5 44	8 2	8 11	6 9	2 27	17
14	2 2	5 50	8 4	8 9	6 3	2 19	16
15	2 10	5 56	8 7	8 7	5 56	2 10	15
16	2 19	6 3	8 9	8 4	5 50	2 2	14
17	2 27	6 9	8 11	8 2	5 44	1 53	13
18	2 36	6 15	8 13	7 59	5 37	1 45	12
19	2 44	6 20	8 15	7 57	5 31	1 36	11
20	2 52	6 26	8 16	7 54	5 24	1 28	10
21	3 1	6 32	8 18	7 51	5 17	1 19	9
22	3 9	6 37	8 19	7 47	5 10	1 10	8
23	3 17	6 43	8 20	7 44	5 3	1 1	7
24	3 25	6 48	8 21	7 40	4 56	0 53	6
25	3 33	6 53	8 22	7 37	4 49	0 44	5
26	3 41	6 58	8 23	7 33	4 42	0 35	4
27	3 49	7 3	8 23	7 29	4 34	0 26	3
28	3 57	7 7	8 24	7 25	4 27	0 18	2
29	4 4	7 12	8 24	7 21	4 20	0 9	1
30	4 12	7 16	8 24	7 16	4 12	0 0	0
Arg. pro.	S. 11.	S. 10.	Sig. 9.	Sig. 8.	Sig. 7.	Sig. 6.	Arg. pro.

VI. EQUATION of the MOON.

Arg. Twice Δ à $\odot$ + Excent. Anom. Δ .

Arg. pro.	Sig. 0.	Sig. 1.	Sig. 2.	Sig. 3.	Sig. 4.	Sig. 5.	Arg. pro.
0	+	+	+	+	+	+	0
0	0 0	0 26	0 44	0 52	0 44	0 26	30
5	0 5	0 30	0 46	0 52	0 42	0 22	25
10	0 9	0 33	0 48	0 51	0 39	0 17	20
15	0 13	0 36	0 50	0 50	0 36	0 13	15
20	0 17	0 39	0 51	0 48	0 33	0 9	10
25	0 22	0 42	0 52	0 46	0 30	0 5	5
30	0 26	0 44	0 52	0 44	0 26	0 0	0
Arg. pro.	S. 11.	S. 10.	Sig. 9.	Sig. 8.	Sig. 7.	Sig. 6.	Arg. pro.

VII. EQUATION of the MOON.

Arg. Twice Δ à $\odot$ — Excent. Anom. $\odot$.

Arg. pro.	Sig. 0.	Sig. 1.	Sig. 2.	Sig. 3.	Sig. 4.	Sig. 5.	Arg. pro.
0	—	—	—	—	—	—	0
0	0 0	0 29	0 49	0 57	0 49	0 29	30
5	0 5	0 33	0 52	0 57	0 47	0 24	25
10	0 10	0 37	0 54	0 56	0 44	0 19	20
15	0 15	0 40	0 56	0 56	0 40	0 15	15
20	0 19	0 44	0 56	0 54	0 37	0 10	10
25	0 24	0 47	0 57	0 52	0 33	0 5	5
30	0 29	0 49	0 57	0 49	0 29	0 0	0
Arg. pro.	S. 11.	S. 10.	Sig. 9.	Sig. 8.	Sig. 7.	Sig. 6.	Arg. pro.

VIII. EQUATION of the MOON.

Arg. Twice Δ à $\odot$ + Excentric Anom. $\odot$.

Arg. pro.	Sig. 0.	Sig. 1.	Sig. 2.	Sig. 3.	Sig. 4.	Sig. 5.	Arg. pro.
0	—	—	—	—	—	—	0
0	0 0	0 21	0 36	0 42	0 36	0 21	30
5	0 4	0 24	0 38	0 42	0 34	0 18	25
10	0 7	0 27	0 40	0 41	0 32	0 14	20
15	0 11	0 30	0 41	0 41	0 30	0 11	15
20	0 14	0 32	0 41	0 40	0 27	0 7	10
25	0 18	0 34	0 42	0 38	0 24	0 4	5
30	0 21	0 36	0 42	0 36	0 21	0 0	0
Arg. pro.	S. 11.	S. 10.	Sig. 9.	Sig. 8.	Sig. 7.	Sig. 6.	Arg. pro.

SIMILAR to VI. above, of Euler, is Mayer's VI. Equation, his Maximum being $1' 30''$ instead of $52''$; and similar to VII. above, of Euler, is Mayer's III. Eq. whose Max. is $1' 2''$ instead of $57''$; and similar to Euler's VIII. above, is Mayer's II. whose Max. is $54''$ instead of $42''$, according to Euler: allowing for the Dif. between the mean Anomalies of Mayer, and excentric Anomalies of $\odot$ and Δ of Euler, on the respective Arguments, the Signs of both these Authors Equations agreeing. But Mr. Mayer has not any Equat. similar to Mr. Euler's V. Equat. above, which Equat. is the only Exception of Difference from all Authors before him that we have seen. The Moon's Place in her Orbit is nearly obtained by the VIII. foregoing EQUATIONS, partly from Mr. MAYER, and partly from Mr. EULER's Plan. After which Orbit Place of the Moon is had, MAYER's XIV. or Reduction-Equation from the Orbit to the Ecliptic may be applied, or any of our own Reduction-equations, for that Purpose. And the Parallax and apparent Semi-Diameter of the Moon to be afterwards determined according to our Tables for that Purpose, at the Beginning of this Work, or else by Mr. MAYER's Supplemental Equation-Tables, for the same Purpose, and not by the Halleian Method; because EULER considers no second EQUATION of the Moon's Apogee for determining the Argument of true Anomaly of the Moon, on which the Parallax, and consequently apparent Semi-Diameter of the Moon, chiefly depends, according to the Halleian Computation.

Mr. EULER has given five Equations of the ascending Node, following the foregoing VIII. of the Moon's Orbit, and with the 3d Equation of the Moon's $\odot$ has given the Inclination of her Orbit, which he corrects by two following Equations, with those of the $\odot$. And with the 4th Equation of the Node, he has given a Reduction of the Orbit-Place of the Moon to that of the Ecliptic, at a mean Quantity, which Equation he makes the same as the 5th Equation of the Node.

He then determines the Distance of the Moon from the Earth, by the following Arguments to VIII. Equations: 1. Mean Anom. Δ ; 2. Excentric An. Sun; 3. Moon twice equated from Sun; 4. Twice Moon à Sun — excentric An. Moon; 5. Twice Moon à Sun — twice excent. An. Δ ; 6. Twice Δ à Sun + excentric An. Δ ; 7. Twice Δ à Sun — excent. An. Sun; 8. Twice Δ à Sun + excent. An. Sun. He corrects the first Distance by the several following Equations of Distances, according to their proper Signs, and having obtained the correct Distance (according to this Method) he takes out the horizontal Parallax, and apparent Diameter of the Moon, according to the Argument of that correct Distance. N. B. He makes the Moon's horizontal Parallax to the apparent Diameter as $60' 40''$ to $33' 20''$. — But these Methods are not so certain and correct as others we have exhibited; yet we would not omit to oblige our Readers with Mr. EULER's ingenious Discoveries.

DIFFERENT

DIFFERENT METHODS

OF SOLVING THE

Keplerian PROBLEM.

A LIMITED CORRECTION of WARD's Hypothesis, by two Terms in Seconds, to be connected with the mean Anomaly, for determining the true Angle, at the upper Focus of a Planet, from Aphelion.

PUT a = Sem. Transf. Orbit, b = Sem. Conjugate, c = Eccentricity, and describing a Circle, on the upper Focus, whose Radius = 1, z = Arch next Aphel. to where a Line intersects it drawn from the upper Focus to the Planet; Sine of that Arch = y , Cos. = x . Whence, by Property of an Ellipsis, the Length of the intersecting Line, or Distance from the upper Focus to the Planet = $\frac{a^2 - c^2}{a + cx} = \frac{bb}{a + cx}$ (because $bb = aa - cc$) and the Planet's

Distance from the Sun = $\frac{aa + cc + 2acx}{a + cx}$; and $\frac{1}{2}bb\dot{z} + \frac{\frac{1}{2}b^2c^2y^2\dot{z}}{a + cz} =$ Fluxion of the Area described by the Planet from Aphelion, by Rays drawn to the Sun, in the lower Focus.

Resolve $\frac{1}{a + cx}$ into $\frac{1}{a^2} - \frac{2cx}{a^2}$, &c. for a Nearness. Whence the Fluxion above becomes $\frac{1}{2}b^2\dot{z} + \frac{\frac{1}{2}b^2c^2y^2\dot{z}}{a^2} - \frac{b^2c^3xy^2\dot{z}}{a^3}$, which, because $y\dot{z} = -\dot{x}$, and $x\dot{z} = \dot{y}$, reduces to $\frac{1}{2}b^2\dot{z} + \frac{b^2c^2}{2aa}x - \dot{x}y - \frac{b^2c^3y^2\dot{y}}{a^3}$, whose Fluent is

$\frac{1}{2}b^2z + \frac{b^2c^2}{2a^2}x - \frac{1}{2}xy - \frac{b^2c^3y^3}{3a^3}$ (the Area of the circular Segment next Aphelion, formed by the Curve z , its Sine y , and $1 - x$, its versed Sine from the Vertex next Aphelion, being $= \frac{1}{2}z - \frac{1}{2}xy$) which Fluent when z Arch = Semi-Circle = p , and y , its Sine, = 0, becomes $\frac{b^2p}{2} + \frac{b^2p}{2} \times \frac{c^2}{2a^2} = \frac{b^2pd}{2}$ (supposing $d = 1 +$

$\frac{c^2}{2a^2}$) the Area of the Semi-Orbit or Ellipsis from Aphelion. Whence, as $\frac{b^2pd}{2}$ Area Semi-Ellipsis : $\frac{b^2dz}{2} - \frac{b^2c^2xy}{4aa} - \frac{b^2c^3y^3}{3a^3}$ Area elliptic Sector from Aphelion :: p Length of an Arch of 180° : $z - \frac{c^2xy}{2a^2d} - \frac{2c^3y^3}{3a^3d} =$

the Length of the correspondent Arch of Mean Anomaly = A . From which Equation, $z = A + \frac{c^2xy}{2a^2d} + \frac{2c^3y^3}{3a^3d} = A + \frac{c^2}{4a^2d} \times \text{Sin. } 2z + \frac{2c^3}{3a^3d} \times \text{Sin. } z$ (because xy , or Cos. $z \times \text{Sin. } z = \frac{1}{2} \text{S. } 2z$). Here, for another Nearness

Different METHODS of solving the KEPLERIAN PROBLEM.

or Approximation, you may consider the *two last* Terms of the Equation, viz. $\frac{c^2}{4a^2d} \times \text{Sin. } 2z + \frac{2c^3}{3a^3d} \times \text{Sin. } z$ of small Value in Respect of the Arch A, and therefore make $z=A$, being nearly so. Whence, for $\text{Sin. } z$ and $\text{Sin. } 2z$, substituting $\text{Sin. } A$ and $\text{Sin. } 2A$, and $z=A + \frac{cc}{4a^2d} \times \text{Sin. } 2A + \frac{2c^3}{3a^3d} \times \text{Sin. } A$, where z the Arch round the Upper Focus, or Angle from Aphelion, is now not to be considered equal to A, but to $A +$ its Correction, by the Quantities $\frac{cc}{4a^2d} \times \text{Sin. } 2A + \frac{2c^3}{3a^3d} \times \text{Sin. } A$, which Quantities are reduced to Seconds of a Degree, by saying

As 3,4159 &c. (Length of an Arch of 180° , Radius being 1) is to 648000 Seconds in that Arch, so is $\frac{cc}{4a^2d} \times \text{Sin. } 2A + \frac{2c^3}{3a^3d} \times \text{Sin. } A$ (the Arch of Correction) to $51566 \times \frac{c^2}{a^2d} \times \text{Sin. } 2A + 137510 \times \frac{c^3}{a^3d} \times$

$\text{Sin. } A$ = the Number of Seconds to be added to the given Mean Anomaly, for the correct Angle of the Planet from Aphelion, at the Upper Focus, which Correction serves in the Planetary Orbits of small Eccentricity, Saturn, Jupiter, the Earth, Venus, and the Moon, but not correctly in the Orbits of Mars and Mercury, while our universal Rule, Page 99, correctly serves for all Orbits whatsoever.

To determine the Value of the foregoing Correction of the Mean Anomaly (according to Ward's Hypothesis) so that the $\angle$ at the Upper Focus of a Planet's Place, from Aphelion, may be truly known according to the above limited Correction?

LOGARITHMICALLY, from the foregoing Expressions.

$$4.7123634 + 2 \text{ Log. } \frac{c}{a} - \text{Log. } d + \text{Log. Sin. } 2A = \text{Log. 1st Term.}$$

$$5.1383343 + 3 \text{ Log. } \frac{c}{a} - \text{Log. } d + 3 \text{ Log. Sin. } A = \text{Log. 2d Term.}$$

Here a , the Sem. Transf. being considered = 1, the Log. of $\frac{c}{a} = \text{Log. } c$ of Eccentricity, and $d = 1 + \frac{cc}{2aa} = 1 + \frac{cc}{2}$, and its Log. is small, and may be rejected or taken at a mean Quantity, for all the Planetary Orbits, except those of Mars and Mercury.

Suppose $c=.06$, then $d=1.0018$, whose Log. = 0.0007810.

Hence, $4.71148 + 2 \text{ Log. } c + \text{Log. Sin. } 2A = \text{Log. 1st Term in Seconds.}$

$3 \times 1.71252 + \text{Log. } c + \text{Log. Sin. } A = \text{Log. 2d Term in Seconds.}$

IN WORDS.

1. To the Sum of the constant Log. 4.71148, and twice the Log. of Eccentricity in Parts of Unity, the Sem. Transverse, add the Log. Sine of twice the Mean Anomaly, and the Sum (rejecting Radius) will be the Logarithm of the first Term in Seconds, to be added to, or subtracted from, the Mean Anomaly, according as it is less or greater than 90° Degrees.

2. To the Sum of the constant Log. 1.71252, and the Log. of Eccentricity, add the Log. Sine of the Mean Anomaly, the Sum (rejecting Radius) being tripled, will be the Log. of the second Term in Seconds, to be always added to the Mean Anomaly.

N. B. The Mean Anomaly is here always reckoned from and to Aphelion, wherein the Seconds may be omitted in computing the Corrections; which being made, Say, as the Aphelion Dist. is to the Perih. Dist. so is the Tangent of half the corrected Mean Anomaly, or true Angle of the Planet at the Upper Focus, from Aphelion, to the Tangent of half the true Anomaly, or Angle, at the lower Focus of a Planet, from Aphelion, required.

EXAMPLE I.

The Eccentricity of the Orbit of Jupiter, being 0.048219 (according to Dr. Halley) and the Mean Anomaly of that Planet 75° , required the true Anomaly?

1st Term.

4.71148 Const. Log.

7.36643 twice Log. Eccy.

2.07791 Sum.

9.69897 Log. S. twice An.

1.77688 . . $59'' 82''$

2d Term.

1.71252 Const. Log.

8.68322 Log. Eccentricity.

0.39574 Sum.

9.98494 Log. Sine Anom.

0.38068

tripled, 1.14204 . . $13'' 86''$

59.82 +

Correction 73.68 +

Hence the $\angle$ of the Planet at the Upper Focus =

$75^\circ 1' 13'' 40'''$

$\frac{1}{2} 37 30 36 50$. . Tang. . . 9.8851410

Const. Log. Orb. 9.9580851

34 52 33 53

Tang. 9.8432261

Doubled 69 45 7 46 the true Anomaly.

75 00 00 Mean Anomaly.

5 14 52 54 True Equation, agreeing with Dr.

Halley's Tables.

Different METHODS of solving the KEPLERIAN PROBLEM.

By Bullialdus's Correction.		Log.
As Sem. Conj. $= \sqrt{1.048219 \times 0.951781}$		0.0005554 co.
To Sem. Transf. $= 1$		0.0000000
So Tang. 75° Mean Anom.		10.5719475
To Tang. 75° $1' 5'' 45'''$ $\angle$ at Upper Focus		10.5725029
$\frac{1}{2}$. . . 37 $30' 33''$. . . Tang. . .		9.8851243
	Conf. Log. Orb.	9.9580851
$34^\circ 52' 30''$. . . Tang. . .		9.8432094
Doubled 69 $45'$ $0''$ The true Anom. nearly.		
75 $00'$ $0''$ M. Anomaly.		
5 $15'$ $0''$ Equation, nearly.		
5 $14'$ $52''$ Above.		
+ $8''$ Error.		

Here it is observable that Bullialdus's is a near Correction.

EXAMPLE II.

Required the true Anom. from the Mean 120° , in the same Orbit?		1st Term.	2d Term.
4.71148	conf. Log.	1.71252	conf. Log.
7.36643	tw. Log. Eccy.	8.68322	Log. Eccy.
2.07791	co. L. for 1. T.	0.39574	conf. Lo. for 2d Ter.
9.93753	Lo. S. 2 An. $=$	9.93755	Log. S. An. $= 120^\circ$.
2.01544	L. $103'' 62''$	0.33329	
		9.95 + trip. 0.99937	Log. $9'' 95''$
- $1' 33'' 40'' = 93.67$ Correction.			
M.A. $120^\circ 0' 00''$			

119 $58' 26''$ the correct $\angle$ at the Upper Focus, from Aphel.	
half 59 $59' 13''$. . . Tangent . . .	10.2383328
	Conf. Log. Orb. 9.9580851
57 $32' 10''$. . . Tangent . . .	10.1964179
doubl. 115 $4' 20''$ The true Anom.	
120 $0' 00''$ M. Anom.	

$4^\circ 55' 40''$ The correct Equation, agreeing with Dr. Halley's Tables.

By Bullialdus's Correction.

As Sem. Conjugate of the Orbit . . .		Log.
To Sem. Transf. 1		0.0005554 co.
So Tangent M. Anom. 120° or T. 60° .		0.0000000
		10.2385606
To Tang. $119^\circ 58' 6''$ or $60^\circ 1' 54''$. . .		10.2391160
$\angle$ at the Upper Focus, nearly.		
Half . . . 59 $59' 3''$. . . Tangent . . .		10.2382834
	Conf. Log. Orb.	9.9580851
Doubled 57 $32' 0''$. . . Tang. 10.1963685		
115 $4' 0''$ True Anom. nearly.		
120 $0' 0''$ M. Anom.		
4 $56'$ $0''$ Equation nearly		
4 $55' 40''$ Correct Equation from above.		

+ $20''$ Error, by Bullialdus.

EXAMPLE III.

Required the Correction of 140° Mean Anomaly of the Moon's Orbit; the Eccentricity being greatest, or 0.066777? Also to find the true Equation and Anomaly?

1. Term.	2. Term.
4.71148 Const. Log.	1.71252 Const. Log.
7.64925 twice Log. Eccentricity.	8.82463 Log. Eccy.
2.36073 Const. Log. for 1st Term	0.53715 Con. Lo. for 2d Ter.
9.99335 Log. S. 2. An. $= 280^\circ$ or 80° .	9.80806 Log. S. An. $= 140^\circ$.
2.35408 Log. $226'' = 3' 36''$	0.34521
2 + tripled 1.03563 Log. $2''$ fere +	

$3' 34''$ Correction, agreeing with Halley's Tables, pro expediendo calculo, &c.

$\frac{1}{2}$ M. Anom. $70^\circ 0' 00''$
1 47 Correction

$\frac{1}{2}$ $\angle$ at Upper Focus 69 $58' 13''$. Tangent .	Log.
	10.4382336
	Log. for Eq. $\angle$ Cent. 9.9419120

67 $22' 37'' 25'''$ Tangent	10.3801456
Doubled 134 $45' 14'' 50'''$ True Anom. $\angle$ required.	
140 $00' 00'' 00'''$ Mean Anom. $\angle$.	

- 5 14 45 10 Correct Equation $\angle$'s Center.
M. Eq. $\angle$ Center.

By p. 68. Arg. $4^\circ 20' = 140^\circ$ Anom. $4^\circ 16' 34''$
For greatest Eccentricity + 58 17 Gr. Dif. Eq. [above.
5 14 51 Eq. $\angle$ Center nearly as above.

By Bullialdus's Correction.

As Sem. Conj. $= \sqrt{1.066777 \times 0.933223}$		Log.
To Sem. Transverse $= 1$		0.0009705 co.
So Tangent M. Anom. 140° or T. 40°		0.0000000
		9.9238135
To Tang. $40^\circ 3' 47''$ or $139^\circ 56' 13''$.		9.9247840
$\frac{1}{2}$ 69 $58' 6''$ Tan. 10.4381875		
	Conf. Log. for Orb.	9.9419120

67 $22' 30''$ Tan. 10.3800995	
Doubled 134 $45' 0''$ The true Anomaly, nearly.	
140 $00' 0''$ M. Anom.	

- 5 15 $0''$ Equation, nearly.
- 5 14 45 Equation fr. above correct.

+ $15''$ Error, by Bullialdus.

EXAMPLE IV.

To find the Angle at the Upper Focus at 110° M. Anomaly in the Orbit of Mars, whose Eccentricity, according to Dr. Halley, is 0.09263936? And likewise to determine the true Anomaly, and the Equation of the Orbit.

By the Logarithmic Equations foregoing,

1st Term.	2d Term.
4.712363 Const. Log.	5.138334 Const. Log.
7.933591 2 Log. Eccy.	6.900387 3 Log. Eccy.
9.998139 co. Log. d.	9.998139 co. Log. d.
2.644093 Sum. For the Orbit $\angle$.	2.036860 Sum.
+ 9.808067 Log. S. 2 A $= 220^\circ$.	9.911957 3 Log. S. A.
2.452160 Log. $283'' 24''$	1.955817 Log. $90'' 334''$
- 4' $43'' 24''$ 1st Term.	1' $30'' 33''$
+ 1 $30' 33''$ 2d Term.	

3 12, 91 Correction.
110 $0' 00'' 00'''$ M. Anom.

109 $56' 47'' 09'''$ $\angle$ at the Upper Focus.

Different METHODS of solving the KEPLERIAN PROBLEM.

Half 54 58 23.5 . Tangent	Log ^s .
Conf. Log. Orb.	10.1543409
	9.9189890
49 48 51.5 . Tangent	10.0733299
Doubled 99 37 43 True Anomaly.	
110 00 00 M. Anom.	
Dif. —10 22 17 Equation Orbit.	
By Dr. Halley's Tab. —10 22 15 Equation δ for 3° 20' M. An.	
+2'' Error.	
By Bullialdus's Correction.	Log ^s .
As Sem. Conj. $\sqrt{1.092639 \times .907361}$.	0.0018716 co.
To Sem. Transf. 1.	0.0000000
So Tang. M. Anom. 110° or 70° its Supp ^t .	10.4389341
To Tan. 109° 55' 15" or its Supp ^t 70° 4' 15" [∠ at the Upper Focus, nearly.]	10.4408057
Half 54° 57' 37" . Tangent . . .	Log.
Conf. Log. Orb.	10.1541348
	9.9189890
49 48 3 . Tangent . . .	10.0731238
Doubled 99 36 6 True Anomaly, nearly.	
110 00 0 M. Anom.	
Dif. —10 23 54 Equation.	
—10 22 15 Equation by Dr. Halley's Tables.	
+1 39 Error, by Bullialdus.	
By RULE. P. 88.	Log ^s .
As Rad.	10.0000000
To Cos. M. An. or S. 20°	9.5340517
So Eccy + its $\frac{1}{2}$ = .1157992 . . .	9.0637055
To .0396057—	8.5977572
As .9603943	0.0175505 co.
To Double Eccy . . .1852787 . . .	9.2678254
So S. M. An. 110° or 70°	9.9729858
To S. 10° 26' 40"	9.2583617
	3
Log. S. Cube 7.7750851	
Log. 3. 4771215—	
— 6' 49"	Log. S. $\frac{1}{3}$ Cube. 7.2979636
Equation 10 19 51.5	
10 22 15 Dr. Halley.	
—2 23.5 Error by this Rule, greater than by Bul-	
lialdus's Correction of Ward.	

N. B. The foregoing limited Correction of Ward's Hypothesis by two Terms only, will give the Angle at the Upper Focus of a Planet, from Aphelion, correct, in the Orbits of the Earth and Venus (of small Eccentricity) by the Use of the first Term only: The second Term in Orbits of small Eccentricity, being but of small Value, in respect of the first Term, may be then safely rejected.

Improvement being as much the Object of our Attention as Things of original Invention, the limited Correction of Ward's Hypothesis, by two Terms, you will find has not always the Readiness of the RULE, p. 88: Similar to which, we hope to see other RULES deduced, for finding, directly and truly, the EQUATIONS for all Orbits whatsoever. An Approximation of this Sort will be less tedious and perplexing than a Number of Terms as follow, employed to find the true Angle at the upper Focus, which being done, (by much Labour and Attention) then there must be a further Operation to determine the true Anomaly, and consequently the EQUATION of the Orbit; which EQUATION would be best arrived at by the direct Method above proposed.

N. B. This Rule is chiefly of Use for the Orbits of the Earth, Venus, Jupiter and Saturn, and sometimes that of the Moon, but never to be depended on for the Orbits of Mars and Mercury. The Truth or Failure of this, or any other Rule for the same Purpose, may be proved by our Universal and Correct Rule, p. 99.

EXAMPLE V.

The Mean Anomaly of Mercury being 115°. (Eccentricity .20589) to determine the Angle at the Upper Focus, True Anomaly and Equation of the Orbit?

1st Term.	2d Term.
4.712363 Conf. Log.	5.138334 Conf. Log.
8.627270 2 Log. Eccy.	7.940906 3 Log. Eccy.
9.990891 Co. Log. d.	9.990891 Co. Log. d.
3.310524 Sum. For Orb. δ	3.070131 Sum.
9.884254 Log. S. 2 An.	9.871827 3 Log. S. A.
3 214778 Log. 1639", 7—	2.941958 Log. 874", 9+
	14' 34", 9
1. Term. 27' 19", 7—	
2. Term. 14' 34", 9+	
12 44, 8— limited Correction by 2 Terms.	
115° 00 00	
114 47 15, 2 ∠ at the Upper Focus, nearly.	
Half 57 23 37, 6 . Tangent . . .	Log ^s .
Conf. Log. Orb. δ	10.1940374
	9.8185730
45° 49' 54", 2 . Tangent . . .	10.0126104
Doubled 91 39 48, 4 The True Anomaly.	
115 00 00 . M. Anomaly.	
Dif. —23 20 11, 6 Equation by the limited Correction.	
Ar. 3° 25' or 115° —23 18 35 Equation in Dr. Halley's Tables.	
+1 36, 6 Error of the limited Correction.	
By Bullialdus's Correction.	Log ^s .
As Sem. Conj. $\sqrt{1.20589 \times .79411}$.	0.0094058 co.
To Sem. Transf. 1.	0.0000000
So Tang. M. Anom. 115° or 65° . . .	10.3313275
To 114° 31' 41" ∠ at Upper Focus] 65° 28' 19" too great an Error.	10.3407333
For $\frac{1}{2}$ 57° 15' 50", 5 . Tangent . . .	10.1918731
Conf. Log. Orb. δ	9.8185730
45° 41' 20" Tan. . . .	10.0104461
doubl. 91 22 49 True Anom. . . .	Error about 19'.
	[in this Orbit, by Bullialdus.

Different METHODS of solving the KEPLERIAN PROBLEM.

Putting $e = \frac{c}{a}$, $f = \frac{b}{a}$, $S = \text{Sine } A$, $s = \text{Sine following Quantity}$, and continuing the Terms of the former

Fluxion of the elliptic Sector, by making $\frac{1}{a^2} - \frac{2cx}{a^3} + \frac{3c^2x^2}{a^4} - \frac{4c^3x^3}{a^5}$, &c. $= \frac{1}{a+cx}$, and considering the Rectangle of the Cosines of any two Angles (Radius being Unity) to be equal to half the Sum of the Cosines of the Sum and

Diff. of the Angles, then, after a prolix Process, $\frac{180 \times 60 \times 60}{3.14159} \times \frac{2}{1+4ee} \times \frac{2}{3} e^3 f S^3 - \frac{37e^5 f}{15} S^5 + \frac{1}{2} e^2 f f. 2A + \frac{5e^4 f^2}{32}$

f. 4 A. will be the Number of Seconds to be added to the mean Anomaly, that the Angle at the upper Focus, (and thence the Angle at the lower one) may be more correctly found. Whence are derived the

FOLLOWING EQUATIONS.

Let E denote the Log. of Eccentricity divided by half the greater Axis, F the Log. of half the less Axis divided by half the greater, G the Log. of the Sum of the Squares of half the greater Axis and twice the Eccentricity divided by the Square of half the greater Axis.

Take $P = 1.71277 + E + \frac{1}{3}F + \frac{1}{3}G$ } Then the Log^s. of 4 Terms
 $Q = 1.14130 + E + \frac{1}{3}F$ } in Seconds to be connected
 $R = 4.71236 + 2E + F$ } with A, will be
 $S = 4.50824$ }

1. $3 \times P + \text{Log. } S. A - \text{Log. Rad.}$
2. $5 \times Q + \text{Log. } S. A - \text{Log. Rad.}$
3. $R + \text{Log. } S. 2A - \text{Log. Rad.}$
4. $S + \text{Log. } S. 4A - \text{Log. Rad.}$

Of which 4 Terms the first is always to be added, the Second always to be subtracted: The other two to be added or subtracted according as the Sines of the respective Arguments, 2 A and 4 A are positive or negative.

The two principal of the Terms (see p. 53. of Misc. Tracts) agree with those before exhibited, being like those two Terms or Equations laid down by Sir ISAAC NEWTON in Schol. to 31. Prop. Book I. of his Principia. If a correcter and conciser Fluent of the Fluxion of an elliptic Sector, formed by Rays drawn from the Curve to the lower Focus can be obtained, a farther and more certain Correction of Ward's Hypothesis will be of Advantage, which otherwise, in numerous unconverging Terms, will be intricate and useless.

As the foregoing Correction of Ward's Hypothesis by four Terms is not perfectly exact in all Parts of the Orbit of Mercury, a farther Correction in this, and other more excentric Orbits, is as follows.

R U L E.

As half greater Axis of the Ellipsis is to the Eccentricity, so is Sine of the mean Anomaly to the Sine of another Angle first found; subtract this first found Angle from the mean Anomaly, and to the Log. Sine of the Remainder (which you may call the eccentric Anomaly) add the Sum of the Log. of the Eccentricity and the constant Log. 1.7581227 (the last being Log. of Degrees 57°.295779 = Rad.) the Sum (rejecting Radius) will be the Log. of an Angle in Degrees and decimal Parts, which being subtracted from the first found Angle, the Difference will leave a Correction to be added to the mean Anomaly.

With this corrected Anomaly, repeat the Operation (if necessary) by always adding the last Correction to the given mean Anomaly.

Then say, As the greater Sem. Axis of the Ellipsis is to the less, so is the Tangent of the corrected Anomaly, to the Tangent of the Angle at the upper Focus of the Ellipsis; whence the Angle at the lower Focus, or true Anomaly, is had by the common Proportion.

The foregoing RULE is true to a Second at one Operation, in all the Orbits of the Planets, except those of Mars and Mercury; in the former of the two last mentioned Orbits, the greatest Error is about three or four Seconds, and in the latter amounts to nearly as many Minutes, in which last Orbit three repeated Operations will be necessary for a true Correction. But, to avoid this Trouble, the following Method of Computation, (not quite so prolix as the former) is true to half a Second, even in the Orbit of Mercury, without repeating the prolix Operation!

A LESS PROLIX RULE.

Add together the Log. Cos. of the Sum of the mean Anomaly once corrected, and the excentric Anomaly, the Log. Secant of the Angle first found (by the former Rule) and twice the Log. of Eccentricity, (in all of which Angles the Seconds may be neglected) the Sum of all (deducting twice the Radius) will be the Log. of a Fraction to be added to Unity, when the Sum of the said Anomalies is between 90° and 270°, but otherwise to be subtracted therefrom.

Then the Log. of this Sum, or Remainder, being subtracted from the Log. of the first Correction in Minutes, you will have the Log. of the true Correction in Minutes, to be added to the given Mean Anomaly.

Different METHODS of Solving the KEPLERIAN PROBLEM.

EXAMPLE.

The mean Anomaly being 70° , in the Orbit of Mercury, whose Eccentricity has been assigned $= .20589$, to determine the true Anomaly, according to the LAST METHOD?

OPERATIONS.		Log ^s .
1. L. Sine M. An. 70°		9.9729858
Log. Eccentricity .20589		9.3136353
Log. $11^\circ 9' .33$ $\angle$ first found		9.2866211
$70^\circ 00' .00$ M. An.		
2. Diff. $58^\circ 50' .67$ Ecc. An: L. S.		9.9323552
Add Log. for the Orbit		1.0717583
Log. $10^\circ 5' .71 = 10^\circ .0952$		1.0041135
$\angle$ f. found $11^\circ 9' .33$		
[Operation repeats. Here the former]		
Dif. $1^\circ 3' .62$ Correction add to M. An.		
+ $70^\circ 00'$ M. Anom.		
+ $58^\circ 50'$ Excentr. Anom.		
[Here the Operation continues, when not repeated.]		
3. $\angle$ Sum $129^\circ 53'$ Cof. $39^\circ 53'$		9.8070
Add Log. Sec. $\angle$ f. found $11^\circ 9'$		10.0083
Add twice Log. Eccy		8.6272
$0.0272 \angle$ Log.		8.4425
+ 1.		
4. Sum 1.0272		0.01187
Log. Min ^s first Correction $63' .62$		1.80359
Log. Min ^s true Correction $61' .9$		1.79172
5. M. An. $70^\circ + 61' .9 = 71^\circ 1' .9$ Log. Tan.		10.4638087
Add Log. Conj. Axis		9.9905940
$\angle$ at Upper Focus, Log. Tan. $70^\circ 38' .82$		10.4544027
6. Half		35
Log. Ratio greatest and least Distances		9.8504350
		9.8185730
$\frac{1}{2}$ true An. $25^\circ 1' .02$ L. Tan.		9.6690080
Doubled $50^\circ 2' .04 = 50^\circ 2' .2''$, the true Anom. required.		

† Sum An^s.
 † Add to } 1. { between 90° and 270°
 sub. fr. } 1. { otherwise.

Compare the foregoing Method with our universal and correct Method at p. 96. directly solving the same.

The Mean Anomaly of Mercury being as before, to find the true Anomaly, by RULE p. 88.

M. An.	M. An.	4th Prop.	
under. } 90° { above. }	above. } 270° { under. }	add to. }	1.
Logarithms.			
As Rad.			10.0000000
To Cof. M. Anom. or S. 20°			9.5340517
So the Eccy. + its $\frac{1}{2} = .2573625$			9.4105452
To .08802314			8.9445969
1. + the Sem. Transf.			
Now, As 1.08802314			9.9633619 co.
To .41178 Doubl. Eccy.			9.6146652
So Sine M. Anom. 70°			9.9729858
To S. $20^\circ 49' 58''$			9.5510129
			3
		Log. S. Cube	8.6530387
		Log. 3.	4771215
		Log. S. $\frac{1}{3}$ Cube	8.1759172
Equation 19 58 25—			
M. Anom. $70^\circ 00' 00''$			
Nearly True An. $50^\circ 1' 35''$ by the Practical Rule, p. 88.			
Correctly True An. $50^\circ 2' 2''$ by universal and correct RULE, p. 96.			
— 27" Error of Rule, p. 88, in this Part of [the Orbit]			

REMARK.

The celebrated Author of the late *Miscellaneous Tracts*, Mr. T. Simpson, in the first Page of his Preface, has denominated his limited Correction (not designed to solve the Keplerian Problem, universally, as it is done at p. 96. of this Work) "a Correction of Ward's circular Hypothesis." But Ward's Hypothesis has, by all others, been considered as elliptical (and not circular) requiring elliptical Correction. And yet, in another Place, at p. 48. of his *mathematical Essays*, the same celebrated Author has examined the Error of Ward's Hypothesis (five or six Minutes in the Orbit of Mars) and corrected it for the elliptical Defect of Bullialdus's Hypothesis; which Hypothesis itself used to be a Correction of Ward's.

He has next examined, (p. 49. same Essays) a different circular Hypothesis to the antient one, for Bullialdus's Hypothesis: And, instead of there considering, as the Antients did, to answer their Purpose, the circular Hypothesis, as having double the elliptic Eccentricity, he has examined and attempted to correct the circular Hypothesis, so called of the Antients, by supposing it of equal Eccentricity with Ward's late elliptical Hypothesis, which he has (as we before observed) denominated the circular Hypothesis.

From all which Novelties, and *Miscellaneous Discoveries*, in his late *Tracts*, aforesaid, such as the Precession of the Equinoxes, double the Quantity to what Sir ISAAC NEWTON found it, from Experience; The Advantage arising, by Chance, from taking the Mean of a Number of Observations, &c. but, above all, his Deduction of all the less Lunar Equations, without the Central, from a mechanic Principle, and compounded analytical Equations, with a VOLUME promised, in Futuro, depending partly on Dr. Bradley's Observations) we may expect to see exhibited every Operation necessary to the forming of a complete Theory of the Moon's Motion, (depending more on physical Hypotheses, than on Experience and Observation) and so behold a New and glorious Light shining on ASTRONOMY! or else, by some Mischance, in using the Mean of a Number of Operations, (more laborious than difficult) we shall find the Subject involved in impenetrable Darkness!

RULES and EXAMPLES

FOR DETERMINING

Other REQUISITES of ELLIPTICAL ORBITS.

PROPOSITION I.

To determine the Distance of a Planet, or Comet, from the SUN, from the true Anomaly and Eccentricity of the Orbit given?

EXAMPLE.

Mercury's true Anomaly being $66^{\circ} 32' 24''$, and Eccentricity of his Orbit ,20589, required his Distance from the SUN?

RULE 1.

Logarithms.

As Square Radius (taking it 1) 0.0000000
To Square Sine half true An. $33^{\circ} 16' 12''$ 9.4784880
So Dist. betⁿ. Foci, or double Eccy. ,41178 9.6146652

To Fourth Proportional $P =$,1239234 9.0931532

Then, As Sum Perih. Dist. +P,9180334 co. 0.0371415

To Perih. Dist. ,794110 9.8998807

So Aphel. Dist. 1.20589 0.0813077

To Dist. $\varnothing$ from $\odot$ 1.043109 0.0183299

RULE 2.

As Radius (taking it 1) 0.0000000

To Cos. true An. $66^{\circ} 32' 24''$ 9.6000017

So the Eccentricity, ,20589 9.3136353

To Fourth Propor^l. $Q =$,0819666 8.9136370

Then, As half greater Axis—Q,9180334 co. 0.0371415

To Perih. Dist. ,794110 9.8998807

So Aphel. Dist. 1.20589 0.0813077

To Dist. $\varnothing$ from $\odot$ 1.043109 0.0183299

N. B. The First Term in the last Proportion is always half greater Axis $\pm Q$.

Sig^a. true Anom^a.

For $\begin{cases} 3. & 4. & 5. & 6. & 7. & 8. & \} + \\ 0. & 1. & 2. & 9. & 10. & 11. & \} - \end{cases} Q +$ half greater Axis is the First Term.

PROPOSITION II.

To determine the Eccentric Anomaly of a Planet, or Comet, from the true Anomaly and Eccentricity given?

EXAMPLE.

Mercury's true Anomaly is given $66^{\circ} 32' 24''$, and Eccentricity of the Orbit ,20589, required the Eccentric Anomaly?

RULE 3.

Logarithms.

As $\sqrt{\text{Perih. Dist. ,79411}}$ co. 0.0500596

To $\sqrt{\text{Aphel. Dist. 1.20589}}$ 0.0406538

So Tang. half true Anom. $\varnothing 33^{\circ} 16' 12''$ 9.8169886

To Tang. half eccentric An. $\varnothing 38^{\circ} 57' 25''$ 9.9077020

Doubled . 77 54 50 excentric Anomaly required.

PROPOSITION III.

To determine the Mean Anomaly and Equation correspondent of a Planet, or Comet, from the Excentric Anomaly and Eccentricity of the Orbit given?

EXAMPLE.

Excentric Anomaly of Mercury is given $77^{\circ} 54' 50''$, and Eccentricity of his Orbit ,20589, required his Mean Anomaly and Equation of his Orbit?

Logarithms.

Radius $57^{\circ} 32' 57''$ 1.7581227

Eccentricity ,20589 9.3136353

See p. 96. Const. Log. for Orbit 1.0717580

RULES for finding the REQUISITES of ELLIPTICAL ORBITS.

RULE 4.

Excentric Anom.	77° 54' 50"	Const. Log. Orbit	1.0717580
		Log. Sine	9.9902651+
Dif. bet. cor.			
Ex. & M. An.	+ 11 32 6 . 11° 53' 51"		1.0620231

Sum, M. Anom. 89 26 56 [By Dr. Halley's Tables,
True Anom. . 66 32 24— [Eqⁿ. 22° 54' 30"

Our Equation & 22 54 32 agreeing with Dr. Halley.

PROPOSITION IV.

To determine the Distance of a Planet, or Comet, from the SUN, from the excentric Anomaly, true Anomaly, and Eccentricity of the Orbit given?

EXAMPLE.

The true Anomaly of Mercury being 66° 32' 24", Excentric Anomaly 77° 54' 50", and Eccentricity of the Orbit .20589, required his Distance from the SUN?

RULE 5.

As Sine true An.	66° 32' 24"	co.	0.0374705
To Sine Exc. An.	77 54 50		0.9902651
So Sem. Conjugate =			
√ Ap. Dist. x Per. Dist. =	√ 1.20589 x .79411		9.9905943

To Dist. & from ☉ . . 1.043109 . . 0.0183299

The same as before.

OTHERWISE.

RULE 6.

As Sine half true Anom.	33° 16' 12"	co.	0.2607560
To Sine ½ Excent. An.	38 57 25		9.7984685
So √ Perih. Dist. .79411			9.9499404

To √ Dist. & from ☉ . . 0.0091649

The Dist. & from ☉ req^d. 1.043109 doubled 0.0183298
The same as before.

Sides Δ in same Ratios.

98923 = ☉ &, from Sun to Mercury
41178 = ☉ F, from Sun to Upper Focus
101077 = & F, from Mercury to Upper Focus

half 241178 = Sum

120589 = H = Half
19512 = H — & F (opposite to ∠ required)
79411 = H — ☉ F
21666 = H — ☉ &

N. B. When the Distances from the Sun are determined by the foregoing Rules in any of the planetary Orbits, a Table of Logarithms may be composed to these Distances, at Sight, from Gardiner's Tables, to six Places of Figures.

PROPOSITION V.

To determine the greatest Equation of a Planet, or Comet, or greatest Difference between its mean and true Anomalies?

EXAMPLE.

The Sem. Transverse of Mercury's Orbit being 1, and Eccentricity .20589, required the greatest Equation?

RULE 7.

From the lower Focus, as a Center with Radius, a mean Proportional between the Sem. Transverse and Sem. Conjugate of the Orbit, describe an Arch to intersect the elliptical Curve, and the Point of that Intersection will be the Planet's true Place, at the Time of the greatest Equation?

Hence, √ Ap. Dist. x Perih. Dist. 1.20589 x .79411 = .9785754 = Semi Conjugate, √ .9785754 x 1 = .98923 a mean Proportional betwixt both Axes, being the Distance of Mercury from the Sun, at the Time of his greatest Equation; Double Eccentricity = .41178 Dist. Foci; and 1.01077, the Distance from the Planet to the upper Focus (by the Nature of an Ellipsis, the two Distances of a Planet from the upper and lower Focus, being always equal to the Transverse Axis of its Orbit) whence, from these three Sides of a given Triangle, the Angle of true Anomaly, at the Sun, or lower Focus, is determined to be 81° 4' 6", 8, as follows.

Sides Δ

. 98923
. 41178
. 1.01077

Sum 2.41178

Ind. Ind.

120589 = H = Half	co. Log.	4.9186923	9	1.20589
19512 = H — & F (opposite to ∠ required)	co. Log.	5.7096982	0	.19512
79411 = H — ☉ F	Log.	4.8998807	9	.79411
21666 = H — ☉ &	Log.	4.3357787	9	.21666

Half 19.8640499 | 29 deduct Rad. fr. [these Indices
40° 32' 3", 4 . . Tang. 9.9320249
Doubled 81 4 6, 8 . . The ∠ at ☉, opposite & F, being the ∠ [of true Anomaly, required.

To determine an Angle opposite to a given Side in any right-lined Triangle, whose three Sides are given?

RULES for finding the REQUISITES of ELLIPTICAL ORBITS.

RULE 8.

• • Add the three Sides together, and take $\frac{1}{2}$ the Sum, and all the Differences betwixt that half Sum and each of the 3 Sides, according to the foregoing Order. Then set down under each other, the *arithmetical Complements of the Logarithms* of the half Sum, and of that Difference between the half Sum and the Side opposite to the Angle sought; then set down (under those co. Log^s.) the *Logarithms* of the Differences betwixt the said half Sum and the other two Sides; the half Sum of which will be the Tangent of half the Angle required.

Now (by Rule 3.)

	Log ^s .
As $\sqrt{\text{Perih. Dist. } .79411}$ co.	0.0500596
To $\sqrt{\text{Aphel. Dist. } 1,20589}$	0.0406538
So Tang. $\frac{1}{2}$ true An. (just found) $40^{\circ} 32' 3'' .4$	9.9320249
To Tang. $\frac{1}{2}$ excentric An. $\varnothing$ at the Time	
of greatest Equation . . . $46^{\circ} 29' 57'' .2$	10.0227383
Doubled . . . $92 59 54 .4$ the excent.	
[Anom. at the Time of the greatest Equation.	
(By Rule 4.) Const. Log. for Orbit	1.0717580
Cor. ex. An. $92^{\circ} 59' 54'' .4$ } . Log. Sine	9.9994050 +
Supplement $87 0 5 6$ }	
Dif. bet. M. and excent.]	
An. . $+11 46 49 .7$. $11^{\circ} 78' 048$.	1.0711630

Sum, M. An. $104 46 44 .1$

Tr. An. $-81 4 6 .8$

Dif the Gr. 2

EQUATION $23 42 37 .3$ required, agreeing with Dr. Halley's Tables; our Equation being greatest but by a Second.

PROPOSITION VI.

To determine the Eccentricity and other Dimensions of a Planet's or Comet's Orbit from the greatest Equation given?

EXAMPLE.

The greatest Equation of Mercury's Orbit being $23^{\circ} 42' 37''$, required his Eccentricity?

RULE 9.

By a Table of natural Sines, the Sine of half the greatest Equation of a Planetary Orbit (*Radius or Semi-Transverse being Unity*) will be very nearly equal to the Eccentricity, in Orbits not much excentric; but in Orbits more or very excentric, as in that of Mercury, and the Orbits of Comets, the natural Sine of half the greatest Equation will be somewhat less than the true Eccentricity. So that the Eccentricity thus determined, for very excentric Orbits, must be augmented, to compute the greatest Equation from thence, by the preceding Example, to correspond with the greatest Equation given. For the greatest Equation computed always augments with the Increase of Eccentricity.

NOW, from the near Eccentricity, found by a Table of natural Sines (*corresponding to the Semi-Transverse = 1*) determine the less Semi-Axis, it being $\sqrt{\text{Ap. Dist.} \times \text{Perih. Dist.}}$; and then determine the mean Proportional betwixt the two Semi-Axes, which is the Distance of the Planet (or Comet) at that Time, of the greatest Equation, from the SUN.

THEN, Three Sides of a plain Triangle being given, (*the Distance of the Planet from the Sun and upper Focus, and Distance of the Foci*) determine the Angle of true A-

nomaly, (by Rule 8, preceeding) then the eccentric Anomaly, and lastly the mean Anomaly, corresponding; from whence the greatest Equation will be determined, (*being the Difference between the true and mean Anomalies*) nearly, according to the assumed Eccentricity.

N O W, Say, as the computed greatest Equation, is to the greatest Equation given, so is the assumed Eccentricity to the true Eccentricity required.

N. B. The greatest Equations and Eccentricities, nearly assumed, are in a very near Proportion to one another.

OTHERWISE, for the very ECCENTRIC ORBITS of COMETS.

RULE 10.

MAKE two Assumptions of Eccentricities, one less and the other greater than the true, (which you may easily do by having Regard to the natural Sine of $\frac{1}{2}$ greatest Equation) and compute the greatest Equations, by the foregoing Rules, to both Eccentricities assumed. Then take their Excess, and Defect, by comparing them with the greatest Equation given; taking also the Differences of Eccentricities assumed.

Then say, As the Sum of Excess and Defect of the computed greatest Equation, above and under the given greatest Equation, is to the Difference of Eccentricities assumed, so is the Defect of the computed greatest Equation under the given greatest Equation, to a correspondent Difference of Eccentricity, to be added to the defective Eccentricity, for the true Eccentricity required.

N. B. IF BOTH the computed greatest Equations should happen to come out too little, or less than the given greatest Equation, you must subtract their Defects from one another, and say,

As the Difference of Defects of Equations, is to the Difference of assumed Eccentricities, so is the greater Defect of Equation, to the Difference of Eccentricity to be added to the less or greater defective Eccentricity, respectively assumed, for the true Eccentricity, required. According to THE METHOD of TRIAL and ERROR.

* • NOW, $\frac{1}{2}$ greatest Equation given of $\varnothing$'s Orbit = $11^{\circ} 51' 18'' .4$, whose natural Sine is .205438 (by TABLE of SINES) for the Eccentricity of $\varnothing$'s Orbit, whereof you may observe that the 3 first Figures, 205 are correct (in Respect of the Semi-Transverse = 1) by what has been given and determined; but, 206 too much. Therefore, Let the Eccentricity be assumed, 2055 about one ten-thousandth Part greater than the Sine of $\frac{1}{2}$ greatest Equation. In which Assumption, from $\frac{1}{2}$ greatest Equation, you may sometimes reject Seconds, and take the Sine of the next whole Minutes greater, as of $11^{\circ} 52'$, when the Orbit, as at present, is much excentric, which Sine would be = .2056349, or, 2056 at four Places.

Eccentricity assumed = .2055, whence $\sqrt{1,2055 \times .7945} = \sqrt{\text{Aph. Dist.} \times \text{Perih. Dist.}} = \text{Sem. Conjugate}$; and $\sqrt{1.2055 \times .7945 \times 1}$ (for 1 equal Sem. Transverse) = $1.2055 \times .7945$ the mean Proportional between the Axes = .989271 = direct Dist. $\varnothing$ fr. $\odot$, hence, 1,010729, = Dist. $\varnothing$ fr. upper Focus, F; and Dist. Foci = doub. Eccentricity = .4110, being the three Sides of a given Triangle to find the $\angle$ opposite to $\varnothing$ F, or $\angle \varnothing \odot F$, the true Anomaly of $\varnothing$ at that Time?

RULES for finding the REQUISITES of ELLIPTICAL ORBITS.

Sides Δ .	
(By Rule 8.) ,989271 = $\odot \frac{1}{2}$	fr. Sun to Mercury
,411000 = $\odot F$	fr. Sun to upper Focus
1,010729 = $\frac{1}{2} F$	fr. Mercury to upper Focus.
2,411000 = Sum.	
$\frac{1}{2} \dots 1,205500$ = Half = H	co. Log. 9.9188328
,194771 = H — $\frac{1}{2} F$ (opposite to Angle required)	co. Log. 0.7104757
,794500 = H — $\odot F$	Log. 9.9000939
,216229 = H — $\odot \frac{1}{2}$	Log. 9.3349139
$\frac{1}{2} \dots$	19.8643163
40° 32' 34",6	Tan. 9.9321581
Doubled 81 5 9,2 the $\angle$ at $\odot$, or true Anomaly.	
NOW (by Rule 3.) As $\sqrt{\text{Perih. Dist.}}, 7945$	co. 0.0499530
To $\sqrt{\text{Aph. Dist.}}, 1,2055$	co. 0.0405836
So Tan. $\frac{1}{2}$ true An. found 40° 32' 34",6	9.9321581
To Tan. $\frac{1}{2}$ eccen. An. $\frac{1}{2}$ at $\frac{1}{2}$	
Time of gr. Equation $\frac{1}{2}$ 46° 29' 47"	10.0226947
Doubled 92 59 34, Eccen. An. at Time of greatest Equation.	
(By Rule 4.) Radius 57°, 2957779	1.7581227
Eccentricity, 2055	9.3128118
See p. 96. Const. Log. for the Orbit	1.0709345
Correct eccentric An. 92° 59' 34",2	
Supplement 87 0 26 } ... Log. Sine	9.9994073 +
Dif. bet. M. and Ecc. Anomalies + 11 45 29,6 . 11°, 75823	1.0703418
Sum, Mean Anom. 104 45 3,6	
True Anom. 81 5 9,2 —	
Dif. The computed gr. Equation . . 23 39 54,4 fr. assumed Eccentricity.	
True gr. Equation . . 23 42 37,3 fr. correct Eccentricity.	
NOW, As 23° 39' 54" : 23° 42' 37" } Defect 2' 43" of Equation fr. assumed Eccentricity.	
Or 85194" : 85357" }	assumed Eccentricity, 2055 : 20589 the true Eccentricity, req ^d .

* * * Hence, it is evident, that by a careful and near Assumption of the Eccentricity, from a TABLE of natural Sines, a very great Accuracy, from Computation, will result in determining the true Eccentricity of the Orbit from the greatest Equation given; and the Trouble of a Second Assumption and Operation, for any of the planetary Orbits, will be thereby avoided.

REMARKS.

REMARK I. Some of the French Authors make the greatest Equation of Mercury to be not less than 24°, and Mons. De la Caille, particularly, in one Place of his Elements of Astronomy, p. 83, makes it to be 24°. 3'. 5". with Eccentricity 211165 to 1011276 the greater Semi-axis, being as 20881 Eccentricity to 1 the Semi-transverse. Street makes the Eccentricity of $\frac{1}{2}$ 7970 to 38710 the Semi-transverse, the same as 20589 to 1, according to Dr. Halley; Street's Mercurial Astronomy being reckoned very correct.

To determine the Principal Diameters of the Orbits of Planets?

They are to be taken in the Subsesquialte (or $\frac{2}{3}$ Power) Proportion of the Periodic Times (by Prop. 15. B. I. Newtonian Principia) and then to be severally augmented in the Proportion of the Sum of the Masses of Matter in the Sun and each Planet to the First of Two Mean Proportionals betwixt that Sum and the Quantity of Matter in the Sun, by Prop. 60. Book I. Principia. They revolve in periodic Times that are in the Sefquialte Proportion of their Distance from the Primary.

RULES

RULES for finding the REQUISITES of ELLIPTICAL ORBITS.

The Eccentricities and Apellions of the Orbits of Planets are determined by Prop. 18. B. I. *Newtonian Principia*. That is, from a Focus and the principal Axis given to describe elliptic Trajectories which shall pass through given Points, and touch right Lines given by Position.

REMARK II. The foregoing Method of computing the Eccentricity of a Planet from the greatest Equation, tho' indirectly, is reckoned better for determining the Dimensions of the Orbits of Planets, than the Method by known Distances and true angular Velocities; because the Distances deduced will be more effected by Errors arising in observing the true angular Velocities than the Eccentricity will be by the Errors made in the greatest Equation. For (according to Monf. De la Caille) the Relation of Mercury's Aphelion and Perihelion Distances is determined by a Comparison of an Arch of $10^{\circ} 3' 35'' \frac{1}{2}$ to another Arch of $27^{\circ} 23''$ whereof one Arch is little more than double the other. Whereas those Distances may be determined from the Comparison of an Angle of 24° the Sum of Inequalities, down to no Angle or Inequality at all: Mercury's Eccentricity causing all that Variation of Inequality from the Mean Motion, from Aphelion or Perihelion (where the Mean and true Motion is equal) to the Points of the two Mean Distances (in respect of the two Semi-axes) which Inequality, nor any Variation from the uniform Motion could happen, but by the Cause of the said Eccentricity of the Orbit.

REMARK III. To deduce, from Observation, the greatest Equation of Mercury, the Time must be determined when Mercury is in those Points of his Orbit, where the Distance (as has been observed) of that Planet from the Sun is a mean Proportional between the two Semi-axes, both before and after the Passage of Mercury through the Apfides.

This Time being when the true and mean Velocities are equal, two Instants, at the Distance of 24 Hours, must be found when Mercury shall have passed through $4^{\circ} 5' 32'' \frac{1}{2}$, by real diurnal Motion; being equal to the mean diurnal Motion of that Planet, according to Monsieur De la Caille.

Now, according to Monf. De la Caille's Observations, 1740, for Paris, $\odot$'s true diurnal Motion about 15th July, and 4th September, is found to be $4^{\circ} 5' 32'' \frac{1}{2}$, therefore interpolating his Differences, $4^{\circ} 14' 33''$; $4^{\circ} 6' 28''$; $3^{\circ} 58' 44''$; between his Observations of 13, 14, 15 and 16th of July, the Time when $4^{\circ} 5' 32'' \frac{1}{2}$ is passed over by Mercury, will be found to be between July 14th 2h 49' and July 15th 2h 49m; the Middle of which Time, will be July 14th 14h 49m, for the Instant when Mercury's true and mean Velocities were in Equality; or when that Planet was at his mean Dist. of the Semi-axes, from the Sun: (the other mean Dist. commonly so called, being the Length of the Semi-Transverse from the Sun, at the End of the Conjugate) Now interpolating Mercury's true Places on the 14, 15 and 16th of July (according to Monf. De la Caille's Observations) Mercury's true Place at the Instant of Equality of mean and true Velocity, July 14th 14h 49m, will be $5^{\circ} 22' 57'' 27''$.

Also interpolating the Differences between the Observations of Mercury's true Places, 3, 4, 5 and 6 of September, (according to the Observations of the same Author) and you will find Sept. 3d 22h 14m $\frac{1}{2}$ for the Time of his mean Dist. (of the Semi-axes) from $\odot$, when his true and mean Velocities are equal, (the Velocities being least and greatest at Aphelion and Perihelion,) and his true Place at that Time, $11^{\circ} 4^{\circ} 49' 10''$.

Now, the Difference between these two Places $5^{\circ} 22' 57'' 27''$
and $11^{\circ} 4^{\circ} 49' 10''$

will be the true Motion, or Sum of Mercury's real Velocities during the Interval of his Passage between his said mean Distances
The Time of which aforesaid Interval is $52^d 7^h 25^m \frac{1}{2}$.
During which Time Mercury would have described by uniform Motion (allowing 12^s in $87^d 23^h 15^m 32^s$)

Dif. $5^{\circ} 11' 51' 43''$

$6^{\circ} 29' 58' 33''$
 $- 5^{\circ} 11' 51' 43''$

The Difference between these two Motions $1^{\circ} 18' 6' 50''$

The Half of which is the Sum of all Mercury's Inequalities, and consequently is his greatest Equation $24^{\circ} 3' 25''$

Differing from the greatest Equation, in another Place of my Author, by $+ 26''$

Interpolation is not necessary for determining the Instant when a Planet is in its mean Distance (of the Semi-axes) if the Orbit is not very eccentric; it being then sufficient to compare its true Motion, at the Time when an Equality of its real and mean Velocities nearly happen, with the Mean Motion corresponding to that Interval.

For, on the 14th of July at Noon, $\odot$'s true Place was $5^{\circ} 14' 43' 43''$
And on the 14th of September at Noon, his Mean Place was $7^{\circ} 2' 48' 7''$

Inequalities nearly in that Interval, the Difference $1^{\circ} 18' 4' 24''$
Half of which is the greatest Equation $24^{\circ} 2' 12''$

Likewise, on July 15th at Noon, the true Place of $\odot$ $5^{\circ} 10' 37' 15''$
And September 4th at Noon, the mean Place $6^{\circ} 18' 42' 35''$

Difference, nearly the Inequalities of Motion in that Interval $1^{\circ} 18' 5' 20''$
The Half of which is the greatest Equation $24^{\circ} 2' 40''$

Which Difference, from the first greatest Equation, derived from the true Motion, is caused by the Inequality of Mercury's Velocity from the 14th to 15th of July.

But, in the planetary Orbits of small Eccentricity, the Inequality from Day to Day is hardly perceptible.

REMARK IV. The two Points of mean Motion in the former Interval, gives the Position of the Line of Apfides, those Points being equi-distant from that Line.

The middle Way between $5^{\circ} 22' 57'' 27''$
and $11^{\circ} 4^{\circ} 49' 10''$

Sum $16^{\circ} 27' 46' 37''$

Will be the Place of Mercury's Aphelion, Half $8^{\circ} 13' 53' 19''$

But the Position of the Line of Apfides, in general, may be had by determining two Points in the Orbit that have equal Velocities, which Method however is only fit for very excentric Orbits, wherein the Inequalities of Velocity, from Day to Day, are considerable enough to determine, with a Degree of Accuracy, the Time and Place, at which the Velocity would be equal to a given Quantity; without being much affected by the Errors of Observation. But, in Orbits of small Eccentricity, where the Velocities from Day to Day are nearly uniform, this Method will be useless.

PRINCIPLES of the PLANETS PLACES and MOTIONS.

The SUN'S Mean RADICAL PLACES and MOTIONS, in REVOLUTIONS SIGNS and DECIMALS.

RADICAL YEARS. Old and New Style.	Sun's mean Place from the Equinox.	Sun's mean Anomaly, from Equinox.	Sun's Apogee, from Equinox.	First * Aries, from Equinox.
Jan. 1. { 1600 } O. S.	9° 20' 12' 31" our	6° 14' 11' 32"	3° 6' 0' 59"	0° 27' 35' 10"
Jan. 1. { 1600 } N. S.	9 10 21 8	6 4 20 11	3 6 0 57	0 27 35 9
Jan. 1. { 1700 } O. S.	9 20 58 3 our	6 13 14 34	3 7 43 29	0 28 59 20
Jan. 1. { 1700 } N. S.	9 10 7 32	6 2 24 5	3 7 43 27	0 28 59 19
Our Places reduce to	Newton's by — 0' 3" Halley's . . — 0 10 Morrison's . . + 0 10 Mayer's . . — 0 8 Euler's . . — 2 12 Bradley's wanting.	by — 0' 3" + 5 10 + 1 50 + 1 8 — 23 23	by 0' 0" — 5 20 + 2 0 + 1 0 + 21 11	N. B. The Eq. forego- ing being connected with those of Motion below, and both joined with our pre- sent Places will give those by others.
Equation of Places.	N. B. Mayer's Paris Places ⊙ are but 23" less than those corresp ^t Places at Greenwich.			
TIME forward.	Mean Motion ⊙.	Mean Motion ⊙ An.	Mean Motion ⊙ Apog.	Mean Motion 1* ♈.
100 Julian Years.	100° 0' 0' 45' 32" our	99° 11' 29' 3' 2"	0° 1' 42' 30"	0° 1' 24' 10"
Our Motions in 100 Ju- lian Years, reduce, from 1600 and 1700, to	Newton's by — 12' 0" Halley's . . + 0 20 Mayer's . . + 14 40 Euler's . . + 28 0 Bradley's wanting.	by + 1' 18' 0" + 1 23 0 + 2 15 20 + 17 8 0	by — 1' 30" — 1 22 40" + 2 30 — 16 40	 — 0 50 + 0 10 + 1 40
Equation of Motion.				
Divide each Quantity above by 100, for the Equation of Motion to our Places for 1 Year, which multiplied by the Years from 1600 or 1700 will be the whole Equation of Motion, and connected with Equation of Place above will be the Equation of our present Places to others.				
4 Julian Years.	Decimals. +4° 0'.00101185185 &c. exactly +1' 49" 16" 48iv	Decimals. 4° — 0'.00126592592 exactly — 2' 16" 43" 12iv	Sign. + ,0022777777 + 4' 6" exactly.	Sign. + ,00187037037 + 3' 22" exactly.
Our				
Halley	+4° 0'.0010119753 exactly +1' 49" 17" 36iv	4° — 0'.0012350617 exactly — 2' 13" 23" 12iv	Sign. + ,002247037037 4' 2" 40" 48iv exactly	Sign. + ,00185185 &c. 3' 20" exactly.
1 Com. Year, { Our	Decimals. + 11°.9920392374 11° 99' 45' 40" 14" &c.	Decimals. + 11°.9914778627 11° 29' 44' 39" 36" &c.	Sign. + ,000561374756 1' 0" 37" 42iv &c.	Sign. + ,0004672725 50' 27" 53" &c.
or 365 Days { Halley	11°.9920392683	11°.9914778925	Sign. + ,0005613758	Sign. + ,0004626558 + 49' 57" 56"
Common Years { 1. — 14' 19" 45"	— 14' 19" 45"	— 15' 20" 23"	+ 1' 0" 37"	+ 0 50' 0" 50" 28"
2. — 28 39 31	— 28 39 31	— 30 40 46	+ 2 1 15	+ 1 40' 1 40 55
3. — 42 59 17	— 42 59 17	— 46 1 10	+ 3 1 53 [Halley.]	+ 2 50' 2 31 23 Our
After Bissextile	Our.			
Day	Sign. + ,0328549021 59' 8" 19" 45 iv	Sign. + ,032853364 59' 8" 9" 47iv	Sign. + ,000001538 9" 58iv	Sign. + ,0000012802 8" 17iv 44v
Hour	Sign. + ,0013689542 2' 27" 50" 11, &c.	Sign. + ,00136889 2' 27" 50" 11 &c.	⊙'s Apogee + his Anom. = his Longitude. There- fore, ⊙ Longitude — his Ap. = his Anomaly.	Therefore when it is Leap- Year, a Day, and Day's Mo- tion must be subtracted for the Months of Jan. and Feb. Being the New and shorter Computation: the Lp.-Yr. not taking Place till 29th of February.
Minute	Sign. + ,000022815904 2" 27" 50iv.	Sign. + ,000022814 2" 27" 50iv	N. B. The Rad. Pls. for the Beginn ^g . of Leap-Yrs are advanced a Day's Mot. +	
Second	Sign. + ,00000038026 2" 27iv 50v, &c.	Sign. + ,00000038024 2" 27iv 50v &c.	Jul. Yrs. Dec. 21073, 1707317 The same as a Revolution of ⊙ from ⊙'s An.	Jul. Yrs. Dec. 25663, 366336 Rev ⁿ ⊙ à *. Dec. 365 ^d , 256532906 [6h 9m 24s 26th 33 rd 365 ^d , 256391047 [6h 9m 12s 11th 18 th &c]
REVOLUTION.	Decimals. 365D ^s .2423006021 365 ^d 5h 48m 54s 46th 19fo. Mean Solar Year.	Decimals. 365D ^s .259727107 365 ^d 6h 14m 0s 25th 19fo. Anomalistical Year.	Sydereal Year, or Revol ⁿ ⊙ from fixed Star.	Day. + ,026131624 36m 25s 46th 20fo
Our				
Halley	Decimals. 365D ^s .2422996627 365 ^d 5h 48m 54s 41th 27fo	Decimals. 365D ^s .259398291 365 ^d 6h 13m 32s 0th 43fo	Our Sydereal Year greater than our Solar by 20m 29s 40th 14fo &c.	Day. + ,025564188 36m 48s 44th &c. *s advance, in 4 Yrs.
4 Years Solar, ano- malistical, or Syde- real, retard or advac of 4 Julian Years, { Our	Day. — ,030797591 44m 20s 54th 42fo	Day. + ,032908428 56m 1s 41th 17fo 26ft	Dr. Halley's Sydereal greater than his Solar by 20m 17s 30th &c.	Dr. Halley's Anomalistical greater than his Solar Year by 24m 37s 19th 16fo &c.
Hall.	Day. — ,0308013493 44m 21s 14th 11fo Seasons go back, in 4 Yrs.	Day. + ,037593164 54m 8s 2th 57fo An ^d advance in 4 Yrs.	Our Anomalistical Yr great- er than our Solar by 25m 5s 39th &c.	
according to —, or +, as to Month-days.		Vernal Equinox. Mar. 9 ^d 2h 16m 44s, 1700 Mar. 8 15 54 30, O. S. Or 19, N. S. Halley [1756]		
Com. Years { 1. + 5h 48m 54s	+ 5h 48m 54s			
2. + 11 37 49	+ 11 37 49			
3. + 17 26 44	+ 17 26 44			
After Bissextile	Seasons go forward each Yr.			

PRINCIPLES and ELEMENTS of the PLANETS PLACES and MOTIONS, according to late IMPROVEMENTS.

The SUN'S Mean MOTIONS continued, in Signs and DECIMALS.				
TIME forward.	Mean Mot. Sun.	Mean Mot. Sun An.	Mean Motion Sun Ap.	Mean Mot. 1 st V.
ANOMALISTICAL YEAR.	Decimals.		Sign.	Sign.
365 ^d 6 ^h 14 ^m 0 ^s } Our	+12 ^s .00057154609	+12 ^s	+0,0005694595	+0,0004676049
25 th 19 th 10 th } Halley	12 ^s .0005617737	+12 ^s	+0,0005617737	+0,00046309122
LUNATION. Dec.	Sign.	Sign. 36292	Sign.	Sign.
29 ^d 53 ^h 50 ^m 8 ^s 11 ⁹ } Our	+0,97022467137	+0,97017923097	+0,0004542100	+0,0003780506
+12 ^h 44 ^m 3 ^s 2 th } Halley	29 ^d 6 ^h 24 ^m 1 ^s &c.	29 ^d 6 ^h 19 ^m 21 ^s &c.	4 ^h 54 ^m 19 ^s &c.	4 ^h 4 ^m 58 ^s &c.
Decimals.	Sign.	Sign.	Sign.	
29 ^d 53 ^h 50 ^m 8 ^s 11 ⁹ } Halley	+0,9702246715135	+0,970179230808	+0,000454184	
+12 ^h 44 ^m 3 ^s 2 th } Halley	29 ^d 6 ^h 24 ^m 1 ^s &c.	29 ^d 6 ^h 19 ^m 21 ^s &c.	4 ^h 54 ^m 19 ^s &c.	

The MOON'S Mean RADICAL PLACES and MOTIONS, in REVOLUTIONS, SIGNS and DECIMALS.

Radical YEARS, Old and New Style.	Moon's mean Places from the Equinox.	M. Pl. Moon's mean Anomaly, fr. Equinox.	M. Pl. Moon's Apogee, fr. Equinox.	M. Pl. Moon's ascending Node fr. Equinox.
Jan. 1. 1600 O. S.	7 ^s 28 ^m 7 ^s 42 ^m	1 ^s 19 ^m 41 ^s 47 ^m	6 ^s 8 ^m 25 ^s 55 ^m	10 ^s 0 ^m 55 ^s 28 ^m
Jan. 1. 1600 N. S.	3 16 21 52	9 9 2 48	6 7 19 4	10 1 27 14
Jan. 1. 1700 O. S.	6 ^s 5 ^m 58 ^s 7 ^m	8 ^s 8 ^m 20 ^s 57 ^m	9 ^s 27 ^m 37 ^s 10 ^m	8 ^s 16 ^m 44 ^s 13 ^m
Jan. 1. 1700 N. S.	1 11 1 42	3 14 38 4	9 26 23 38	5 17 19 10
Our Places reduce to . . .	Newton's by -0 ^s 10 ^m Halley's . . -1 10 Mayer's . . 0 0 Euler's . . +0 20 Bradley's wanting.	by . . -3 ^s 10 ^m . . -4 10 . . +2 0 . . -1 14 . . +10 50	by . . +3 ^s 0 ^m . . +3 0 . . -2 0 . . +0 54 . . -10 30	by . . -0 ^s 10 ^m . . -1 10 +4 1 . . +1 50
Equation of Places.	N.B. Mayer's Paris-Places were reduced to Greenwich.			
TIME forward.	Mean Motion Moon.	Mean Motion D's Anom.	Mean Motion D's Apogee.	Mean Motion D's Node.
100 Julian Years.	1336 ^r 10 ^s 7 ^m 50 ^s 25 ^m	1325 ^r 6 ^s 18 ^m 39 ^s 10 ^m	11 ^r 3 ^s 19 ^m 11 ^s 15 ^m	5 ^r 4 ^s 14 ^m 11 ^s 15 ^m
Our Motions in 100 Julian Years from 1600 reduce to . . .	Newton's . . 0 ^s 0 ^m Halley's . . 0 0 Mayer's . . 0 0 Euler's . . +1 55 Bradley's wanting.	by . . . 0 ^s 0 ^m . . . 0 0 . . . 0 0 . . . +1 55 . . . 0 0	by . . . 0 ^s 0 ^m . . . 0 0 . . . 0 0 . . . 0 0 . . . 0 0	. . . 0 ^s 0 ^m . . . 0 0 . . . 0 0 . . . 0 0 . . . 0 0
Equation of Motion.				

Divide any Quantity, variable from our own Motⁿ, by 100, for the Eq. of Mot. to our Pls for 1 Yr, which X'd for the Yrs from 1600 to 1700, will be the whole Equat. of Motion, and connected with Equat. of Places above will be the absolute Eqⁿ. of our Present Places to those of others

Julian Years . . .	Decimals.	Decimals.	Decimals.	Decimals.
	+53 ^r 5 ^s 69 ^m 45 ^s 37 ^m 03 ^s	+53 ^r 05 ^s 26 ^m 48 ^s 70 ^m 37 ^s 03 ^s	+5 ^s 42 ^m 55 ^s 33 ^m 33 ^s &c.	-2 ^s 57 ^m 89 ^s 16 ^m 66 ^s &c.
	+5 ^s 20 ^m 42 ^s 49 ^m exactly.	+0 ^s 7 ^m 56 ^s 46 ^m exactly.	+5 ^s 12 ^m 46 ^s 3 ^m exactly.	-2 ^s 17 ^m 22 ^s 3 ^m exactly.
365 Days	Decimals.	Decimals.	Decimals.	Sign.
	+13 ^r 4 ^s 31 ^m 28 ^s 10 ^m 13 ^s 14 ^m	+13 ^r 2 ^s 95 ^m 73 ^s 42 ^m 7	+1 ^s 35 ^m 54 ^s 67 ^m 43 ^s 14 ^m	-6442878735
	13 ^r 4 ^s 9 ^m 23 ^s 3 ^m 29 ^s &c.	13 ^r 2 ^s 28 ^m 43 ^s 13 ^m &c.	1 ^s 10 ^m 39 ^s 50 ^m 29 ^s &c.	19 ^m 19 ^s 43 ^m 11 ^s &c.
Com. Years, { 1	+4 ^s 9 ^m 23 ^s 3 ^m	+2 ^s 28 ^m 43 ^s 13 ^m	+1 ^s 10 ^m 39 ^s 50 ^m	-0 ^s 19 ^m 19 ^s 43 ^m
{ 2	+8 18 46 7	+5 27 26 26	+2 21 19 41	-1 8 39 26
{ 3	+0 28 9 10	+8 26 9 39	+4 1 59 32	-1 27 59 10
Day.	Sign.	Sign.	Sign.	Sign.
	+4392131784	+435499569	+0037136094	-0017651722
	13 ^r 10 ^s 35 ^m 1 ^s &c.	13 ^r 3 ^s 53 ^m 57 ^s &c.	6 ^s 41 ^m 4 ^s &c.	3 ^s 10 ^m 38 ^s &c.
Hour.	+018300549	+018145815	+000154734	-0000715487
	32 ^m 56 ^s 27 ^m &c.	32 ^m 39 ^s 44 ^m &c.	16 ^m 43 ^s	7 ^m 56 ^s 35 ^m
Minute.	+000305009	+00030243	+000002579	-0000012258
	32 ^m 56 ^s 27 ^m &c.	32 ^m 39 ^s 44 ^m &c.	16 ^m 43 ^s	7 ^m 56 ^s
Second.	+000005083	+0000050405	+000000043	-00000002043
	32 ^m 56 ^s &c.	32 ^m 39 ^s &c.	16 ^m 43 ^s	7 ^m 56 ^s
Meridical REVOLUTION, of	Decimals.	Decimals.	Ds. Decimals.	Ds. Decimals.
	27 ^d 32 ^m 58 ^s 45 ^m 72 ^s	27 ^d 55 ^m 45 ^s 22 ^m 84 ^s	3231.357611277	6798.20338173
	D. +7 ^h 43 ^m 4 ^s 54 ^m	An. D. +13 ^h 18 ^m 34 ^s 11 ^m	Ap. D. +8 ^h 34 ^m 57 ^s 36 ^m	D. +4 ^h 52 ^m 52 ^s 11 ^m
Meridical REVO- { 1	Decimals.	Decimals.	Ds	Ds
lution. { 2	29 ^d 53 ^h 50 ^m 8 ^s 11 ⁹	29 ^d 58 ^h 02 ^m 64 ^s 68 ^m 00 ^s 5	411.78632769861 &c.	146.6197060918 &c.
{ 3	D. +12 ^h 44 ^m 3 ^s 2 th	An. +23 ^h 31 ^m 34 ^s 52 ^m &c.	Ap. +18 ^h 52 ^m 18 ^s 42 ^m &c.	D. +14 ^h 52 ^m 22 ^s 13 ^m &c.
LUNATION. Mot.	Decimals.	Decimals.	Sign.	Sign.
	+12 ^s .8602246554 &c.	+12 ^s .860559558 &c.	+10966509748	-052126578 &c.
	D. 29 ^d 6 ^h 24 ^m 1 ^s &c.	An. 25 ^d 49 ^h 0 ^m 25 ^s &c.	Ap. 3 ^h 17 ^m 23 ^s 49 ^m	D. 1 ^h 33 ^m 49 ^s 40 ^m &c.
ANOMALISTICAL YEAR.	Decimals.	Decimals.	Decimals.	Decimals.
	+13 ^r 4.42 6741279	+13 ^r 35.07031055	+0 ^r 1 ^s 355430729	-0 ^r 0 ^s 6447457501
	12 ^d 48 ^m 8 ^s 3 ^m	2 ^d 6 ^m 33 ^s 32 ^m	100 3 ^s 58 ^m 39 ^s	190 20 ^s 32 ^m 32 ^s &c.
in An. Year.	+12 ^r 48.4261795051	{ 0 à 0 }	1 ^s 02235124049	
	12 ^d 47 ^m 7 ^s 23 ^m &c.	{ in a Lunation. }	+00 40 ^s 13 ^m 59 ^s	

ELEMENTS of the PLANETS PLACES and MOTIONS; according to late IMPROVEMENTS.

In REVOLUTIONS, Signs, and DECIMALS.

The Mean PLACES and MOTIONS of the MOON from the SUN, and Time of the Mean NEW MOON in JANUARY, &c.

Our Radixes.			According to Dr. HALLEY.		
RADICAL YEARS.	Mean Place Moon à Sun.	Mean New Moon.	Mean Diff. Moon à Sun.	Mean New Moon.	
Jan. 1. { 1600 } O.S.	10° 7' 55' 11"	Jan. 5 ^d 6 ^h 31 ^m 52 ^s	10° 7' 54' 11"	Jan. 5 ^d 6 ^h 33 ^m 50 ^s	O.S. 1600
Jan. 1. { 1600 } N.S.	6 6 0 44	Jan. 15 6 31 52	6 5 59 44	Jan. 15 6 33 50	N.S. 1600
Jan. 1. { 1700 } O.S.	8 15 0 4	Jan. 9 ^d 14 ^h 42 ^m 44 ^s	8 14° 59' 4"	Jan. 9 ^d 14 ^h 44 ^m 42 ^s	O.S. 1700
Jan. 1. { 1700 } N.S.	4 0 54 10	Jan. 19 14 42 44	4 0 53 11	Jan. 19 14 44 42	N.S. 1700
TIME forward.	Mean Mot. Moon à Sun.	Advce. & Retrt. of Luna.	Mean Mot. Moon à Sun.	Mean Lunations.	
100 Julian Years.	1236 ^r 10 ^s 7' 4' 53"	+4 ^d 8 ^h 10 ^m 52 ^s 16 th 29 ^{fo}	1236 ^r 10 ^s 7' 4' 52" 40"	1236,853	
4 Julian Years.	49 ^r 5 ^s 20 ^o 40' 59" 43" 12iv	— 14 ^d 0 ^h 1 ^m 30 ^s 34 th	49 ^r 5 ^s 20 ^o 40' 59" 42" 24iv	49,474 12	
365 Days.	+12 ^r 4 ^s 320770894	— 14 ^d 0 ^h 10482961	+12 ^r 4 ^s 3207708631	12,36 nearly.	
Common Yrs. { 1.	+4 ^s 9' 37' 23" 15"	— 10 ^d 1 ^h 11 ^m 23 ^s , &c.	+4 ^s 9' 37' 23" 15"	12.	
2.	+8 19 14 46 30	— 21 6 22 47	+8 19 14 46 30	25.	
3.	+0 28 52 9 45	— 2 8 50 7	+0 28 52 9 45	37.	
After Biffextile.		our.			
Day.	Sign. + ,4063582763 12° 11' 26" 41", &c.	Mayer. Sun's Mot.	Mayer. Moon's Mot. 4 Years. [53 ^r 5 ^s 20 ^o 42' [53 ^r 36"	Mayer's Solar Year.	Equa Motn. — 20", for 100 Years, fr. 1600 or 1700. And Equation of Diff. Moon à Sun, — 1', being connect- ed with our D à ☉, will cor- respond with Halley's D à ☉ very nearly.
Hour.	+ ,016931595 30' 28" 36", &c.	100 Julian Years. 100 ^r 0 ^s 0 ^o 45' 46" 40"	Sun's Motion. = 1200 ^s 0254320987	Decimals. 365 ^d 24225926899 5 ^h 48 ^m 51 ^s 12 th	
Minute.	+ ,000282193 30" 28" 36iv	4 Julian Years. 4 ^r 0 ^s 0 ^o 1' 49" 52"	Sun's Mot. = 48 ^s 0010172839	Moon's Mot. Mayer. 27 ^d 3215827586 Perl. Rev.	
Second.	+ ,000004703 30" 28iv 36v, &c.	363 Days. 11 ^s 29 ^o 45' 40" 23", &c.	Sun's Mot. = 11 ^s 9920405945	Synodical Revn with Sun. 29 ^d 53 ^o 5890025	
1. Lunation.	Sun à Moon's Apogee. ,86055959174 25° 49' 0" 26", &c. Halley.	Sun à Moon's Node. 1 ^s 022351251199 1 ^s 0 ^o 40' 13" 56"	Accurately. In 365 Days . . 12.360064241 4 Jul. Yrs. . . 49.474120153 223 Lunations . . Halley	Lunations. Sun à Moon's Node. 12 ^s 6363271109 11 ^s 984329017	

The Mean PLACES and MOTIONS of Mercury, Venus, Mars, and Jupiter.

RADICAL YEARS.	Mean Places ☿, from Equinox.	M. Places Apb. ☿.	M. Pls. Node ☿.	Mean Places ♀, from Equinox.	M. Places Apb. ♀.	M. Pls. Node ♀.
Jan. 1. { 1600 } O.S.	0 ^s 9' 10' 8"	8 ^s 11 ^o 15' 48"	1 ^s 13 ^o 17' 0"	4 ^s 5' 57' 42"	10 ^s 4' 57' 16"	2 ^s 13 ^o 6' 15"
Jan. 1. { 1600 } N.S.	10 28 14 43	8 11 15 47	1 13 16 59	3 19 56 24	10 4 57 14	2 13 6 12
Jan. 1. { 1700 } O.S.	2 ^s 23' 12' 1"	8 ^s 12 ^o 43' 25"	1 ^s 14 ^o 40' 20"	10 ^s 25' 9' 34"	10 ^s 6' 31' 29"	2 ^s 13 ^o 57' 55"
Jan. 1. { 1700 } N.S.	1 8 11 4	8 12 43 24	1 14 40 19	10 7 32 8	10 6 31 27	2 13 57 52
TIME forward.	Mean Motion ☿.	M. Mot. Ap. ☿.	M. Mot. Node ☿.	Mean Motion ♀.	M. Mot. Ap. ♀.	M. Mot. Node ♀.
100 Julian Years.	415 ^r 2 ^s 14 ^o 2' 13"	0 ^s 10' 27' 37"	0 ^s 10' 23' 20"	162 ^r 6 ^s 19 ^o 11' 52"	0 ^s 10' 34' 13"	0 ^s 0 ^o 51' 40"
4 Julian Years.	16 ^r 7 ^s 8 ^o 57' 41" 19" 12iv	3' 30" 16" 48iv	0 ^s 0 ^o 3' 20"	6 ^r 6 ^s 0 ^o 46' 4" 28" 48iv	3' 46" 7" 12iv	0 ^s 0 ^o 2' 4"
See Phil. Transact. No. 386.	Decimals. +199 ^s 29871593592	Degrees. +05841111, &c.	Degrees. +0555555, &c.	Decimals. +78 ^s 02559703701	Degrees. [&c. +0628111111	Degrees. +03444444, &c.
365 Days.	Decimals. +49 ^s 79057584733	+52" 34"	+50"	Decimals. +19 ^s 49304717215	+56" 31"	+2 ^s 31"
Common Yrs. { 1.	+1 ^s 23' 43' 2" 11"	+0' 52"	+0' 50"	+7 ^s 14' 47' 29" 5"	+0' 56"	+0' 31"
2.	+3 17 26 4 22	+1 45	+1 40	+2 29 34 58 10	+1 53	+1 2
3.	+5 11 9 6 33	+2 37	+2 30	+10 14 22 27 15	+2 49	+1 33
After Biffextile.						
Day.	Sign. + ,1364125365 4° 5' 32" 33" 14iv	Seconds. + ,144 8" 38" 24iv	Seconds. + ,137 8" 13" 12iv	Sign. + ,05340560869 1° 36' 7" 48" 20iv	Seconds. + ,155 9" 18" 11iv	Seconds. + ,085 5" 6" 11iv
Hour.	Sign. + ,0056838517 10' 13" 51" 22iv	Seconds. + ,006 21" 36iv	Seconds. + ,00228333, &c. 8" 13iv 12v	Sign. + ,00222523369 4' 0" 19" 30iv	Seconds. + ,00645833, &c. 23" 15iv	Seconds. + ,00354166, &c. 12" 49iv
Minute.	Sign. + ,0000947308 10" 13" 51iv, &c.			Sign. + ,00003708722 4" 0" 19iv, &c.		
Second.	Sign. + ,0000015788 10" 14iv, fere.			Sign. + ,000006182 4" 0iv, &c.		

P. 120. 2d Col. ☉ M. Motn.

Com. Yrs. { 1. — 14' 19" 45"
2. — 28 39 31
3. — 42 59 17
EMENDATIONS as above.

P. 119. 1. 2. For Trajectories read Trajectories.

1. 11. For, where the m. and true Motions are equal, read, where the m. and true Anomalies are equal.

ELEMENTS of the PLANETS PLACES and MOTIONS, according to our late IMPROVEMENTS. In REVOLUTIONS, SIGNS and DECIMALS.

The Mean PLACES and MOTIONS of MARS and JUPITER.

RADICAL YEARS.	Mean Places δ from Equinox.	M. Pl. Aph. δ	M. Pls. Node δ	Mean Places η from Equinox.	M. Pls. Aph. η	M. Pls. Node η
Jan. 1. { 1600 } O.S.	3 ^s 25' 26' 49"	4 ^s 28' 36' 40"	1 ^s 16' 21' 22"	4 ^s 10' 20' 24"	6 ^s 9' 34' 24"	3 ^s 5' 34' 10"
Jan. 1. { 1600 } N.S.	3 20 12 22	4 28 36 38	1 16 21 21	4 9 30 31	6 9 34 22	3 5 34 9
Jan. 1. { 1700 } O.S.	5 ^s 27' 9' 9"	5 ^s 0' 33' 20"	1 ^s 17' 24' 42"	9 ^s 16' 48' 35"	6 ^s 9' 33' 48"	3 ^s 7' 34' 10"
Jan. 1. { 1700 } N.S.	5 21 23 15	5 0 33 18	1 17 24 41	9 15 53 43	6 9 33 46	3 7 34 9
TIME forward.	M. Mot. δ from Eq.	M. Mot. Aph. δ	M. Mot. Node δ	Mean Motion η	M. Mot. Ap. η	M. Mot. Node η
100 Julian Years.	53 ^r 2 ^s 1' 42' 20"	0 ^s 1' 56' 40"	0 ^s 1' 3' 20"	8 ^r 5 ^s 6' 28' 11"	0 ^s 2' 0' 0"	0 ^s 1' 23' 20"
	2 ^r 1 ^s 15' 40' 5" 36"	0 ^s 0' 4' 40"	0 ^s 0' 2' 32"	0 ^s 4 ^s 10' 27' 31" 38" 24iv	0 ^s 0' 4' 48"	0 ^s 0' 3' 20"
4 Julian Years.	Decimals.	Deg.	Degs.	Decimals.	Degs.	Degs.
	+25 ^s .52227407407	+0.7777777 &c.	0.0422222 &c.	+4 ^s .04862629629	+0.8	+0.555555 &c.
365 Days.	Decimals.			Decimals.		
	+6 ^s .37620125738	+1' 10"	+38"	+13 ^s .01146479045	+1' 12"	+50"
Com. Years. { 1.	+6 ^s 11' 17' 19" 44"	+1' 10"	+0' 38"	+1 ^s 0' 20' 38" 5"	+1' 12"	+0' 50"
2.	+10 22 34 19 28	+2 20	+1 16	+2 0 41 16 10	+2 24	+1 40
3.	+7 3 51 29 12	+3 30	+1 54	+3 1 1 54 15	+3 36	+2 20
After Biffextile.						
Day	Sign. +0.1746904454 31' 26" 39" 24iv.	Secs. +192 11" 31" 12iv	Secs. +104 6" 14" 24iv	Sign. +0.0277113367 4' 59" 16" 56iv	Secs. +197 11" 47" 12iv	Secs. +137 8" 13" 12iv
Hour	Sign. +0.0072787685 1' 18" 6" &c.	Secs. 0.008 28" 48iv	Secs. 0.00433 &c. 15" 34iv	Sign. +0.0001154639 12" 28" 12iv	Secs. 0.00820833 &c. 29" 32iv &c.	Secs. 0.00570833 &c. 20" 33iv
Minute	Sign. +0.00001213128 1" 18" 36iv			Sign. +0.0000019244 12" 28iv.		
Second	Sign. +0.00000020218 1" 18iv &c.			Sign. +0.00000032076 12iv 28v &c.		

The Mean PLACES and MOTIONS of SATURN.

RADICAL YEARS.	Mean Places η	M. Pls. Node η	M. Pls. Node η
Jan. 1. { 1600 } O.S.	6 ^s 16' 10' 32"	8 ^s 28' 31' 56"	3 ^s 18' 51' 46"
Jan. 1. { 1600 } N.S.	6 15 50 26	8 28 31 54	3 18 51 45
Jan. 1. { 1700 } O.S.	11 ^s 9' 16' 32"	8 ^s 28' 33' 10"	3 ^s 21' 5' 6"
Jan. 1. { 1700 } N.S.	11 8 54 26	8 28 33 18	3 21 5 5
TIME forward.	Mean Motion η	M. Mot. Ap. η	M. Mot. Node η
100 Julian Years.	3 ^r 4 ^s 23' 6' 0"	0 ^s 2' 13' 20"	0 ^s 0' 30' 0"
	1 ^s 18' 55' 26" 24"	0 ^s 0' 5' 20"	0 ^s 0' 1' 12"
4 Julian Years.	Decimals.		Degs.
	+1 ^s .6308	+0.88888 &c.	+0.2
365 Days.	Decimals.		
	+0 ^s .4074209445	+1' 10"	+18"
Com. Years. { 1.	0 ^s 12' 13' 21" 27"	+1' 20"	+18"
2.	0 24 26 42 54	+2 40	+36
3.	1 6 40 4 21	+4 0	+54
After Biffextile.			
Day	Sign. +0.00111162217 2' 0" 33" 6iv	Secs. +219 12" 8" 24iv	Secs. +049 2" 56" 24iv
Hour	Sign. +0.0000465092 5" 1" 22iv	Secs. +0.009125 32" 57iv	Secs. +0.002041666 7" 20iv
Minute	Sign. +0.0000007751 5" 11iv 22v		
Second	Sign. +0.0000000129 5iv 1v &c.		

The PLANETS Mean Periodical REVOLUTIONS.

	Ds Decimals.	Ds h m s th
δ	87.968454378	87 23 14 34 27
η	224.695493091	224 16 41 30 36
η	686.929383687	686 22 18 18 45
η	4330.357686018	4330 8 35 4 4
η	10750.551876380	10750 13 14 4 7

The PLANETS Mean Synodical REVOLUTIONS with each other.

	Ds Decimals.	Ds h m s th
δ	39.6300348836	39 15 7 15 1
η	31.1035889370	31 2 29 10 5
η	29.5305908511	29 12 44 3 3
η	28.4532706809	28 10 52 42 35
η	27.4950595246	27 11 52 53 8
η	27.3911968963	27 9 23 19 24
δ	144.5662497937	144 13 35 23 59
η	115.8775010497	115 21 3 30 10
η	100.8882436086	100 21 19 4 14
η	89.7925323996	89 10 1 12 38
η	88.6942117894	88 16 39 39 54
δ	583.9214751501	583 22 0 55 27
η	333.9217217235	333 22 7 16 45
η	236.9926725095	236 23 49 26 54
η	229.4920685698	229 11 48 34 43
δ	779.9370265795	779 22 29 19 5
η	398.8861976230	398 21 16 7 28
η	378.0875534780	378 2 6 4 37
δ	816.4425616661	816 10 37 17 19
η	733.8182627679	733 19 38 17 54
η	7251.1412534284	7251 3 23 24 17

TO determine the Place of a Planet at any Time by its periodical Revolution, and the Place of that Planet given for some one Time Before or After?

RULE. Divide the Time, in Days and Decimals, after or before the radical Time of the Place given, by the Time of the Planet's periodical Revolution, in Days and Decimals, and the Quotient will be the Number of Revolutions performed in that Interval, which turned into Signs, Degrees, Minutes, &c. (neglecting whole Revolutions) and added or subtracted to or from the mean radical Place, according as the Time of the Place sought is after or before the Time of the Place given, and you will have the mean Place required: Which Operation may be readily performed by a Table of Logarithms to many Places. See Examples for finding the Mean Places from those given, further on.

ELEMENTS

**ELEMENTS of the PLANETS' PLACES and MOTIONS; according to late IMPROVEMENTS.
In REVOLUTIONS, SIGNS and DECIMALS.**

MEAN DISTANCES of the PLANETS from the SUN, Eccentricities and Inclinations of their Orbits, with their Aphelions and Nodes from 1° .

PLANETS.	By Mr. STREET's Tables.		Dr. HALLEY's Tables			Mr. STREET's Tables.		
	Mean Dist. from Sun.	Eccentricity.	Mean Dist. from Sun.	Eccentricity.	greatest Equan.	Aph. à 1° .	Q à 1° .	Incl. Orb.
h	953800 As 1	54700 to, 0573495	954002 1	54386 , 05700322	$6^{\circ} 32' 4''$	$75^{\circ} 28' 30''$	$25^{\circ} 22' 30''$	$2^{\circ} 30'$
u	520110 1	25050 , 048163	520093 1	25072 , 0482192	$5^{\circ} 31' 36''$	$5^{\circ} 9' 50''$	$2^{\circ} 8' 00''$	$1^{\circ} 20'$
δ	152369 1	14100 , 0925385	152369 1	14170 , 09263936	$10^{\circ} 40' 2''$	$4^{\circ} 1' 12''$	$0^{\circ} 19' 10''$	$1^{\circ} 52'$
φ	72333 1	517 , 0071475	72334 1	504 , 0069813	$0^{\circ} 48' 0''$	$9^{\circ} 5' 00''$	$1^{\circ} 15' 16''$	$3^{\circ} 24'$
$\varnothing$	38710 1	7970 , 20589	38710 1	7970 , 20589	$23^{\circ} 42' 36''$	$7^{\circ} 13' 48''$	$0^{\circ} 15' 42''$	$6^{\circ} 14'$

THE FIXED STARS being at Rest, the periodic Times of the Five Primary Planets, and of the Earth, about the Sun, (describing Areas proportional to the Times of Description) are in the Sefquipluate (or $1\frac{1}{2} = \frac{1}{2}$ Power) Proportion of their mean Distance from the SUN. Princip. B. III.

AUTHORITIES.	The Periodic Times, in Respect of the Fixed Stars, of the Planets and Earth revolving about the SUN.					
	h	u	δ	$\ominus$ Orb.	φ	$\varnothing$
According to Sir Isaac Newton.	Days 10759,275	Days 4332,514	Days 686,9785	Days 365,2565	Days 224,6176	Days 87,9692
	The Mean Distances of the Planets and Earth from the SUN.					
	Parts.	Parts.	Parts.	Parts.	Parts.	Parts.
To Kepler	951000	519650	152350	100000	72400	38806
To Bullialdus	954198	522520	152350	100000	72398	38585
To periodic Times	954006	520096	152369	100000	72333	38710

The MEAN PLACES and MOTIONS of the FOUR SATELLITES of JUPITER.

RADICAL YEARS.	Mean Place 1, from Equinox.	Mean Place 2, from Equinox.	Mean Place 3, from Equinox.	Mean Place 4, from Equinox.	M. Place Apfis 4, from Equinox.
Jan. 1. { 1600 } O.S.	$9^{\circ} 7' 16' 20''$	$118^{\circ} 24' 9' 29''$	$98^{\circ} 25' 17' 11''$	$8^{\circ} 19' 51' 40''$	$8^{\circ} 27' 48' 6''$
Jan. 1. { 1600 } N.S.	$1^{\circ} 12' 22' 57''$	$2^{\circ} 0' 24' 36''$	$5^{\circ} 2' 6' 37''$	$1^{\circ} 14' 9' 0''$	$8^{\circ} 27' 47' 6''$
Jan. 1. { 1700 } O.S.	$5^{\circ} 2' 16' 20''$	$38^{\circ} 18' 31' 59''$	$118^{\circ} 16' 48' 51''$	$3^{\circ} 14' 41' 40''$	$10^{\circ} 27' 48' 6''$
Jan. 1. { 1700 } N.S.	$2^{\circ} 13' 53' 37''$	$2^{\circ} 13' 24' 37''$	$5^{\circ} 3' 19' 13''$	$7^{\circ} 17' 24' 44''$	$10^{\circ} 27' 47' 0''$
TIME forward.	Mean Motion 1.	Mean Motion 2.	Mean Motion 3.	Mean Motion 4.	M. Motion Apf. 4.
100 Julian Years.	$20645^{\circ} 7' 25''$	$10285^{\circ} 38' 24' 22' 30''$	$510^{\circ} 18' 21' 31' 40''$	$2188^{\circ} 6' 24' 50''$	$2^{\circ} 0' 0' 0''$
4 Julian Years.	$825^{\circ} 9' 27' 24' 0''$	$411^{\circ} 48' 28' 34' 30''$	$204^{\circ} 2' 14' 3' 40''$	$87^{\circ} 6' 15' 23' 36''$	$0^{\circ} 2' 24' 0''$
365 Days.	$+825^{\circ} 9' 91333333$	$+411^{\circ} 48' 9525$	$+204^{\circ} 2' 46870378$	$+87^{\circ} 6' 51311111$	$+08$
Com. Years { 1	$+206^{\circ} 35' 782591832$	$+402^{\circ} 9' 593335044$	$+51^{\circ} 05' 19786234$	$+21^{\circ} 10' 448518518$	$+02$ fere
After Bissextile. { 2	$+38^{\circ} 23' 28' 40''$	$+98^{\circ} 11' 48'$	$+0^{\circ} 5' 56' 9''$	$+10^{\circ} 13' 27' 20''$	$+00$ 36'
{ 3	$+7^{\circ} 16' 57' 20''$	$+6^{\circ} 23' 36''$	$+0^{\circ} 1' 52' 18''$	$+8^{\circ} 26' 54' 40''$	$+1^{\circ} 12'$
	$+11^{\circ} 10' 26' 0''$	$+4^{\circ} 5' 24''$	$+0^{\circ} 17' 48' 27''$	$+7^{\circ} 10' 22' 0''$	$+1^{\circ} 48'$
Day	$+68.782966$ $68^{\circ} 23' 20' 20'' 19'''$	$+38.379159822$ $38^{\circ} 11' 22' 29' 15'''$ &c.	$+18.67725441$ $18^{\circ} 0' 19' 3'' 28'''$ &c.	$+719037037$ $08^{\circ} 21' 34' 16'''$ &c.	$+00005479$ $5'' 55'''$
Hour	$+2826235$ $8^{\circ} 28' 43' 20'''$ &c.	$+140798325$ $4^{\circ} 13' 26' 13'''$ &c.	$+0698856$ $20^{\circ} 5' 47' 38'''$ &c.	$+029959876$ $53'' 55''' 40iv$	In Jan. and Feb. Leap Year, compute for Day and Day's Motion less than the given time See Tab. farther on of Ds. fr. Jan. 1, include.
Minute	$+00471039$ $8' 28'' 43'''$ &c.	$+002346638$ $4' 13'' 26'''$ &c.	$+00116476$ $2' 5'' 47'''$ &c.	$+000499331$ $53''' 55iv$	
Second.	$+00007850$ $8'' 28''' 43iv$ &c.	$+00003911$ $4'' 13''' 26iv$ &c.	$+00001941$ $2'' 5''' 47iv$ &c.	$+000008322$ $53iv 55v$ &c.	
Time of a Revolution.	128 $1^d. 769137506$ $1^d. 10^h. 27^m. 33^s. 29th$ $9^fo. 57fi$	128 $3^d. 5511785863$ $3^d. 13^h. 13^m. 41^s. 49th$ $47^fo. 28fi$	128 $7^d. 1545496471$ $7^d. 3^h. 42^m. 33^s. 5th$ $22^fo. 14fi$	128 $16^d. 6889873287$ $16^d. 16^h. 32^m. 8^s. 30th$ $18^fo. 43fi$	128 600 Julian Years.

OF JUPITER'S FOUR SATELLITES.

THE fixed Stars being at rest, the periodic Times of the Circumjovial Satellites, or MOONS (describing Areas round Jupiter, proportional to the Times of Description) are in the sesquuplicate (or $1\frac{1}{2} = \frac{3}{2}$ Power) Proportion of their mean Distances from the Center of Jupiter.

AUTHORITIES.	Periodic Times of the Satellites of Jupiter.			
	d h m s	d h m s	d h m s	d h m s
According to Sir Isaac Newton	1 18 27 34	3 13 13 42	7 3 42 36	16 16 32 9
to Dr. Halley's Tables	1 18 27 33	3 13 13 42	7 3 42 33	16 16 32 8
	Distances of Jupiter's Satellites from his Center.			
From the Observations of Bonelli	5 $\frac{2}{3}$ Rad. $\frac{1}{4}$	8 $\frac{2}{3}$ Rad. $\frac{1}{4}$	14 Rad. $\frac{1}{4}$	24 $\frac{2}{3}$ Rad. $\frac{1}{4}$
Townley, by Micrometer	5,52	8,78	13,47	24,72
Cassini, by the Telescope	5	8	13	23
Cassini, by Eclipses of Satellites	5 $\frac{2}{3}$	9	14 $\frac{2}{3}$	25 $\frac{1}{3}$
From the periodic Times.	5,667	9,017	14,384	25,299

REMARKS.

MR. POUND determined, by the Help of excellent Micrometers, the Diameters of Jupiter, and the Elongation of his Satellites, thus. The greatest heliocentric Elongation of the 4th Satellite was taken with a Micrometer, in a 15 Foot Telescope, and, at a mean Dist. of $24\frac{1}{2}$ $\odot$, was found about $8' 16''$. The Elongation of the 3d Satellite was taken with a Micrometer in a Telescope of 123 Feet, at the same Distance of Jupiter from the Earth, and was found $4' 42''$. The greatest Elongation of the other Satellites, at the same Dist. of $24\frac{1}{2}$ $\odot$, are determined, from the periodic Times, to be $1' 56'' 47'''$ and $1' 51'' 6'''$.

The Diameter of Jupiter, taken with the Micrometer, in an 123 Foot Telescope, several Times, and reduced to Jupiter's mean Distance from the Earth, proved always less than $40''$, never less than $38''$, generally $39''$. This Diameter in shorter Telescopes is $40''$, or $41''$. For Jupiter's Light is a little dilated by the unequal Refrangibility of the Rays, which Dilatation bears a less Ratio to the Diameter of Jupiter in the longer and more perfect Telescopes than in those which are shorter and less perfect.

The Times in which the 1st and 3d Satellites passed over Jupiter's Body, were observed from the Beginning of the Ingress to the Beginning of the Egress, and from the complete Ingress to the complete Egress, with the long Telescope. And from the Transit of the first Satellite, the Diameter of Jupiter, at its mean Distance from the Earth, came out $37\frac{1}{2}''$, and from the Transit of the third $37\frac{1}{2}''$. There was observed also the Time in which the Shadow of the first Satellite passed over Jupiter's Body, and thence the Diameter of Jupiter, at its mean Distance from the Earth, came out about $37''$. If we suppose its Diameter to be $37\frac{1}{2}''$, very nearly, then the greatest Elongation of the

will be respectively equal to . . . 5,965 . . . 9,494 . . . 15,141 . . . 26,63 . . . Radii of $\frac{1}{4}$.

See Principia, B. III. Phenomenon 1.

Of the celebrated Dr. BRADLEY's Observations, concerning the SITUATIONS and MOTIONS of the SATELLITES of JUPITER.

THIS indefatigable and accurate Observer of the Heavens, has compared the oldest and most accurate Observations he could procure, with his own modern Observations of Jupiter's Satellites taken at Wansled. Who has likewise compared the Observations at the Distances of one, two, or three Revolutions of Jupiter, and sometimes found very remarkable Differences in the Motions of the three inferior Satellites; especially in the Second, or that next but one to Jupiter.

He doubts whether these Inequalities do not partly arise from the Eccentricities of the Orbits, and the Motion of their Apfides. And, by what he can collect from the Motions of the Second Satellite, he thinks it probable that the said Inequalities may proceed from the mutual Action of the Satellites on one another. Sometimes (he says) the Motion of the Second Satellite varies so much, in so small a Time, from its mean Quantity, that a small Eccentricity will not account for it; while the Rest of the Observations will not admit of a large Eccentricity — He finds that the Period of these Errors nearly answers to the Time the three inferior Satellites take up in returning to the same Situation, in respect of each other, and to the Axis of Jupiter's Shadow; which Period is performed in 437 Days, or in 123 Revolutions of the Second Satellite.

After which Period, our judicious Author says, that the like Errors return nearly in the same Order. But, in the intermediate Time, after 60 Revolutions, this Second Satellite will deviate 10, 20, 30, and sometimes 40 Minutes of Time from its common Rate of Motion, during the 7 foregoing, or 7 following Months. That, because the Satellites of Jupiter are not found in the same Place in the Heavens after the Completion of the aforesaid Period, it is possible (and probable) these Errors may vary somewhat on that Account. That if the Orbit of this Second Satellite be likewise eccentric, as late Observations persuade that it is, the Inequalities arising from both Causes must be very intricate, and not easily to be distinguished by Observation only.

The Errors of the 1st and 3d Satellite are not so great as those of the 2d, but appear to arise from the same Causes with those Errors: not wholly depending on the Eccentricity.

Our judicious Author has likewise observed a sensible Difference between the Durations of the Eclipses of the 1st Satellite made at the different Nodes, which (he says) were longer and shorter alternately. That is, the Durations of the Eclipses of the 1st Satellite, at the descending Node, in Leo, in the Years 1682-3, 1694-5, and 1718, were, at least $2^h 20^m$; whereas, at the ascending Node, in Aquarius, in the Years 1677, 1689, the Length of those Eclipses, of the 1st Satellite, did not exceed $2^h 14^m$; which was confirmed by comparing together as many Observations of such Immersions and Emergences, as could be procured, and such as were near together. Hence it is manifest, this Difference of Length of Time in the Eclipses of the 1st Satellite, did not wholly arise from the Eccentricity of the Orbit, if it has any; but to what other Cause it may be assigned Dr. Bradley pretends not to determine.

And hence we may observe, how precarious or uncertain is the Method of determining the Long. of Places from the Eclipses of $\frac{1}{4}$'s Satellites.

Dr. Bradley, however, wishes, till we can get more Light in this Matter from future Observations, that some great Geometer, (such as the celebrated PROFESSOR, MR. T. SIMPSON) in Imitation of the GREAT NEWTON, would apply himself to investigate these Irregularities, or Inequalities, from the certain and demonstrative Principles of Gravity: which, however, we will suppose, depend on physical Causes and Hypotheses, as well as on the Mean of a Number of Operations; in which we are the more confirmed, by the celebrated Author of the late miscellaneous Tracts.

From the Observations of the 4th Satellite, its Orbit is confirmed to be elliptical: And all Dr. Bradley's late Observations of the Motion of that Satellite will be correctly represented by supposing the greatest Equation of that Satellite equal to that of the Planet Venus, $48'$: That its higher Apfide was in $\propto 8^o 0'$ at the Beginning of the Year 1717. Though by comparing this Hypothesis with the older Observations of the Years 1671, 1676, and 1677, the Computations were found to differ greatly from the Observations; yet putting back the Apfide to $\propto 14^o 0'$, at the Beginning of the Year 1677, those Differences mostly vanished. Allowing therefore an equable Motion to this 4th Satellite of Jupiter, of 60 forward in ten Years, the Hypothesis aforesaid was found to agree with the intermediate Observations. The Numbers on that Principle, were found every where to agree with the Heavens (excepting only two Observations, both justly to be suspected) within $10'$ of a Degree.

As to the Latitudes of the Satellites of Jupiter, it was found, from Observations, that the Nodes of the 4th Satellite were, in the Year 1718, in $\propto$ and $\Omega 110\frac{1}{2}$, the Nodes of the 3d Satellite, lying then very near them. The same Places are therefore assigned by Dr. Bradley to the Nodes of the two inferior Satellites, as Observation has not yet discovered any Thing to the contrary. Who takes Notice that if the Nodes of the Satellites were, 40 Years before that Time, in $\propto$ and $\Omega 15^o$, according to Cassini. (whose Authority is of great Weight) that they will appear to have gone back about 1^o in each of Jupiter's Revolutions, by which the present Place of their Nodes may be nearly assigned.

Cassini's Inclination of the Orbits of the 3 inferior Satellites to the Plane of Jupiter's Orbit being $2^o 55'$, is still retained; but the Inclination of the Orbit of the 4th Satellite, $2^o 42'$, Dr. Bradley has considered somewhat less than Cassini's. And as it is a very difficult Matter to determine, with Accuracy, the Situation of Circles so extremely small, so Dr. Bradley is of Opinion that it ought not to be undertaken but with most excellent Telescopes. Mr. Short, an excellent Optician in the Strand, we hear is a Competitor for this Honour.

ELEMENTS of the PLANETS PLACES and MOTIONS, according to our late IMPROVEMENTS.
In REVOLUTIONS, SIGNS and DECIMALS.

The Mean PLACES and MOTIONS of the Five SATELLITES of SATURN.

RADICAL YEARS.	Mean Place 1, from Equinox.	Mean Place 2, from Equinox.	Mean Place 3, from Equinox.	Mean Place 4, from Equinox.	Mean Place 5, from Equinox.
Jan. 1. { 1600 } O. S.	2° 12' 45"	3° 28' 55"	5° 13' 15"	8° 7' 11"	0° 25' 45"
Jan. 1. { 1600 } N. S.	10 25 47	8 3 34	2 26 21	0 21 25	11 10 22
Jan. 1. { 1700 } O. S.	0° 18' 56"	7° 14' 38"	8° 10' 38"	3° 23' 28"	6° 5' 1"
Jan. 1. { 1700 } N. S.	2 21 16	7 7 45	3 4 2	7 15 7	4 15 5
TIME forward.	Mean Motion 1.	Mean Motion 2.	Mean Motion 3.	Mean Motion 4.	Mean Motion 5.
100 Julian Years.	19347° 10' 6" 11"	13345° 3' 15" 44"	8085° 2' 27" 24"	2290° 7' 16" 17"	460° 5' 9" 15"
4 Julian Years.	773° 10' 29" 2' 50" 24"	533° 9' 22" 13' 45" 36"	323° 4' 27" 29' 45" 36"	91° 7' 15" 3' 4"	18° 5' 0" 22' 12"
365 Days	+9286°.968244444	+6405°.74097777	+3880°.916533333	+1099°.5017037037	+221°.0123333333
Common Years } 1.	+193° 4' 152° 152786	133° 4' 339121758	+80° 9' 565047684	22° 10' 687283950	4° 7' 215264658
2.	+4° 4' 35' 14" 51"	+4° 10' 10' 25" 9"	+9° 16' 57' 5" 9"	+10° 20' 37' 6" 40"	+7° 6' 27' 28" 35"
3.	+8 9 10 29 42	+8 20 20 50 18	+7 3 54 10 18	+9 11 14 13 20	+2 12 54 57 10
After Bissextile.	+0 13 45 44 53	+1 0 31 15 27	+4 20 51 15 27	+8 1 51 20 0	+9 19 22 25 45
Day	+6°.3565833295 6° 10' 41" 51"	+4°.384490744 4° 11' 32" 5"	+2°.656342596 2° 19' 41" 25"	+752567901 22° 34' 37" 20"	+151274697 4° 32' 17" 40"
Hour	+264857638 7° 56' 44" 37" &c.	+182687114 5° 28' 50" 12" &c.	+110680941 3° 19' 13" 32" &c.	+031356996 56' 26" 33" &c.	+0063031123 11' 20" 44" &c.
Minute	+0044142939 7' 56" 44" &c.	+003044785 5' 28" 50" &c.	+0018446823 3' 19" 13" &c.	+0005226166 56" 26" 33iv &c.	+00010505187 11" 20" 44iv &c.
Second	+0000735715 7" 56" 44iv &c.	+000050746 5" 28" 50iv &c.	+0000307447 3" 19" 13iv &c.	+0000087102 56" 26iv 33v &c.	+00000175086 11" 20iv 44v &c.
Time of a Revolution of 1/2's Satellites.	12 ^s 1d.8878066058 1d 21h 18m 26s 29th 26fo 40fi	12 ^s 2d.7369199068 2d 17h 41m 9s 52th 47fo 48fi	12 ^s 4d.5174895799 4d 12h 25m 11s 5th 48fo 55fi	12 ^s 15d.9454050329 15d 22h 41m 22s 59th 41fo 26fi	12 ^s 79d.3258898076 79d 7h 49m 16s 52th 45fo 45fi

The Mean MOTIONS of the Five SATELLITES of SATURN, deduced from the Periodic Times, as given, and not computed from the Motion, (*De Tabulis Satellitum Saturni Kkkk 3*, Halley's *Astronomical Tables*) shewing the Difference from the True Motion and Times of Revolutions deduced from that Motion, above—Hence the correct Revolutions (as above) should be always deduced from the Motion, for a Length of Time, and not the Motion from the Revolutions, in a limited short Time, as Below:

TIME forward.	Mean Motion 1.	Mean Motion 2.	Mean Motion 3.	Mean Motion 4.	Mean Motion 5.
100 Julian Years.	19347° 9' 14' 26" 12"	13344° 7' 9' 30' 21"	8085° 2' 20' 41' 10"	2290° 7' 22' 51' 32"	460° 5' 12' 33' 38"
4 Julian Years.	773° 10' 28' 10' 39"	533° 9' 12' 22' 48"	323° 4' 27' 13' 38"	91° 7' 15' 18' 51"	18° 5' 0' 30' 8"
365 Days.	+773° 10' 939248468	+533° 9' 4126745472	+323° 4' 907581626	+91° 7' 5104785992	+18° 5' 0167473552
Com. Years. } 1.	193° 4' 145671236	+133° 4' 257102127	+80° 9' 5628112884	+22° 10' 6894761725	+4° 7' 2163674084
2.	+4° 4' 22' 12"	+4° 7' 42' 47"	+9° 16' 53' 3"	+10° 20' 41' 3"	+7° 6' 29' 27"
3.	+8 8 44 24	+8 15 25 34	+7 3 46 6	+9 11 22 6	+2 12 58 55
After Bissextile.	+0 13 6 36	+0 23 8 21	+4 20 39 9	+8 2 3 9	+9 19 28 22
Day	+6°.356563482 6° 10' 41' 48" 51" &c.	+4°.3842660354 4° 11' 31' 40" &c.	+2°.656336473 2° 19' 41' 24" 20" &c.	Sign. +7525739093 22° 34' 37" 58" &c.	Sign. +1512777216 4° 32' 17" 59"
Hour	Sign. +264856811 7° 56' 44" 32" &c.	Sign. +182677751 5° 28' 49" 11" &c.	Sign. +110680686 3° 19' 13" 30" &c.	+031357246 56' 26" 34" &c.	+0063032384 11' 20" 44" &c.
Minute	+00441428 7' 56" 44" &c.	+003044629 5' 28" 49" &c.	+001844678 3' 19" 13" &c.	+0005226207 56" 26" 34iv &c.	+0001050539 11" 20" 44iv &c.
Second	+00007357 7" 56" 44iv &c.	+0000507438 5" 28" 49iv &c.	+000030744 3" 19" 13iv &c.	+00000871034 56" 26iv 34v &c.	+0000017509 11" 20iv &c.
Time of a Revolution.	12 Signs. 1d 21h 18m 27" = 1d.8878125	12 Signs. 2d 17h 41m 22" = 2d.737060185	12 Signs. 4d 12h 25m 12" = 4d.5175	12 Signs. 15d 22h 41m 12" = 15d.945277777	12 Signs. 79d 7h 47m 0s = 79d.324305555

Of JUPITER'S SATELLITES.

Rad. 1/2's Orbit to Rad. Sat's. Orb.	Revolution 1/2 to Revolution Sat's.	Velocities of Sat's. to Velocity 1/2.
1 1964 to 1	2449 to 1	2449 to 1964
2 1236 to 1	1220 to 1	1220 to 1236
3 778 to 1	606 to 1	606 to 778
4 443 to 1	260 to 1	260 to 443
5 275 to 1	121 to 1	121 to 275

Of SATURN'S SATELLITES.

Rad. 1/2 Orb. to Rad. Sat's. Orb.	Revolution 1/2 to Revolution Sat's.	Velocities of Sat's. to Velocity 1/2.
1 2905 to 1	5702 to 1	5702 to 2905
2 2292 to 1	3932 to 1	3932 to 2292
3 1719 to 1	2382 to 1	2382 to 1719
4 717 to 1	675 to 1	675 to 717
5 238 to 1	135 to 1	135 to 238

De la Caille. See further on.

OF SATURN'S FIVE SATELLITES.

THE fixed Stars being at Rest, the periodic Times of the Circum-Saturnial SATELLITES, or MOONS (describing Areas round Saturn proportionally to the Times of Description) are in the Sefquuplicate (or $1\frac{1}{2} = \frac{3}{2}$ Power) Proportion of their mean Distances from the Center of Saturn.

AUTHORITIES.

AUTHORITIES.	Periodic Times of the Satellites of Saturn.				
	1	2	3	4	5
	d h m s	d h m s	d h m s	d h m s	d h m s
According to Sir Isaac Newton	1 21 18 27	2 17 41 22	4 12 25 12	15 22 41 14	79 7 48 0
To Cassini and Halley	1 21 18 27	2 17 41 22	4 12 25 12	15 22 41 12	79 7 47 0
	Distances of Saturn's Satellites from his Center.				
	1 $\frac{1}{2}$ Rad. Ring	2 $\frac{1}{2}$ Rad. Ring	3 $\frac{1}{2}$ Rad. Ring	8 Rad. Ring	24 Rad. Ring
According to Sir Isaac Newton	1,93	2,47	3,45	8	23,35
To periodic Times	2,097	2,686	3,752	8,698	25,348
From Mr. Huygen's Telescope, recited also by Dr. Halley. See his Tables.	4,893 Rad. $\frac{1}{2}$	6,268 Rad. $\frac{1}{2}$	8,754 Rad. $\frac{1}{2}$	20,295 Rad. $\frac{1}{2}$	59,154 Rad. $\frac{1}{2}$

REMARKS.

The first of the foregoing Motions, are esteemed the best Improvements, preferable to those in *Philos. Transact.* N. 356, and sufficient for determining the Places of the Satellites of Saturn, in Respect of that Primary. From whence a more perfect Theory, by future and diligent Observations of the Motions of these Satellites, can only be expected.

N. B. The Forces tending to the Center of Saturn are reciprocally in the duplicate (or Square) Proportion of the Distances. And the Cubes of the Distances of the Satellites, from the Center of Saturn, as the Squares of the periodic Times.

The celebrated Mr. Pound found the Ratio of the Fourth, or Huygenian Satellite, (the biggest of all the 5) to the Diam. of Saturn's Ring, as 374 to 43, or as 8,7 to 1 nearly. And likewise found the Ratio of the Diameter of the Ring to the Diameter of $\frac{1}{2}$, to be as 7 to 3: whence, by the foregoing Proposition, the Distances above are computed from the Distance and Period of the Fourth Satellite, as discovered by Mr. Pound, by Means of the Royal Society's Telescope of 123 Feet long, fitted with a curious Micrometer.

Anno 1719, May 29^d 10^h Julian Style, Mr. Pound observed, with the same Instrument, its greatest easterly Digression, 3' 7" from the Center of Saturn, whose Place was then η 7° 41'. From whence, by Computation, the Ratio of the Dist. of this 4th Satellite from Saturn is to the Dist. of the Sun from the Earth, as 8,25 to 1000; whence the Distances of the other Satellites may be estimated.

The Diam. of Saturn's Ring by the same Telescope was found (at the Time aforesaid) to be 43", consequently, when $\frac{1}{2}$ is at his mean Dist. $\frac{1}{2}$ is 42", and Diam. of $\frac{1}{2}$ 18". These Appearances are in very long and excellent Telescopes, because in such Telescopes, the apparent Magnitudes of the heavenly Bodies bear a greater Proportion to the Dilatation of Light, in the Extremities of those Bodies, than in shorter Telescopes. If all the spurious Light be rejected, the apparent Diameter of Saturn will not amount to more than 16". Cassini supposed the four inferior Satellites to move in the Plane of the Ring, or that their Orbits are inclined to the Plane of Saturn's Orbit, in an Angle of 30°. For Saturn being about the middle of Gemini and Sagittarius, the greater Axis of the Ring (then of the greatest Width) is found to be nearly double the Length of the less Axis; and these Satellites seem to describe elliptic Orbits, always similar to the elliptic Ring. In their greatest Digressions from Saturn, they are always found in the greater Axis of the Ring produced. These Phenomena could not happen unless the Orbits of the Satellites had nearly the same Situation with the Plane of his Ring.

The same excellent Astronomer since discovered that the 5th and outward Satellite is carried round Saturn in an Orbit very different from the Rest: For its ascending Node was found to be η 5° 0', and the Inclination to the Ecliptic an Angle of 18° only, or half of the Inclination of the former Satellites. Therefore, a Table of Inclination and Reduction is accommodated to the 5th Satellite, among our Tables further on.

The famous Astronomer, Miraldus, has (since the Discoveries of Cassini) by Means of the best Telescopes, and piercing Eyes, sought the Nodes of Saturn's Ring (See *Mem. Roy. Acad. of Paris* for 1715 and 1716). He determines from subtil Observations, that the Plane of the said Ring, in the Year 1715, intersected the Plane of Saturn's Orbit, in η and χ 19° 45'; now, granting the Angle of Inclination of the said Ring to be 30°, the same Plane of the Ring intersected the Earth's Orbit, in η and χ 16° $\frac{1}{2}$, with an Inclination to the Plane of that Orbit in an Angle of 31° 20'. Therefore, at any given Time, to determine, exactly, the Position, Species and Points of the Apogee and Perigee, both of the Ellipses of the Ring, and of the Ellipses described by the Satellites, an oblique-angled Spherical Triangle must be resolved, as is shewn in our Precepts for computing the Latitudes of the Satellites of Jupiter.

The Earth's Latitude, in Respect of the Orbit of Saturn, scarcely ever exceeding 15', it may (as this Affair is not sufficiently known) be neglected, as if Saturn and the Earth moved in the same Orbit. Therefore, from the geocentric Place of $\frac{1}{2}$, deduct 5° 19° 45', and the Remainder will be the Argument of Latitude; with which, in the Table of Latitude of the Satellites of Saturn, 1, 2, 3, 4, take the Inclination, which is the Angle that a visual Ray, from the Earth to Saturn, is inclined to the Satellite-Orbits. The Sine of which Angle is to the Radius, as the less Diameters of these apparent Orbits are to the greater. Let the Ellipsis of the Ring be of the same Species: The Apogee Semi-circle lies towards the North, when the Argument of Latitude is less than six Signs; but lies towards the South, when the Argument is greater.

A Table of Reduction is therefore added, (hereafter) requisite in so great an Inclination of the Planes of the Satellites, that we may know their true Apogee.

A General View of the ELEMENTS of the THEORY of OUR PLANETARY SYSTEM.

According to Sir ISAAC NEWTON'S Principles of Matter and Motion, and other Authorities.

Planetary BODIES.	Apparent Diameters seen at a mean Distance from $\odot$.	Apparent Diameters seen at a mean Distance from $\odot$.	Tr. Diameters. $\odot$'s Parallax $10\frac{1}{2}''$.	Revolutions on their Axes, to same fixed $\star$.	Periodical (rather Synodical) Revolutions through their Orbits, to the same fixed Star.	Inclination of their Axes to Axis of Ecliptic.	Inclination of their Orbits to the Ecliptic.	Proportion of mean Distances fr. the Sun.	Places of their Nodes from Equinox 1760, Beginning N. S.
Names.	' "	' "	Diam. $\odot$.		Days, Decr. Y. D. H. M. S.	' "	' "	As	S. o ' "
SUN	32 12	...	91,7431	25° 12h	...	7 30	...	0,00006	...
MERCURY	0 20	0 21	0,3670	unknown.	87,9692	unknown.	6 59 20	0,38770	8 16 6 29
VENUS	1 20	0 30	1,0275	23h 20m	224,6176	0 24 14 49 20	75 0 2 23 5	0,72333	11 14 28 49
EARTH	...	0 21	1,0000	23h 56m	365,2565	1 0 6 9 21	23 28 $\frac{1}{2}$ 0 0 0	1,00000	...
MOON	31 16 $\frac{1}{2}$	0 6	0,2740	27d 7h 43m	27,3216	0 27 7 43 0	2 10 5 to +18'	1, $\pm$	11 26 54 26
MARS	0 30	0 12	0,5229	24h 40m	686,9785	1 32 1 23 29 2	0 0 1 52 0	1,52369	8 18 2 41
JUPITER	0 39	0 37	9,1468	9h 56m	4332,5140	11 31 5 12 20 9	0 0 1 20 0	5,20096	25 8 24 9
SATURN	0 16	0 16	7,2661	unknown.	10759,2750	29 16 7 6 36 0	unknown. 2 33 30	9,54006	25 21 23 2
	observed.				1st Year com.				

N. B. When the Satellites have greater Velocities than their Primaries, they must be retrograde in their inferior Conjunction, but direct in their superior. The Velocity of Jupiter's second Satellite being nearly equal to that of its Primary, it must appear stationary in the inferior Conjunction. When the Velocity of the Primary is the greater, the Satellite appears direct. See Bottom of the foregoing Table, and of Page 128.

A General VIEW continued of the ELEMENTS of the THEORY of our PLANETARY SYSTEM.

Bodies.	Surfaces.	Magnitudes.	Quantities of Matter.	Densities.	Weight on their Sur- face.	Light and Heat, on Surfaces.	Ratio of their Polar to Equatorial Dia- meters.	Eccentricities of their Orbits.	Places of their Appella, from E- quinox, 1760, Be- gin. N. S.
	Newton.	Newton.	Newton.	Newton.	Newton.	Newton.	Newton.	Halley.	Halley.
	As sq. Diam.	As cub. Diam.	As	As	As	As	As	Sem. Transf. r.	s o ' "
SUN	8416,7963	772183.	169282.	0,25	22,9885	45025.		...	...
MERCURY	0,1347	0,0494	unknown.	unknown.	unknown.	6,6737		0,20589	12 35 58
VENUS	1,0557	1,0848	unknown.	unknown.	unknown.	1,91146		0,0069813	7 27 56
EARTH	1,0000	1,0000	1,0000	1,0000	1,0000	1,0000		0,01685 our	21 21 11
MOON	0,0751	0,0206	0,02513	1,2225	0,3333	1, ±		fr., 0,066777 } to, 0,43323 }	7 50 42
MARS	0,3780	0,1976	unknown.	unknown.	unknown.	0,43166		0,0926393	1 43 18
JUPITER	83,6639	765,2574	158,652	0,23625	2,16781	0,03696	.. 12 to 13	0,0482192	10 45 46
SATURN	52,7962	383,6124	56,0350	0,1675	1,21609	0,01098.		0,05700322	29 51 18

Bodies.	Periodical Revolu- tions.	Proportion of mean Dist- ances fr. ☉.	Period Time, as Cubes M. Dist.	Diameters.	Magnitudes.	Diameters in English Miles.	Mean Distan- ces fr. the Sun, in English Miles.	Progres- sive Vels- cities per Minute in their Orbs.	Rotary Vels- cities of their Equator per Minute.
	Street.	Street.		11" ☉ Parx.	Keil.	Keil.	Whiston.	Whiston.	Miles.
	d h m s								
SUN	...	...	...	...	...	...	763460	...	63.
MERCURY	87 23 15 53	0,38710	...	0,3333	0,037	4240	32,000,000	1605	unknown.
VENUS	224 16 49 24	0,72333	...	1.	1.	7906	59,000,000	1146	0,71
EARTH	365 6 9 50	1,00000	...	1.	1.	7935	81,000,000	968	17,3
MOON	...	1, ±, 003	...	0,2740	0,0205	2175	± 81,000,000	38	0,158
MARS	686 23 27 30	1,52369	...	0,5000	1,25	4444	123,000,000	779	9,26
JUPITER	4332 12 20 25	5,20110	...	15,0833	3431,5	81155	424,000,000	425	432.
SATURN	10759 6 36 26	9,53800	...	11,4166	1488.	67870	777,000,000	315	unknown.

Bodies.	Would descend to the Sun in Days.	The projectile Forces being itopic, D would descend to ☉ in 4 Days.	Time of their Passage thro' Aphelion.	Appellion fr. 1 * V.	Periodical Re- volns. to *s.	Lat. M. Parx.	Gr. Parx.	M. Parx.	Least Par.
			Monsieur De la	Caille.		☉. D.	D.	D.	D.
			Yr. Mo. d h m s	s o ' "	d h m				
SUN	...	...	...	...	...	0 57 20	61' 7"	57' 18"	53' 29"
MERCURY	15 1/2	...	1740 Aug. 9 6 27 40	7 13 54 30	87 23 15 1/2	30 57 16	Gr. Ap. Diam. D.	M. Ap. Diam. D.	Least Ap. Diam. D.
VENUS	39 1/2	...	1743 Jan. 23 15 24 10	9 7 49 20	224 16 48 1/2	38 57 14	33' 37"	31' 31"	29' 25"
EARTH	64 1/2	...	1744 Dec. 29 2 51 40	8 8 42 45	365 6 9 1/2	45 57 12	Our Tables.		
MOON	64 1/2	...	...	...	...	52 57 10	Saturn's Ring; Diamrs. 9 to 4.		
MARS	121	...	1745 Jan. 12 8 31 40	4 1 49 50	686 23 30 1/2	60 57 8	h X m	19° Ring	least visi- ble.
JUPITER	290	...	1744 Apr. 9 12 50 40	5 10 30 40	4332 12 0	90 57 4	h II f		
SATURN	767	...	1723 Sept. 4 23 50 40	7 29 26 15	10759 8 0	Newton.			

N. B. D from ☉ = 240,000 Eng. Miles at a mean Distance.

Bodies.	Diameters.	Surfa- ces.	Magni- tudes.	Distances from Sun.	Revolutions on their Axes.	Periodical Revolu- tions through their Orbits to the same Point of the Eclip- tic. 1st Com. Yr.	Lat. M. Parx.	Gr. Parx.	M. Parx.	Least Par.
		Connoissance des Temps.		Greatest. Least.		Yr. d h m s	☉. D.	D.	D.	D.
	Dis. ☉.	D. ☉sq.	Ds ☉cub.	Sem. D. ☉	Sem. D. ☉					
SUN	100.	10000	1000000	...	...	25d 12h	...	...	...	...
MERCURY	1/2	1/9	1/27	10274	6754	unkn.	0 87 23 14 34			
VENUS	1	1	1	16016	15796	23h 20m	0 224 16 41 30			
EARTH	1	1	1	22374	21626	23h 56m	1 0 5 48 54			
MOON	1/2 +	1/3	1/5	62 fr. ☉.	54 fr. ☉.	27d 7h	0 27 7 43 5			
MARS	3/5	1/3	1/5	36630	30426	24h 40m	1 321 22 18 19			
JUPITER	10+	106	1170	119900	108900	9h 56m	11 313 8 35 4			
SATURN	10-	99	980	221870	197802	unkn.	29 158 13 14 42			

Most of the above Differences of Proportions arise from the Difference of the Sun's Parallax, taken 10 $\frac{1}{2}$, 11, 12 $\frac{1}{2}$, or more Seconds of a Degree, (under 15") which will be truly determined, to a 500 Part of the whole, on May 26, 1761, when Venus will transit the Sun's Disk. The Difference of Proportion in the Revolutions of the same Bodies, their Places and Points of Passage, in Respect of Time, is owing to a greater or less Accuracy of Instruments, and also Diligence and Ability of the Observers. Much Improvement herein is expected from the accurate Observation of the Discoverer of the Aberration of Light, our present Astronomer Royal, Dr. Bradley.

* * * Each Satellite's Motion as seen from the Sun, being compounded of its own proper Motion, and that of its Primary, two Sorts of Re- volutions must be distinguished among them; the one periodical, being the Time the Satellite takes in describing 360°, seen from the Primary's Center; the other Synodical, being the Time the Satellite takes in returning to the same Phase, or Aspect, in Respect of the Sun. The Synodical therefore exceeds the periodical Revolution by the Time the Satellite spends in running over an Arc of its Orbit similar to that run over by the Primary in the Time of the Satellite's periodical Revolution.

A TABLE shewing, at Sight, the NUMBER OF DAYS, from the 1st of January to any Month-Day of the Year following in a common or Leap Year.

Com. Year.	Leap Year.	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
M. Ds	M. Ds	No Ds	No Ds	No Ds	No Ds	No Ds	No Ds	No Ds	No Ds	No Ds	No Ds	No Ds	No Ds
1	0	1	32	60	91	121	152	182	213	244	274	305	335
2	1	2	33	61	92	122	153	183	214	245	275	306	336
3	2	3	34	62	93	123	154	184	215	246	276	307	337
4	3	4	35	63	94	124	155	185	216	247	277	308	338
5	4	5	36	64	95	125	156	186	217	248	278	309	339
6	5	6	37	65	96	126	157	187	218	249	279	310	340
7	6	7	38	66	97	127	158	188	219	250	280	311	341
8	7	8	39	67	98	128	159	189	220	251	281	312	342
9	8	9	40	68	99	129	160	190	221	252	282	313	343
10	9	10	41	69	100	130	161	191	222	253	283	314	344
11	10	11	42	70	101	131	162	192	223	254	284	315	345
12	11	12	43	71	102	132	163	193	224	255	285	316	346
13	12	13	44	72	103	133	164	194	225	256	286	317	347
14	13	14	45	73	104	134	165	195	226	257	287	318	348
15	14	15	46	74	105	135	166	196	227	258	288	319	349
16	15	16	47	75	106	136	167	197	228	259	289	320	350
17	16	17	48	76	107	137	168	198	229	260	290	321	351
18	17	18	49	77	108	138	169	199	230	261	291	322	352
19	18	19	50	78	109	139	170	200	231	262	292	323	353
20	19	20	51	79	110	140	171	201	232	263	293	324	354
21	20	21	52	80	111	141	172	202	233	264	294	325	355
22	21	22	53	81	112	142	173	203	234	265	295	326	356
23	22	23	54	82	113	143	174	204	235	266	296	327	357
24	23	24	55	83	114	144	175	205	236	267	297	328	358
25	24	25	56	84	115	145	176	206	237	268	298	329	359
26	25	26	57	85	116	146	177	207	238	269	299	330	360
27	26	27	58	86	117	147	178	208	239	270	300	331	361
28	27	28	59	87	118	148	179	209	240	271	301	332	362
29	28	29	60	88	119	149	180	210	241	272	302	333	363
30	29	30		89	120	150	181	211	242	273	303	334	364
31	30	31		90	121	151	182	212	243	274	304	335	365
	31			91	122	152		213	244		305		366

EXAMPLE I.

To find the Number of Days from January 1, to October 31, following, in a common and also Leap-Year?

Against 31 in Month-Day Columns { in the common Year } Column you find under Oct. { 304 } reqd.
 { in Leap-Year } { 305 }

EXAMPLE II.

To find the Number of Days from the 1st of January, to the 23d of February, in a common or Leap-Year?

Against 23 in the first Month-Day Column (serving for January and February in common and Leap-Years) you find under February 54 Days, for both Years, required.

N. B. The above Table is first made according to the common Year, or First Column, serving equally for January and February in common and Leap Years. And therefore a Day's Place advanced in Second Column, serves for Leap-Year, after February (of 29 Days) by adding a Day to that Month.

Hence, the Reason for our adding a Day's Motion to the Epochs, or Places, for the Beginning of Leap Years, to make them correspond with Places for January 1st, at Noon. For thus contrived, you see the Day's Motion ordered to be added, in other Astronomical Tables, for all Months after February, in Leap-Year, is already done in our Epochs for the Beginning of that Year, whilst there is only a Day and Day's Motion to be taken out left for the two Months of January and February, for each Leap-Year.

* * In like Manner the Day's Difference between Old and New Stile do not take Place till February the 29th of the Julian Century.

A TABLE for reducing Julian Years to Days, and the contrary.

Julian Years.	Days.	Julian Years.	Days.
* 1	365	600	219150
* 2	730	700	255675
* 3	1095	800	292200
4	1461	900	328725
8	2922	1000	365250
12	4383	2000	730500
16	5844	3000	1095750
20	7305	4000	1461000
24	8766	5000	1826250
28	10227	6000	2191500
32	11688		
36	13149		
40	14610		
44	16071		
48	17532		
52	18993		
56	20454		
60	21915		
64	23376		
68	24837		
72	26298		
76	27759		
80	29220		
84	30681		
88	32142		
92	33603		
96	35064		
100	36525		
200	73050		
300	109575		
400	146100		
500	182625		

CONSTR. After the 1st 3 Yrs, 1461 Days are added for each 4 Yrs, from the 1st 4 Yrs, continually, as far as 100 Yrs; then 36525 Days for 100 Yrs are continually added to the Days for the last hundred Years.

Arithmetically.

Multiply the No. of Julian Yrs by 365 Ds, and to the Product add the Quotient of those Yrs divided by 4, and the Sum will be the Ds in those Yrs, when the 4th from the 1st is a Leap-Year.

But if the 1st, 2d, or 3d is a Leap-Yr, then a Day must be added to the said Sum for the exact No. of Days required.

Ex. To find the Number of Days in 18 Years, the famous Eclipse-Period?

365 Ds
18 Years

2920

365

Product 6570

+4=18÷4

Sum 6574 Days required,

when 3d or 4th Yr. fr. 1st is a Leap-Yr.

but 6575 Days,

when 1st or 2d Yr. from 1st is Leap-Yr.

223 Lunations = 6585 Ds 7^h 43^m 20^s.

Yrs. Beg.

= { 18 Yrs 11^d 7^h 43^m 20^s } 3 4 LP-Yr.

{ 18 10 7 43 20 } 1 2

For 16 Yrs = 5844 Days

2 Yrs

730 + 1

— 6574, and 6575

6585 . . 6585

11 10 Ds.

JULIAN

JULIAN YEARS.		SOLAR YEARS. Our.		SOLAR YEARS. Mayer.		Gregorian YEARS.	Sydereal YEARS.	Gregorian greater and less than So- lar Years. Our.	Sydereal greater and less than So- lar Years. Our.																		
Years.	Days.	EQUATION of Julian Years.				EQUATION of Julian Years.				Equation of Julian Years. Days.	EQUATION of Julian Years.				+ Gr. — less.				20 ^m 29 ^s 40 th 16 ^{fo}								
		5 ^h 48 ^m 54 ^s 46 th 19 ^{fo}	Days.	H	M	S	5 ^h 48 ^m 51 ^s 12 th 19 ^{fo}	Days.	H		M	S	6 ^h 9 ^m 24 ^s 26 th 35 ^{fo}	Days.	H	M	S	Days.	H	M	S						
*1	365	+	..	5	48	55	+	..	5	48	51		+	..	6	9	24	+	..	5	48	55	+	..	0	20	30
*2	730	+	..	11	37	49	+	..	11	37	42	Hundred	+	..	12	18	49	+	..	11	37	49	+	..	0	41	0
*3	1095	+	..	17	26	44	+	..	17	26	34	Years	+	..	18	28	13	+	..	17	26	44	+	..	1	12	29
4	1461	—	..	0	44	21	—	..	0	44	35	— 1/4	+	..	0	37	38	—	..	0	44	21	—	..	0	6	43
8	2922	—	..	1	28	42	—	..	1	29	10	of them,	+	..	1	15	16	—	..	1	28	42	—	..	0	13	26
12	4383	—	..	2	13	3	—	..	2	13	46	= N ^o . of	+	..	1	52	53	—	..	2	13	3	—	..	0	20	10
16	5844	—	..	2	57	24	—	..	2	58	21	Days new	+	..	2	30	31	—	..	2	57	24	—	..	0	26	53
20	7305	—	..	3	41	44	—	..	3	42	56	is forward	+	..	3	8	9	—	..	3	41	44	—	..	0	33	35
40	14610	—	..	7	23	29	—	..	7	25	52	of old	+	..	6	16	18	—	..	7	23	29	—	..	1	7	11
60	21915	—	..	11	5	14	—	..	11	8	48	Style.	+	..	9	24	27	—	..	11	5	14	—	..	1	40	47
80	29220	—	..	14	46	58	—	..	14	51	44		+	..	12	32	35	—	..	14	46	58	—	..	2	14	23
100	36525	—	0	18	28	43	—	0	18	34	40	— 1	+	0	15	40	44	+	..	5	31	17	—	..	2	47	59
200	73050	—	1	12	57	26	—	1	13	9	20	— 2	+	1	7	21	29	+	..	11	2	34	—	..	5	35	57
300	109575	—	2	7	26	8	—	2	7	44	0	— 3	+	1	23	2	13	+	..	16	33	52	—	..	8	23	55
400	146100	—	3	1	54	51	—	3	2	18	40	— 3	+	2	14	42	57	—	..	1	54	51	—	..	11	11	54
500	182625	—	3	20	23	34	—	3	20	53	20	— 4	+	3	6	23	41	+	..	3	36	26	—	..	13	59	53
600	219150	—	4	14	52	17	—	4	15	27	59	— 5	+	3	22	4	26	+	..	9	7	43	—	..	16	47	51
700	255675	—	5	9	21	0	—	5	10	2	39	— 6	+	4	13	45	10	+	..	14	39	0	—	..	19	35	50
800	292200	—	6	3	49	42	—	6	4	37	19	— 6	+	5	5	25	54	—	..	3	49	42	—	..	22	23	48
900	328725	—	6	22	18	25	—	6	23	11	59	— 7	+	5	21	6	39	+	..	1	41	35	—	1	1	11	46
1000	365250	—	7	16	47	8	—	7	17	46	39	— 8	+	6	12	47	23	+	..	7	12	52	—	1	3	59	55
2000	730500	—	15	9	34	16	—	15	11	33	18	— 15	+	13	1	34	46	—	..	9	34	16	—	2	7	59	30
3000	1095750	—	23	2	21	24	—	23	5	19	57	— 23	+	19	14	22	9	—	..	2	21	24	—	3	11	59	15
4000	1461000	—	30	19	8	32	—	30	23	6	37	— 30	+	26	3	9	32	—	..	19	8	32	—	4	15	59	0
5000	1826250	—	38	11	55	40	—	38	16	53	16	— 38	+	32	15	56	55	—	..	11	55	40	—	5	19	59	5
6000	2191500	—	46	4	42	48	—	46	10	39	55	— 45	+	39	4	44	18	—	1	4	42	48	—	6	23	58	30
7000	2556750	—	53	21	29	56	—	54	4	26	34	— 53	+	45	17	31	41	—	0	21	29	56	—	8	3	58	11
8000	2922000	—	61	14	17	4	—	61	22	13	13	— 60	+	52	6	19	5	—	1	14	17	4	—	9	7	57	59
9000	3287250	—	69	7	4	12	—	69	15	59	52	— 68	+	58	19	6	28	—	1	7	4	12	—	10	11	57	44
10000	3652500	—	76	23	5	120	—	77	9	46	32	— 75	+	65	7	53	51	—	1	23	5	120	—	11	15	57	29

* When 1, 2, or 3d Year is Bissextile 1 Day must be added.

The above Equation to be applied to the Days of Julian Years in 2d Column, for the Days in the same Number of Solar Years.

Example.

To find the Number of Days in 673 Solar Years?

Yrs.	Ju.	Days	Eqn.	d	h	m	s
Ag ^t . 600	..	219150	..	4	14	52	17
60	..	21915	..	0	11	5	14
12	..	4383	..	0	2	13	3
1	..	365	..	+	0	5	48
2d Year Biss.	..	1	..	+	0	0	0
Ys. Ju. 673	..	245814	..	5	22	21	39
				5	22	21	39

Sol. Y. 673 .. 245808 1 38 21

[Days required.]

And the same Method for Mayer's Equations.

N. B. The above Equations being made from the 2d Column of Days when 1st, 2d, or 3d Year is Bissextile, and 1 Day adds to Julian Years, and subtracts from Equations, you may find the Days to any Number of Solar Years from the Equations, as they stand for a Day less of Julian Years; amounting both Ways to the same.

The above Equations to be applied to the Days in Ju. Yrs for Ds in the same No of Syderal Yrs.

Ex. To find the Number of Days in 673 Syderal Years.

Yrs.	Ju.	Eqn.	d	h	m	s
Ag ^t . 600	+	3	22	4	26	
60	+	0	9	24	27	
12	+	0	1	52	53	
1	+	0	6	9	24	
J. Y. 273	+	4	15	31	10	
Ds. 245813	0	0	0	0	0	
						found before.

Days h m s

Dif. 245808 8 28 50

in 273 Solar Years, req^d.

N. B. The Equations — + shew how much the Solar and Syderal Years respectively retreat or advance in the Month-days of the Yrs. Julian: Or — and + shew how much the Month-day by each Account is forward or back of the Month-day by the Julian Account.

Example.

To find the Difference between 250 Gregorian and Solar Years?

Y. Gr.	h	m	s
200	+	11	2
40	—	7	23
8	—	1	28
2	+	11	37
250	+	13	48

Ho. &c.

more in 250 Years Gregorian than in as many Years Solar, required.

So for the Rest.

Example.

To find the Difference between 3400 Years Syderal and Solar?

S. Yrs.	d	h	m	s
3000	—	3	11	59
400	—	0	11	11
3400	—	3	23	11

d^s ho. &c.

less in 3400 Syderal than in as many Solar Years.

So for the Rest.

Of the PRECESSION of the EQUINOXES.

SIR ISAAC NEWTON (Prop. 39. B. 3. of the *Principia*) by a Deduction from the Principles of Gravity, (contrary to the *Simpsonian Hypothesis*) demonstrates, that the annual Regress of the Equinoxes, would be $9'' 56''' 50iv$, at a mean Rate, in a circular Orbit, if the Plane of the Equator, and Ecliptic coincided. But, because they are inclined to one another in an Angle of about $23^\circ 30'$, this Recession of Motion must be diminished in the Proportion of the Sine Complement of $23^\circ 30'$ to Radius, or as 91706 to 100000, reducing it to $9'' 7''' 20iv$, which is the Annual Precession of the Equinoxes arising from the Action of the Sun only.

But, the Force of the Moon to move the Sea (and consequently Equinoxes) to that Force of the Sun is proved (Prop. 37. B. 3.) to be as 44815 to 1, whence $1:44815::9'' 7''' 20iv:40'' 52''' 52iv$. The Precession of the Equinoxes from the Action of the Moon; and the Sum of this and the other Precession is $50'' 0''' 12iv$ for the Mean annual Precession of the Equinoxes, by the joint Actions of the Moon and Sun on the Earth, to move the Equinoxes; supposing the Height of the Earth at the Equator to exceed the Height at the Poles by $17\frac{1}{2}$ Miles only. If the Height of the Earth at the Equator exceeds the Height at the Poles by more than $17\frac{1}{2}$ Miles, the terrestrial Matter will be more rare near the Surface, than at the Center of the Earth; and the Precession of the Equinoxes will be increased by the greater Excess of Height, and diminished by the greater Rarity of Matter.

Year	Long. Pleiades.	Year	Long. $\star$ Aries:
In 294 before Christ	$0^\circ 27' 54'' 0'''$ Timocbaris.	In 1601 . . .	$0^\circ 27' 36'' 50'''$ Tycho Brahe.
1727 since Christ	$1^\circ 26' 10'' 58'''$ See Sherburne, p. 12. Wing, [p. 56.]	1700 . . .	$0^\circ 28' 59'' 20'''$ Flamsteed.

Yrs 2021 Distance. $0^\circ 28' 16'' 58'''$ Dif. Motion.

Whence $50'' 22''' 48iv$ Mean Annual Precession.

But the celebrated Dr. Bradley, our Astronomer Royal, has, with great Diligence and Judgment, accurately determined the Quantity of this variable Precession through an intire Revolution of the Moon's Node, (performed in 18 Years, 11 Days, 7 Hours and 45 Minutes) on which its Variation, as well as that of the Ecliptic Obliquity depends: Being both at a Mean in 3° Longitude of the Node, and at their greatest and least Quantities in 0 and 6° Longitude thereof, respectively. See Table, p. 33.

The famous Mr. Mayer, of Gottingen, in his *Astronomical Tables*, determines from the Paris Observations, the mean Equinoctial Precession to be $1^\circ 24' 20''$ for 100 Julian Years, which is at the Mean Rate of $50'' 36'''$ a Year. But from Dr. Bradley's more accurate Observations we find it to be $1^\circ 24' 10''$ only, in 100 Julian Years, or $50'' 30'''$, a Year that the Fixed Stars go forward of the Equinox, at a Mean Rate, or which comes to the same Thing, that the Equinox recedes from the Fixed Stars, in those Times, respectively.

The renowned T. Simpson physically finds, on antineutonian Principles, the Quantity of Precession (supposing uniform Density of the Earth) to be $21'' 7'''$ by the Sun's Force only, (instead of $9'' 7''' 20iv$) which he artificially deducts from $50''$, the Precession, from Newtonian Principles, by the Sun and Moon's Force, and so determines the Precession by the Moon's Force, to hide Mistake, explode Newtonian, and establish Simpsonian Reasoning. See p. 33. Misc. Trans.

Of the PROGRESSION of the Aphelions and Nodes of the Orbits of the Planets, in Respect of the FIXED STARS, and of the Progression of the APHELION of the Earth's Orbit.

SIR ISAAC NEWTON (Prop. 14. B. 3.) proves (from Prop. 11. B. 1.) that the Aphelions of Planetary Orbits are immoveable, as are likewise the Planes of those Orbits, by Prop. 1, same Book, (That if the Planes are fixed, so must the Nodes) except so far as they may be disturbed and suffer some Inequalities by the mutual Actions of the Planets and Comets, in their Revolutions, upon one another: which, however, produce but small Motions and Inequalities.

That the Fixed Stars are immoveable, because they keep the same Positions to the Aphelions and Nodes of the Planets.

That since these Stars are liable to no sensible Parallax, from the annual Motion of the Earth; they can have no Force on our System; because of their immense Distances.

That, if they had any Effect upon the Bodies of our System, yet, by their being dispersed every where round the Heavens, their contrary Attractions would destroy their Mutual Actions, by Prop. 70. B. 1.

That, since the Planets near the Sun, (Mercury, Venus, the Earth, and Mars) are so small, that they cannot act but with little Force on each other, therefore their Aphelions and Nodes must remain fixed, except so far as they are disturbed by the Actions of Jupiter, (the greatest of all the Planets) Saturn, and the higher Bodies. Whence, we find, by the Theory of Gravity, that the Aphelions of the lower Planets move a little in Consequencia, or according to a following Order of Signs, in respect of the Fixed Stars.

That this Motion of the Aphelions of the Planets is in the Sesquuplicate Ratio, or one and half Power (betwixt the simple and duplicate Ratios) of their several Distances from the Sun.

That, if the Aphelion of Mars, in an 100 Julian Years, is carried $33' 20''$ forward in respect of the Fixed Stars, the Aphelions of the Earth, of Venus, and of Mercury, according to the said Sesquuplicate Proportion, will, in an 100 Julian Years, move forward $17' 40''$, $10' 53''$, and $4' 16''$, respectively.

Now, to $1^\circ 24' 20''$ Mayer's Motion of the Fixed Stars beyond the Equinox,
Add $17' 40''$ Sir Isaac Newton's Motion of the Earth's Apogee beyond them,

} in 100 Julian Years.

Sum $1^\circ 42' 0''$ Motion of the Earth's Apogee, beyond the Equinox,
But, to $1^\circ 24' 10''$ Dr. Bradley's Sydereal Motion, beyond the Equinox,
Add $17' 40''$ Sir Isaac Newton's Apogee Progression beyond the Stars,

} in 100 Julian Years.

Sum $1^\circ 41' 50''$ Motion of the $\odot$ or $\oplus$'s Apogee beyond the Equinox,
And, to $1^\circ 23' 20''$ Dr. Halley's Motion of $\star$'s forward
Add $17' 40''$ Sir Isaac Newton's Apogee Progression beyond them,

} in 100 Julian Years.

Sum $1^\circ 41' 0''$ Motion of the $\odot$ or $\oplus$'s Apogee from Equinox,
 $1^\circ 41' 7' 20''$ Dr. Halley's Motion of Apogee for the same Time. See his Tables.

But, as neither of these Quantities are found to agree with Observation, we have settled the mean Motion of $\odot$'s Apogee at $1^\circ 42' 30''$ from the Equinox in 100 Years; being $40''$ more than by the Theory of Gravity, which Sir Isaac always refers to Observation, as the Proof of Theory, In the Time of Hipparchus, 140 Years before Christ, the Sun's Apogee was in $5^\circ 30'$ II, 1884 Years, since which Time in 1744, it has advanced 33° to the Colures, which is at the mean Rate of $1' 3''$ yearly; as the Intervals of some other Observations, loosely taken, ascertain. But a Mean being taken of the Quantities deduced from several remote Observations shews the Motion of the Sun's Apogee to be, as we have before determined it, $61'' 30'''$, a Year: so small a Motion of the Earth's Aphelion, in respect of the Stars, distinguished from the Precession of the Equinox, being hard to determine by the physical Laws of Gravity only.

Of the INEQUALITIES of the EARTH'S MOTIONS, caused by the GRAVITATION of the MOON.

Whence are produced the Precession of the Equinoxes, Nutation, and different Motion of the Earth's Axis; and also Change of the Ecciptic Obliquity.

IF the Earth had no Moon, it would only be impressed with a Central Force towards the Sun. But being attended with a Moon, gravitating towards it, the Earth's Motion, round the Sun, is thereby changed and accelerated from the Moon's Reaction, when the Moon is retarded, and the contrary.

These Inequalities, in the Earth's Motion, are insensible, in respect of the Sun, whose apparent Motions are ever equal, and in contrary Direction to the real Motions of the Earth. The Moon's Vicinity to us is the Reason of these Inequalities being seen from the Earth. For, if the Moon went round the Sun at the same Distance, and in like Manner, as it goes about the Earth, the Inequalities seen from the Earth would then appear, at least, 275 Times less than they appear at present. And the Moon's greatest Inequality, $2^{\circ} 41'$, caused by her Combination of Gravity towards the Sun and Earth, would then appear not above $35''$, very difficult to be observed. Besides, the Inequalities, caused by the $\odot$, in the $\ominus$'s Motion, cannot be near so considerable as the Inequalities of the Moon's Motion; because the Earth is about 50 Times bigger than the $\odot$; and therefore the Inequalities caused in the $\ominus$'s Motions, by the Gravitation of the Moon, will be almost imperceptible with regard to the Sun, and consequently the reciprocal Gravitations of the Earth and Moon, can no more affect the Computation of the true Place of the Earth, seen from the Sun, (or of the Sun's true Place seen from the Earth) than if the Earth had no $\odot$, or Satellite.

Nevertheless, The Moon's Gravitation towards the Earth, acting on the Earth's Spheroidal Figure, produce two very sensible Effects. The one the ebbing and flowing of the Tide; the other a Variation of the Planes of the Ecciptic and Equator: causing the Intersections of those Planes, and consequently Points of the Equinoxes, to move retrograde.

For, the Moon's Orbit being inclined about $5^{\circ} 9'$ to the Ecciptic, and its ascending Ω making a Revolution in about 18 Years, through the Signs of the Ecciptic, it follows, that the Inclination of the Moon's Orbit is continually changing its Position with the terrestrial Equator. When the Moon's ascending Ω coincides with the first Point of Aries, which is the ascending Node of the Equator, then the Inclination of the Moon's Orbit to the Plane of the Equator will be the Sum of $5^{\circ} 9'$, and $23^{\circ} 28' = 28^{\circ} 37'$; but when the ascending Node meets with the first Point of Libra, or descending Node of the Equator, then the Moon's Orbit will be inclined to the Earth's Equator by only the Difference of $23^{\circ} 28'$ and $5^{\circ} 9' = 18^{\circ} 19'$. Therefore the Inclination of the Moon's Orbit to the Plane of the Ecciptic, increases, in about 9 Years, from $18^{\circ} 19'$ to $28^{\circ} 37'$; and decreases from $28^{\circ} 37'$ to $18^{\circ} 19'$ in about 9 Years, more; when a Revolution of the Moon's Node is completed.

Were the Earth an homogeneous Sphere, its central Force towards the Sun and Moon would cause no Change in the Position of its Axis, on which it revolves: But its Figure being a flattened Sphere next its Poles (resembling a Spheroid generated by the Rotation of an Ellipsis on its less Axis) its Matter is raised or accumulated about the Equator, whereon the Sun and Moon acting obliquely, its Equinoxes are made to retrograde, and its Ecciptic Obliquity drawn to a nearer and nearer Coincidence with the Equator: besides having a periodical Variation with the Moon's Node.

For, suppose the Excess of the Spheroidal Matter, above the Spherical, to be collected about the Earth's Equator into a Ring, adhering to the Plane of the globular Earth, inscribed in the Spheroidal one, the two Points where this Ring intersects the Plane of the Ecciptic, or the two terrestrial Nodes, are the Points of Aries and Libra. Let all the Particles of Matter, of which the said protuberant Ring is composed, be considered as so many little Moons, (See Newtonian Princip. B. 1. Prop. lxvi. Cor. 20.) revolving about the Earth in the same Space of Time as the Points of its Surface revolves, viz. in $23^h 56^m 4^s$; then the supposed Ring of Moons will be affected by 3 Central Forces of Gravity.

The 1st infinitely great in respect of the other two, is the Gravity towards the Earth's Center.

The 2^d is their Gravitation towards the real Moon, in the Heavens.

The 3^d their Gravitation towards the Sun, the weakest of the three Forces; the Sun being vastly farther from the Earth than the Earth is from the Moon.

By combining the 1st of these Forces with each of the other, it follows,

That the Nodes of the Ring must always tend to move retrograde; and therefore the Line where the Planes of the terrestrial Equator and Ecciptic intersect, must also tend to move retrograde; causing a Precession of the Points of the Equinoxes.

Now, because the Plane of the Ecciptic, wherein the Sun is situated, always makes an $\angle$ of about $23^{\circ} 28'$ with the Plane of the Equator, it follows, that the Quantity of the Precession of the Equinoxes, resulting from the combined Gravitation of the aforesaid Ring, about the Earth, and its Gravitation towards the Sun, is apparently equal at each of the $\ominus$'s periodical Revolutions: Sir Isaac Newton making it $9'' 7''' 20iv$, arising from the Force of the Sun only.

But the Inclination of the $\odot$'s Orbit to the Plane of the Equator, being sometimes $18^{\circ} 19'$, and at other Times $28^{\circ} 37'$, the Retrogradation on the Nodes of the said Ring, or Precession of the Equinoxes, from the Ring's combined Gravity towards the Earth and Moon, must vary in Quantity: being greatest when the $\angle$ of Inclination is greatest, and the contrary. Hence, The Precession of the Equinoxes, from these two Causes, varies during a Period of about 18 Years; being greatest, or about $58''$ in a Year, when the $\odot$'s ascending Ω is in Υ , and least, or about $43''$ in a Year, when her Node is in $\cap$. The mean Precession is about $50''$ in a Year, when the Moon's Node is in $\overline{\alpha}$ or φ .

Hence, it appears, that the Inclination of the said Ring, and consequently of the Plane of the Ecciptic to that of the Equator, must be subject to a periodical Variation: Their Variation from the Sun's Action being too small for Observation. But the Variations from the Moon's Action are apparent. For, the Obliquity of the Ecciptic is observed to be sensibly greater, when the Moon's ascending Ω is in Υ , and sensibly less when in the opposite Sign of $\cap$.

RULES for determining the QUANTITY of PRECESSION, NUTATION, and CHANGES, thereby, of DECLINATION, and RIGHT ASCENSION, in respect of the STARS. According to Table, p. 33.

To find the present Equation of PRECESSION?

As Rad. is to Sine Ω 's Distance from the nearest Equinoctial Point, so is the greatest Equation of Precession to the present Equation, sought.

To be added to the Mean Precession when the ascending Ω is in any of the 6 Southern Signs, but to be subtracted therefrom, when in any of the 6 Northern Signs.

N. B. The Decrease of Inclination of the Equator to the Ecciptic, from the Ω in the Beginning of Υ , is proportional to the versed Sine of the Node's present Distance, from Υ .

Whence, the said Inclination will be in its mean Quantity when the Ω is in the Solstice of $\overline{\alpha}$ or φ . Consequently, the Difference between the Mean and true Inclination will be, as the Difference between the versed Sine of the Ω 's present Distance from Υ , and the versed Sine of 90° . That is, As Cosine of Node's Distance from Υ . Therefore,

RULES for determining the QUANTITY of PRECESSION, NUTATION, &c.

To find the Nutation at any Time?

As Radius is to the Cosine of the Node's Distance from the nearest Equinoctial Point, so is the greatest Nutation (or Equation of Ecciptic Obliquity) to the present Nutation.

To be added to the Mean Obliquity of the Ecciptic when the Node is in any of the 6 ascending Signs ♈, ♊, ♋, ♌, ♍, ♎, but to be subtracted therefrom, when the Node is any of the 6 descending Signs ♏, ♏, ♐, ♑, ♒, ♓.

To find the CHANGE of a Star's DECLINATION and Right Ascension, arising from the Nutation of the Earth's Axis?

As Radius is to the Sine of the Star's Right Ascension, so is the Nutation (or given Change of the Equator's Inclination to the Ecciptic) to the Change of the Star's Declination caused by that Nutation. And, As Cotang. of the Star's Declination is to the Cosine of its Right Ascension, so is the Nutation in the Table, to the Change of the Star's Right Ascension answerable.

To find the CHANGE of the Star's DECLINATION and Right Ascension, arising from the Precession of the Equinox?

As Coscant of Obliquity of the Ecciptic is to the Cosine of the Star's Right Ascension, so is the Precession of the Equinox (or Change of the Star's Longitude) to the Change of the Star's Declination caused by Precession. And, As the Cosine of the Star's Declination is to the Cotang. of the Angle of Position, so is the Change of Declination (found by the last Proportion) to the Change of the Right Ascension, answerable. Which Changes of Declination and Right Ascension are additive or subtractive according to the Nature of the Declination and Right Ascension, from which they vary.

N. B. When the Precession, in the second foregoing Case, amounts to some Minutes, the Mean Right Ascension must be used at the Middle of the Interval to make the Conclusion exact. For though the given Right Ascension of the Interval may be sufficient for the Purpose in common Cases, without the Trouble of a Computation for the middle Time, yet where the greatest Accuracy is required, the Operation must be repeated; or the mean Result taken of a Number of Operations. See Misc. Tracts.

A small Motion of Nutation and Precession (see the said Tracts) arises from the Moon's Declination. The greatest Quantity of which is to the greatest Quantity of that depending on the Sun, in a Ratio compounded of the Ratio of the Densities of the two Bodies, that of their periodic Times, and that of the Sines of the Inclination of their respective Orbits to the Plane of the Equator, nearly; amounting in no Circumstance to more than $\frac{1}{4}$ of a Second. And, there is another Nutation and Precession arising from more than all that have been considered by the same celebrated Author; being the mean Result of a Number of Operations, to be explained farther on.

OBLIQUITY of the ECLIPTIC.

To determine the OBLIQUITY of the ECLIPTIC with the EQUATOR, from OBSERVATION.

It is always equal to the Sun's greatest Declination, in either of the Tropics. Therefore,

RULE I. The Difference taken between the Sun's Meridian Altitude, on either Day of the Solstice, and the Height of the celestial Equator at the Place of Observation, will be nearly equal to the Sun's greatest Declination, or Obliquity of the Ecciptic, required.

RULE II. The Sun's Meridian Altitude being taken, at each Tropic, or Solstice, the Difference of these two Altitudes will be correctly equal to the Distance of the two Tropics; the half of which Distance will be the true Distance of each Tropic from the Equator, correctly equal to the Ecciptic Obliquity, required.

EXAMPLES.

1st Observation. The Sun's Meridian Altitude at the Winter Solstice, 1743, observed at Paris $17^{\circ} 40' 7''$
By Page 2. Height of the celestial Equator, at Paris, (Lat. being $48^{\circ} 50' 10''$) $41^{\circ} 9' 50''$

Ecciptic Obliquity, nearly, The Difference

2^d Observation. The Sun's Meridian Altitude, at the Summer Solstice, 1744, observed at Paris $64^{\circ} 37' 17''$
Height of the Equator at Paris, as before $41^{\circ} 9' 50''$

Ecciptic Obliquity, nearly, The Diff.

Former Diff. $23^{\circ} 27' 27''$
Sum $23^{\circ} 29' 43''$

Or, 1st Meridian Altitude $17^{\circ} 40' 7''$
2^d Meridian Altitude $64^{\circ} 37' 17''$

Diff. $46^{\circ} 57' 10''$

Mean Ecciptic Obliquity, correctly.

Sum $46^{\circ} 57' 10''$

Half $23^{\circ} 28' 35''$

$\frac{1}{2}$. The Ecc. Obv. $23^{\circ} 28' 35''$, correctly, as before.

From the Mean of two Operations only.

N. B. This Mean Quantity of the Ecciptic Obliquity is found to decrease less than 1 Min. of a Degree in a Century, besides having a periodical Variation of $18''$, in a Revolution of the Moon's $\odot$; varying from its greatest Obliquity $23^{\circ} 28' 39''$, the $\odot$'s $\odot$ being in Υ , to $23^{\circ} 28' 21''$, its least, the $\odot$ in $\cap$; and so by Rotation of the Node's Longitude, according to Table, p. 33.

Of the MEASURE of TIME.

The SYDEREAL and SOLAR DAY.

THE Rotation of the Earth, on its own *Axis* to the Left, or Easterly, being *uniform*, (causing an apparent uniform Revolution of the *fixed Stars*, about the Earth, to the Right, or Westerly) is the *equal Measure of Time*.

For, the Diameter of the Earth's Orbit being but a *Point* in Proportion to the immense Distance of the *fixed Stars*, their Position cannot be altered to us by any other Motion of the Earth than that of its Rotation on its own *Axis*, except a small *apparent Change* caused by the progressive Motion of Light, and of the Earth in its Orbit. And therefore the *real* Rotation Easterly of the same Point, or Meridian of the Earth, to the same fixed Point in the Heavens, (or the apparent Revolution, Westerly, of the same *fixed Star*, to the same Spot, or Meridian, of the Earth) through an entire Revolution, is *always the equal Measure of what is called the Sydereal Day*.

But, the Earth having an *accelerated* or *retarded* Motion, through its Orbit, to the Left, round the Sun, and a Rotation the same Way, or to the Left, on its *Axis*, causes an apparent Motion of the Sun to the Right. The Interval of these two Motions (*real or apparent*) at the Sun's next Return to the same Meridian, is what the *Astronomers* call the *Astronomical, or mean Solar Day*, which they make to begin and end at Noon. For if they reckoned it to begin and end from and to any other Time of the Day, (as from and to Morning or Evening) the *Inequalities of the solar Days* would be much greater, on Account of the *Increase and Decrease of artificial Days*. And the Sun being the *bigger Light*, and the Stars revolving in Succession through the Year, the solar Day, for that Reason, is found more convenient than the *sydereal*, for the *common Measure of Time*.

If the Sun had no other apparent Motion than that of its *diurnal* Revolution round the Earth, it would, every Day, appear to describe the same *Parallel*, through the Heavens, (from Rising to Setting) and be accompanied with the same *fixed Stars* at its Return to the same Meridian. But by the Earth's Progression in its Orbit, *as before described*, the Sun, every Day, appears to be removed as much to the Left, or Eastward, of the Sun's Place on the former Day, as the Earth has really moved the same Way, in the opposite Part of the Orbit or Ecliptic. And therefore, *apparently*, the Sun returns to the same Meridian each Day, about $59' 8''$ (the Earth's mean diurnal Motion in right Ascension) later than on the former Day, or Return of the same *fixed Star* to the same Meridian. For the Stars appear to advance, each Day, about $59' 8''$, (or $\frac{1}{365}$ Part of a Revolution) of the Sun.

Hence 1 mean astronomical or solar Day is measured by the Sum of 1 Revolution of the Earth on its *Axis*, and $\frac{1}{365}$ Part of another Revolution, = $365^\circ + 59' 8''$, OR

An ASTRONOMICAL or SOLAR DAY, AT ALL TIMES,

Is accurately measured by the Sum of 360° of the Equator, and an Arc of the Equator correspondent to the Arc of the elliptic Orbit, described by the Earth (or Sun apparently) in that Day, i. e. $360^\circ + \text{Diff. R. A. in that Day}$.

For, when the Earth has described a Revolution of 360° to the *fixed Stars*, on its own *Axis*, it must still revolve to the Left, that Day, as much as its *annual* Departure in R. A. the same Way in its Orbit, to bring the Meridian of the Earth under the Sun; then *apparently* removed to the Left, or Eastward, as much as the Earth has advanced in the contrary Part of her elliptic Orbit.

The like is also evident by the apparent Motion of the Sun and Stars exhibited on the celestial Globes.

Therefore, if a *fixed Star* comes to the Meridian with the Sun, at Noon, the same *fixed Star*, after one Month or $\frac{1}{12}$ of a Year, will come to the same Meridian exactly $\frac{1}{12}$ of 24 Hours, or 2 Hours, sooner, and so, in Proportion, for any Number of Months after. In 6 Months, or Half a Year, it will come to the Meridian at Midnight, 12 Hours preceding the Sun; and in 365 Days, being almost a solar Year, it will come to the Meridian about one Day sooner, or nearly return to it again with the Sun; in which last Interval, the same *fixed Star* will have return'd to the Meridian about 366 Times, or made nearly that Number of Revolutions to 365 Returns to the Meridian, or Revolutions, of the Sun.

Hence, $\frac{365 \text{ Days}}{366 \text{ Revs.}} = 23^h 56^m 3^s 56^{th} 3^{rd} 56^{th}$, &c. = Time of 1 Revolution of the Earth or *fixed Stars*

being nearly the Quantity of the Sydereal Day, in mean solar Time, but not correctly. See Fergusson's Astronomy (P. 95, 96) erroneously making 366 Sydereal Revolutions, exactly, in 365 solar Days.

The above Computation does not consider the Earth's Motion (or Sun's apparently) through the whole Orbit, or Ecliptic in right Ascension; but only provides for 365 Days of a Revolution. Therefore the compleat Year must be divided by itself + 1, the Revolution gained in that Time, for the correcter Length of the Sydereal Day.

Hence, $\frac{365.2423006021 \text{ Days.}}{366.2423006021 \text{ Revs.}} = .9972695617 \text{ Day} = 23^h 56^m 4^s 5^{th} 26^{th}$, &c. correcter Time of the

SYDEREAL DAY.

Or,

MEASURE of TIME.

Or, As 365 D^s.243006021 : 360° passed over in R. A. :: 1 D. : 59'.1388236, &c.

Sun's mean diurnal Motion in R. A. = 59' 8" 19''' 45iv, &c.

Now, As 360° + 59'.1388236 = 360°.98564706 apparent diurnal Mot. ☉ : 24 Ho. } : 23^h.9344696, &c.
:: 360° apparent diurnal Mot. * }

The SYDEREAL DAY, in mean Solar Time, as before, 23^h 56^m 4^s 5th 26^{fo}, &c.

Hence, any fixed Star comes to the Meridian each } by . . 3^m 55^s 54th 34^{fo}
Night, sooner than on the former Night }

[Time by a regulated Clock.

N. B. As a Turn of the Earth on its Axis measures one syderal Day, there must be one Turn, or syderal Day more, in the Course of the Earth through her Orbit, than the Number of solar Days in which that Course is compleated, whatever Number those Days may be. And the same holds with respect to any other Planet revolving on its Axis to the Stars, and the Number of Days in which it goes through its Orbit.

The SYDEREAL DAY deduced correctly.

A syderal Year, or Revolution of the Sun with the same Star, or ☉ à *, is not till 365 D^s.256532906 (because the Stars go forward 50" 30''' a Year) consequently, the Number of syderal diurnal Revolutions of that Star to the same Meridian, in that Time = 366.256532906.

Whence, $\frac{365.256532906 \text{ Days.}}{366.256532906 \text{ Revs.}} = .9972696733 \text{ Day.}$

The SYDEREAL DAY, in mean Solar Time, = 23^h 56^m 4^s 5th 59^{fo} 10^{fi}, &c. correctly.

Hence, any fixed Star comes to the same Meridian each } by . . . 4^m 55^s 54th 0^{fo} 50^{fi}, Time by a regulated
Night, sooner than on the former Night, correctly, }

[Clock.

* * One apparent Revolution of the Sun to the Meridian will be lost by a Planet moving round him; in the same Manner that a Traveller would lose a Day going round the Earth the same Way with the apparent Motion of the Sun. Who would reckon one Day less at his Return than the Inhabitants remaining at the Place of his Setting out; whatever Number of Days, or Time, he took to go round the Earth.

Hence, the Number of solar Days that any Planet takes to go round the Sun in its Orbit, will be always 1 less than the Number of syderal Days, or Revolutions, of that Planet to the same fixed Star, in that Time. And if the Earth turned but once round its Axis in a Year, the same Way that it goes round the Sun in its Orbit, there would be continual Day on that Side of the Earth next the Sun at its Setting out, and continual Night on the opposite Side: As the Moon keeps the same Face towards us by turning once round her Axis while she goes once round the Earth.

THE TABLE, Page 21, is used for the Regulation of Clocks and Watches, it shews the Difference in Succession, between a Number of solar and syderal Days. The Difference between 1 solar and 1 syderal Day is shewn to be 3^m 55^s 54th, as aforegoing, that the fixed Stars daily accelerate, or come to the Meridian sooner each Night than on the Night before.

To regulate a CLOCK or WATCH.

If, through a small Hole, in a Window-Shutter, or in a Piece of Metal fixed to a Window, you observe at what Time any Star disappears or vanishes behind a Chimney, or Corner of a House, at a short Distance, and you further observe the same Star vanish or disappear the Night following 3 Minutes, 56 Seconds, sooner, by the Clock or Watch, and on the Second Night 7 Minutes, 52 Seconds, sooner; the Third Night 11 Minutes, 48 Seconds, sooner, and so on every Night, sooner, in that Proportion, for several Days, according to the Table, in Page 21. of the Stars Acceleration, then the Clock or Watch will be found to go true, according to mean or uniform Time. Otherwise, it must be altered to go by that Rule, and may be regulated by it to half a Second of Time.

The Time-measurer being thus well regulated according to the uniform Motion of the Earth on its Axis, or equal Measure of the syderal Day, it will shew you the different Inequality of the solar Day from Noon to Noon; to be sometimes greater and sometimes less than 24 Hours by the Clock. The Time by a regulated Clock, and that by a true Sun-Dial will never be the same, except about April 15, June 16, August 31, and December 24, each Year. For, by the equal Movement of the Machine, all the Year, the unequal Time by the Sun will be found slower than the Time by the Clock, from December 24 to April 15; and, from thence, till June 16, faster; from thence to August 31, slower; and thence to December 24, faster than the Clock: The Sun's Time being the unequal Motion:

[N. B.

MEASURE of TIME.

N. B. The foregoing Days are those, on which the equal and unequal Movements meet, or shew the same Time; but are not the Days, whereon the Sun's equal and true Motion is alike, for 24 Hours, hereafter shewn.

The EQUATION-TABLES, at p. 19 and 20, exactly shew the Difference between the Time by the Sun, and that by an equal moving Clock, for the Year. Especially the Table at Page 19, giving that Equation readily from the Sun's Place, or Longitude only.

The Longitude of the fixed Stars being known, when any Star, whose Relation to the Ecliptic is known, comes to the Meridian at Midnight, the Sun's Place, or Longitude, in the Ecliptic being then opposite to the Point of the Ecliptic coming to the Meridian, will be also known.

EQUATION of TIME.

MEAN and TRUE DAYS.

TO measure Time by a Clock two Sorts of Time or Days must be necessarily distinguished.

The one true or apparent, known by the Interval between the Instant of the Sun's Center passing the Meridian, and that of its next Return to the same Meridian.

The other, a mean Day, or Time, known by the Interval from one Noon to the next, such as would be daily observed, if the Sun's Motion was uniform like that of the Equator.

Mean Time is shewn by well regulated Clocks; and true Time is deduced from Observations of the Sun.

The true and mean Day are each divided into 24 Hours, or 86400 Seconds of Time. The Measure of the mean Day, is the Sum of 360° of the Equator, and $59' 8''$, annual Motion in R. A. passing under the Meridian; which, divided by 86400, gives $15' 28''$, passing under the Meridian in each Second of Time.

The Measure of the true Day is the Sum of 360° of the Equator, and the Motion or Increase of right Ascension for that Day.

When the Sun is in Apogee, about June 30, Leap-Year, its Motion in R. A. in a Day, is about $1^\circ 2' 6''$; therefore $361^\circ 2' 6''$ is then the Measure of the true Day. Say, As $360^\circ 59' 8'' : 24^h$ mean Time :: $361^\circ 2' 6'' : 24^h 0^m 12^s$, true Time, corresponding; the true Day being then 12^s longer than the mean Day.

The true and mean Day are of equal Length when the Sun's Increase of R. A. is $59' 8''$ for a Day; happening about February 10, May 15, July 26, November 2. Leap-Year, N. S.

The above Days respect only the mean Length of the solar Days, when the mean and true Motion of the Sun, for 24 Hours is alike, or the Clock (going sometimes before and sometimes after the Sun) marks nearly the same Difference of Time, or has no Increase for a Day; the Difference between the Clock and Sun-Dial at 12, being nearly the same as at all other Times of the Day. These Days are distinguished from the Days before observed, on which the Time marked by the Clock, going always equal, before and after the Sun, meet with Noon, or 12, marked by the Sun, moving unequally.

The Clock and Sun-Dial on those Days (April 15, June 16, August 31, and December 24) go nearly together, or mark the same Time for 24 Hours. At other Times of the Year there is a considerable Difference between the true Noon, or Instant, of the Sun's Center passing the Meridian, and the mean Noon, or Instant, that the Sun's Center would have passed it, had its Motion in R. A. continued uniform, from any one of those last mentioned Days.

The Differences of each mean and corresponding true Day, successively, through the Year, from any one Day of the Year, accumulate insensibly; yet, at the End of many Days, form a considerable Variation between the Time of the Day, by the Clock, (going equally from the same Time of the Day or Noon by the Sun) and Time of the Day, by the unequal Motion of the Sun. This Difference, or Equation, from whatever Epocha, at Noon, or Time of the Day by the Sun, the mean Movement begins its Reckoning, is at all Times, successively, equal to the Difference between the Sun's mean Place, and its true right Ascension, reduced to Time. For the Sun's mean Place would be advanced as far in the Equator, as in the Ecliptic, by uniform Motion, and consequently the Difference between that Distance, and the true R. A. in the Equator, is the Arc of Equation between mean and true Time; being the very Principle on which the Difference between the mean and true Time depends. But if the Sun's mean Progress in R. A. or the Earth's Rotation on its Axis, be not uniform, this Equation of natural Days must be defective.

The aforesaid Arc of Equation of Time is reducible to two Arcs; one of which is the Difference of the true Place of the Sun and its R. A. and the other is the Difference of the true and mean Place of the Sun, and therefore the Sum of Differences, of which Arcs, converted into Time, will produce the Equation of Time, on its first Principles.

Tycho had a Suspicion of the Acceleration or Retardation of the Earth's Motion on its Axis, and was for ballancing the Part of the forementioned Equation of the Difference between the Sun's true and mean Place, (depending on the Sun's mean Anomaly) by only using the other Part of the Equation or Difference between the Sun's true Place and right Ascension, (depending on the ecliptic Obliquity) which he called the *Empyric*; because it served him to find, by Calculation, the Eclipses he had observed.

EQUATION of TIME.

Street imagined that this supposed *Inequality* of the Earth's Rotation, or first Mover, was compensated, by making Use of the Part of Equation neglected by Tycho, under a *contrary Title*, or Sign, as he has given it in his *Astronomia Carolina*; though we find no Reason for any *Inequality* of the Earth's Rotation.

Besides the common Equation of Time, giving the Hour by the Clock when it is Noon by the Sun, for each Day of the Year, there is annually published, (in an excellent *Ephemeris*, intitled the *Connoissance des Temps*, by Monsieur Miraldi, of France, sent for into England for Want of Astronomy being more encouraged among us) another Equation of Clock-Time, correspondent to the Increase of Time by the Sun. This last Equation begins its *Epocha* with the Time by the Sun, November 2, when the equal and true Motion of the Sun for a Day is alike: and not from the common *Epocha's*, of April 15, June 16, August 31, or December 24, when the Sun and Clock mark the same Time. It shews the Increase, or Difference, of Clock-Time from November 2, for each Day of the Year, correspondent to the Time by the Sun. The Difference of any two of which *Increases*, for any two Days of the Year, shews the Increase or Decrease of Clock-Time, correctly, correspondent to the Increase or Decrease of Time by the Sun in that Interval. This Equation is exactly conformable to the other common Equation of Time, and its Increase and Decrease in the same Interval.

EXAMPLE. On November 2, (the *Epocha* of the increasing Clock-Time Equation) Noon by the Sun, according to the common Equation of Time, is at 11^h 43^m 51^s by a Clock, well regulated; and therefore this Clock being set at 12, on the same Day, when it is 12 or Noon by the Sun; it will continue to go 16^m 9^s forward of correct mean Time, each Day round the Year.

Required the Increase of Clock-Time, correspondent to the Sun's Time, from November 12, (1st after Leap-Year) to December 10 following, and the correct Times by a regulated Clock, when it is Noon by the Sun, on each of those Days?

	Incr. of Clock-Time	Correct M. Noon.	
November 12	0 ^h 42 ^m	11 ^h 44 ^m 33 ^s	} By the Connoissance des Temps.
December 10	9 37	11 53 27	
Increase of Clock-Time in that Interval } correspondent to Time by the Sun.	8 55 Dif.	same as 8 54 Dif.	
But, November 12	12 ^h 0 ^m 42 ^s Noon by Clock, from above.		
	— 16 9 too fast.		
	11 44 33 correct mean Noon, by a regulated Clock.		
December 10	12 9 37 Noon by the Clock, from above.		
	— 16 9 too fast.		
	11 53 28 correct mean Noon, by a regulated Clock.		
	So for the Rest.		

A Clock may be regulated by making it go from its Time of 12 by the Sun, with the Increase or Decrease of Time, in a Day, or for a longer Interval, according to the EQUATION-TABLES of Time following, for each Day of the Year.

The other Method of regulating a Clock, is by making it measure 23^h 56^m 4^s from the Instant of any Star's Passage through the Meridian, to its next Return to the same Meridian, as before observed. What its Return exceeds that Time it goes too fast, what it wants, it goes too slow, of correct mean Time.

To find the true Time, or Noon, by a regulated Clock or Watch?

RULE. Observe the Time marked by a Clock, when the Sun's Center is on the Meridian, being the Instant that the Sun ceases to rise any higher, and is going to descend. Or, you may observe the two Times marked by the regulated Clock, when the Sun has an equal Altitude in the Fore and After Noon; the Middle of which two Instants will be the true Noon marked by the Clock. Whatever the Time, marked by the Clock, is short of, or exceeds, 12, at the true Noon, must be added to, or subtracted from, the Hours, Minutes and Seconds, marked by the Clock at all other Times of the Day, for the apparent or true Time, by the Sun, required.

EXAMPLE. Let the Time of observing a given Altitude of the Sun, marked by the Clock, in the Forenoon, be 9^h 20^m 34^s, and the Time of observing the same Altitude on the Afternoon be 2^h 59^m 36^s, required, from thence, the Time marked by the Clock when it is Noon, by the Sun?

Time 1st Observation	9 ^h 20 ^m 34 ^s —	
	12 0 0	
1st Observation	2 39 26 before 12.	
Time, 2d Observation	2 59 36 after 12.	
Sum	5 39 2 Interval.	
1/2	2 49 31	} + Time 1st Observation. — fr. Time 2d Observation.

Hence, Noon by the Sun, is at . . . 0 10 5 after 12 by the Clock.

Which Time must be subtracted from all other Hours of the Day by the Clock, for the Time apparent by the Sun.

EQUATION of TIME.

The *Instant* of the Passage of a *Star*, or *Planet*, over the *Meridian*, is determined by the like *METHOD* of *CORRESPONDING ALTITUDES*. The farther the *Sun* or *Star* is from the *Meridian*, the *faster* it ascends or descends, and is therefore the fitter for *Observation*, if equal *Altitudes* can be observed.

This *Method* is of most use for observing the *Stars*, or *Sun* in his *Solstices*, having no *Change* of *Declination* in the *Interval* of the two *Observations*. For, when the *Sun* has a *Change* of *Declination*, whereby it approaches the elevated *Pole*, and *increases* the *Meridian Altitude* with the *Length* of *Days*, and consequently makes the *Afternoon Arc* greater than the *Forenoon Arc*, described between the same *Altitudes*, then the *Mean*, between the two observed *Instants*, will be *later* than it really happened. And the *contrary* follows in the *Decrease* of *Days*: For then the *Change* of the *Sun's Declination* being towards the depressed *Pole*, the *Mean* of the two observed *Instants* will be *sooner* than it really was.

The greatest *Error* in the true *Time*, thus computed, occasioned by the *Sun's Change* of *Declination*, does not exceed 30^s . But, to compensate for that *Inaccuracy*, we have given a *Table* for the different *Latitudes* and *Intervals* between the two *Observations*, exhibiting the *Mean Instant*, or *Noon* of the *Sun*, marked by the *Clocks*, as correctly as if no *Change* of the *Sun's Declination* ever happened. See this *Table* farther on.

Of the NATURE and DIVISION of TIME.

TIME is the *Succession* of *Duration*, measured by the *Motion*, or *Change* of *Place*, of the *Celestial Bodies*, in their respective *Revolutions* through their *Orbits*; particularly, by the apparent *Motion* of the *Sun*, or the real *Motion* of the *Earth* in its *Orbit*. And the *Standard Measure* of *Time* to which all other *Measure* thereof refers by *Comparison*, is the *Mean Solar Year* (subdivided into *Months*, *Weeks*, *Days*, *Hours*, *Minutes*, and *Seconds*) during one *Revolution* of the *Earth* (or *Sun* apparently) through its *Orbit*.

The *Mean Solar Day*, and its *Parts*, differ from the *variable Solar Day* (from *Noon* to *Noon*) and its *Parts*, as has been shewn before.

The *Mean Solar Year* is determined by dividing the *Number* of *Days* by the *Number* of the *Mean Solar Year*. *Revolutions* apparent, or real, of the *Sun* or *Earth*, in that *Time*, as derived from the *Sun's Mean Motion*.

Thus, $\frac{36525 \text{ Days in } 100 \text{ Julian Years}}{\text{Rev}^s \text{ Dec}^s} = 365,2423006021$, or $365^d 5^h 48^m 54^s 46^{th} 19^{to} 16^{fl}$, &c.
 $100,00210802469$, in 100 Julian Years
 Length of the *Mean Solar Year*. See p. 7.

The *Length* of the *Year* between other like *Points* of the *Ecliptic* differs by many *Seconds* of *Time* more or less, according to the following *Computations* from *Dr. Halley's Tables*, (See *Palladium* for 1756) where the *Equation* of *Time*, and *Decrease* of the *Sun's Mean Anomaly*, for a *Revolution* to different *Points* of the *Ecliptic* are considered.

		Apparent Time.	Eq. of m. Yr.	Dif. Length Year.	
		March 19 ^d 15 ^h 54 ^m 30 ^s	+ 0 ^s	365 ^d 5 ^h 48 ^m 55 ^s	
Variable Solar Year.	Dr. Halley's mean Solar Year, 365 ^d 5 ^h 48 ^m 55 ^s nearly, between the next vernal Equinoxes; nearly according to our Mean Year.	♈ April 19 4 58 33	— 23	365 5 48 32	Since the Sun's Entrance into the same Sign to the same Sign again.
		♉ May 20 5 55 7	— 49	365 5 48 6	
		♊ June 20 14 55 25	— 61	365 5 47 54	
		♋ July 22 1 54 9	— 58	365 5 47 57	
		♌ Aug. 22 8 12 43	— 41	365 5 48 14	
		♍ Sept. 22 4 25 34	— 16	365 5 48 39	
		♎ Oct. 22 12 1 17	+ 8	365 5 49 3	
		♏ Nov. 21 7 53 25	+ 9	365 5 49 4	
		♐ Dec. 20 19 57 56	+ 40	365 5 49 35	
		♑ Jan. 20 0 26 4	+ 38	365 5 49 33	
		♒ Feb. 18 15 17 10	+ 24	365 5 49 19	

N. B. Though the *Length* of the above *Mean Astronomical Year* is sometimes determined by the *Number* of *Revolutions* betwixt a given *Number* of *Solstices*, containing a certain *Number* of *Days*, on which account the *mean Year* is commonly called the *Tropical Year*; yet the *Length* of the *Year* so determined is properly the *Distance* of *Time* between the next *vernal Equinoxes*, or *Mean Year*, before ascertained, *falsely distinguished* by the *Name* of *Tropical*. For, according to the foregoing *Computation*, the *Tropical Year* of *Cancer* is seen to be less than the *Mean Year*, by 61 *Seconds* of *Time*; and the *Tropical Year* of *Capricorn* more by 40 *Seconds* of *Time* than the *Mean Year*; as the *Year* of the *Autumnal Equinox* is seen to be less than the *Year* of the *Vernal Equinox*, by about 16 *Seconds* of *Time*; owing to the *progressive Motion* of the *Sun's Apogee*, and *regressive one* (or *Precession*) of the *Equinox*, in respect of the *fixed Stars*, each *Julian Year*; making together $1' 1'' 50'''$, in a *Julian Year*, that the *Sun's Apogee* goes forward of the *Equinox*.

Hence,

DIVISION of TIME.

Hence, To $365^d \ 5^h \ 48^m \ 54^s \ 46^{th}$, the *Solar Mean Year*.
Near Sydereal Year. Add $20 \ 29 \ 42\frac{1}{2}$, answring $50'' \ 30'''$ Mot. $\ast$ s in a Year *nearly*.
 Sum, $365 \ 6 \ 9 \ 24 \ 28\frac{1}{2}$, the *Sydereal Year*, or Time in which the Sun returns to the same *Star*.
 Again, To $365^d \ 5^h \ 48^m \ 54^s \ 46^{th}$, the *Mean Year*.
Near Anomalifical Year. Add $25 \ 5 \ 41$, answring to $61'' \ 50'''$ nearly Mot. $\odot$ Apog. in a Mean Year.
 Sum, $365 \ 6 \ 14 \ 0 \ 27$, the *Anomalifical Year*, *nearly*, or Time in which the Sun returns to the
 [same *Anomaly*.]

Page Twenty.

Correct

Sydeareal Year. But correctly,

12 ^s	0	0'	27"	19 ^m	12iv exactly = ☉'s Mean Motion	} in 1 Julian Year.
	—	50	30	00	= M. Motion Fixed Stars	
				D ^s	D ^s	

As Dif. 11 29 59 36 49 12 : 365,25 :: 12^s : 365,256532906 = 365^d 6^h 9^m 24^s 26^th 35^fo 5^fi fere, the Length
[of the Sydeareal Year, or Revolution ☉ à *.

12	0	0	27	19	12 ☉'s M. Motion.	} in 1 Julian Year.
	—	1	1	50	0 M. Mot. ☉'s Apog.	
				D ^s	D ^s	

As Dif. 11 29 59 25 29 12 : 365,25 :: 12^s : 365,259727107, &c. = 365^d 6^h 14^m 0^s 25^th 19^fo 21^fi &c. the
Length of the Anomalifical Year, correctly.

Correct N. B. The above correct Proportions are new ; there being no Instance, that we have seen, in any former Anomalistical Year. Astronomy, where the Sydereal and Anomalistical Year are correctly deduced on their true Principles. The Time of a Mean Lunar Year is measured by twelve Lunations, or Revolutions, of the Moon to the Sun, from any Synodical Conjunction, to the next, of the Moon and Sun. Now, a Mean Lunation, or one mean Lunar Month, is thus determined.

[illegible]

The *civil* or *political* Year consist of a certain Number of Days, according to the Usage or Custom of any particular Nation; of which some reckon according to the Days in the *Lunar*, but most according to the Days in the *Solar* Year.

The *civil* Solar Year with us contains 365 Days for three Years together, and every *fourth* Year, commonly called *Leap-Year*, or *Bissextile*, contains 366 Days. This Account of Time, or reckoning, by the Year, is called the *Julian Account*, from *Julius Cæsar*, who first instituted it, as a Means to make the *civil* and *correct solar* Year, and its Seasons keep Pace together. He ordered, in each *Leap-Year*, the *Intercalary Day*, to be added to the 23d of *February*, and for the 23d and 24th of that Month to be reckoned as *one Day*, which 23d being the 6th of the *Calends* of *March*, and twice reckoned, called *Bis sextus Dies*, gave the Name of *Bissextile* to that Year. In our English Almanacs this *Intercalary Day* is added at the End of *February*, making the 29th Day of that Month, in each *Bissextile* or *Leap-Year*.

Hence, a *Mean Julian Year* consists of 365^d 6^h 0^m 0^s 0th 0^{fo} 0^{fi}
 But a *Mean Solar Year* consists of 365 5 48 54 46 19 16 &c.

Difference, a mean Julian Year is greater than a mean Solar, }
Or, Time that the Seasons go back each Julian Year } . . . 11^m 5^s 13th 40^{fo} 44^{fi}

Hence, $\frac{1 \text{ Day}}{,0076993979} = \text{Yrs Decs \&c.}$ $\left\{ \begin{array}{l} \text{Or, } 365^{\text{d}}, 25 \\ 365,2423006021 \\ \hline ,0076993979 \end{array} \right. \begin{array}{l} \text{Julian Year.} \\ \text{Solar Year.} \\ \text{Difference.} \end{array}$

Or, $129^{\text{yrs}} 23^{\text{h}} 55^{\text{m}} 56^{\text{s}} 11^{\text{th}} 13^{\text{fo}} 25^{\text{fl}}$ &c. or about 130 Years, in which the Seasons, or Equinoxes, fall back, or happen *earlier* by 1 Day in the *Julian Account*.

This falling back of Seasons causing great Confusion in settling the Time of *Easter*, according to its original Institution, Pope *Gregory XIII.* took in Hand to reform the Calendar, in 1582; whose Example was followed in 1752, by the *British* Parliament.

In the Time of the *Nicene Council*, Anno Christi 325, the *Vernal Equinox* is said to have happened (but was rather erroneously determined) on the 21st of *March*: For, by *Halley's Astronomical Tables* it happened on *March* the 20th, 11^h 25^m in the Forenoon; which, among Astronomers, is on *March* the 19th 23^h 25^m P. M. and not on the 21st of *March*, according to *Error established*. Besides, we find the *Vernal Equinox*, according to *Halley*, fell on *March* the 19th, 15^h 54^m 30^s apparent Time, in the Year 1756 (*See Palladium for that Year*) which corresponds with

DIVISION of TIME.

March the 20th, 3^h 54^m 30^s, Morning; and fell about 44^m 21^s, later in 1752, that Year for which the Old Stile was corrected to correspond with the new one, or foreign Account of Time, and likewise adapted to the Error of the Equinox, falling on the 21st, instead of on the 20th of March, as it fell at the Nicene Council, and falls at this Time, though the Alteration of our Stile to correspond with the foreign Account was the principal View. Pope Gregory XIII. first settled the New Stile 1582, who called 5th October the 15th for that Year, thereby striking 10 Days out of the Calendar. For, dividing 1257 Years since the Nicene Council, by 130 Years, in which the Seasons fall back a Day, the Quotient will be above 9½ Days, the Seasons were then fallen back. From hence, 1257 Solar Years, since the Nicene Council appear to have been compleated, besides 9½ Days, in 1257 Years of the Julian Reckoning; for which Reason 10 Days lost in the solar Account of Time, were first struck out of the Julian Calendar, since made to correspond with solar Time.

And to prevent (as much as possible) the falling Back of the Seasons in the Month Days for Alteration of Style. the future, and to keep a Conformity between the Gregorian and Solar Account of Time, (the one in whole Days, and the other in Days and Parts, to the Year) Pope Gregory farther ordered, that every Fourth Hundred Year, from 1600, should consist of 366 Days, as usual; but that every Three Hundred Years or Centuries, in Succession from even Hundreds, should consist of 365 Days only (similar to three successive Years after each Bissextile) and not contain 366 Days each, as formerly. Thus he provided for 3 Days, that the Seasons fall back in every 400 (instead of 390) Years. See p. 30, 31.

In this NEW ACCOUNT of Time, called the GREGORIAN or NEW STILE, from Pope Gregory it's Author, the Seasons fall back but one Day in 5200 Years.

For, they fall back in 5200 Years by Julian Account = $\frac{5200}{130} = 40$ Days.

They go forward in 5200 Years, by the Gregorian Account = $\frac{5200}{400} \times 3 = 39$ —

Difference, they fall back in 5200 Years Gregorian Account . . . 1 Day.

Or, because in 400 Gregorian Account, but 1 Day of the Season's falling back in the Julian Account is considered, and provided for, which happens in 390 Years, 10 Years, in every 400 Gregorian, are unprovided for, in which the Seasons fall Back; and therefore $\frac{130}{10} = 13$, Times 400 Years = 5200 Years Gregorian, as before, in which the Seasons fall back 1 Day.

Anno Chr.
The Old Stile was corrected in England to the New, for . . . 1752
The Year of the Nicene Council 325

Difference. The Years from Nicene Council to our Alteration of Stile . . 1427

Now, $\frac{1427}{130} = 11$ Days, nearly, the Seasons fall back from the Time of the Nicene Council, to 1752, the Time when the Stile was altered in England.

Which 11 Days lost in the solar Account of Time, were supplied in the Gregorian Account, in our Calendar, by calling the 3d of September 1752, the 14th of that Month. This Alteration of Style reduced the Seasons (with regard to the Month Days) as they stood at the Council of Nice: For 1427 solar Years and 11 Days over are nearly compleated in that Time instead of 1427 Years only, reckoned by the Julian Account.

Hence it appears, that reducing the same Number of Days into different Years of any Kind, does not give a different Value to those Years in either Account of the same Time; which is contrary to common Opinion that those Days are lost. For, whether we reckon the same Number of Shillings, in Guineas, at Twenty Shillings and Six Pence, or at 21 Shillings each Guinea, the different Number of Guineas by either Account, in the same Number of Shilling Pieces, will have but the same Value, similar to the same Number of Days in a different Number of Solar, Julian, and Gregorian Years.

Of YEARS.

The civil or common Year, is of different Length, according to the Custom among different Nations. Some Nations reckon this Year by the Solar and some by the Lunar Motion.

The Civil Year, in most Parts of Europe, contains 365 Days for three Years successively, called common Years, and every 4th Year contains 366 Days, called Leap Year, or Bissextile.

These Civil Years are called also Julian Years, from Julius Cæsar, who added a Day every 4th Year to make (as he expected) the Civil and Solar Account of Time keep Pace together; and keep the Seasons, by that Means, nearly to the same Days of the Month.

The civil or common lunar Year is likewise Complete, or Vacant. The Complete consists of 354 Complete. Vacant. Days, at the End of which the Year begins again. The Vacant, or Embolismic Year, is that wherein a Month is added to make the Lunar correspond with the solar Account of Time. By this Method, the Jews kept their Account of Time, according to the Lunar Motion. But, by adding no more than a Month of 30 Days, called Ve-Adar, every third Year, in their Account, it fell Short of the solar Reckoning, in that Time, by about 3½ Days.

The

DIVISION of TIME.

The first Romans used the *Lunar, Vacant, or Embolimic, Year*, as it was settled by *Romulus* the first King, and Founder of *Rome*, who made it consist of ten *Lunar Months*, or *Lunations*, and therefore terminated 61 Days short of the solar Year; by which the Seasons in the *Roman Year* were unfixed, and wandering, in the Month days on which they fell: Whence they had a Table published by their *High Priest* to appoint the Time of their Spring and other Seasons. *Julius Cæsar*, considering this Difficulty, reformed the *Calendar* (as aforesaid) by making the Year consist of 365 Days and Six Hours, reckoning the Hours every fourth Year only, consisting of 366 Days. But the *Julian Account* of Time, thus stated, required a still farther Correction, by the *Equinoxes* falling back about a Day in 130 Years, and unfixing the Times of the Seasons, and the Fall of *Easter*, and consequently of the other *Feasts* thereon depending. Therefore *POPE GREGORY XIII.* took the *Calendar* in Hand, and, in the Year 1582, ordered (as before observed) the 5th of *October* to be called the 15th. By this Means he restored 10 Days the Seasons had retreated, since the *Nicene Council, Anno Christi 325*, and reduced the *Julian* to the *Gregorian Account* of Time. Who, by reckoning 3 Days less in 400 *Gregorian Years*, for the Time to come, made the *civil* or *political*, nearly to keep Pace with the *Solar Years*.

Of MONTHS.

Month. The MONTHS are of two Sorts, *astronomical* and *civil*.

Astronomical. The *astronomical Month* is the Time in which the Moon goes through the *Zodiac*, and beyond; and is either *periodical* or *synodical*.

Civil. The *Periodical Month* is the Time in which the Moon makes a complete Revolution from any Point in the *Ecliptic* to the same again, which, at a Mean Rate of Motion, is in $27^d 7^h 43^m 5^s$, as has been computed.

Periodical. The *Synodical Month*, called also a *Lunation*, is the Interval of Time between two next Conjunctions of the Moon and Sun, being $29^d 12^h 44^m 3^s$, at a mean Rate of Motion, as has been computed.

Synodical. The *Civil Months* are limited for the Purpose of civil Life; and differ in their Names, Number of Days, also Beginning and Ending, according to the Custom in different Nations.

Civil. The first Month of the *Jewish Year* happened according to the Moon, in our *August* and *September*; the Second *Jewish Month* in *September* and *October*, &c. O. S.

Antient. The First Month of the *Egyptian Year* began on the 29th of our *August*, O. S.

The First Month of the *Arabic and Turkish Year* began the 16th of our *July*, O. S.

The First Month of the *Grecian Year* happened, according to the Moon, in our *June* and *July*; the Second in *July* and *August*, &c. O. S. as set down in the following Table, useful in *antient Astronomy*.

DIVISION of the YEAR by the ANTIENTS.

The JEWISH YEAR.			The EGYPTIAN YEAR.			The ARABIC and TURKISH YEAR.			The ANTIENT GRECIAN YEAR.		
No.	Jewish Months.	Fell on our Ms, O. S. Days	No.	Egyptian Ms.	Fell on our Ms, O. S. Days	No.	Arabian Mths.	Fell on our Ms, O. S. Days	No.	Old Grecian Months.	Fell on our Ms, O. S. Days
1	Tisri . . .	Aug. Sept. 30	1	Tbotb . . .	August 29 30	1	Muharram . .	July 16 30	1	Hecatombæon	June July 30
2	Marchesvan . .	Sep. Oct. 29	2	Paopbi . . .	Sept. 28 30	2	Sapbar . . .	August 15 29	2	Metagitnion	July Aug 29
3	Casseu . . .	Oct. Nov. 30	3	Atbir . . .	Oct. 28 30	3	Rabia I. . .	Sept. 13 30	3	Bædromion	Aug. Sept. 30
4	Tebeth . . .	Nov. Dec. 29	4	Cbojac . . .	Nov. 27 30	4	Rabia II. . .	Oct. 13 29	4	Pyaneption	Sept. Oct. 29
5	Sbebat . . .	Dec. Jan. 30	5	Tybi . . .	Dec. 27 30	5	Jomada I. . .	Nov. 11 30	5	Mæmactætion	Oct. Nov. 30
6	Adar . . .	Jan. Feb. 29	6	Mecbir . . .	Jan. 26 30	6	Jomada II. . .	Dec. 11 29	6	Pofideon	Nov. Dec. 29
7	Nisan, or Abib	Feb. Mar 30	7	Phamenoth . .	Feb. 25 30	7	Rajab . . .	Jan. 9 30	7	Gamelion	Dec. Jan. 30
8	Jiar . . .	Mar. Apr. 29	8	Parmutbi . .	March 27 30	8	Shaftan . . .	Feb. 8 29	8	Antbresterion	Jan. Feb. 29
9	Sivan . . .	Apr. May 30	9	Pachon . . .	April 26 30	9	Ramadan . .	March 8 30	9	Elapheobolion	Feb. Mar. 30
10	Tamuz . . .	May June 29	10	Payni . . .	May 26 30	10	Sbarwal . . .	April 8 29	10	Munichion	Mar. Apr. 29
11	Ab . . .	June July 30	11	Epiphi . . .	June 25 30	11	Dulbaadab . .	May 7 30	11	Tbargelion	Apr. May 30
12	Elul . . .	July Aug. 29	12	Mefori . . .	July 25 30	12	Dulbeggia . .	June 5 29	12	Schirrophorion	May June 29
Days in the Year 354			Epagomenæ, or Days added 5			Days in the Year . . 354			Days in the Year 354		
In the Embolimic Year, after Adar, they added a Month called Ve-Adar, of 30 Days. But the other 12 Months last 30			Days in the Year . . 365			They add 11 Days at the End of each Year to keep the same Months to the same Seasons.			N. B. We would have inserted the Names of Week-Days used by each Nation; but wanted good Authority.		

Common. A common Month is divided into 4 Weeks; of which Kind of Months there are 13, or 52 Weeks, in a *Julian Year*; besides 1 Day over, or 2 in *Bisextile*.

Week. A WEEK is divided into 7 Days, whose Names are known. The antient *Gentiles* gave Names to the Days of the Week, from the Names of the *Planets*, which they supposed to preside over them. The 1st Day of the Week they denominated *Sun*, the 2d *Moon*, the 3d *Mars*, 4th *Mercury*, 5th *Jupiter*, 6th *Saturn*.

DIVISION of TIME.

Of DAYS.

- Day.** A DAY is either *natural* or *artificial*, in Propriety of Distinction.
- Natural.** The *Natural* Day is divided into 24 Hours, from the same to the same Time.
- Artificial.** The *Artificial* Day is the Time from the Sun Rising to the Sun Setting of that Time.
- The *Natural* Day is *Astronomical* or *Civil*.
- Astronomical.** The *Astronomical* Day begins at Noon and ends the Noon following. Astronomers find this Limit, of Beginning and Ending Day, fittest for their Purpose of Computation, by the diurnal Revolution of the Sun from Meridian to Meridian. For the Days would be much more *unequal* (as was observed) if they begun and terminated at Sun Rising or Setting, or any other Time of the Sun's diurnal Revolutions, except at *Midnight*; which yet would not, so well as Noon to begin Time at, answer the Purpose of Calculation.
- The *British, French, Dutch, Germans, Spanish, Portuguese, and Egyptians*, in common Reckoning, begin their Day at *Midnight*. The *ancient Greeks, Jews, Bohemians, Silesians, and modern Italians and Chinese*, did, and now begin Day at Sun Setting.
- The *ancient Babylonians, Persians, Syrians, and modern Greeks*, did, and now begin Day at Sun Rising, according to the civil Custom of each Nation.
- N. B. The *Italians* reckon the *natural* Day by 24 Hours of the Clock. Other Nations, by reckoning twice twelve of the Clock, prevent their Clocks striking 24, (like the *Italian* Clocks) at one Time.

Of HOURS.

- Hour.** An HOUR is divided into 60 Parts called *Minutes*.
- The Hour is either equal or unequal.
- Equal.** An *equal* Hour is the 24th Part of a mean Day, measured by a well regulated Clock, or Watch.
- Unequal.** An *unequal* Hour of a *natural* Day, is measured by the Sun's *unequal* Motion in the *Ecliptic*, inclined to the Equator in an Angle of $23^{\circ} 28' \frac{1}{2}$, while 15° , and the $\frac{1}{24}$ of the Increase of R.
- A. for a Day, counted on the Equinoctial, successively passes the Meridian.
- An *unequal* Hour is otherwise the $\frac{1}{24}$ Part of the *Artificial* Day from Sun Rising to Setting; and is also the $\frac{1}{24}$ Part of the Night, from Sun Setting to Rising, as reckoned by the *Jews*. These *unequal* or *Jewish* Hours, increase and decrease with the Length of Days and Nights. For Length of Jewish or unequal Hours, see Ladies Diary for former Years, by the late Mr. Henry Beighton, F. R. S.

Of MINUTES and SECONDS.

- Minute.** A MINUTE of Time is divided into 60 equal Parts, called *Seconds*, and each *Second* is divided into 60 equal Parts called *Thirds*, and so on, in a *Sexagesimal* Subordination.
- Second.** The *Jews, Chaldeans, and Arabians* divide the Hour into 1080 equal Parts, or *Scruples*, containing 18 Times 60; whereby one Minute contains 18 *Scruples*.

Of CYCLES, or PERIODS.

- A Cycle, or Period, is a Revolution of Time.
- Sun's Cycle.** The Cycle of the Sun, or *Dominical-Letter-Cycle*, is a Revolution of 28 Years, in which Interval of Time, the Days of the Week in the *Julian* Account return to the same Days of the Month. And the Sun's Place never varies every 7 Times 4, or 28 Years, above 7 Times $1' 53'' = 13' 11''$, on the same Month-day, not 50' in 100 *Julian* Years, according to *Ferguson's Astronomy*. After the Completion of this Cycle, the Week days begin the same Order over again, with respect to the same Month-days of the *Julian* Reckoning, which is only interrupted by the common Hundredth *Gregorian* Year, coming between. See Tables further on, of this Cycle for finding the *Dominical Letter* at Sight, according to both *Stiles*, for any Year before and since *Christ*.
- The Cycle of the Moon, commonly called the *Golden Number*, is a Revolution of 19 Years, in which Interval of Time, according to the *Julian* Account, (including the 6 odd Hours yearly) the Conjunctions, Oppositions, and other Aspects of the Moon with the Sun, are within an Hour and a Half of the same Situation as they were on the same Month-days in the preceding Cycle. But, this Cycle is interrupted a Day, like the Sun's Cycle, by the common Hundredth *Gregorian* Year coming between. See p. 31. And Tables farther on, for finding the *Golden Number* at Sight, according to both *Stiles*, for any Year before and since *Christ*.
- Indiction.** The *INDICTION-CYCLE* is a Revolution of 15 Years, only; used by the *Romans* for determining the Times of particular Payments by the Subjects to the Republic: Which Cycle was founded by *Constantine*, in the Year of *Christ* 312.
- See Rules for determining the Numbers of each of these Cycles, arithmetically, farther on.

Of the FIXED STARS.

They are so called, because of the *Fixed Distance* they are generally observed to keep from one another: Their apparent diurnal Revolution in the Heavens to the Right arising only from the Earth's Rotation to the Left on its *Axis*. They appear of different *Magnitudes* from their different, but immense, Distances, at which they are placed, in the infinite Space, from our Sight, and are seen much bigger by the naked Eye than through a *Telescope*, magnifying 200 Times, taking off the scattered Rays, causing a *Radiation* or *Scintillation* to our Sight; and thereby renders their true Appearance so many distinct Points of native Light. If the Stars shone with borrowed and not (as they do) with native Light, they would be as invisible without the Use of *Telescopes*, as the Satellites of *Jupiter*, any one of which appear bigger, when viewed by a *Telescope*, than the greatest of the *Fixed Stars*.

The Number of *Fixed Stars* discoverable by the naked Eye, in either the *Northern* or *Southern Hemisphere*, exceeds not a *Thousand*, while they appear, at first Sight, to be innumerable; but the Deception is in looking confusedly, and not distinctly, upon them.

The *BRITISH CATALOGUE*, consisting of a large Number of Stars, which cannot be seen without the Help of the *Telescope*, counts not above 3000 in both Hemispheres.

The Light of the *Moon* to this Earth in the Night-Time, being much superior to the faint glimmering Light of the Stars, it follows from thence, that the Stars were not designed for giving Light in the Night only; because many more than are visible to the naked Eye require the Help of a *Telescope* to discover them. And, as our *Sun* is surrounded with a System of Planets and Comets, which would be invisible from the nearest fixed Star, it is probable that every Star is a *Sun* to a remote System of planetary Bodies, though invisible to us at this Distance. This Plurality of innumerable Worlds quite confists with the infinite Power, Wisdom, and Greatness of the CREATOR OF ALL WE BEHOLD!

The Stars on account of their different Appearance, have been distributed into Six Classes, or Magnitudes; the 6th being the least Magnitude visible to the naked Eye. This Distribution being made long before the Invention of *Telescopes*, those Stars which cannot be seen without the Help of that Instrument are distinguished by the Name of *Telescopic Stars*.

Of CONSTELLATIONS.

The *Antients*, to distinguish the Stars, divided the *Starry Sphere* into different *Constellations*, or Collections of Stars, situated next one another, in the Heavens, by drawing about them the *Out-lines* of different Things and Animals, according as they supposed a Collection of Stars to represent. Those Stars, which could not be brought under any designed Forms, were denominated *unformed Stars*.

Those *Constellations*, and likewise *unformed Stars*, are now represented on a *Celestial Globe*, by which any particular Star, and its Situation, in Respect of another, is readily known and distinguished, by giving the most remarkable Stars the most remarkable Places.

The Number of the *antient Constellations* is 48, and the Number on our *modern Globes* about 70. On *MARTIN'S Globes*, being an Improvement on *SENEC'S*, are inserted *Bayer's Letters*. The first Letters of the Greek Alphabet are put to denote the Magnitudes of the Stars in each Constellation, nearly according to the Order of Magnitude from α , denoting the Star of the first Magnitude; β denoting that of the second, &c. where-by the particular Magnitudes, and Names of the Stars are as truly defined and distinguished, as if they were shewn us in the Heavens. See *MARTIN'S GLOBES*, in Fleet-street, London.

The *Starry Heavens* are divided into three Parts, 1st, the *Zodiac*, of 16 Degrees Breadth, quite round the Heavens, taking all the planetary Orbits, and Orbit of the *Moon*: Containing 12 Constellations; *Aries, Taurus, Gemini, Cancer, Leo, Virgo, Libra, Scorpio, Sagittarius, Capricorn, Aquarius, and Pisces*. In the middle of this Zone, or Belt, is the *Ecliptic*, or Circle which the Earth annually describes, as seen from the Sun; and which the Sun appears to describe, as seen from our Earth. 2d, The Region of the Heavens on the North Side of the *Zodiac*, containing 21 Constellations. 3d, All the Region on the South Side, containing 15 Constellations.

The *Antients* divided the *Zodiac* into twelve Parts by the equal Dropping of twelve Parts of Water, (at each of which Parts dropped out, they noted a particular Star's Rising) the whole 12 Parts having first dropped through a small Vessel into a Receiver, from the observed rising of the same Star to its observed Rising the Night following.

The Names of the *Constellations*, and different Stars, observed in each Hemisphere, by different Astronomers, since the Improvement of Astronomy began, are as follow.

ANTIENT CONSTELLATIONS.

Numb.	Latin Names.	English Names.	Ptolemy.	Tycho.	Hevelius.	Flamsteed.
1	Ursa Minor	The Little Bear	8	7	12	24
2	Ursa Major	Great Bear	35	29	73	87
3	Draco	Dragon	31	32	40	80
4	Cepheus	Cepheus	13	4	51	35
5	Bootes, Arctophilax	Bootes	23	18	52	54
6	Corona Borealis	Northern Crown	8	8	8	21
7	Hercules	Hercules kneeling	29	28	45	113
8	Lyra	Harp	10	11	17	21
9	Cygnus, Gallina	Swan	19	18	47	81
10	Cassiopea	Lady in her Chair	13	26	37	55
11	Perseus	Perseus	29	29	46	59
12	Auriga	Waggoner	14	9	40	66
13	Serpentarius, Ophiuchus	Serpentarius	29	15	40	74
14	Serpens	Serpent	18	13	22	64
15	Sagitta	Arrow	5	5	5	18
16	Aquila, Vultur	Eagle	15	12	23	71
17	Antinous	Antinous	10	3	19	18
18	Delphinus	Dolphin	4	4	6	10
19	Equulus, Equisetio	Horse's Head	20	19	38	89
20	Pegasus, Equus	Flying Horse	23	23	47	66
21	Andromeda	Andromeda	4	4	12	16
22	Triangulum	Triangle				

ANTIENT

FIXED STARS.

ANTIEN'T CONSTELLATIONS continued.

Numb.	Latin Names.	English Names.	Ptolemy.	Tycho.	Hevelius.	Flamsteed.
23	Aries	Ram	18	21	27	66
24	Taurus	Bull	44	43	51	141
25	Gemini	Twins	25	25	38	85
26	Cancer	Crab	23	15	29	83
27	Leo	Lion	35	30	49	95
28	Coma Berenices	Berenices Hair	35	14	21	43
29	Virgo	Virgin	32	33	50	110
30	Libra, Chelæ	Scales	17	10	20	51
31	Scorpius	Scorpion	24	10	20	44
32	Sagittarius	Archer	31	14	22	69
33	Capricornus	Goat	28	28	29	51
34	Aquarius	Water Bearer	45	41	47	108
35	Pisces	Fishes	38	36	39	113
36	Cetus	Whale	22	21	45	97
37	Orion	Orion	38	42	62	78
38	Eridanus, Fluvius	The River	34	10	27	84
39	Lepus	Hare	12	13	16	19
40	Canis Major	Great Dog	29	13	21	31
41	Canis Minor	Little Dog	2	2	13	14
42	Argo Navis	Ship	45	3	4	64
43	Hydra	Hydra	27	19	31	60
44	Crater	Cup	7	3	10	31
45	Corvus	Crow	7	4		9
46	Centaurus	Centaur	37			35
47	Lupus	Wolf	19			24
48	Ara	Altar	7			9
49	Corona Australis	Southern Crown	13			12
50	Pisces Australis	Southern Fish	18			24

2777

NEW SOUTHERN CONSTELLATIONS, discovered by MARINERS.

Numb. Const.	Latin Names.	English Names.	Stars Flamst.
1	Columba Noachi	Noah's Dove	10
2	Robur Carolinum	The Royal Oak	12
3	Grus	Crane	13
4	Phœnix	Phœnix	13
5	Indus	Indian	12
6	Parvo	Peacock	14
7	Apus, Avis Indica	Bird of Paradise	11
8	Apis, Musca	Bee or Fly	4
9	Chamaeleon	Chameleon	10
10	Triangulum Australis	South Triangle	5
11	Piscis volans, Passer	Flying Fish	8
12	Dorado, Xiphias	Sword Fish	6
13	Toucan	American Goose	9
14	Hydrus	Water Snake	10

Sum 137

HEVELIUS's CONSTELLATIONS composed of unformed Stars.

Numb.	Latin Names.	English Names.	Hevelius.	Flamst.
1	Lynx	The Lynx	19	44
2	Leo Minor	Little Lion		53
3	Asterion and Cbara	Greyhound	23	25
4	Cerberus	Cerberus	4	
5	Vulpecula & Anser	Fox and Goose	27	35
6	Scutum Sobieski	Sobieski's Shield	7	
7	Lacerta	Lizard	10	16
8	Camelopardus	Camelopard	32	58
9	Monoceros	Unicorn	19	31
10	Sextans	Sextant	11	41

303

Milky Way.

The Luminous Track, round the Heavens, appearing single in some Places, and double in others, called the Milky Way, from its Whiteness, is caused by the numerous Stars scattered therein, (powdered with Stars as Milton terms it) none of which can be distinctly seen but by the Means of good Telescopes.

Of LUCID and OTHER SPOTS in the HEAVENS.

Several luminous Spots, in the Heavens, appear magnified, and more luminous, being seen through good Telescopes; but are void of Stars. One of which Spots is seen in Andromeda's Girdle, first observed in the Year 1612, by Simon Marius, with whitish Rays near its Middle. This luminous Spot is subject to several Changes, and is sometimes invisible. Another of these Spots is seen near the Ecliptic, between the Head and Fore of Sagittary; being small, but very luminous. A Third is seen on the Back of Cancer, being too South to be visible in England. A Fourth, of a smaller Size, is seen before Antinous's Right Foot; with one Star in it, by which it appears the brighter. A Fifth is seen in the Constellation of Hercules, between the Stars of Bayer, ζ , and η , being visible, though appearing but small, to the naked Eye, when the Sky is clear, and the Moon absent.

There

FIXED STARS.

There are (besides the Appearances spoke of) *cloudy Stars* in the Heavens, so called from their appearing of a dim and misty Light to the naked Eye; but appear through a Telescope, to be broad illuminated Parts of the Sky, containing one, or more Stars. Five of these cloudy Stars are mentioned by Ptolemy. 1. One, at the Extremity of the Right-hand of Perseus. 2. One, in the midst of the Crab. 3. One, unformed, near the Sting of the Scorpion. 4. The Eye of Sagittary. 5. One, in the Head of Orion. In the first of these more Stars appear, through the Telescope, than in any of the rest; though 21 have been reckoned in Orion's Head; and above 40 in the Crab. Two are visible in the Eye of Sagittary, without a Telescope; but several more appear by the Use of that Instrument. Flamsteed observed a cloudy Star in the Bow of Sagittary, containing many small Stars. Cassini and Flamsteed observed one between the Great and Little Dog, appearing full of Stars, only visible by the Telescope. The two whitish Spots near the South Pole, called the Magellanic Clouds. Magellanic Clouds, by Sailors, resembling the Milky Way, to the naked Eye, appear through Telescopes to be a Mixture of small Clouds and Stars.

But, the most remarkable of all the cloudy Stars is in the middle of Orion's Sword, where Seven Stars (three of which are close together) seem to shine through a Cloud, very lucid near the middle; but faint and imperfect about the Skirts: looking like a Gap in the Sky, through which appears Part of a brighter Region. Though most of the cloudy Spaces are but a few Minutes of a Degree in Breadth, yet, as they are among the Fixed Stars, they must be every one larger Spaces than our Solar System occupies; as in which there seems to be a perpetual uninterrupted Day, among innumerable Worlds.

OF NEW PERIODICAL STARS.

Several Stars are described by *antient Astronomers* not now to be found; and others are now visible to the naked Eye not recorded in *antient Catalogues*. Hipparchus observed a new Star about 150 Years before Christ, who has not mentioned in what Part of the Heavens it was seen; though it occasioned his making a Catalogue of the Stars the most antient of any that we now have.

The first new Star that we have any certain Account of was discovered by Cornelius Gemma, on the 11th of November, 1572, in Cassiopea's Chair. It surpassed Sirius in Magnitude and Brightness, and was seen for 16 Months in Succession. It appeared, at first, bigger than Jupiter to some Eyes, by which Means it was seen at Mid-Day; afterwards it decayed gradually in its Lustre and Magnitude, till March 1573, when it became invisible.

On the 13th of August, 1596, David Fabricius observed the *Stella Mira*, or wonderful Star, in the Neck of the Whale, since disappearing and appearing, periodically, seven Times in six Years, continuing in its greatest Lustre, for 15 Days together, which is never quite extinguished.

In the Year 1600, William Jansenius discovered a changeable Star in the Neck of the Swan, which, in Time, became so small, as made it thought to have disappeared till the Years 1657, 1658, and 1659, when it recovered its former Lustre and Magnitude; but soon decayed in both, and is now of the smallest Size.

In the Year 1604, Kepler, and several of his Friends, saw a new Star near the Heel of the Right Foot of Serpentarius, so bright and sparkling, that it exceeded any Thing of the Kind they before had seen, who observed that it was every Moment changing into some of the Colours of the Rainbow, except when it was near the Horizon, and generally appeared white. It exceeded Jupiter in Magnitude, which it was near during the Month of October, but distinguished from Jupiter by his steadier Light. This Star disappeared between October 1605 and February following, and has not since appeared.

In the Year 1670, July the 15th, Hevelius discovered a new Star, which, in October following was so decayed as to be hardly perceptible. In April following, it regained its former Lustre; but wholly disappeared in August. In March 1672 it appeared again very small, disappearing ever since.

In the Year 1686, a new Star was discovered by Kirch, disappearing and returning periodically in 404 Days.

In the Year 1672, Cassini observed a Star in the Neck of the Bull, which he judged was not visible in Tycho's Time, nor yet when Bayer made his Catalogue.

OF CHANGES in the HEAVENS.

Many Stars, besides those before-mentioned, have been observed to change their Magnitudes and Appearances; but as none of them were ever observed to have Tails, 'tis concluded they could none of them be Comets; having no Parallax in their greatest Lustre and Magnitudes. It appears probable, that these periodical Stars, having vast Clusters of dark Spots, make slow Rotations on their own Axes; by which Means they disappear when the Side covered with Spots is turned towards us. And those Stars, breaking out of a sudden, with such resplendent Brightness, are probably Suns of other Systems, whose Fuel being much exhausted, acquire their Blaze and Splendor, continuing for some Time by the Accession of the Comets of that System, lodging on their Surface. For, this (according to Sir Isaac Newton) appears to be the greatest Use and End of the Cometary Part of a System.

And M. Maupertuis, in his Dissertation of the Figures of the Celestial Bodies (p. 61 to 63) is of Opinion, that some Stars, by their prodigious swift Rotation on their Axes, assume not only the Figures of flatted Globes next their Axes, but, by the great centrifugal Force, arising from such swift Rotations, may become of the Figure of Mill-stones, or even be reduced to flat circular Planes, so thin, as to be quite invisible when their Edges are turned towards us: As Saturn's Ring is invisible in such Positions. That, when very excentric Planets, or Comets, go round any flat Star in Orbits much inclined to its Equator, the Attraction of the Planets or Comets, in their Perihelion, must alter the Inclination of the Star; on which Account, it will appear more or less large and luminous, as its Broadside is more or less turned towards us. And thus he thinks we may account for the apparent Changes of the Magnitude and Lustre of those Stars; as likewise for their appearing and disappearing at certain Intervals.

Some of the Stars, particularly Arcturus, have been observed to change their Places, in the Heavens, above a Minute of a Degree, in Respect of the Situation of other Stars. But whether this is owing to any real Motion of the Stars themselves is the Business of many Ages to determine.

If our Solar System changes its Place, with Regard to absolute Space, it must, in Time, occasion an apparent Change in the Distances of the Stars from one another. And, in such a Case, the Places of the nearest Stars to us being more affected than those at a greater Distance, their relative Positions will seem to alter, though the Stars themselves were really immovable. On the other hand, if our System remains at Rest, and any of the Stars of other Systems, have real Motion, in respect of infinite Space, their Positions will be changed thereby, and apparent Places, more or less, according as those Stars are nearer or further from us; and the swifter or slower those Motions are, and their Directions more or less suited to our Perception. And the same will happen from real and different Motions in the Systems themselves.

The Obliquity of the Ecliptic to the Equinoctial is, at present, about one third part of a Degree less than Ptolemy determined it. For most of the Astronomers after him found this Obliquity to decrease gradually down to Tycho's Time. It is objected that we cannot depend on the Observations of the antient Astronomers, on Account of the Incorrectness of their Instruments, it may be urged, that Tycho and Flamsteed were both able Observers, and yet Flamsteed makes the Ecliptic Obliquity about $2\frac{1}{2}$ Min. of a Degree less than Tycho did, about 100 Years before; which Difference can hardly arise from the different Cor-

FIXED STARS.

rectness of the Instruments with which they observed. For, Ptolemy lived 1324 Years before Tycho; and the gradual Decrease of Ecciptic Obliquity, nearly corresponds to the Difference of Time between these three Great Astronomers. Though the Variation of the Obliquity is shewn by Dr. Bradley, to depend on the Place of the Moon's Node through its Revolution, (See Tab. p. 33.) Yet, if we consider that the Earth is not a perfect, but a flattened Sphere, whose shorter Axis is less than its Equatorial Diameter, and that the Sun and Moon are continually and obliquely acting upon the greater Mass of Matter about the Equator, and attracting it to a nearer and nearer Coincidence with the Ecciptic, it will appear in some Degree probable, that the Attractions constantly diminish the Angle between the Planes of the Ecciptic and Equator. Nor is it less probable, (as the World is to be destroyed, or come to a Period, according to the Scriptures) that the mutual Attractions of all the Planets have a Tendency to bring the Planes of all their Orbits to a final Coincidence: Though the Effect be so small as not to produce a sensible Change in many Ages. See Ferguson's Astronomy.

—The Stars shall fade away, the Sun Himself
Grew dim with Age, and Nature sink in Tears,
But thou shalt flourish in immortal Youth,
Unhurt amidst the War of Elements,
The Wrecks of Matter and the Crust of Worlds!

CATO.

As Astronomers begin their Computations from certain fixed Points in the Heavens, called *Epochs*, or Radical Places; so Historians begin their Reckonings from certain fixed Points of Time, called *Æras*, or Radices of Time. The most remarkable Radices of Time are the *Æras* of Creation, the Greek *Olympiads*, the Building of Rome, of Nabonassar, the Death of Alexander the Great, the Birth of Christ, the Arabian Hegyra, and Persian Yesdegerd. These, and others of less Note, are determined according to the Julian Period, Age of the World, and Years before and since Christ, as in the following Table.

REMARKABLE ÆRAS and EVENTS.
According to the Years of the Julian Period, of Creation, and of Christ.

N ^o .	Remarkable Æras, or Events.	Yrs. Jul. Per.	Yrs. of Cre.	Yrs. bef. Chr.	N ^o .	Remarkable Æras, or Events.	Yrs. Jul. Per.	Yrs. of Cre.	Yrs. bef. Chr.
THE					THE				
1	Creation of the World, by Strauchius . . .	764	1	3949	26	Persian Monarchy founded by Cyrus . . .	4154	3390	559
2	Deluge, or Noah's Flood . . .	2420	1656	2293	27	Battle of Maratbon . . .	4224	3460	489
3	Assyrian Monarchy, by Nimrod . . .	2537	1733	2176	28	Beginning of the Reign of Art. Longimanus . . .	4249	3485	464
4	Birth of Abraham . . .	2712	1948	2001	29	Beginning of Daniel's 70 Weeks . . .	4250	3492	457
5	Beginning of the Kingdom of Argives . . .	2856	2092	1857	30	Beginning of the Peloponnesian War . . .	4282	3518	431
6	Beginning of the Kingdom of Athens, by Cecrops . . .	3157	2393	1556	31	DEATH of ALEXANDER the GREAT . . .	4390	3626	323
7	Departure of Israelites from Egypt . . .	3216	2452	1497	32	Restoration of the Jews . . .	4548	3784	129
8	Their Entrance in Canaan, or Jubilee . . .	3256	2492	1457	33	Correction of the Calendar by Julius Cæsar . . .	4669	3905	47
9	Destruction of Troy . . .	3529	2865	1184	34	Beginning of the Reign of Herod . . .	4673	3909	40
10	Beginning of King David's Reign . . .	3653	2889	1060	35	Spanish Æra . . .	4675	3911	38
11	Foundation of Solomon's Temple . . .	3696	2932	1017	36	Battle at Actium . . .	4683	3919	30
12	Argonautic Expedition . . .	3776	3012	937	37	Taking of Alexandria . . .	4683	3919	30
13	Arbaces, the I. King of the Medes . . .	3838	3074	875	38	Æra of the Title of Augustus . . .	4686	3922	27
14	Mandaucus the II. . .	3865	3101	848	39	TRUE ÆRA of CHRIST'S BIRTH . . .	4709	3945	4
15	Sofarmus III. . .	3915	3151	798	40	Death of Herod . . .	4710	3946	3
16	Artica IV. . .	3945	3181	768	41	Reputed Æra of CHRIST, by Dionysius exiguus . . .	4713	3949	0
17	Cardica V. . .	3996	3232	718					since
18	Pbraortes VI. . .	4057	3293	656	42	True Year of CHRIST'S DEATH . . .	4746	3982	33
19	Cyaxares VII. . .	4080	3316	633	43	Destruction of Jerusalem . . .	4783	4019	70
20	BEGINNING of the Greek OLYMPIADS . . .	3938	3174	775	44	Dioclesian Persecution . . .	5015	4251	302
21	Cætonian Æra of building ROME . . .	3961	3179	752	45	Æra of Constantine the GREAT . . .	5019	4255	306
22	ÆRA of NABONASSAR . . .	3967	3202	746	46	Council of Nice . . .	5038	4274	325
23	Destruction of Samaria . . .	3990	3226	723	47	ÆRA of HEGIRA . . .	5335	4571	622
24	Babylonish Captivity . . .	4133	3349	600	48	ÆRA of YESDEGERD . . .	5344	4580	631
25	Destruction of Solomon's Temple . . .	4124	3360	589	49	Jallalæan Æra . . .	5791	5027	1078
					50	Æra of the Reformation . . .	6230	5466	1517

Of ÆRAS, CYCLES, and PERIODS.

THE Solar and Lunar Cycles of 28 and 19 Years, and Roman Indiction of 15 Years, multiplied together, produce the great Julian Period of 7980 Years; having its Beginning 764 Years before the supposed Æra of Creation, when all the three Cycles began together. And as the said Julian Period is not yet completed, it therefore comprehends all other Cycles, Æras, and Periods. And as there is but one Year in the whole Julian Period, having the same Numbers or Parts of the three whole Cycles, of which the said Period is composed, therefore, if Historians had marked, in their Writings, the Numbers of the three Cycles for the Year in which any remarkable Event, or memorable Transaction happened, there could have been no Dispute about the Time of its happening.

To find the Year of the Julian Period since Christ? The Dionysian, or reputed, Æra of Christ's Birth, being about the End of the 4713 Year of the Julian Period.

RULE. To the Current Year of Christ add 4713, the Sum will be the Julian Period.

To find the Year of the Julian Period before the reputed Coming of Christ?

RULE. Subtract the Number of the given Year before Christ from 4713, and the Remainder will be the Julian Period.

EXAMPLE

ERAS, CYCLES, and PERIODS.

EXAMPLE I. 1759 Years since Christ.
4713 Year of Julian Period at Christ's Coming.

Sum, 6472 Year of Jul. Period 1759 Yrs since Christ.

EXAMPLE II. 1759 Years before Christ.
4714 Year Julian Period at Christ's Coming.

Dif. 2956 Year Jul. Per. 1759 Years before Christ.

To find the Numbers of the three Cycles of the Sun, Moon, and Indiction for a given Year of the Julian Period.

RULE. Divide the given Year by 28, 19, and 15, respectively, and the three Remainders will be the Numbers of the Cycles sought. The three Quotients will be the Cycles completed since the Julian Period began.

Yr. Per.
Examples, 28) 4713 (168 Cycles since.

Rem. 9 Cycle of Sun,
or Dominical Letter,

Yr. Per.
19) 4713 (248 Cycles since.

Rem. 1 Cycle of Moon,
or Golden Number.

Yr. Per.
15) 4713 (314 Cycles since.

Rem. 3 Cycle of Indiction.
All at the Birth of Christ.

Of the BIRTH and DEATH of CHRIST.

THE common Era of Christ is 4 Years later than the true Era. For Christ was born before the Death of Herod the Great, who sought to kill him as soon as he heard of his Birth. And according to the Testimony of Josephus (B. 15. Ch. 8.) there was an Eclipse of the Moon in the Time of Herod's last Sickness, a little before his Death. Which Eclipse some Astronomical Tables shew to have happened in the 4710 Year of the Julian Period, March 13^d 3^h 21^m after Midnight, at Jerusalem. Now, Christ must have been born some Months before Herod's Death, because in the Interval (betwixt his Birth and Herod's Death) he was carried into Egypt, for Preservation of his Life. The latest Time therefore at which the true Era of the Birth can be fixed is about the End of the 4709th Year of the Julian Period, which is four Years before the reputed Era.

Christ died in the 4746 Year of the Julian Period on Friday, April 3, in the 33d Year of his Age, according to the reputed Era, or 37th Year according to the true Era, discovered by the Eclipse of the Moon happening a little before the Time of Herod's Death. He was put on the Cross, at Jerusalem, at 12 o'Clock (as we reckon Time) and expired about three in the Afternoon: the whole Neighbourhood of Jerusalem, being overspread with a miraculous Darkness, during the Time of his Suffering.

This miraculous Darkness, (on the 4th Year of the 202d Olympiad) Ptolemy, the Trallian (an Hebræan Writer) mentions as an Eclipse of the Sun that then happened, through his Want of Knowledge in Astronomy. For it was full Moon about that Time, by Astronomical Tables, and consequently there could be no Eclipse of the Sun.

For ample Account of the Time of Christ's Crucifixion and Age, at his Death, see p. 42. Palladium 1753.

The reputed Era of Christ's Birth was never settled till the Year of Christ 527: when Dionysius Exiguus, a Roman Abbot, fixed it at the End of the 4713th Year of the Julian Period; being four Years too late, for the Reason before given.

The Birth of CHRIST, according to Nicholas Man, Esq; in his Chronology, is proved to be just 6 Years before the said vulgar, or reputed, Era; the Eclipse of the Moon, a little before Herod's Death, happening on March the 13th, at 3 in the Morning.

But, that the ERAS of CHRONOLOGY may be examined and settled on the Foundation of Truth; we here give the Reader Sir ISAAC NEWTON'S ASTRONOMICAL PRINCIPLES OF CHRONOLOGY; by which he determines the GRAND ERA of the Argonautic Expedition as the Foundation of all his Chronology.

He observes, that Eudoxus, in his Account of the Spheres of the Antients, placed the Solstices and Equinoxes in the Middle of the Constellations of Aries, Cancer, Chelæ and Capricorn; and that this Sphere or Globe was first made by Museus, and the Asterisms delineated upon it by Chiron, two Persons of the Argonauts.

It has been shewn, by the Precession of the Equinoxes, contrary to the Order of Signs, that the fixed Stars appear to go forward, in the Signs, 50" a Year. And because at the End of the Year 1689, the Equinoctial Colure, passing through the middle Point, between the first and last Star of Aries, then did cut the Ecliptic in $8^{\circ} 6' 44''$, it is thence evident, that the Equinox had then gone back $36^{\circ} 44'$, and therefore, As 50" is to 1 Year, so $36^{\circ} 44'$, to 2645 Years, for the Time since the Argonautic Expedition, to the Beginning of the Year 1690, or 955 before Christ.

But, Sir Isaac Newton (in order to determine the Time more correctly) finds the mean Place of the Colure of the Equinoxes, and Solstices, by considering the several Stars they passed through among the different Constellations, according to Eudoxus, as follows.

In the Back of Aries is a Star of the 6th Magnitude, marked γ , by Bayer, in the End of the Year 1689, its Longitude was $8^{\circ} 9' 38' 45''$; and the Equinoctial Colure passing through, cut the Ecliptic, according to Eudoxus, in $8^{\circ} 6' 58' 57''$.

In the Head of Cetus are two Stars of the 4th Magnitude, called ν and ξ by Bayer: Eudoxus's Colure passing in the Middle between them, cut the Ecliptic in $8^{\circ} 6' 58' 51''$, at the End of the Year 1689.

In the extreme Flexure of Eridanus, there was formerly a Star of the 4th Magnitude, (of late referred to the Breast of Cetus) being the only Star in Eridanus, through which the Colure can pass. Its Longitude was, at the End of the Year 1689, $7^{\circ} 25' 22' 10''$, and the Colure of the Equinox passing through it cut the Ecliptic in $8^{\circ} 7' 12' 40''$.

In the Head of Perseus, rightly delineated, is a Star of the 4th Magnitude, called τ by Bayer; its Longitude was $8^{\circ} 23' 25' 30''$, at the End of the Year 1689; and the Colure of the Equinox, passing through it, cut the Ecliptic in $8^{\circ} 6' 18' 57''$.

In the Right Hand of Perseus, rightly delineated, is a Star of the 4th Magnitude, whose Longitude, at the End of the Year 1689, was $8^{\circ} 24' 25' 27''$, and the equinoctial Colure, passing through it, cut the Ecliptic in $8^{\circ} 4' 56' 40''$.

The Sum of all these	$8^{\circ} 6' 58' 57''$
Five Places of the	$8^{\circ} 6' 58' 51''$
Colure	$8^{\circ} 7' 12' 40''$
	$8^{\circ} 6' 18' 57''$
	$8^{\circ} 4' 56' 40''$

Is . . . $1^{\circ} 2' 26' 5''$

The 5th Part of which . . . $8^{\circ} 6' 29' 13''$, is therefore the mean Place, in which the Colure in the End of the Year 1689, did cut the Ecliptic: Taking the Mean of a Number of Observations.

After this Manner Sir Isaac Newton determines the Mean Place of the Solstitial Colure to be in $9^{\circ} 6' 28' 46''$, which being very nearly 90° from the other Colure, shews the Deduction to be rightly made.

The Equinoxes, at this Time, having departed $1^{\circ} 6' 29'$ from the Cardinal Points of Chiron, shew that 2628 Years have elapsed, since that Time, which is correcter than the former Number of Years, (though less by only 17) for the Time, since the Argonautic Era to the Year of Christ 1689; making the Argonautic Era about 938 Years before Christ.

By other similar Methods Sir Isaac Newton shews, that this Era of the Argonauts ought to be placed in that Age of the World. And having determined this most Antient Era, he makes his future Computations in Chronology with Reference thereto. Who, with great Skill and Penetration, has suited his Chronology to agree with the Course of Nature, the Principles of Astronomy, with Sacred History, with Herodotus, the Fathers of profane History, and with itself. Which great Author is therefore not to be contradicted but by his EQUALS in Judgment, or such, at least, as understand him.

CHRONOLOGICAL TABLES.

DOMINICAL LETTERS for Julian or OLD STYLE, for ever.

YEARS before CHRIST.										YEARS since CHRIST.									
CENTURIES										CENTURIES									
YEARS above		CENTURIES		YEARS before		CENTURIES		YEARS since		CENTURIES		YEARS above		CENTURIES		YEARS before		CENTURIES	
CHRIST.		CHRIST.		CHRIST.		CHRIST.		CHRIST.		CHRIST.		CHRIST.		CHRIST.		CHRIST.		CHRIST.	
DC		CB		BA		AG		GF		FE		ED		DC		CB		BA	
1	29	57	85	E	D	C	B	A	G	F	E	D	C	B	A	G	F	E	D
2	30	58	86	F	E	D	C	B	A	G	F	E	D	C	B	A	G	F	E
3	31	59	87	G	F	E	D	C	B	A	G	F	E	D	C	B	A	G	F
4	32	60	88	BA	AG	GF	FE	ED	DC	CB	BA	AG	GF	FE	ED	DC	CB	BA	AG
5	33	61	89	C	B	A	G	F	E	D	C	B	A	G	F	E	D	C	B
6	34	62	90	D	C	B	A	G	F	E	D	C	B	A	G	F	E	D	C
7	35	63	91	E	D	C	B	A	G	F	E	D	C	B	A	G	F	E	D
8	36	64	92	GF	FE	ED	DC	CB	BA	AG	GF	FE	ED	DC	CB	BA	AG	GF	FE
9	37	65	93	A	G	F	E	D	C	B	A	G	F	E	D	C	B	A	G
10	38	66	94	B	A	G	F	E	D	C	B	A	G	F	E	D	C	B	A
11	39	67	95	C	B	A	G	F	E	D	C	B	A	G	F	E	D	C	B
12	40	68	96	ED	DC	CB	BA	AG	GF	FE	ED	DC	CB	BA	AG	GF	FE	ED	DC
13	41	69	97	F	E	D	C	B	A	G	F	E	D	C	B	A	G	F	E
14	42	70	98	G	F	E	D	C	B	A	G	F	E	D	C	B	A	G	F
15	43	71	99	A	G	F	E	D	C	B	A	G	F	E	D	C	B	A	G
16	44	72		CB	BA	AG	GF	FE	ED	DC	CB	BA	AG	GF	FE	ED	DC	CB	BA
17	45	73		D	C	B	A	G	F	E	D	C	B	A	G	F	E	D	C
18	46	74		E	D	C	B	A	G	F	E	D	C	B	A	G	F	E	D
19	47	75		F	E	D	C	B	A	G	F	E	D	C	B	A	G	F	E
20	48	76		AG	GF	FE	ED	DC	CB	BA	AG	GF	FE	ED	DC	CB	BA	AG	GF
21	49	77		B	A	G	F	E	D	C	B	A	G	F	E	D	C	B	A
22	50	78		C	B	A	G	F	E	D	C	B	A	G	F	E	D	C	B
23	51	79		D	C	B	A	G	F	E	D	C	B	A	G	F	E	D	C
24	52	80		FE	ED	DC	CB	BA	AG	GF	FE	ED	DC	CB	BA	AG	GF	FE	ED
25	53	81		G	F	E	D	C	B	A	G	F	E	D	C	B	A	G	F
26	54	82		A	G	F	E	D	C	B	A	G	F	E	D	C	B	A	G
27	55	83		B	A	G	F	E	D	C	B	A	G	F	E	D	C	B	A
28	56	84		DC	CB	BA	AG	GF	FE	ED	DC	CB	BA	AG	GF	FE	ED	DC	CB

USE. The Dominical Letter stands under the Column of Centuries, or whole Hundreds, and against the Year (on the Left) above Centuries, before Christ, for the Letter required.

Construction. There being a Revolution of the Julian Dominical Letter in every 700 Julian Years, any Number of Julian Years before Christ, will have the same Julian Dominical Letter as the Complement of those Years to any Number of 700 Years, will have for Years since Christ.

Example. $\left\{ \begin{array}{l} 7 \text{ Hundreds} \dots 3500 \\ \text{Before Christ} \dots -2563 \\ \hline \text{Comp. since Chr.} \dots 937 \end{array} \right.$ Yrs. Dom. Let. $\left\{ \begin{array}{l} G \ 3501 \\ \dots A \ 2563 \\ \hline G \ 937 \end{array} \right.$

USE. The Dominical Letter stands under the Column of Centuries, and against the Year (on the left) above Centuries, since Christ, for the Letter required.

Construction. From the foregoing Argument reverted, any Number of Julian Years since Christ, will have the same Julian Dominical Letter as the Complement of those Years to any Number of 700 Years, will have for Years before Christ.

Example. $\left\{ \begin{array}{l} 7 \text{ Hundreds} \dots 5600 \\ \text{Since Christ} \dots -3666 \\ \hline \text{Comp. bef. Chr.} \dots 1934 \end{array} \right.$ Yrs. Dom. Let. $\left\{ \begin{array}{l} \dots E \ 5601 \\ \dots F \ 3666 \\ \hline \dots F \ 1934 \end{array} \right.$

N. B. By increasing the Columns of Hundreds by Sevens, the Dominical Letter may be found for any Number of Julian Years before or since CHRIST; though never so great.

In Leap-Year the 1st Dominical Letter serves to the 24th of February, exclusive, (24th being twice reckoned, or 24 and 25 the same Month-days, in the Roman Calendar in Bissextile, have the same Letter) the Second Dominical Letter serves for the Rest of the Year. But in our English Calendar this extraordinary Day is added on the 29th of February.

This Table requires 1 to be ded from given year and also the Table BC next Page

vide Pa
155. 157
170.

CHRONOLOGICAL TABLES.

DOMINICAL-LETTERS for Gregorian or New Style, for ever.

YEARS before CHRIST.					YEARS since CHRIST.				
CENTURIES.					CENTURIES				
221									

USE. The Dominical Letter stands under the Column of Centuries, or whole Hundreds, and against the Year (on the Left) above Centuries, before Christ, for the Letter required.

Construction. There being a Revolution of the Gregorian Dom. Letter every 400 Greg. Years, any No. of Years before Christ will have the same Greg. Dom. Letter as the Complement of those Years to any No. of 4 Hundred Years will have for Years since Christ.

Ex. { 4 Hundreds . . . 3600
Before Christ . . . 1791 D C B
Comp. since Christ 1807 D C B

So for the Rest.
N. B. Tho' there was no N. Style before Pope Gregory ordained it, in 1582, by calling the 5 Oct. O. S. the 15 N., at the Years may be thus computed in conformity to our present Style.

N. B. By increasing the Columns of Hundreds by Fours, the Dom. Letter may be found for any No. of Years before or since Christ.

USE. The Dominical Letter stands under the Column of Centuries, and against the Year (on the Left) above Centuries, since Christ, for the Letter required.

Construction. In every 4 Hundred Gregorian Years there are 20871 Weeks just, or 3 Days less than in the same No. of Jul. Years; hence a Revolution of the Dominical Letter every 4 Hundred Gregorian Years is evident.

N. B. The Dominical Letter for any Number of Years since Christ will be the same, as the Dominical Letter for the Complement of those Years to any Number of 4 Hundreds, before Christ.

Ex. { 4 Hundreds . . . 4000
Since Christ . . . 2333 A
Comp. before Christ 1667 A

So for the Rest.

WEEK and MONTH-DAY. Shewing, by the Sunday Letter, of either Style, the Day of the Week to any Day of the Month, in that Year, for ever.

MONTHS.	SUNDAY-LETTERS.						
	A	B	C	D	E	F	G
JANUARY 31	1	2	3	4	5	6	7
OCTOBER 31	8	9	10	11	12	13	14
	15	16	17	18	19	20	21
	22	23	24	25	26	27	28
	29	30	31	1	2	3	4
FEBR. 28. 29	5	6	7	8	9	10	11
MARCH 31	12	13	14	15	16	17	18
NOVEMBER 30	19	20	21	22	23	24	25
	26	27	28	29	30	31	1
	2	3	4	5	6	7	8
APRIL 30	9	10	11	12	13	14	15
JULY 31	16	17	18	19	20	21	22
	23	24	25	26	27	28	29
	30	31	1	2	3	4	5
	6	7	8	9	10	11	12
AUGUST 31	13	14	15	16	17	18	19
	20	21	22	23	24	25	26
	27	28	29	30	31	1	2
	3	4	5	6	7	8	9
SEPTEMBER 30	10	11	12	13	14	15	16
DECEMBER 31	17	18	19	20	21	22	23
	24	25	26	27	28	29	30
	31	1	2	3	4	5	6
	7	8	9	10	11	12	13
MAY 31	14	15	16	17	18	19	20
	21	22	23	24	25	26	27
	28	29	30	31	1	2	3
	4	5	6	7	8	9	10
JUNE 30	11	12	13	14	15	16	17
	18	19	20	21	22	23	24
	25	26	27	28	29	30	

USE. Under the Sunday-Letter, for the Year, and against the Month, is the Sunday-Column, of all the Sundays in that Month, for that YEAR. Next to which, on the Right, is the Monday-Column, next Tuesday-Column, &c. Whence all the Week-Days of that Month are immediately known.

Example 1. To find on what Week the 12th of June happened 455 Years since Christ, N. S.

The Dominical Letter was (by the foregoing Table) C. — Now, under C, against June 6, 13, 20, 27 is the Sunday-Column; next to which (on the Left) is the Saturday-Column, wherein is found 12, which therefore was on a Saturday, required.

Example 2. Required the Week-day of September 19, 1777 Years before Christ, according to N. S. ?

The Dominical Letter is found B. D C

Hence, under E, against September (to the Left) the Sunday-Column is 7, 14, 21, 28, all the Sundays in that Month; and to the Left the 19th Day is in the Friday-Column, which therefore was on a Friday, required.

Hence Apr. 3d, 33 Ch. O. S. was on a Friday.

And May 28th, 585 bef. Chr. on Monday, after a Battle between the Medes and Lydians, Peace was made; not on a Wednesday, according to Ferguson's Author mistakes the 1st of Chr. f. the Year before) Christ: W. Beg. of (or End of chronological Eras.

Greg. Years before or since Christ,

CHRONOLOGICAL

use
p 157
170

CHRONOLOGICAL TABLES.

MONTHS.	Golden Numbers	Advance Days	A TABLE, to find the Place or Sign of the Moon, nearly, by the Golden Number and Days of the Month, till 1899; which will serve for ever, by adding the Days the Moon's go forward in the Month-Days, for a given Period, beyond that Time.																									The Moon's Place or Sign			
			According to the Gregorian or New Style. London Meridian.																												
Jan. Oct.	7	1	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	♈		
		2	1	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	♉	
	18	3	2	1	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	♊	
	10	4	3	2	1	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	♋	
Fe. Mar. No.	2	5	4	3	2	1	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	♌	
		6	5	4	3	2	1	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	♍	
December	13	7	6	5	4	3	2	1	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	♎	
	5	8	7	6	5	4	3	2	1	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	♏	
April		9	8	7	6	5	4	3	2	1	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	♐	
	16	10	9	8	7	6	5	4	3	2	1	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	♑	
	8	11	10	9	8	7	6	5	4	3	2	1	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	♒	
May		12	11	10	9	8	7	6	5	4	3	2	1	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	♓	
	19	13	12	11	10	9	8	7	6	5	4	3	2	1	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	♈
	11	14	13	12	11	10	9	8	7	6	5	4	3	2	1	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	♉
	3	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	27	26	25	24	23	22	21	20	19	18	17	16	15	14	♊
June		16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	27	26	25	24	23	22	21	20	19	18	17	16	15	♋
	14	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	27	26	25	24	23	22	21	20	19	18	17	16	♌
July		6	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	27	26	25	24	23	22	21	20	19	18	♍
		19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	27	26	25	24	23	22	21	20	19	18	♎
	17	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	27	26	25	24	23	22	21	20	19	♏
	9	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	27	26	25	24	23	22	21	20	♐
August		1	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	27	26	25	24	23	22	♑
		23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	27	26	25	24	23	22	♒
	12	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	27	26	25	24	23	♓
	4	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	27	26	25	24	♈
September		26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	27	26	25	♉
	15	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	27	26	♊

EXAMPLE of the USE.

To find the Place or Sign of the Moon, for July 10, 1764, the Golden Number being 17?

Against July stands 1, under or above which, stands the Day of the Month, viz. 10, against which, on the Left, stands 27, Advance Days, and the Golden Number being 17, against which stands 1, to the right Hand, under, or above, which, stands the Advance-Days, viz. 27; against which, to the Right, stands the first Part of Sagittarius, for the Place or Sign the Moon will be in, required.

By the Reverse Method the Day of the Month may be found to any Sign.

To find the Place or Sign of the Moon, for April 29, 1765, the Golden Number being 18?

Against April stands 1, under which stands 2, for the 29th Day, (the 1st being always 28, when the Day of the Month is above 27) against which, to the Left, stands 10, for the Advance-Days. But against the Golden Number 18, stands 1, under, or above which, stand those same 10 Advance-Days, against which, to the Right, stands the latter Part of Leo for Answer. And the like for all other Cases.

CONSTRUCTION.

Any Golden Number is chosen to stand against any Month, (to begin with January is best) so as the Number of Advance-Days, standing to the Left, against the Day of that Month, being found under 1, placed against the Golden Number, shall shew the fore or latter Part of the Sign the Moon is in, to the Right Hand of that Number of Advance-Days; which being determined for any one Day of any Month, the Rest of the Months must be placed forward or backward, in such Order of Succession, or Retrogression, that the 1st Day, next to 27 Days, be reckoned 28; the 2d, the 29th, and so on, to the present Month's End: observing, that as but 27 Days are reckoned for the Moon's going through all the Signs, instead of 27 Days, 7 Hours, and about 43 Minutes, that each third succeeding Month begins a Day later, or at 82 Days End, by counting forward: otherwise the Sign will go on too fast by 7 Hours and 43 Minutes every 27 Days, or in about a Month.

When the Moon's Place is ascertained to the Golden Number for any Year, and the Moon's Mean Place being known to go forward by 13 Days in each Leap Year, or by 10 Days in each Common Year, every Golden Number advanced by 4, or 1, must be placed 13 or 10 Days forward of the former Golden Numbers respectively, which will always point out, (by carrying the Advance-Days so much forward) the Place or Sign of the Moon, for any Year and Month-Day, required.

CHRONOLOGICAL

CHRONOLOGICAL TABLES.

MONTHS.	Golden Numbers	Advance Days.	A TABLE, shewing the Moon's Age by the Golden Number, to any Day of the Month, till 1899. Which will serve for ever, by subtracting the Days the Moon's Age retreats, or the New, Full, or Quarter Moons go forward, in the Month-Days, for a given Period, beyond that Time.																														Moon's Age	Moon's Phases																		
			According to the Gregorian or New Style. London Meridian.																																																	
	7	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1																				
	18	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2																				
		3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3																				
	10	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4																				
		5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5																				
	2	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6																				
Jan. March	13	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7																				
		8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8																				
Fe. Ap. May	5	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9																				
June	16	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10																				
July		11	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11																				
		8	12	11	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12																			
August	19	13	12	11	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12																			
September		14	13	12	11	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12																		
October	11	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12																	
November		16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12																
December	3	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12															
	14	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12														
		19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12													
	6	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12												
	17	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12											
		22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12										
	9	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12									
		24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12								
	1	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12							
	12	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12						
		27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12					
	4	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12				
	15	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12			
		30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	30	29	28	27	26	25	24	23	22	21	20	19	18	17	16	15	14	13	12		

EXAMPLE of the USE.

TO find the Age of the Moon, the 14th of August, 1760, the Golden Number being 13?

Against August stands 1, above, or under which find the Day of the Month, viz. 14; against which to the left Hand stands 26, the No. of Advance Days; Against 13, the Golden Number for the Year, stands 1, under, or above which, stands the No. of Advance-days, viz. 26, against which to the right Hand stands the 2d Day of the Moon in the first Quarter, for Answer. — The Converse finds the Day of the Month to any New, Full, or Quarter Moon.

TO find on what Day the Moon will be at Full in April 1759, the Golden Number being 12?

Against 12 the Golden Number for the Year, stands 1, under, or above, which, against the Full Moon, stands the No. of Advance Days, viz. 20, against which, of the Advance Day Column, in the Month-Day Column for April, to the Right, stands the 12th Day for Answer. And so for other Cases.

CONSTRUCTION.

Any Golden Number is chosen for any Month (to begin with January is best) so as the Number of Advance-Days, standing to the Left, against the Day of that Month, being found under 1, placed against the Golden Number, shall shew the Day of the Moon, to the Right Hand of that Number of Advance-Days; which being determined for any one Day of any Month, the Rest of the Months must be placed forward or backward, in their Order of Succession, or Retrogression; but so as the first Day, next to 30 Days, be reckoned the 31st Day of that Month; observing, that as 30 Days are reckoned for the Time of a Mean Lunation, instead of 29 Days, 12 Hours, 44 Minutes, and about 3 Seconds, the each Second Succeeding Month begins a Day sooner, or at 59 Days End, by counting forward; otherwise the Lunations will proceed too slow by 12 Hours, 44 Minutes, and 3 Seconds in every 30 Days, or about a Month.

When the Moon's Age is ascertained for any Year, to the Golden Number for that Year, the Lunations, or Lunar-Cycles, being known to retreat 14 Days every four Years, or 11 Days every succeeding Year, the Moon's Age comes forward as much; whereby the Golden Numbers for four Years, and one Year forward, must be advanced 14 and 11 Days, respectively, of the former Golden Numbers; which will always point out, (by carrying the Advance-Days forward with them, answerable to any Day of the Month) the Day of the Moon's Age required.

CHRONOLOGICAL TABLES.

A TABLE of the Sun's Place throughout the Year; by which the Moon's Place may be found, and her Rising and Setting nearly, for London, or any Part of the known World. According to New Style.

Days of Mth	Jan.	Feb.	March.	April.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	For finding the Moon's Place.	For finding the Moon's Place.	High Water at London.	No. of the Sign.
													D Age.	D's Sig. and Degree.	D Souths.	
1	☉ 11°	☉ 13°	☉ 11°	☉ 12°	☉ 11°	☉ 11°	☉ 9°	☉ 14°	☉ 9°	☉ 11°	☉ 10°	☉ 9°	1	0 13	12 A 48	☉
2	12	14	12	13	12	12	10	13	10	9	10	10	2	0 26	1 36	0
3	13	15	13	14	13	13	11	14	11	10	11	11	3	1 9	2 24	8
4	14	16	14	15	14	14	12	15	12	11	12	12	4	1 23	3 12	1
5	15	17	15	16	15	15	13	16	13	12	13	13	5	2 6	4 0	
6	16	18	16	17	16	16	14	17	14	13	14	14	6	2 19	4 48	II
7	17	19	17	18	17	17	15	18	15	14	15	15	7	3 2	5 36	2
8	18	20	18	19	18	18	16	19	16	15	16	16	8	3 15	6 24	
9	19	21	19	20	19	19	17	20	17	16	17	17	9	3 28	7 12	III
10	20	22	20	21	20	20	18	21	18	17	18	18	10	4 12	8 0	3
11	21	23	21	22	21	21	19	22	19	18	19	19	11	4 25	8 48	Ω
12	22	24	22	23	22	22	20	23	20	19	20	20	12	5 8	9 36	4
13	23	25	23	24	23	23	21	24	21	20	21	21	13	5 21	10 24	η
14	24	26	24	25	24	24	22	25	22	21	22	22	14	6 4	11 12	5
15	25	27	25	26	25	25	23	26	23	22	23	23	15	6 17	12 0	
16	26	28	26	27	26	26	24	27	24	23	24	24	16	7 0	12 M 48	6
17	27	29	27	28	27	27	25	28	25	24	25	25	17	7 13	1 36	7
18	28	30	28	29	28	28	26	29	26	25	26	26	18	7 26	2 24	
19	29	31	29	30	29	29	27	30	27	26	27	27	19	8 10	3 12	III
20	30		30		30	30	28		28	27	28	28	20	8 23	4 0	7
21	1	3	1	2	1	1	29	1	28	28	29	29	21	9 6	4 48	8
22	2	4	2	3	2	2	30	2	29	29	30	30	22	9 19	5 36	
23	3	5	3	4	3	3	1	3	30	30	1	1	23	10 2	6 24	9
24	4	6	4	5	4	4	2	4	1	31	2	2	24	10 15	7 12	
25	5	7	5	6	5	5	3	5	2	2	3	3	25	10 28	8 0	
26	6	8	6	7	6	6	4	6	3	3	4	4	26	11 12	8 48	10
27	7	9	7	8	7	7	5	7	4	4	5	5	27	11 25	9 36	
28	8	10	8	9	8	8	6	8	5	5	6	6	28	0 8	10 24	11
29	9		9		9	9	7	9	6	6	7	7	29	0 21	11 12	
30	10		10		10	10	8	10	7	7	8	8	30	1 4	12 0	
31	11		11		11	11	9	11	8	8	9	9				

USE. The Sun's Place is expressed in Signs and Degrees under each Month. On the left Hand of the Degrees of the Sun's Place stand the Golden Numbers, respectively answering to the Days of the Month, whereon the New Moons happen; when the Moon's Place is the same with the Sun's. From whence, the Days of her Age must be reckoned forward to the Month Day, whereon the Moon's Place is required: Adding the Signs and Degrees gone forward, by the Moon, according to the Number of Days of her Age, in the annexed Table. When the Sign and Degree of the Moon's Place are found, observe against what Day of the Month the same Sun's Place stands in the Table, and the Moon's half Time above the Horizon, will be the same as that half Day: known by the nearest Month-Day in Table p. 156, of Sun's Setting, which must be deducted from the Time of the Moon's Southing for her Rising, or added thereto for her Setting, at London, or in England.

In other Parts of the World, use the Semi-Duration Arc, according to the Place's Latitude, and Moon's Place or Declⁿ, considered without Latitude. See Tab. of Semi-Duration Arcs farther on.

To find the Moon's Place and Rising, July 16, 1760? The Golden No. being 13, stands against 13th July at New Moon, when the Sun and Moon's Place will be 3^h 21^m
For 3 Days of the Moon add 1 9

Sum 5 0

Or 11^h 00, to which answers Aug. 23, when the Half-Day, (equal the half Continuance of Moon above the Horizon) is 7^h 3^m which taken from 2^h 24^m D's Southing, leaves 7^h 21^m Morning her Rising. And so for other Cases.

HIGH WATER, at the following PLACES, in Hours and Minutes, before or after the Time of High Water at London.

Note, a stands for after, and b for before.

Amsterdam, a 0 ^h 30 ^m	Fowey, a 3 10	Harwich, b 3 30	Ostend, b 2 0	Spithead, b 3 0
Brest, a 1 0	Flushing, b 1 30	Hastings, b 4 0	Orfordness, b 4 0	Shoreham, b 4 0
Bidlington Pier, a 2 0	Gouries Gut, b 0 5	St. Helens, b 5 30	Plymouth, a 3 10	Tinmouth Haven, a 0 30
Bridgewater, a 4 40	Gravesend, b 1 20	Havre-de-Grace, b 5 30	Portland, a 5 50	Tinmouth, a 3 40
Buoy of Nore, b 1 30	Gunfleet, b 4 0	Loo, a 3 10	Portsmouth, b 2 0	Torbay, a 3 40
Calais, b 3 0	Hartlepool, a 0 30	Lime, a 3 50	Rochester, b 1 20	Topsham, a 3 50
Dartmouth, a 3 30	Humber, a 2 0	Leigh, b 0 5	Rammekins, b 1 20	Texel, a 4 40
Dover, b 3 0	Hull, a 3 30	Mountsbay, a 1 15	Redsand, b 2 0	Weymouth, a 4 25
Dieppe, b 4 0	Harborough, a 3 30	Maze, b 0 5	Scilly, a 1 45	Yarmouth Pier, b 4 40
Exmouth, a 3 50	Harflew, a 5 50	Needles, b 4 40	Shoe Bacon, b 2 0	

EXAMPLE. To find the Time of High-Water at Rochester, on the 10th Day of the Moon's Age.

At London, on that Day, it is high Water — 9^h 53^m
subtract 1 20

High Water at Rochester — 8 33

And the like for other Places, by adding or subtracting to or from High Water at London.

CHRONOLOGICAL TABLES.

A TABLE, to find the Time of the *Paschal* or *Easter* Full Moon, and consequently *Easter*, for ever, by the Golden Numbers, and Days the *Lunar* Cycle of 19 Years goes forward in any given Period, *New Style*.

Paschal Full Days	Dom. Letter.	GOLDEN NUMBERS.																		
		i	ii	iii	iv	v	vi	vii	viii	ix	x	xi	xii	xiii	xiv	xv	xvi	xvii	xviii	xix
Mar. 21	C	8	19	0	11	22	3	14	25	6	17	28	9	20	1	12	23	4	15	26
22	D	9	20	1	12	23	4	15	26	7	18	29	10	21	2	13	24	5	16	27
23	E	10	21	2	13	24	5	16	27	8	19	0	11	22	3	14	25	6	17	28
24	F	11	22	3	14	25	6	17	28	9	20	1	12	23	4	15	26	7	18	29
25	G	12	23	4	15	26	7	18	29	10	21	2	13	24	5	16	27	8	19	0
26	A	13	24	5	16	27	8	19	0	11	22	3	14	25	6	17	28	9	20	1
27	B	14	25	6	17	28	9	20	1	12	23	4	15	26	7	18	29	10	21	2
28	C	15	26	7	18	29	10	21	2	13	24	5	16	27	8	19	0	11	22	3
29	D	16	27	8	19	0	11	22	3	14	25	6	17	28	9	20	1	12	23	4
30	E	17	28	9	20	1	12	23	4	15	26	7	18	29	10	21	2	13	24	5
31	F	18	29	10	21	2	13	24	5	16	27	8	19	0	11	22	3	14	25	6
Apr. 1	G	19	0	11	22	3	14	25	6	17	28	9	20	1	12	23	4	15	26	7
2	A	20	1	12	23	4	15	26	7	18	29	10	21	2	13	24	5	16	27	8
3	B	21	2	13	24	5	16	27	8	19	0	11	22	3	14	25	6	17	28	9
4	C	22	3	14	25	6	17	28	9	20	1	12	23	4	15	26	7	18	29	10
5	D	23	4	15	26	7	18	29	10	21	2	13	24	5	16	27	8	19	0	11
6	E	24	5	16	27	8	19	0	11	22	3	14	25	6	17	28	9	20	1	12
7	F	25	6	17	28	9	20	1	12	23	4	15	26	7	18	29	10	21	2	13
8	G	26	7	18	29	10	21	2	13	24	5	16	27	8	19	0	11	22	3	14
9	A	27	8	19	0	11	22	3	14	25	6	17	28	9	20	1	12	23	4	15
10	B	28	9	20	1	12	23	4	15	26	7	18	29	10	21	2	13	24	5	16
11	C	29	10	21	2	13	24	5	16	27	8	19	0	11	22	3	14	25	6	17
12	D	0	11	22	3	14	25	6	17	28	9	20	1	12	23	4	15	26	7	18
13	E	1	12	23	4	15	26	7	18	29	10	21	2	13	24	5	16	27	8	19
14	F	2	13	24	5	16	27	8	19	0	11	22	3	14	25	6	17	28	9	20
15	G	3	14	25	6	17	28	9	20	1	12	23	4	15	26	7	18	29	10	21
16	A	4	15	26	7	18	29	10	21	2	13	24	5	16	27	8	19	0	11	22
17	B	5	16	27	8	19	0	11	22	3	14	25	6	17	28	9	20	1	12	23
17	B	5	16	27	8	19	0	11	22	3	14	25	7	18	29	10	21	2	13	24
18	C	6	17	28	9	20	1	12	23	4	15	26	7	18	29	10	21	2	13	24
18	C	7	18	29	10	21	2	13	24	5	16	27	8	19	0	11	22	3	14	25

EXAMPLE. To find the Full Moon before Easter-Sunday, in the Year 1765?

Against the Period 1700, in TABLE below stands 1, and the Golden Number for 1765 being 18, against 1, the Days gone forward in that Period, under the Golden Number xviii, in this Table stands April 6, in the first Column for Answer. And the Dominical Letter for 1765 being F, reckoning on from April 6, to F, Apr. 7 is Easter-Sunday, and so for other Cases.

NOTE. The Days the Cycle is gone forward, standing under the given Golden Number, and against a certain Month-day, serve for a given Period, according to TABLE below.

A TABLE of the Number of Days, the *New*, *Full*, or *Quarter* Moons in the *Gregorian* Account, go forward in the Months, or the *Lunar* Cycle, (of 19 Years, or 235 Synodical Months) advances of the Cycle in the Beginning of 1600.

Bisextile Years.	Gregor. Years of Chr.	Lu. Cyc. go forw. Days.	Gregor. Years of Chr.	Lu. Cyc. go forw. Days.	Gregor. Years of Chr.	Lu. Cyc. go forw. Days.	Gregor. Years of Chr.	Lu. Cyc. go forw. Days.	Gregor. Years of Chr.	Lu. Cyc. go forw. Days.	Gregor. Years of Chr.	Lu. Cyc. go forw. Days.
B	1600	0										
	1700	1	2900	6	4100	11	5300	16	6500	21	7700	26
	1800	1	3000	6	4200	12	5400	17	6600	22	7800	27
	1900	2	3100	7	4300	12	5500	17	6700	23	7900	28
B	2000	2	3200	7	4400	12	5600	17	6800	22	8000	27
	2100	2	3300	7	4500	13	5700	18	6900	23	8100	28
	2200	3	3400	8	4600	13	5800	18	7000	24	8200	29
	2300	4	3500	9	4700	14	5900	19	7100	24	8300	29
B	2400	3	3600	8	4800	14	6000	19	7200	24	8400	29
	2500	4	3700	9	4900	14	6100	19	7300	25	8500	0
	2600	5	3800	10	5000	15	6200	20	7400	25	&c.	
	2700	5	3900	10	5100	16	6300	21	7500	26		
B	2800	5	4000	10	5200	15	6400	20	7600	26		

See the Reason, Page 30, why the Lunar Cycle goes forward in the Months, according to a certain Number of Days, as here inserted.

CHRONOLOGICAL TABLES.

A TABLE, shewing the MOVEABLE FEASTS, by the DOMINICAL LETTER and GOLDEN NUMBER, from 1700 to 1899, according to the Gregorian or New Style.

D. L.	GOLDEN NUMBERS.	From Xmas to Sh. Sund.	Shrov. Sund.	Easter Sunday	Rog. Sund.	Afc. Day.	Whit. Sund.	Trin. Sund.	Adv. Sund.
A	3.11.14	6 ^w 0 ^d	Fe. 5	Mar. 26	Ap. 30	May 4	May 14	May 21	Dec. 3
	5.8.13.16.19	7	12	Ap. 2	May 7	11	21	28	3
	2.7.10.18.	8	19	9	14	18	28	Jun. 4	3
	1.4.9.12.15	9	26	16	21	25	Jun. 4	11	3
	6.17	10	Mar. 5	23	28	Jun. 1	11	18	3
B	3.11.14.19	6 1	Feb. 6	Mar. 27	May 1	May 5	May 15	May 22	No. 27
	2.5.8.13.16	7 1	13	Apr. 3	8	12	22	29	27
	7 10.15.18	8 1	20	10	15	19	29	Jun. 5	27
	1.4.9.12	9 1	27	17	22	26	Jun. 5	12	27
	6.17	10 1	Mar. 6	24	29	Jun. 2	12	19	27
C	3.8.11.14.19	6 2	Feb. 7	Mar. 28	May 2	May 6	May 16	May 23	28
	2.5.13.16	7 2	14	Apr. 4	9	13	23	30	28
	4.7.10.15.18	8 2	21	11	16	20	30	Jun. 6	28
	1.9.12.17	9 2	28	18	23	27	Jun. 6	13	28
	6.	10 2	Mar. 7	25	30	Jun. 3	13	20	28
D	14.	5 3	Feb. 1	Mar. 22	Ap. 26	Ap. 30	May 10	May 17	29
	3.8.11.19	6 3	8	29	May 3	May 7	17	24	29
	2.5.10.13.16	7 3	15	Apr. 5	10	14	24	31	29
	4.7.15.18	8 3	22	12	17	21	31	Jun. 7	29
	1.6.9.12.17	9 3	Mar. 1	19	24	28	Jun. 7	14	29
E	3.14.	5 4	Feb. 2	Mar. 23	Ap. 27	1	May 11	May 18	30
	8.11.16.19	6 4	9	30	May 4	8	18	25	30
	2.5.10.13	7 4	16	Apr. 6	11	15	25	Jun. 1	30
	4.7.12.15.18	8 4	23	13	18	22	Jun. 1	8	30
	1.6.9.17	9 4	Mar. 2	20	25	29	8	15	30
F	3.14.	5 5	Feb. 3	Mar. 24	Ap. 28	2	May 12	May 19	Dec. 1
	5.8.11.16.19	6 5	10	31	May 5	9	19	26	1
	2.10.13.18	7 5	17	Apr. 7	12	16	26	Jun. 2	1
	1.4.7.12.15	8 5	24	14	19	23	Jun. 2	9	1
	6.9.17	9 5	Mar. 3	21	26	30	9	16	1
G	3.11.14	5 6	Feb. 4	Mar. 25	Ap. 29	3	May 13	May 20	2
	5.8.16.19	6 6	11	Apr. 1	May 6	10	20	27	2
	2.7.10.13.18	7 6	18	8	13	17	27	Jun. 3	2
	1.4.12.15	8 6	25	15	20	24	Jun. 3	10	2
	6.9.17	9 6	Mar. 4	22	27	31	10	17	2

EXAMPLE. What Time do the Moveable Feasts fall in 1762? The Dominical Letter being C, and Golden Number 15; Shrove Sunday is February 21; Easter, April 11; Rogation, May 16; Ascension-Day, May 20; Whitsunday, May 30; Trinity Sunday, June 6; Advent Sunday, November 28; so for other Years.

Note. The above Table will serve from 1900 to 2199, except when the Easter Moon falls on a Sunday; then Easter falls a Week after the Time, by the Tables.

A TABLE, shewing the Moveable Terms, from 1700 to 1899.

Easter Term begins.	Ends.	Trinity Term begins.	Ends.
Ap. 12	May 8	May 26	Jun. 14
19	15	Jun. 2	21
26	22	9	28
May 3	29	16	July 5
10	Jun. 5	23	12
Ap. 13	May 9	May 27	Jun. 15
20	16	Jun. 3	22
27	23	10	29
May 4	30	17	July 6
11	Jun. 6	24	13
Ap. 14	May 10	May 28	Jun. 16
21	17	Jun. 4	23
28	24	11	30
May 5	31	18	July 7
12	Jun. 7	25	14
Ap. 8	May 4	May 22	Jun. 10
15	11	29	17
22	18	Jun. 5	24
29	25	12	July 1
May 6	Jun. 1	19	8
Ap. 9	May 5	May 23	Jun. 11
16	12	30	18
23	19	Jun. 6	25
30	26	13	July 2
May 7	Jun. 2	20	9
Ap. 10	May 6	May 24	Jun. 12
17	13	31	19
24	20	Jun. 7	26
May 1	27	14	July 3
8	Jun. 3	21	10
Ap. 11	May 7	May 25	Jun. 13
18	14	Jun. 1	20
25	21	8	27
May 2	28	15	July 4
9	Jun. 4	22	11

Easter Term begins April 28, ends May 24. Trinity Term begins June 11, ends June 30, for the same Year 1762.

Note. Easter Term begins 17 Days, and ends 43 Days after Easter. Trinity Term begins 5 Days, and ends 24 Days after Trinity. Therefore Easter Term continues 27, and Trinity 20 Days, inclusive.

Emendation. P. 148. For F, (against 4 Years current since Chr. O. S. under 700) make FE with the Pen. Same Pa. For FD (against 12 Yrs. bef. Chr. O. S. under 700) make ED.

In the Dom. Letter and Golden No. Tables, O. and N. Style, the Yrs. since Christ, were first considered as completed, or 1, 2, 3, &c. Years of Chr. reckoned after completed, (instead of current) by which the 0 Year was considered Bissextile, fr. the Beginning to the End of the 1 Year since the Birth of Christ; the 1st Year from thence to the End of 2d, &c. whereby 1st Year before Christ was considered 1st after Bissextile, which is Bissextile, considering the Years since Chr. current, as they are. Therefore, to find the D. let. or Golden No. for any Yr. before Chr. 1 must be deducted from the true Year, before Christ, and then find them, in the Tables from the Date remaining; because 1 Year before Chr. should stand in the Place of 0, Year; 2 in the Place of 1, Year; 3 in Place of 2 Years, &c. in the Tables for Years before Christ.—While the similar Tables, for Years since Christ, are used as they stand.

CHRONOLOGICAL TABLES.

A TABLE, shewing, at Sight, the MOVEABLE FEASTS, and TERMS, for ever, by the DOMINICAL LETTER, and GOLDEN NUMBER, according to Old Style.
Of Use in HISTORY and CHRONOLOGY.

D. L.	GOLDEN NUMBERS.	From Xmas to Sh. Sund.	Shrove Sund.	Easter Sunday	Rog. Sunday	Ascen. Day.	Whit-Sunday	Trin. Sunday	Advent Sunday	Easter Term begins.	Ends.	Trinity Term begins.	Ends.
A	2.5.13.16	6 ^w 0 ^d	Feo. 5	Mar. 26	Apr. 30	May 4	May 14	May 21	Dec. 3	Apr. 12	May 8	May 26	Jun. 14
	7.10.15.18	7	12	Apr. 2	May 7	11	21	28	3	19	15	Jun. 2	21
	1.4.9.12	8	19	9	14	18	28	Jun. 4	3	26	22	9	28
	3.6.11.14.17	9	26	16	21	25	Jun. 4	11	3	May 3	29	16	July 5
	8.19	10	Mar. 5	23	28	Jun. 1	11	18	3	10	Jun. 5	23	12
B	2.5.13.16	6 1	Feb. 6	Mar. 27	May 1	May 5	May 15	May 22	No. 27	Apr. 13	May 9	May 27	Jun. 15
	4.7.10.15.18	7 1	13	Apr. 3	8	12	22	29	27	20	16	Jun. 3	22
	1.9.12.17	8 1	20	10	15	19	29	Jun. 5	27	27	23	10	29
	3.6.11.14	9 1	27	17	22	26	Jun. 5	12	27	May 4	30	17	July 6
	8.19	10 1	Mar. 6	24	29	Jun. 2	12	19	27	11	Jun. 6	24	13
C	2.5.10.13.16	6 2	Feb. 7	Mar. 28	May 2	May 6	May 16	May 23	28	Apr. 14	May 10	May 28	Jun. 16
	4.7.15.18	7 2	14	Apr. 4	9	13	23	30	28	21	17	Jun. 4	23
	1.6.9.12.17	8 2	21	11	16	20	30	Jun. 6	28	28	24	11	30
	3.11.14.19	9 2	28	18	23	27	Jun. 6	13	28	May 5	31	18	July 7
	8	10 2	Mar. 7	25	30	Jun. 3	13	20	28	12	Jun. 7	25	14
D	16	5 3	Feb. 1	Mar. 22	Apr. 26	Apr. 30	May 10	May 17	29	Apr. 8	May 4	May 22	Jun. 10
	2.5.10.13	6 3	8	29	May 3	May 7	17	24	29	15	11	29	17
	4.7.12.15.18	7 3	15	Apr. 5	10	14	24	31	29	22	18	Jun. 5	24
	1.6.9.17	8 3	22	12	17	21	31	Jun. 7	29	29	25	12	July 1
	3.8.11.14.19	9 3	Mar. 1	19	24	28	Jun. 7	14	29	May 6	Jun. 1	19	8
E	5.16	5 4	Feb. 2	Mar. 23	Apr. 27	1	May 11	May 18	30	Apr. 9	May 5	May 23	Jun. 11
	2.10.13.18	6 4	9	30	May 4	8	18	25	30	16	12	30	18
	1.4.7.12.15	7 4	16	Apr. 6	11	15	25	Jun. 1	30	23	19	Jun. 6	25
	6.9.14.17	8 4	23	13	18	22	Jun. 1	8	30	30	26	13	July 2
	3.8.11.19	9 4	Mar. 2	20	25	29	8	15	30	May 7	Jun. 2	20	9
F	5.16	5 5	Feb. 3	Mar. 24	Apr. 28	2	May 12	May 19	Dec. 1	Apr. 10	May 6	May 24	Jun. 12
	2.7.10.13.18	6 5	10	31	May 5	9	19	26	1	17	13	31	19
	1.4.12.15	7 5	17	Apr. 7	12	16	26	Jun. 2	1	24	20	Jun. 7	26
	3.6.9.14.17	8 5	24	14	19	23	Jun. 2	9	1	May 1	27	14	July 3
	8.11.19	9 5	Mar. 3	21	26	30	9	16	1	8	Jun. 3	21	10
G	5.13.16	5 6	Feb. 4	Mar. 25	Apr. 29	3	May 13	May 20	2	Apr. 11	May 7	May 25	Jun. 13
	2.7.10.18	6 6	11	Apr. 1	May 6	10	20	27	2	18	14	Jun. 1	20
	1.4.9.12.15	7 6	18	8	13	17	27	Jun. 3	2	25	21	8	27
	3.6.14.17	8 6	25	15	20	24	Jun. 3	10	2	May 2	28	15	July 4
	8.11.19	9 6	Mar. 4	22	27	31	10	17	2	9	Jun. 4	22	11

A General TABLE of the SUN'S Rising and Setting, for England and Ireland, NEW STYLE. To be used with Table, Page 153.

Days	January.		February.		March.		April.		May.		June.		July.		August.		September.		October.		November.		December.																											
	rises.	sets.	rises.	sets.	rises.	sets.	rises.	sets.	rises.	sets.	rises.	sets.	rises.	sets.	rises.	sets.	rises.	sets.	rises.	sets.	rises.	sets.	rises.	sets.																										
3	8	33	57	7	23	4	37	6	29	5	32	5	27	6	34	4	32	7	29	3	50	8	10	3	46	8	14	4	22	7	37	5	17	6	42	6	16	5	43	7	15	4	44	7	59	1				
6	8	13	57	7	18	4	43	6	23	5	38	5	21	6	40	4	27	7	34	3	47	8	13	3	48	8	12	4	27	7	32	5	23	6	36	6	22	5	37	7	20	4	39	8	2	3	58			
9	7	58	4	2	13	4	48	6	17	5	44	5	16	6	45	4	22	7	39	3	46	8	14	3	51	8	9	4	31	7	28	5	29	6	30	6	28	5	31	7	25	4	34	8	4	3	59			
12	7	55	4	5	8	4	53	6	11	5	50	5	10	6	51	4	17	7	44	3	45	8	15	3	54	8	6	4	36	7	23	5	35	6	24	6	53	5	26	7	31	4	28	8	6	3	54			
15	7	51	4	9	7	2	4	59	6	5	56	5	4	6	57	4	13	7	48	3	44	8	16	3	57	8	3	4	42	7	17	5	41	6	18	6	39	5	20	7	36	4	24	8	7	3	53			
18	7	48	4	12	6	5	5	59	6	2	53	5	2	6	58	7	3	4	7	53	3	43	8	17	4	0	8	0	4	47	7	12	5	47	6	12	6	45	5	14	7	41	4	19	8	8	3	52		
21	7	44	4	16	6	5	10	5	53	6	8	4	53	7	8	4	4	7	57	3	43	8	17	4	4	7	5	6	4	53	7	6	5	52	6	7	6	51	5	8	7	45	4	15	8	8	3	52		
24	7	39	4	21	6	4	5	16	5	47	6	14	4	8	7	13	4	0	8	1	43	8	17	4	8	7	5	2	4	58	7	1	5	58	6	16	5	55	5	2	7	49	4	11	8	8	3	52		
27	7	34	4	26	6	3	5	21	5	41	6	20	4	2	7	19	3	5	7	8	4	44	8	16	4	12	7	4	5	48	5	4	55	6	4	5	55	7	2	4	57	7	52	4	8	7	3	53		
30	7	30	4	30					5	35	6	26	4	3	7	24	3	5	4	8	7	3	45	8	15	4	17	7	4	5	43	5	9	6	50	6	10	5	49	7	7	4	52	7	56	4	8	5	3	55

N. B. The Time of the Sun's Setting, in the above Table, being doubled, will shew the Length of that Day, in the Year.

REMARKS

REMARKS on the ÆRAS, or Points of TIME.

THE most famous Æra is that of *Jesus Christ*, beginning at the *Kalends of January*, immediately following the *Birth*. Though this Æra is generally received by *Christians*, yet the *English* and *Irish* used an Æra later by a whole Year in all public and ecclesiastical Writings, before the *Alteration of the Style*. They did not begin the Year with the 1st of *January* following the *Nativity*, but with the 1st of the *Conception*, or *Incarnation*, observed on the 25th of *March*; and therefore the *English* reckoned from the *Feast of Lady-day*, 1748, that there were completed 1747 Years. But, from the *Birth* of our Lord to the *Feast* of the *Nativity*, in the Year 1747, they numbered only 1746 Years elapsed; whereas all the Rest of the *Christian World* reckoned 1747.

Dionysius Minor, Author of the *Christian Æra*, made the *Conception* of *Christ* to be on the 8th of the *Kalends of April*, on the first Year of this Æra, and born the Winter following, at the End of the 46th Year of the Reformation of the *Calendar* by *Julius Cæsar*.

This Way of *Computation* was universally accepted, at first, but universally receded from by Degrees; and did not take Place in *England*, and some other Places, before the End of the old *Style*.

The common Opinion is, that *Christ* was born the Winter preceding the Time in which *Dionysius* reckoned the *Conception*; yet the *English* and the Rest of the *Christian World*, agreed about the Time since his *Birth* for the greater Part of the Year; but for three Months, from the *Kalends of January* to the 8th of the *Kalends of April*, they wrote one Year less.

CONTROVERSIES about ÆRAS.

There are many *Controversies* about the Æra of *CREATION*, among the *Chronologists*, (and always will) some affirming the World to be created 3950 Years before *Christ*; others affirm 3984. The *Greek-Church*, and *Eastern Emperors* Æra of *Creation* was 5509 Years before *Christ*. But the *Chinese Date* of the World is much older; and such *chronological Disputes* will never be settled.

The antient Æra of the *Olympiads* begun 775 Years before *Christ*, on the *Kalends of July*, according to some; but 776 according to the *Greeks*, who yet make the Year of the *Julian Period* 3938.

The Æra of the Building of *Rome* is either the *Varronian*, 752 Years, or the *Capitolian*, beginning *April 21*, 751 Years before *Christ*.

The Æra of *Nabonassar*, famous among *Astronomers*, began the 26th of *February* of the *Julian Year*, reckoned backward to 746 before *Christ*: That Day being the 1st of the *Egyptian Year*. *Ptolomy*, and, after him, *Copernicus*, computed the Places of the *cælestial Bodies* from it.

The Æra of *Alexander* begins 324 Years before *Christ*, *November 12*, being on a *Sunday*, and according to *Chronology*, which was also the 1st Day of the *Egyptian Year*. Between the Æra of *Nabonassar* and *Alexander*, there are 424 *Egyptian Years*. The *Abissines* reckoned by the Æra of *Martyrs*, called *Dioclesian*, beginning *August 29*, in the Year of *Christ* 284.

The *Turks* and *Arabians* Æra, call'd *Hegira*, takes its Beginning from *Mahomet's Flight*, *July 16*, Anno *Christi* 622.

The *Persian Æra* is called *Jesdegerd* begins *June 16*, Anno *Christi* 632.

The *Julian Period*, containing all other Æras, has been described, Page 146; which Æras, when reduced to it, differ, according to different Authorities.

The Antients distinguished their Time by other Revolutions; such as the *Jubilee* of 49 or 50 Years; an Age of 100; and an *Olympiad* of 4 Years, among the *Greeks*, beginning at the *New Moon*, next after the *Summer Solstice*, 775 Years before *Christ*.

The *Persians* observe the *Egyptian Account* of 365 Days in a Year to this Day. In which Reckoning the *Equinoxes* and *Seasons* move forward through the Year in the Space of 1506 *Egyptian Years*, 4 Hours, 27 Minutes, 8 Seconds, according to the *Newtonian Mean Solar Year*, when the same Season begins again. This Year is extremely convenient for *Astronomical Computation*; because of no Motion to be added for *intercalary Days* in the Reckoning; there being none of those Days taking Place.

The Difference of a Year, or more, between some of these Æras, and those we have given, at Page 146, (which we find disputable) we leave to be settled by *Chronologists*, making this Subject their chief Study; as there is no Likelihood for Men of Science to agree exactly in these Matters, not capable of Demonstration.

The Week-day correspondent to the Month-day, for the reputed Year, many Years before *Christ*, is another Obstacle lying in the Road to Truth, in *chronological Computation*. For, before the Time of *Julius Cæsar*, the Form of the Year, or Year of Confusion, was never exactly and regularly settled, by which it will be hard to know the Week-day, on which every successive Year, since *Creation*, began; as likewise the Week-day, correspondent to the Month-day of any Year, since *Creation*: The same depending on the settled Form, or known Changes, of the Year, from some particular Date. The Antient Week and Month-Days, before *Christ*, the most certainly known, are reduced to the *Julian Form* of Reckoning, whereby they become intelligible.

When *Julius Cæsar* reformed the *Calendar* in the 45th Year before *Christ*, he ordered that the 1 Day of the *Julian Year*, should begin on *Jan. 1*, immediately after the old *Roman Year* of *Confusion*, which happened, according to *Chronology*, on a *Friday*. Now (by *Tab. p. 148.*) the *Dom. Let.* for 45 Years before *Christ*, (deducting 1 from 45 to find it) was D, old *Style*. Whence (by *Tab. p. 149.*) the 1st of *Jan.* in that Year was on a *Friday*, agreeing with *Chronology*. Likewise, we find, on *Retrospection*, that 28th *May*, in the 585 Year before *Christ*, being *Leap-Year*, was on a *Wednesday* and not on a *Monday*, when the Battle between the *Medes* and *Lydians* was decided. Our Mistake against *Ferguson's Astronomy*, at p. 149, we therefore freely give up to Truth and Justice. At the same Time we observe, if his Æra, 24 Years bef. *Chr.* be true, when the *Calendar* was corrected, that *Jan. 1st* happened, by our and his Tables, on a *Sunday*, and not on a *Friday*, according to *Chronology*. That the 24th, and not 23d *Feb.* (see p. 10, *Ferguson's Appendix*) is on the 6th of the *Kalends of March* — and 24 and 25th of *Feb.* (not 23 and 24) are reckoned the same Day, in *Bisextile*.

Mistakes in the Æras or Dates of *Chronology*, may in general be rectified by making them agree with the Week-Days to which they are fixed; the Week-Day being a material Circumstance in all *Chronologies* and *Histories* of EVENTS.

REMARKS on the NATURE and MEASURE of TIME.

TIME, by our Conception, proceeds with a constant equable Flux, or uniform Succession. The Measure of which must therefore be referred to Motion; and that of the apparent uniform Motion of the fixed Stars is best. But the variable Motion of the Sun and Moon have, by the Consent of all Ages, (as more conspicuous and observable than that of the Planets and Stars) been made Choice of for this Purpose; which Bodies seem to have been intended by the divine Architect, besides their other Uses, for perpetual *Chronometers*, to distinguish and mark out the Seasons, Days, and Years, as they are.

Astronomers observe that the Days, and therefore the Hours, and Minutes, are unequal, and that by the Earth slackening her Pace, and sometimes quickening it again in her Orbit; besides her uniform Rotation from West to East, the same Way, on her Axis, in a Day's Time. This they infer from the Sun's different apparent Motion, through the same Length of Year, nearly.

Therefore, to measure Time, or refer it to one Standard of uniform Motion, (as they imagine) they have fixed the Length of the solar Year (or that Time wherein the Sun and Earth finish their several Positions with each other, and begin again) to the constant Period of 365^d 5^h 48^m 57^s 39th, &c. according to *Sir Isaac Newton*; but different by others.

REMARKS on the NATURE and MEASURE of TIME.

But one Way or the other, this Length of the Year is no more certain or permanent than the mean Length of a Day. Since, in this Computation, they divide the Distance of Time between any two distant Equinoxes, or Solstices, observed, by the Number of Revolutions between those Observations, whereby they but determine the mean Length of the variable solar Period, which they afterwards suppose permanent or equal.

To do which, as Sir Isaac Newton went no farther back than 20 Years, betwixt the first and last Observation, he could not determine the Length of the mean Year to so great an Exactness, as if he had gone as far back into Antiquity, as Ptolemy's Time, or before, for his first Observation. For the mean solar Year this Way determined, by the great Distance of Time divided by the great Number of Revolutions, would have come so near the Truth (except an Acceleration of the mean Motion of the Sun or Earth be supposed) that if there had been a considerable Error committed in the Time of the 1st Observation, or elsewhere, yet, it would have become insignificant for one Year, by its Division into so great a Number of Parts or Revolutions.

Hence, $365^d.24234027 : 365^d :: 12^s : 11^s.99203814$, &c. = $11^s.29^m.45^s.40''$, &c. Sun's Motion in 365 Days, according to Sir Isaac Newton. From whence the mean diurnal Motion in the Ecliptic = $59' 8''$, &c. (as before observed in the Tables) whereas the apparent or real Motion amounts to $61'$ sometimes, and at other Times scarcely to $57'$: The mean and true Time being proportional to the mean and true Motions.

But, considering the annual Precession of the equinoctial Points at the mean Rate of $50''$, according to some, but different by others, the Variation of the Ecliptic's Obliquity, the different Distances of the observed Equinoxes, or Solstices, by which the Solar Year is differently determined, ASTRONOMICAL TABLES, formed according to these different Principles (as has been the Custom) must vary more or less from the Truth in Computation; verifying Dr. Halley's Observation to his Friend that "all Astronomical Tables are made to be mended." And the present superior Accuracy we pretend to, in our Tables, is by a Comparison with, and Improvement of, the latest and best Authorities.

Of REST and MOTION of BODIES, and of ANTIENT and MODERN HYPOTHESES.

RELATIVE Rest we observe daily in the Masses of inanimated Matter about us; but ABSOLUTE REST in any Thing we observe, is as hard to distinguish, as an absolute Vacuum in infinite Space.

Dr. Bradley, by his more diligent and correct Observations of the Heavens, was the first who discovered the Aberration of the fixed Stars, (caused by the Motion of Light and of the Earth in its Orbit) who has also made a Discovery of another apparent Motion among the fixed Stars, in his Letter to the Earl of Macclesfield. And it has been the Opinion of the antient, as well as it is now of the modern, Astronomers, that different Worlds, or distant Systems, have a relative Motion among one another, at the same Time they are relatively moving forward (perhaps in Orbits round a central World) in infinite Space. This Notion of a Plurality of Worlds has prevailed ever since the Time of the Pythagoreans, who maintained (as we now do) the Probability of the several Planets being inhabited by a particular Kind of Beings, suited to those Regions. Who likewise maintained that the Planets were retained in their Orbits by the Power of Gravity. That each Planet moved round its Orbit like a Stone whirled round at the End of a String, by Centripetal and Centrifugal Forces. And Lucretius, taught by Democritus, supposed Worlds without Number, in the infinite Space, counterpoising each other, by some general Law of Gravitation. That if Bodies were bounded, those, within the Limit, would, by the Attraction of one another, in Time, unite in the middle Space.

And herein it is observable, that Limits of Space appear as impossible to Conception, as the Limits of Time, or of Existence. For, there is no Thought can imagine when Time and Existence were not; nor can Imagination represent, to a rational Mind, a Possibility, when Time and Existence shall be no more; except in the Sense of all Things (after a Dissolution of our Earth and System) being swallowed up in an eternal Duration and Immensity of Existence, in their several new Modes. Therefore, an ETERNAL and INFINITE EXISTENCE ARE OF UNAVOIDABLE NECESSITY. But the infinite Succession of Time, and the Points of it, from whence all new Forms of Things, Beings, and their astonishing Changes, take their Place and Effect, who but an INFINITE GREAT BEING, (eternally existing before all Worlds) can comprehend!

The Ancient Astronomers first imagined, that the Celestial Bodies had a circular and equal Motion round the Body which they supposed at Rest, but not finding this Circular Hypothesis to agree with their Observation (the Sun spending near eight Days more in the Northern, than in the Southern Semi-Circle of the Ecliptic) to reconcile Appearance better to Observation, fixed the Sun, or Earth, at Rest, 3450 such Parts from the Center of the supposed Circular Orbit, as the Radius of it is 100000, which Distance from the Center is called Eccentricity, whence they readily calculated the Sun's Place, at any Time. This Theory, though agreeing pretty well for the Place of the Sun, the Motion of other Planets could not be accounted for by it: Therefore it was changed by Kepler, for the Elliptical Theory; which supposes the Sun in the lower Focus common to the Elliptical Orbits of all the Planets, with a rotary Motion only about his own Axis, they circulating round him, and describing constantly equal Areas in equal Times (or Areas proportional to the Times) by Rays drawn from the Sun, to each respective Planet. By the same Hypotheses are represented the Motions of all the Secondary Planets round their Primary ones, placed in the lower Focus of each Secondary's Orbit: And this Hypothesis answers to the Appearances in the Heavens beyond all others hitherto invented; though Kepler himself at first doubted whether some Orbits vary from true Ellipses: Thinking it is not improbable that some might have a Form like the Section of an Egg, by the Force of Gravity dilating one Part of the Orbit more than another. But this Theory follows the Track of the first Circular Theory, which supposed the Sun at Rest in the Center of the planetary Orbits, and Planets moving circularly round Him, whereby they must describe equal Areas in equal Times; as they are now found to do in the Elliptical Theory, by Observation.

The Sun's least apparent Diameter in Apogee has been observed $31' 29''$, and his greatest in Perigee $32' 33''$, by which it should result, that the Sun's greatest and least Distances are as 1953 to 1889, as 101661 to 98339 (vid. Keil's Astron. p. 277) and the Eccentricity only 1661 of such Parts, as the Radius of the Circular Orbit, Sem-Transf. (or mean Dist.) of the Elliptical Orbit is 100000: But the antient Eccentricity 3450 is above double 1661, from whence the Circular Theory was long ago concluded to be false. For, admitting but 1725, one half of the antient Eccentricity, it would better agree with the Sun's apparent Diameters observed: But then would not, so well as the whole, account for the Appearances of the Sun's unequal Motion round the Year. And (as the Ancients did) making the Center of the Circular Orbit, the Center of equal Motion, 1725 Eccentricity will not account for the annual Inequalities. For the Prosthaphereses, or Differences between the Sun's mean and true Places are thus twice as much as what they will amount to with this half of antient Eccentricity. This Defect of Eccentricity of the Circular Orbit was afterwards nearly compensated for, by placing half of it each Way from the Orbit's Center, and making the Center of equal Motion at the contrary Extreme to that where is placed the Sun, in the elliptical Orbit, which is supposed, by Ward and Bullialdus, similar to the Center of equal Motion in the Circular Theory. Mr. Street's Eccentricity to this Theory of the Earth's Motion is 1732, Mr. Flamsteed's 1692, Mayer's 1680, Morris's 1683, Our's 1685, which make a small Difference in finding the Sun's true Place from his Mean; nor is this Difference likely to be adjusted, while different Observators of the Heavens, like different Surveyors of the same GENTLEMAN'S ESTATE, vary, in their Quantities, from one another.

Kepler having observed, That the Squares of the Periodical Times of revolving Bodies, are as the Cubes of the Distances from the Centers of the Orbits, about which they are supposed to perform their equal Motions, (i. e. as the Cubes of the mean Distances from the Body about which they revolve.) This UNIVERSAL Theory of MOTION, examined and confirmed by Sir Isaac Newton, and verified by Observation itself, is only contradicted by supposing the Sun's Motion, and the Earth at Rest.

PRACTICAL QUESTIONS

IN

CHRONOLOGY answered.

Of USE in HISTORY.

QUESTION I.

AN Y Year, Month, and Day of the Month, being given, to find the Day of the Week, correspondent?
EXAMPLE. What Day of the Week was the 12th of March, in the 24th Year before the reputed Year of Christ, the Day where-
 in the Eclipse of the Moon happened, a little before the Death of King Herod, at the real Birth of Christ.
 By Tab. p. 148, the Dominical Letters, for the Year, are found, at Sight, to be BA, which last Letter, A, is used after the 24th of Februa-
 ry. Consequently by Tab. p. 149, March 12th, that Year, was on a Sunday, required.

QUESTION II.

The 1st, 2d, 3d, or 4th Week-Day (Sunday, Monday, Tuesday, &c.) being given in any Month of a given Year, to find from thence the Day of the Month?
 This being the Reverse of the former Question is answered from the same Tables, directly, at Sight.

QUESTION III.

The Day of the Month that Easter falls on, in any Year, being given, to find in what Years it shall happen so again, or has so before happened viz. on the same Day of the Month, either according to Old or New Style?

EXAMPLE. Let the 25th of March be the Time of Easter happening, when, according to the Proverb, *My Lord falls in my Lady's Lap?* Here, March the 25th being on a Sunday, the Dominical Letter, by Tab. p. 149, will be always G, for Old or New Style.
 Then, by Tab. p. 155, find March 25th for the Time of Easter happening, against the Dominical Letter G, New Style, and you will find 3. 11. 14. for the Golden Numbers, standing against the Time, when Easter can possibly happen, with the Dominical Letter G, for that Time, New Style.

In Old Style, it happened when the Golden Numbers were 5. 13. 16. and Dominical Letter G (See Tab. p. 156). Now, a Dominical Letter Cycle is 28 Years, and a Golden Number Cycle, 19 Years, (in which Periods the Dominical Letters and Golden Numbers return to the same Order, O. S.) therefore 28 by 19 produce 532 Years, called the Dionysian Period, being the Cycle after which Easter, and all the Moveable Feasts, began and followed the same Order, again, Old Style. But, this Order is interrupted in the New Style, by the Day reckoned less in every succeeding odd hundredth Year, than is reckoned by the Old Julian Account, and there will be intermediate Years, in both Styles, wherein the Golden Number and Dominical Letter will happen the same over again, before the Expiration of the said Dionysian Period; because of the Rotation of 7 Letters 4 Times in the 28 Year-Cycle. For, by Tab. p. 150, the Years when the Golden Numbers are 3. 11. and 14. are as follows.

Golden Number 3.				Golden Number 11.				Golden Number 14.			
1769	1807	1902	2016	1777	1815	1910	2005	1780	1818	1913	2008
88	26	21	G. 35	96	34	29	24	99	37	32	27
Examine these	45	40	54	Adding 8	53	48	43	Adding	56	G. 51	G. 46
Years by Dom.	64	59	73	to the	72	67	62	3 to the	75	70	65
Let. Tab. for	G. 83	78	G. 91 &c.	former.	91	86	81	former.	G. 94	89	84
N. S. p. 149.		97					99 &c.				2102

In this Manner are the Years backward examined for N. S. by Centuries. And likewise the Years backward or forward, for O. S. are thus examined, correspondent to the Golden Numbers 5. 13. 16. by the Dominical Letter Table, Old Style, at p. 148, for any Years (before or since 1759) to which the said Golden Numbers belong.

According to N. S. *Easter* falls on March 25, in the Years 1883, 1894, 1951, 2035, 2046, 2091, &c. from above.

QUESTION IV.

A LADY was born on an Ash-Wednesday, 23 February, N. S. and was under 30 Years of Age, 1759, required her Age?
 By Perpetual Time-Tables, (see farther on) When Ash-Wednesday was on the 23d of February Easter fell on April 10. And, by Tab. p. 155, the Dominical Letter B, and Golden Numbers 7. 10. 15. 18. when Easter can so happen.
 Now, by examining the Dates for those Golden Numbers, from Table, p. 150, between 1729 and 1759 (the Difference 30 Years, being above the Age) and that Dominical Letter for those Dates, by Tab. p. 149, You will find only the Year 1735, for the Time of her Birth, when the Golden Number was 7, and Dominical Letter B. Consequently, the Lady's Age is 24 Years, required.

QUESTION

QUESTIONS in CHRONOLOGY answered.

QUESTION V.

A PERSON above 58 Years of Age, in the Year 1759, was born on a Friday, July 14, about the Year 1700, Old or New Style: Required his Age?

The Dominical Letter, according to Old or New Style, by Tab. p. 149, was A; the nearest Year to which 1699, O. S. or 1702, N. S. When the Dominical Letter was A, for both Styles, by Tab. p. 148, and Tab. p. 149. Whence, his Age, in 1759, must be 60, according to Old Style, or 57, according to New, therefore his Age is 60, required.

In the 1 Year of the reputed Birth of Christ, the Golden Number was 2, the Cycle of the Sun 9, and Roman Indiction 3, Year of the Dionysian Period 457, of the Julian Period 4713. Hence, the Reason of the RULES, p. 146.

QUESTION VI.

The Cycles of the Sun and Moon being given, to find the Year of the Dionysian Period, according to Old Style?

RULE. Multiply the Cycle of the Sun by 57, and the Cycle of the Moon, or Golden Number, by 476; divide the Sum of the Products by 532, the Remainder, after the Quotient, will be the Year of the Dionysian Period, required.

EXAMPLE. The Golden Number is 1, and Cycle of the Sun 9, at the reputed Birth of Christ, to find the Year of the Dionysian Period, at that Time?

476	57
1 Golden Number, or Cycle of the Moon.	9 the Cycle of the Sun.
476	513
513	
532)989(1	
457 Remainder, Dionysian Period, required.	

QUESTION VII.

The Cycle of the Sun, Moon, and Indiction, being given to find the Year of the Julian Period, according to Old Style?

RULE. Multiply the Number 4845 by the Year of the Cycle of the Sun, the Number 4200 by the Year of the Cycle of the Moon, and the Number 6916 by the Year of the Indiction Cycle: Divide the Sum of these Products by 7980 (the Number of Years in the Julian Period), and the Remainder, after the Quotient, will be the Year of the Julian Period, required.

EXAMPLE. To find the Year of the Julian Period, at the reputed Birth of Christ, when the Cycle of the Sun was 9, the Cycle of the Moon, or Golden Number, was 1, and the Indiction 3?

4845	4200	6916
9 Cycle ☉.	1 Cycle ☾.	3 Indiction.
43605	4200	20748
		4200
		43605
		7980)68553(8
		63840
		Remainder 4713 Julian Period required.

QUESTION VIII.

To find the Year of the Dionysian Period, according to Old Style, for any Year before or since Christ?

RULE I. Add 75 (the Complement of 457 to the whole Period 532) to the Number of Years before Christ, and divide that Sum by 532, and subtract what remains from 532, and the last Remainder will be the Year of the Dionysian Period, for the Number of the Year before Christ, required?

N. B. If the Sum of 75, and the Years before Christ, is less than 532, subtract that Sum from 532, and the Remainder will be the Year of the Period, required.

RULE II. Add 457 to the Number of Years since Christ, and divide that Sum by 532, and the Remainder will be the Year of the Dionysian Period, for the Year, since Christ, required.

N. B. If the Sum of 457, and the Years since Christ, is less than 532, that Sum will be the Year of the said Period?

EXAMPLE I. 1000 Years before Christ.

$$\begin{array}{r} +75 \\ 532)1075(2 \\ 1064 \\ \hline -11 \\ 532 \end{array}$$

Remainder 521 Year of the Dionysian Period, required.

EXAMPLE II. 2000 Years since Christ.

$$\begin{array}{r} +457 \\ 532)2457(4 \\ 2128 \end{array}$$

Remainder 329 Year of Dionysian Period, required.

QUESTION IX.

The Year of the Dionysian Period being given to determine from thence the Year before or since Christ?

This being the Reverse of the former Question, let the Dionysian Period be 329, some Year since Christ, or 521, some Year before Christ.

EXAMPLE

QUESTIONS in CHRONOLOGY answered.

EXAMPLE. —521 Number Dionysian Period before Christ.
 532 Number of the wh. Period.
 11 Yrs from Begin. of that Period before Christ.
 457 No. Period the Yr. bef. Christ.
 Sum 468 Years before Christ, when Period was 521.
 Add 532 continually.
 1000 Years before Christ, when Period was 521, &c.
 N. B. In the foregoing Example, before Christ, 1 should be added to the Result, making 522, or else 1 should have been subtracted from 1000, the Yr. before Christ, for the Result of this Operation, then 522, to agree with Chronology. See our Chronological Scale, farther on. *P. 170*

329 No. Period since Christ.
 +75 Years since Christ to Beginning of the 1st Period.
 404 Years since Christ, when Period was 329.
 Add 532 continually.
 936 Years since Christ, when Period was 329.
 532
 1468 Years since Christ, when Period was 329.
 532
 2000 Years since Christ, when Period was 329, &c.

QUESTION X.

To find the Year of the Julian Period, for any Year before or since Christ?
 RULE I. The Number of Years before Christ, subtracted from 4714 (the Number of the Period, in the 1st Year of the reputed Birth of Christ) the Remainder will be the Year of the Julian Period for the Year before Christ, required.
 RULE II. The Number of Years, since Christ, added to 4713, the Number of the Period the Year before Christ, the Sum will be the Year of the Julian Period, for the Year, since Christ, required.

EXAMPLE I. 4714 Year of the Period in 1st Yr. bef. Christ.
 —1759 Years before Christ.
 2955 Year of the Period, required.

EXAMPLE II. 1760 Years since Christ.
 4713 Year of the Period at Christ.
 6473 Year of the Period, required.
 N. B. 7980 the whole Period.

QUESTION XI.

To find the Year before or since Christ correspondent to the given Year of the Julian Period?
 This QUESTION is the Reverse of the former.
 RULE I. The given Year of the Julian Period must be subtracted from 4714 for the Year before Christ.
 RULE II. 4713 must be subtracted from the given Year of the Julian Period for the Year since Christ.
 EXAMPLE. The given Years of the Julian Period, are 2955 and 6473, required the Years of Christ?
 subtracted 4714 subtracted 4713
 Remainder 1759 Yr before Christ, reqd. Rem. 1760 Year since Christ, required.
 N. B. If the Number of the Julian Period is equal to, or less than 4713, the Year is before, if greater, since Christ.

QUESTION XII.

A Person since the Year 1740 said, that 49 Times the Years of his Age was equal to the Date of the Year on which he was born: What Age was he?
 PUT x for the Years of the Person's Age. Then $49x = 1758 - x$. By Supposition. But $49x + x = 1758$, or $x = \frac{1758}{50} = 35\frac{8}{44}$ Years of his Age. But this Age being subtracted from 1758 can leave no whole Number for the Date: whence it could not be 1758 to which his Age reckoned. Let $y =$ the Date, then $x = \frac{y}{50}$; here $y = 1750$, &c. For the Time to which his Age was reckoned, being divisible by 50. Whence $\frac{1750}{50} = 35$, his Age, and 1715 was the Date of his Birth, required.

A GENERAL PROPOSITION.

To find a whole Number, which being divided by other given whole Numbers, given whole Numbers shall remain.
 EXAMPLE. To find a whole Number, which being divided by 2, 3, 4, 5, and 6; 1, 2, 3, 4, and 5, shall respectively remain; but be divided by 7, Nothing shall remain.
 AXIOM. Any whole Numbers being subtracted from any other whole Numbers, a whole Number, or Nothing, shall remain.
 Conditions 1. 2. 3. 4. 5. 6.
 Put $x =$ the whole Number, sought: Then, $\frac{x-1}{2}, \frac{x-2}{3}, \frac{x-3}{4}, \frac{x-4}{5}, \frac{x-5}{6}, \frac{x}{7}$, will be the 6 Conditions of the Question, expressing the same whole Number.
 Let 1st Condition, $\frac{x-1}{2} = a$, Then $x = 2a+1$. Substitute which Value of x , in 2d Condition. Then $\frac{2a-1}{3}$, a whole Number.
 { But, $\frac{3a-3}{3}$, a whole Number.
 Whence, $x = 6b+5$. The 1st and 2d Conditions. . . . Here, $a = 3b+2$ (in lowest Terms) Dif. . . . $\frac{a-2}{3} = b$, a whole Number.

QUESTIONS in CHRONOLOGY answered.

Substitute this Value of x , in the 3d Condition Then, $\frac{6b+2}{4}$, or $\frac{3b+1}{2}$, a whole Number.

$$\left\{ \frac{4b}{2}, \text{ a whole Number.} \right.$$

Whence, $x = 12c + 11$. The 1st, 2d, and 3d Conditions. . . . Here $b = 2c + 1$. . . Dif. $\frac{b-1}{2} = c$, a whole Number.

Substitute this Value of x , in the 4th Condition. Then $\frac{12c+7}{5} = 2c+1, + \frac{2c+2}{5}$, a whole Number.

$$\left\{ \frac{5c}{5}, \text{ a whole Number.} \right.$$

1st Dif. . . . $\frac{3c-2}{5}$, a whole Number.

Whence, $x = 60d + 59$. The 1st, 2d, 3d, and 4th Con. . In lowest Terms, Here, $c = 5d + 4$. . . 2d Dif. $\frac{c-4}{5} = d$, a whole Number.

Also 5th Condition, as will appear.

For, substitute this Value of x , in the 5th Condition. Then, $\frac{60d+54}{6}$ is a whole Number by Consequence.

Therefore, substitute the aforesaid Value of x , in 6th Condition, Then, $\frac{60d+59}{7} = 8d+8, + \frac{4d+3}{7}$, a whole Number.

$$\left\{ \text{But } \frac{7d+7}{7}, \text{ a whole Number.} \right.$$

1st Dif. $\frac{3d+4}{7}$, a whole Number.

Whence, $420e + 119$. The 1. 2. 3. 4. 5. and 6th Conditions. . . . Here, $d = 7e + 1$. . . 2d Dif. $\frac{d-1}{7} = e$, a whole Number.

The GENERAL AFFIRMATIVE THEOREM, in the lowest Terms; answering all the Conditions of the Question, let e be what it will.

Let $e = 0$, then 119 is the least whole Number.

$e = 1$, then 539 is the next whole Number.

$e = 2$, then 959 the next, &c. ad infinitum.

All other Questions of this Nature are answered in this Manner: the chief Business being to reduce the required Conditions, to Fractions in the lowest Terms, as the Operation proceeds, in order that the final Theorem may take in all the Cases that can happen; viz. from 0, 1, 2, 3, 4, &c. the successive Values of the assumed whole Number. And this Theorem will be either affirmative or negative.

OTHER WISE negatively.

PUT x for the Number as before, and Conditions the same.

Let $\frac{x-1}{2} = a$, 1st Condition. Hence $x = 2a + 1$, as before. Substitute in 2d Condition. Then, $\frac{2a-1}{3}$, a whole Number.

$$\left\{ \text{But } \frac{3a}{3}, \text{ a whole Number.} \right.$$

Whence, $x = 6b - 1$. The 1st and 2d Conditions. . . . Here, $a = 3b - 1$. . . Dif. . . . $\frac{a+1}{3} = b$, a whole Number.

Substitute this Value, in the 3d Condition. . . . Then $\frac{6b-4}{4} = \frac{3b-2}{2}$, a whole Number.

Whence, $x = 12c - 1$. The 1. 2. 3. Conditions. . . . Here $b = 2c$. . . viz. $-1 + b, + \frac{b}{2} = c$, a whole Number.

Substitute this Value of x , in 4th Condition. Then, $\frac{12c-5}{5} = -1 + 2c, + \frac{2c}{5}$, a whole Number.

Whence, $x = 60d - 1$. The 1. 2. 3. 4 Conditions. . . . Here $c = 5d$ $\frac{c}{5} = d$, a whole Number.

And also 5 Conditions, as appears.

For, Substitute this Value of x , in the 5th Condition. Then, $\frac{60d-6}{6}$, a whole Number by Consequence.

Therefore,

QUESTIONS in CHRONOLOGY answered.

Therefore, substitute the said Value of x , in the 6th Condition. Then $\frac{60d-1}{7}$, a whole Number.

$$\text{viz. } 3d + \frac{4d-1}{7}, \text{ a whole Number.}$$

$$\text{1st Dif. } \left\{ \frac{7d}{7}, \text{ a whole Number.} \right.$$

$$\text{2d Dif. } \left\{ \frac{3d+1}{7}, \text{ a whole Number.} \right.$$

General affirmative THEOREM.
Whence, $x = 420e + 119$, as before,
answering all the Conditions. }

$$\text{Here, } d = 7e + 2 \dots \frac{d-2}{7} = e, \text{ a whole Number.}$$

$$\text{But } \frac{7d-7}{7}, \text{ a whole Number.}$$

$$\text{From above. 1st Dif. } \frac{3d-6}{7}, \text{ a whole Number.}$$

General negative THEOREM.
Whence, $x = 420e - 301$, answering
all the Conditions. }

$$\text{Here, } d = 7e - 5 \dots \text{2d Dif. } \frac{d+5}{7} = e, \text{ a whole Number.}$$

Let $e=1$, then 119 is the least whole Number.
 $e=2$, then 539 is the next whole Number.
 $e=3$, then 959 the next, &c. ad infinitum, as before.

QUESTION XIII.

REQUIRED the YEAR of CHRIST when the Golden Number is 5, and the Cycle of the Sun 6?
In the 1st Condition, 1 is to be added to the Year, to be divided by 19 for the Golden Number 5 to remain.
In the 2d Condition, 9 is to be added to the Year, to be divided by 28 for the Cycle of the Sun 6 to remain.

Conditions 1. 2. 1. 2.

Put $x = \text{No. Year sought}$: Then $\frac{x+1-5}{9}$, $\frac{x+9-6}{28}$, Or $\frac{x-4}{19}$, $\frac{x+3}{28}$ will be the two Conditions of the Question, expressing the same whole Number.

Let 1st Condition $\frac{x-4}{19} = a$, Then $x = 19a + 4$. Substituting which in 2d Condition $\frac{19a+7}{28}$, a whole Number.

$$\text{1st Dif. } \left\{ \text{But } \frac{28a}{28}, \text{ a whole Number.} \right.$$

$$\text{2d Dif. } \frac{9a-7}{28}, \text{ a whole Number.}$$

$$\text{3d Dif. } \frac{10a+14}{28}, \text{ a whole Number.}$$

$$\text{Here, } a = 28b - 21 \dots \frac{a+21}{28} = b, \text{ a whole Number.}$$

$$\text{But } \frac{28a+28}{28}, \text{ a whole Number.}$$

$$\text{From above, 1st Dif. } \frac{9a+21}{28}, \text{ a whole Number.}$$

$$\text{2d Dif. } \frac{10a-14}{28}, \text{ a whole Number.}$$

$$\text{Here, } a = 28b + 7 \dots \text{3d Dif. } \frac{a-35}{28} = -1 + \frac{a-7}{28} = b, \text{ a whole Number.}$$

QUESTION XIV.

IN what Year of the Dionysian Period was it in the Year before Christ, when the Golden Number was 1, and Cycle of the Sun 9?
The FIRST Conditions of this Question is to find the least Number, which divided by 19 and 28, shall leave 1 and 9 respectively remaining.

Conditions 1. 2.

Put $x = \text{that Number}$, Then $\frac{x-1}{19}$, $\frac{x-9}{28}$ will be the 2 Conditions of the Question, expressing the same whole Number.

QUESTIONS in CHRONOLOGY answered.

Let 1st Condition $\frac{x-1}{19} = a$, Hence $x = 19a + 1$.

NEGATIVE THEOREM.
Whence, $x = 532b - 75$.

AFFIRMATIVE THEOREM.
 $x = 532b + 457$.

Let b be
assumed so
as the No.
comes out less
than 532.

Substitute this Value in 2d Condition, Then, $\frac{19a-8}{28}$, a whole Number.

But $\frac{28a-28}{28}$, a whole Number.

1st Dif. $\frac{9a-20}{28}$, a whole Number.

2d Dif. $\frac{10a+12}{28}$, a whole Number.

Here, $a = 28b - 4 \dots$ 3d Dif. $\frac{a+32}{28} = 1 + \frac{a+4}{28} = b$, a whole Number.

By the Negative Theorem.
Let $b=1$, then 457 the Number of the Period.

By the Affirmative Theorem.
 $b=0$, then 457 is the same Number required.

But $\frac{28a+28}{28}$, a whole Number.

From above, 1st Dif. $\frac{9a+36}{28}$, a whole Number.

2d Dif. $\frac{10a-44}{28}$, a whole Number.

N. B. The Dionysian Period limits the
Answer to the least whole Number, or
Number in that Period.

Here, $a = 28b + 24 \dots$ 3d Dif. $\frac{a-80}{28} = -2 + \frac{a-24}{28} = b$, a whole Number.

N. B. Either of the above Theorems find as many conditional Numbers as you please.

QUESTION XV.

THE Golden Number being one, in the Year before Christ, the Cycle of the Sun 9, (as before) and the Roman Indiction 3, what was then the Year of the Julian Period?

The Conditions here are to find the least whole Number, which being divided by 19, 28, and 15; 1, 9, and 3 shall respectively remain.

Conditions 1. 2. 3.

Put $x =$ the Number sought, Then $\frac{x-1}{19}$, $\frac{x-9}{28}$, $\frac{x-3}{15}$, will be the 3 Conditions of the Question, expressing the same whole Number.

Two of which Conditions were determined above, viz. $x = 532b + 457$.

Substitute which Value of x , in the 3d Condition, Then $\frac{532b+454}{15}$, a whole Number.

Viz. $35b + 30 + \frac{7b+4}{15}$, a whole Number.

$\left\{ \frac{15b-15}{15} \right.$, a whole Number.

1st Dif. $\frac{8b-19}{15}$, a whole Number.

Here, $b = 15c + 8 \dots$ 2d Dif. $-1 + \frac{b-8}{15} = c$, a whole Number.

But $\frac{15b+15}{15}$, a whole Number.

From above, 1st Dif. $\frac{8b+11}{15}$, a whole Number.

Here, $b = 15c - 7 \dots$ 2d Dif. $\frac{b+7}{15} = c$, a whole Number.

NEGATIVE THEOREM.

Whence $x = 7980c - 3267$.

And when $c=1$, then 4713 is the Number
of the Period, as before.

N. B. Either of the above Theorems find as many conditional Numbers as you please.

QUESTION XVI.

THE Golden Number being 11, Cycle of the Sun 2, and Indiction 9, required the Date of the Year, O. S. when these happened?

QUESTIONS in CHRONOLOGY answered.

Conditions 1. 2. 3. 1. 2. 3.
Put x = Year fought. Then $\frac{x+1-11}{19}$, $\frac{x+9-2}{28}$, $\frac{x+3-9}{15}$, or $\frac{x-10}{19}$, $\frac{x+7}{28}$, $\frac{x-6}{15}$, will be the 3 Conditions of the Question, expressing the same whole Number.

Let 1st Condition, $\frac{x-10}{19} = a$, then $x = 19a + 10$.

Substitute this Value of x in 2d Condition. Then, $\frac{19a+17}{28}$, a whole Number.

Whence, $x = 532b + 105$. The 1st and 2d Conditions.

{ But $\frac{28a+28}{28}$, a whole Number.

1st Dif. $\frac{9a+11}{28}$, a whole Number.

2d Dif. $\frac{10a+6}{28}$, a whole Number.

Here, $a = 28b + 5$ 3d Dif. $\frac{a-5}{28} = b$, a whole Number.

Substitute, the foregoing Value x , in the 3d Condition. Then $\frac{532b+99}{15}$, a whole Number.

Viz. $35b + 6 + \frac{7b+9}{15}$, a whole Number.

But $\frac{15b+15}{15}$, a whole Number.

1st Dif. $\frac{8b+6}{15}$, a whole Number.

Here, $b = 15c + 9$ 2d Dif. $\frac{b-9}{15} = c$, a whole Number.

N. B. The above Numbers can happen but once in a Julian Period of 7980 Julian Years, as appears by the Theorem: Notwithstanding which, as many conditional Numbers of the same Properties may be found by the Theorem as you please.

The GENERAL PROPOSITION (Page 161) OTHERWISE Answered.

LET the Quotient of the same whole Number when divided by another whole Number be called a , when divided by a second whole Number be called b , by a third c , by a fourth d , by a fifth e , by a sixth f , &c.

FORMER EXAMPLE. To find a whole Number, which being divided by 2, 3, 4, 5 and 6; 1, 2, 3, 4, and 5 shall, respectively, remain; but being divided by 7, Nothing shall remain.

Conditions 1. 2. 3. 4. 5. 6.
Here, $2a+1 = 3b+2 = 4c+3 = 5d+4 = 6e+5 = 7f$, which are the 6 Conditions of the Question, expressing the same whole Number.

N. B. The chief Thing here to be considered is to find the Value of the different Quotients a , b , c , d , &c. of the same whole Number, when divided by different whole Numbers, so as Nothing shall remain, according to the several Expressions, and Conditions, as above.

AXIOM. Any whole Number taken from the same, or a different whole Number, a whole Number, or Nothing, shall remain.

1. 2. Conditions.
In the 1st and 2d Conditions. $2a+1 = 3b+2$, a whole No. sought; a and b conditional whole Nos. required.
Subtract 1 from each Side, $2a = 3b+1$, a conditional whole Number, divisible by 2.
Subtract $2b$. . . a whole Number, divisible by 2.

Whence, $3b+2=5$, the 1st and 2d Conditions. . . Here $b=1$, least. $b+1$, a whole Number, divisible by 2.

NOW, 6 being the least Multiple, divisible by 2, 3, the Divisors respecting the 2 first Conditions,

3. 1. 2. Conditions.
Therefore, . . . $4c+3 = 6a+5$, a whole Number sought, c and a whole Numbers, required.
Subtract 3 from each Side . . . $4c = 6a+2$, a whole Number, divisible by 4.
Subtract $4a$. . . a whole Number, divisible by 4.

Whence, $6a+5=11$, The 1. 2. 3. Conditions. . . Here $a=1$, least. . . $a+1$, a whole Number, divisible by 2.

NOW, 12 being the least Multiple, divisible by 2, 3, 4, the Divisors respecting the 3 first Conditions,

4. 1. 2. 3. Conditions.
Therefore, . . . $5d+4 = 12a+11$, a whole Number sought, d and a whole Numbers, required.
Subtract 4 from each Side . . . $5d = 12a+7$, a whole Number, divisible by 5.
Subtract $10a+5$, a whole Number, divisible by 5.

Here a may be discovered $= 4$. . . $2a+2$, a whole Number, divisible by 5.

$5a$. . . a whole Number, divisible by 5.

$3a-2$, a whole Number, divisible by 5.

Whence, $12a+11=59$, The 1. 2. 3. 4. Conditions. . . Here $a=4$, least. $a-4$, a whole Number, divisible by 5.

NOW,

QUESTIONS in CHRONOLOGY answered.

NOW, 60 being the least Multiple, divisible by 2, 3, 4, 5, the Divisors respecting the 4 first Conditions,

1. 2. 3. 4. Conditions.

Therefore, $6a+5 = 60a+59$, a whole Number sought, a and a whole Numbers.

Subtract 5 from each Side, $6a = 60a+54$, a whole Number, divisible by 6.

Hence, $60a+59=59$, The 1. 2. 3. 4. 5. Conditions. . . Here $a=0$, least . . which is already divisible by 6.

NOW, 60 being the least Multiple, divisible by 2, 3, 4, 5, 6, the Divisors respecting the first 5 Conditions,

1. 2. 3. 4. 5. Conditions.

Therefore, $7f = 60a+59$, a whole Number sought, divisible by 7, f and a whole Numbers.

Subtract $56a+56$, a whole Number, divisible by 7.

$4a+3$, a whole Number, divisible by 7.

$7a$. . . a whole Number, divisible by 7.

$3a-3$, a whole Number, divisible by 7.

$3a-3$, a whole Number, divisible by 7.

Hence, $60a+59=119$, The 1. 2. 3. 4. 5. 6. Con. Here $a=1$, at least $a+6$, a whole Number, divisible by 7.

NOW, 420 being the least Multiple divisible by 2, 3, 4, 5, 6, 7, the Divisors respecting the 6 Conditions,

1. 2. 3. 4. 5. 6. Conditions.

Therefore, $420a+119$, will be an AFFIRMATIVE GENERAL THEOREM, by which you may determine as many Numbers as you please, having the aforesaid 6 Conditions, where $a=0, 1, 2, 3, 4$, &c. assumed at Pleasure.

The LEAST MULTIPLE, divisible by several Numbers, is had by a continual Multiplication of the lowest Parts, and least Multiples of those Numbers, that will compose all the different Wholes; viz. make up as many of the following Divisors as you can, out of the preceding Ones, and their Multiples, for determining the least Multiple of all Numbers, universally.

Example. 2, 3, being different and not divisible by each other have $1 \times 2, \times 3=6$ for the least Multiple.

But 2, 3, 4, being the same as 2, 3, 2^2 , multiplied together, therefore the different Numbers multiplied, taking the square, cube, &c. Roots, viz. $2 \times 3 \times 2=12$, is the least Multiple of 2, 3, 4.

Again, 2, 3, 4, 5 being the same as 2, 3, 2^2 , 5, the different Numbers multiplied, taking in the Roots, $2 \times 3 \times 2 \times 5=60$, the least Multiple of 2, 3, 4, 5.

Again, 2, 3, 4, 5, 6, being the same as 2, 3, 2^2 , 5, (2, 3) multiplied, the different Numbers multiplied, taking in the Roots, and excluding former Similitudes, $2 \times 3 \times 2 \times 5=60$, the least Multiple of 2, 3, 4, 5, 6.

And, if 2, 3, 4, 5, 6, 7, 8, 9, were the Divisors, then 2, 3, 2^2 , 5, (2, 3) 7, 2^3 , 3^2 , will be the different Numbers of them multiplied together, taking in their Roots, viz. $2 \times 3 \times 2 \times 5 \times 7 \times 2 \times 3=2520$, the least common Multiple of them all.

So for other Cases.

QUESTION XVII.

THE Golden Number being 12, Cycle of the Sun 4, and Roman Indiction 7, to determine the Year of the Julian Period, and likewise the Year of Christ?

Let a, b, c , be the 3 Quotients after dividing the Year of the Period, by 19, 28, and 15, for 12, 4, and 7, respectively, to remain.

1. 2. 3. Conditions.

Therefore, $19a+12 = 28b+4 = 15c+7$.

Subtract $12 \dots 19a = 28b-8$, a whole Number divisible by 19, a and b whole Numbers.

Subtract $19b \dots$ a whole Number divisible by 19.

Remainder $9b-8$, a whole Number divisible by 19.

1st Dif. $10b+8$, a whole Number divisible by 19.

Here, $b=3$, the least. 2d Dif. $b+16$, a whole No. divisible by 19. Whence $28b+4=88$. 1. & 2. Condit.

NOW, 53 being the least Multiple, of which 19 and 28 are Divisors, respecting the first 2 Conditions.

Therefore, $15c+7 = 532a+88$

Subtract $7 \dots 15c = 532a+81$, a whole Number, divisible by 15, c and a whole Numbers.

$15 \times 35a+5 \times 15$. Subtract $525a+75$, a whole Number, divisible by 15.

$7a+6$, a whole Number, divisible by 15.

$15a+15$, a whole Number, divisible by 15.

1st Dif. $8a+9$, a whole Number, divisible by 15.

Here $a=12$, the least, 2d Dif. $a+3$, a whole No. divisible by 15. . . . Whence $532a+188=6476$

Year of Julian Period 1st Yr. bef. Chr. -4713

Year of Christ, required . . . 1759

N. B. The last Quantity divisible by a whole Number should (for finding the least Number sought) be brought out affirmative.

QUESTION XVIII.

REQUIRED a PERSON's Age, on December 4, 1759, N. S. born March 8, (since 1700) in the Year when the Dominical Letter O. S. B, and Golden Number 19? And required also the Date of the Year, and Day of the Week, when the same Person was born?

Gol. No. 19. Dom. Letters.

By Tab. p. 150. $\left\{ \begin{array}{l} 1709 \dots B \\ 28 \text{ with} \dots GE \\ 1700 \text{ and } 1759 \dots 47 \dots D \end{array} \right\}$ By Tab. p. 148.

Hence, the Person was born in the Year 1709, O. S. Therefore, on March 8, O. S. . . . 1759

The Year of the Person's Age will be Dif. Years 50

But, December 4, 1759, N. S. answers to November 23, 1759, O. S.

BIRTH-DAY, March 8, 1759, O. S.

By Table, Page 32. The Difference of Days betwixt which Times, are $245+15=260$ Days.

For, by Table, Page 129, From January to November 23. . . 327 Days. . . To March 8. . . 67 Days. . . Diff. 260

So that the PERSON'S AGE will be 50 Years, and 260 Days, required.

ARITHMETICAL OPERATIONS.

TO turn Time into Degrees and Parts of the Equator, arithmetically ?

RULE. Take the several Sums of the Hours, Minutes, and Seconds, and the Halves thereof respectively. Then take in or consider a Decimal—Place to the Right Hand of each of which Sums, as Units Place, and you will have the Degrees, Minutes, and Seconds of a Degree answering thereto; which being collected will be the Degrees and Parts of the Equator, required.

E X A M P L E S.

$\frac{1}{2}$. . 2	4 Hours 5 Hours.	$\frac{1}{2}$. . 2.5
Sum 6 Hours = 60°.		Sum 7.5 = 75°

$$\begin{array}{r} 7 \text{ Hours} \\ \frac{1}{2} \dots 3.5 \\ \hline \text{Sum } 10.5 = 10^{\circ} 30' \\ \phantom{\text{Sum }} 11 \text{ } 45' \end{array} \qquad \begin{array}{r} 47 \text{ Minutes} \\ \frac{1}{2} \dots 23.5 \\ \hline 70.5 = 70^{\circ} 30' \\ \text{Divide by } 6 \dots 11^{\circ} 45' \end{array}$$

Or, $\begin{array}{r} \text{H M} \quad \text{H} \\ 7 \quad 47 = 7 \cdot 78333, \text{ etc. see p.} \\ \frac{1}{2} \quad \cdot \quad 3 \cdot 89166 \quad 14. \end{array}$

Sum 11.67499
= 1160,75
45' as before.

Collected Sum 116 45 Degrees and Minutes, *required.*

$13 \text{ Hours} \dots 33 \text{ Minutes} \dots 51 \text{ Seconds}$
 $\frac{1}{2} \dots 6.5 \qquad \frac{1}{2} \dots 16.5 \qquad \frac{1}{2} \dots 25.5$

Sum $19.5 = 19^{\circ}$ Sum $49.5 = 49^{\circ} 15'$ Sum $76.5 = 76^{\circ} 15''$
 $\begin{array}{r} 8 \text{ } 15' \\ 12 \text{ } 45'' \end{array}$ $\div 6 \dots 8^{\circ} 15'$ $\div 6 \dots 12' 45''$

Collected Sum $203^{\circ}27'45''$ required.

9 Hours	19 Minutes . . .	27 Seconds
$\frac{1}{2} \dots 4.5$	$\frac{1}{2} \dots 9.5$	$\frac{1}{2} \dots 13.5$
Sum 13.5 = 135°	28.5 = 28 5'	40.5 = 40 5"
4 45'	÷ 6 .. 4 45'	÷ 6 .. 6' 45"
	6 45"	

Collected Sum $139^{\circ} 51' 45''$, required.

$$\begin{array}{r} \text{H M S} \\ \text{Or, } 13 \ 33 \ 51 \\ \qquad \qquad \qquad ,85 \\ = 13.564166, \text{ etc.} \\ \frac{1}{2} \dots 6.782083 \end{array}$$

20.346249 = 203°.46249 . . see p. 14.
as before, 27'.7494
nearly. 44".964

TO turn Degrees and Parts of the EQUATOR into TIME, arithmetically?

RULE. Cut off a Figure to the Right Hand, from the Degrees, Minutes, and Seconds, and take one Third away from each respective Quantity, (after such a Separation of Integers, and the Decimal of each to the Right Hand) what remains will be Hours, Minutes, and Seconds of TIME, respectively; which being collected into a Sum will be the Time sought.

E X A M P L E S.

Hours.
195° or 19,5°
Subtract $\frac{1}{3}$. . 6,5
Remainder 13 Hrs.

80 15' = 80,25 Or, 825
Subtract $\frac{1}{3}$. . ,275

Remainder ,55 Hr.
or 33 Min. reqd.

$$\begin{array}{r} \text{Min}^s. \\ 12' 45'' = 12', 75 \text{ Or } 1,275 \\ \text{Subtract } \frac{1}{3} \text{ . . } .425 \\ \hline \text{Remainder, Min}^s. ,85 \\ 51 \text{ Seconds, reqd.} \end{array}$$

Secs.
56⁴ or 5,6

Subtract $\frac{1}{3}$. . 1,8666

Remainder. Secs. 3,7334
44 Thirds, reqd.

Or, 13,9^o 5,1' 4,5"
 Subtract $\frac{1}{3}$. . . 4,63333 Sub. $\frac{1}{3}$ 1,7 Sub. $\frac{1}{3}$. . . 1,5
 Rem. Hours 9,26667 Rem. M. 3,4 Rem. 3 Secs.
 9^h 16^m 0002 24^s
 3 24^s
 + 3

Collected Sum 9^h 19^m 27^s, required.

$\frac{1}{3} \dots \overset{20,3^0}{6,76666} \dots$ Sub. $\frac{1}{3} \dots \overset{2,7'}{,9} \dots$ Sub. $\frac{1}{3} \dots \overset{4,5''}{1,5} \dots$
 $\frac{13^h}{32^m} \overset{13,53333}{1} \dots$ Rem. $1,8 = 1^m 48^s$ Rem. 3 Secs.
 $\overset{1}{1} \overset{48^s}{3} \dots$

Collected Sum $13^h 33^m 51^s$, required.

Or, $27', 45'' = 27', 75$, or $2, 775$
 Subtract $\frac{1}{3}$. . $.925$
 Remainder. Min^s. $1, 850$
 51^s as before

A TABLE for finding the NUMBER of DAYS advanced and retreated by the EPACTS and LUNATIONS, in the MONTH DAYS, (from 6000 Years before to 6000 Years after Christ) from Old to New Style, from 1800, and the contrary. For determining the EPACT, and Fall of EASTER, according to New Style, as settled by Pope Gregory. See Tab. p. 30. Also to find the Epact, O. S. for the same Centuries.

No. Col. 1.	2.	3.	4.	5.	6.	7.	1.	2.	3.	4.	5.	6.	7.	1.	2.	3.	4.	5.	6.	7.
Centuries before CHRIST, N. S.	Days Diff. from Old to New Style	Days retr. of Cyc. in O. Style	Sum, Days Adv. and retr. of Cyc. from 1800	Days Adv. and Epact from Old to New Style	Golden Nos.	Epact N. S.	Centuries before and since CHRIST, N. S.	Days Diff. from Old to New Style	Days retr. of Cyc. in Old Style	Sum, Days Adv. and retr. of Cyc. from 1800	Days Adv. and Epact from Old to New Style	Golden Nos.	Epact N. S.	Centuries since CHRIST, N. S.	Days Diff. from Old to New Style	Days retr. of Cyc. in O. Style	Sum, Days Adv. and retr. of Cyc. from 1800	Days Adv. and Epact from Old to New Style	Golden Nos.	Epact N. S.
	X		+	+				X		+	+				+		-	-		
B. 6000	47	0	24	23	6	29	B. 2000	17	1	11	6	16	2	B. 2000	13	0	1	12	6	24
5900	46	0	24	22	11	23	1900	16	0	11	5	2	27	2100	14	1	2	12	11	19
5800	45	0	24	21	16	17	1800	15	0	11	4	7	21	2200	15	0	2	13	16	13
5700	44	*1	23	21	2	13	1700	14	1	10	4	12	16	2300	16	0	2	14	2	8
B. 5600	44	0	23	21	7	8	B. 1600	14	0	10	4	17	11	B. 2400	16	1	3	13	7	4
5500	43	0	23	20	12	2	1500	13	0	10	3	3	6	2500	17	0	3	14	12	28
5400	42	1	22	20	17	27	1400	12	1	9	3	8	1	2600	18	0	3	15	17	22
5300	41	0	22	19	3	22	1300	11	0	9	2	13	25	2700	19	1	4	15	3	18
B. 5200	41	0	22	19	8	17	B. 1200	11	0	9	2	18	20	B. 2800	19	0	4	15	8	13
5100	40	1	21	19	13	12	1100	10	1	8	2	4	16	2900	20	0	4	16	13	7
5000	39	0	21	18	18	6	1000	9	0	8	+1	9	10	3000	21	1	5	16	18	2
4900	38	0	21	17	4	1	900	8	0	8	-0	14	4	3100	22	0	5	17	4	27
B. 4800	38	1	20	18	9	27	B. 800	8	0	8	0	19	29	B. 3200	22	0	5	17	9	22
4700	37	0	20	17	14	21	700	7	*1	7	0	5	25	3300	23	1	6	17	14	17
4600	36	0	20	16	19	15	600	6	0	7	1	10	*	3400	24	0	6	18	19	11
4500	35	1	19	16	5	11	500	5	0	7	2	15	13	3500	25	0	6	19	5	6
B. 4400	35	0	19	16	10	6	B. 400	5	1	6	1	1	10	B. 3600	25	1	7	18	10	2
4300	34	0	19	15	15	*	300	4	0	6	2	6	4	3700	26	0	7	19	15	26
4200	33	1	18	15	1	26	200	3	0	6	3	11	28	3800	27	0	7	20	1	21
4100	32	0	18	14	6	20	100	3	1	5	3	16	23	3900	28	1	8	20	6	16
B. 4000	32	0	18	14	11	15	B. 0	2	0	5	3	1	8	B. 4000	28	0	8	20	11	11
3900	31	1	17	14	16	10	100	1	0	5	4	6	2	4100	29	0	8	21	16	5
3800	30	0	17	13	2	5	200	+0	1	4	4	11	27	4200	30	0	8	22	2	*
3700	29	0	17	12	7	29	300	1	0	4	5	16	21	4300	31	*1	9	22	7	25
B. 3600	29	1	16	13	12	25	B. 400	1	0	4	5	2	17	B. 4400	31	0	9	22	12	20
3500	28	0	16	12	17	19	500	2	1	3	5	7	12	4500	32	0	9	23	17	14
3400	27	0	16	11	3	14	600	3	0	3	6	12	6	4600	33	1	10	23	3	10
3300	26	0	16	10	8	8	700	4	0	3	7	17	*	4700	34	0	10	24	8	4
B. 3200	26	*1	15	11	13	4	B. 800	4	1	2	6	3	27	B. 4800	34	0	10	24	13	29
3100	25	0	15	10	18	28	900	5	0	2	7	8	21	4900	35	1	11	24	18	24
3000	24	0	15	9	4	23	1000	6	0	2	8	13	15	5000	36	0	11	25	4	19
2900	23	1	14	9	9	18	1100	7	1	1	8	18	10	5100	37	0	11	26	9	13
B. 2800	23	0	14	9	14	13	B. 1200	7	0	1	8	4	6	B. 5200	37	1	12	25	14	9
2700	22	0	14	8	19	7	1300	8	0	1	9	9	*	5300	38	0	12	26	19	3
2600	21	1	13	8	5	3	1400	9	1	0	9	14	25	5400	39	0	12	27	5	28
2500	20	0	13	7	10	27	1500	10	0	+0	10	19	19	5500	40	1	13	27	10	23
B. 2400	20	0	13	7	15	22	B. 1600	10	0	-0	10	5	15	B. 5600	40	0	13	27	15	18
2300	19	1	12	7	1	18	1700	11	0	0	11	10	9	5700	41	0	13	28	1	13
2200	18	0	12	6	6	12	1800	12	*1	1	11	15	4	5800	42	1	14	28	6	8
2100	17	0	12	5	11	6	1900	13	0	1	12	1	29	5900	43	0	14	29	11	2
														6000	43	0	14	29	16	27

N. B. As 0 signifies the first Year current before Christ, so 100, 200, 300, &c. signify the 101, 201, 301 Years current before Christ, being all Bissextile. So that 1 must be added to all Years before Christ in the Table for the current Years. The Tabular Years being the Years completed. The Years after Christ are let down current (not completed) as they are used in Chronology.

For the MEMORY.

CONSTRUCTION. RULE I. Before Christ, take the Fourth from the Hundreds left one, Then add two: and the Difference of Style will be known.

Year
Ex. 5500 before Christ.

RULE II. From the Hundreds since Christ, take their Fourth, and more two, And the Days will remain betwixt Old Style and New.

54
13—

D.

41+2=43, to be subtracted from the Old Style for the New.

Year
3900 since Chr.

2+1=11—

28 Days to be added to the old Style, for the New.

N. B. The D's Age and Epacts retr. (—) the same No. of Days that the Lunations adv. (+) from O. to N. S. and the contrary, explaining 5. Col. being the Dif. or Sum of 2d and 4. Columns, according to Signs.

To find the Retreat of the Lunar Cycle, or Moon's Age, in the Month-Days of the Old Style, according to its established Rate of Anticipation of 1 Day in every 300 Years (in round Numbers) or 8 Days correctly in every 2500 Years (the mean Rate being 1 Day in 110.7 Years) from whence the Retreat or Advance of the Lunations from Old to New Style, and consequently of the Gregorian Epacts, as settled by the Pope's Authority, are rightly determined. 310.7

RULE I. Subtract 17 from the Centuries since Christ, and divide the Remainder by 25, multiply the Quotient by 8, and reserve the Product.—Divide the last Remainder by 3, and add the Quotient to the Product reserved, if there is now no Remainder, it will be the Retreat of the Cycle, if there is a Remainder that Sum more 1 will be the Cycle's Retreat, required from 1800.

Ex. I. Years 5700 since Christ. 17 25)40(1 by 8 = 8 Product reserved. 25 3)15(5 . . . +5	}	Days 41 Adv. N. S. —13 Retr. of Cyc. Diff. 28 Lunations or Epacts retr. [for that Period.]
---	---	---

Rem^r. 0 Sum 13 Days Retreat of the Cycle, to be deducted from the No. of Days advanced from Old to New Style, for the Cycle's Age and Epact retreated that Dif. from those of Old Style, from 1800.

Ex. II. Years 3100 since Christ. 17 25)14(0 3)14(4, +1 = 5 retr. of Cyc. required. 12	}	31 7 24 2 22-5 = 17 2 Remainder, add 1.
--	---	--

To find the Advance of the Lunar Cycle, or Moon's Age, in Old Style, from 1800, backward to the Year 6000 before Christ?

RULE II. Proceed exactly as in the former RULE for finding the Retreat of Lunations from 1800 to the Year 6000 since Christ; only deduct an Unit from the Result of the same Operation; taking the Distance of Centuries back from the 17th Century in this backward Computation, instead of the Distance of Centuries forward from the same 17th Century, in the forward Computation.

Ex. I. Years 5700 before Christ. +17 Centuries since Christ. 25)74(2, by 8 = 16 Product reserved. 50 3)24(8 . . . 8	Ex. II. Years 3000 since Christ. 17 Centuries since Christ. 25)14(0 by 8 = 0 3)14(4 . . . 4 12 1 2, add 1. Sum 5, by 1. Rule.
--	--

45 Days Advance of the Moon's Cycle, to be connected with the 44 Days retreated from Old to New Style, which being subtracted therefrom, 21 Days will remain that the Cycle's Age and Epacts are advanced, or to be added to the Old Style's Cycle's Age and Epacts, for those of the New Style, from 1800.

Days Advance of the Moon's Cycle, in the Old Style, to which 1 being added, the Advance from Old Style to New, the Sum will be 5 Days retreated by the Cycle's Age or Epacts in the Old Style, or to be subtracted from those of the Old for the New.

The Difference between the Advance of the Days from Old to New Style, and the Retreat of the Lunar Cycle being rightly determined, (by the two foregoing Rules) that Difference being the Retreat of the Cycle's Age or Epacts for a certain Period, in the Month-Days, O. S.—To find from thence the Epacts for N. S.?

RULE. Multiply the Golden Number for the Year by 11, and deduct the said Retreat of Cycle's Age or Epacts from the Product, (adding 30, if it cannot be deducted otherwise) what remains will be the Epact for that Year required.—Or, when (without adding 30) the said Epact Retreat for Years since Christ is deducted from the said Product of the Golden Number by 11, let the Remainder be divided by 30, and what remains will be the required Epact.

Ex. I. Yrs since Chr. G.N. Ds 2300 2 14 Epact Retreat by 11 3000 . . 18 . . 16 E.Ret. 11 18 18 198 16 30)182(6 Rem. 2 Epact, N.S.	22 —14 8 Epact, N.S. right 30 23 22 5 16
--	---

Yr. f. Ch. G.N. Ds 5700 1 28 Epact Retreat. 11 11+30=41 Diff. 13 Epact, N.S. right	41 13 28 41 see above
---	--

CHRONOLOGICAL RULES and REMARKS.

Note
REMARK. The same numeral Quantities, expressing any Number of Years current since Christ, will not express the same Number of Years current before Christ, by the Nature of Chronology.

The numeral Quantities, 0, 100, 200, 300, &c. Years current, since Christ, are found to express

1, 101, 201, 301, &c. Years current, before Christ, and therefore

RULE I. 1 must be added to the numeral Years 0, 100, 200, 300, &c. before Christ, in our Astronomical Tables, to make them correspond with 1, 101, 201, 301, &c. Years Bissextile, before Christ, in Chronology: Or, which is the same, you must take out the Places from the Astronomical Tables aforesaid, for one Year less than any Year current before Christ, in Chronology, to correspond with Truth.

Mr. Mayer's Astronomical Tables required the like Correction in their Use, with our Tables, to make all the solar and lunar Places, for any Number of Years he has expressed, before Christ, to correspond with the true Years, current before Christ, according to Chronology.

Hence, it is doubtful, if Mr. Mayer did not first make this necessary Correction, in taking out his Computations, whether the Principles be correct on which he founds his Quantities of Acceleration of the Moon's Motion. For when he examined and compared the lunar Places in his Tables with those recorded of antient Eclipses and Observations, for Years current before Christ, they might nearly agree, and yet an Error in Chronology might compensate for another in the Quantities of Acceleration; if this should happen to be the Case.

Mr. Ferguson, in his Astronomy, has mentioned the foregoing Paradox (without explaining it) by determining the 28th of May 585 Years current before Christ (when Peace ensued at the Eclipse, putting an End to the Battle betwixt the Medes and Lydians) to be on a Wednesday, as it happen'd, and not on a Monday, according to the vulgar Mistake of Chronologers. Who consider the 0 Year for the Beginning of the 1st Year current of Christ (beginning at 0) instead of considering this 0 Year, to be the 1st Year current before 1st Year of Christ, as it numerically denotes, when 100 is put for 100 Years current before Christ, that should be signified by 101, &c.

CHRONOLOGICAL SCALE.

Tabular Years bef. Chr.										Chr.†.	Tabular Years since Chr†.																					
400	300			200			100			0	100	200			300			400														
401	301			201			101			0	100	200			300			400														
Chronological Years before Chr†.											Chronological Years since Chr†.																					
Tabular less than chronological Yrs. before Chr†, by a Year.											Tabular Years before Chr†.										Chr.†.	Tabular Years since Chr†.										Tabular and chronological Years since Chr†, the same.
											8	7	6	5	4	3	2	1	0	1	2	3	4	5	6	7	8					
											9	8	7	6	5	4	3	2	1	0	1	2	3	4	5	6	7	8				
Chronological Years before Chr†.											Chronological Years since Chr†.																					

From above it is evident that 1 Year less must be used than the Chronological Year current, before Christ, to correspond with the Tabular Year, in seeking the true Answer from Astronomical or Chronological Tables, or from Arithmetical Operations: Being a Correction in Astronomy and Chronology hitherto unknown, or unobserved.

From above it is evident, that the same chronological with the tabular Year since Christ must be used, in seeking the true Answer from astronomical or chronological Tables, or from arithmetical Operations.

Since therefore the Quantities 0, 100, 200, &c.

Numerically denote the 1, 101, 201, &c. Years chronologically, or current, before Christ, in our chronological Tables, Therefore,

RULE II. Deduct 1 from the chronological Date, or Number of the Year current before Christ; then, with the Year remaining, 0, or N. S. find the Dominical Letter, Golden Number, or any other Result, from our Tables, for which they are fitted to determine; and likewise use this remaining Date, in arithmetical Operations concerning Chronology. But never subtract or add any Thing to the Dates, in the Use of our Tables, or any arithmetical Operations about Chronology, for Years current since Christ.

Of the LEAP-YEAR, or YEAR since.

TO find, in any Year before Christ, if it was Leap-Year, or what Year since, according to N. S.

RULE III. Divide Hundreds remaining* by 4, or Years over, What remains take from 4, will the sought-Year discover.

* Even Hundreds after 1 subtracted from the Date.

† 4. 1. 2. 3. Years since Leap-Year.

Yrs.	Yrs.
Examples. 1501 bef. Chr.	1462 before Christ,
— 1	— 1
1500 . . 4) 15(3	4) 61(15
— 3	— 1
4	4
Rem. 1 Year	Rem. 3 Yr. since Leap-Year,
[since Leap-Year, N. S.]	[N. S.]
When 0 remains it is Leap-Year.	

TO find, in any Year before Christ, if it was Leap-Year, or what Year since, according to O. S.

RULE IV. Dividing by 4, after 1 from the Date, The Remainder from 4, shews from Leap-Year how late.*

* 4. 1. 2. 3. Years after Leap-Year.

Yrs.	Yrs.
Examples. 1701 before Christ.	756 before Christ.
— 1	— 1
4) 1700(425	4) 755(188
10	35
20	35
— 0 Leap-Year.	— 3
4	4
Rem. 4 since Lp-Yr. O. S.	Rem. 1 Yr. since Lp-Yr,
[being Leap-Year.]	[O. S.]

TO find, in any Year since Christ, if it be Leap-Year or what Year since, according to N. S.

RULE V. By 4 divide Hundreds, or Years more, and then The Remainder will shew if 'tis Leap-Year, or when.*

* 0. 1. 2. 3. Years since Leap-Year.

Yrs.	Yrs.
Examples. 1600 since Christ.	1775 since Christ.
4) 16(4	4) 75(18
	35
0 Leap-Year, N. S.	3 Yr. since Leap-
	[Year, N. S.]

CHRONOLOGICAL RULES.

TO find, in any Year since Christ, if it be Leap-Year, or what Year since, according to O. S. ?

RULE VI. If the Date of the Year you by 4 should divide, What remains will the Number* since Leap-Year decide.

*O. 1. 2. 3. Years since Leap-Year.

Yrs. since Chr. Yrs. since Chr.
Examples. 4)1751(437 4)1800(45
15 20
31
Rem. 3 Year since Leap- [O. S.
Year, O. S.

Of the DOMINICAL LETTER.

TO find the Dominical Letter for any Year before Christ, according to N. S. ?

RULE VII. Hundreds Left* quote by 4, and the Letter's Place had, By what's left short of 4, doubled; if 1 you add: Then add once and the fourth of odd Years, made 1 less, Quoting Sevens: What remains will the Letter express.

* Even Hundreds after 1 subtracted from the Date.
Example. To find the Dominical Letter for 1535 Years current before Christ according to N. S. ?

1501 35 odd Years.
— 1 — 1
15|00 . . 4)15(3 1/4 . . 34 made 1 less,
— 3 — 8
4 Rem. 42
1 left short of 4 . . 1 3 for 1501.
Doubled 2 7)45[6
Add 1 3 . . C, for 1535, N. S.
3 Letter's Place A B C D E F G.
[C, for 1501 before Christ, N. S. 1. 2. 3. 4. 5. 6. 7.

N. B. In Leap-Years before Christ this Rule gives the 1st Dominical Letter of the Two.

TO find the Dominical Letter for any Year before Christ, O. S. ?

RULE VIII. The Year Left* its Fourth and 4 divide by 7, From what remains you'll find the Letter given, * After 1 subtracted.

Yrs. Yrs.
Examples. 3740 before Chr. 2325 before Christ,
— 1 — 1
1/4 . . 3739 1/4 . . 2324
934 581
4 4
7)4677(668 7)2909(415
47 10
57 39
1 . . A reqd. O. S. 4 . . D reqd. O. S.

N. B. When it is Leap-Year this Rule finds the 1st of the 2 Dominical Letters.

TO find the Dominical Letter for any Year since Christ, N. S. ?

RULE IX. Divide Hundreds by 4, and to twice what remains, One added, the Letter for that Year explains; Take from which (or 7 added) what is Left, after 7, Divides the Years since, and their Fourth — And That's given.*

N. B. When 0 remains for the Letter's Place, after Subtraction, the Letter is G, and shews that 7 also remains.

*A. B. C. D. E. F. G.

1. 2. 3. 4. 5. 6. 7.

Ex. For 17|59 . 4)17(4 1/4 . . 59 above Hundreds.

1 Rem. 14
Doubled 2 7)73(1
Add 1 7
Sum 3 C the Dom. Let. for 1700. 3 Rem. to be sub-
— 3 [since Chr. N. S. [tracted.
O . . or G, Dominical Letter for 1759 since [Christ, N. S.

N. B. When it is Leap-Year this Rule finds the last of the two Dominical Letters.

TO find the Dominical Letter for any Year since Christ, O. S. ?

RULE X. The Year, its Fourth, and 4, by 7 divided, What's Left deduct from 7, the Letter is decided.*

*A. B. C. D. E. F. G.

1. 2. 3. 4. 5. 6. 7.

Yrs. Yrs.
Examples. 1/4 . . 1759 since Christ. 1/4 . . 1785 since Christ.
439 446
4 4
7)2202(314 7)2235(319
10 13
32 65
— 4 Rem. — 2
7 7
Rem. 3 Letter C reqd. Rem. 5 Letter E reqd.

N. B. When it is Leap-Year this Rule finds the last Dominical Letter of the Two.

Of the WEEK-DAY, FOREVER, O. or N. S.

TO find, by the Dominical Letter, O. or N. S. for any Year, on what Day of the Week any Day of the Month happened, or shall happen, in that Year ?

George Brown

AT DUBLIN DWELLS GREGORY BLAND ESQUIRE, GEN'ROUS CHARLES FREEMAN, AND DANIEL FLYER.

USE. In the Roman Kalendar, the first Day of each Month is denoted as follows:

A. 1.	D. 1.	D. 1.	G. 1.	B. 1.	E. 1.
January.	February.	March.	April.	May.	June.
G. 1.	C. 1.	F. 1.	A. 1.	D. 1.	F. 1.
July.	August.	September.	October.	November.	December.

Correspondent to the Letter beginning each Word of the foregoing Memorable Couplet.

EXAMPLE. To find the Week-Day answering to May 24, 1759, since Christ, N. S. ?

The Dominical or Sunday-Letter for the Year is G.

Consequently A denotes Monday; B, Tuesday; C, Wednesday; D, Thursday; E, Friday; F, Saturday.

The 1st of May being always denoted by B, now on a Tuesday, therefore (continually adding 7) the 8th, 15th, and 22d Days are all Tuesdays, and consequently, the 24th Day is on a Thursday, required. The same Method is to be observed for finding the Week-Day on any Month of the Year (by Memory) which the Chronological Tables, p. 149. exhibit at Sight; serving (like the Memorial Rule above) for any Year before or since Christ, either according to New or Old Style.

TO find the Golden Number, or Prime, before Christ, O. or N. S.

RULE. Deduct 2 from the Date, and divide by 19, What remains take therefrom and the Prime will be seen.

right

CHRONOLOGICAL RULES.

Yrs.
Example. 1611 before Christ.

$$\begin{array}{r} 1611 \\ - 2 \\ \hline 1609 \\ 19 \overline{) 1609} \\ \underline{152} \\ 89 \\ \underline{76} \\ 13 \\ \underline{19} \end{array}$$

Compare this Example with that p. 150, where I should have been deducted from the Year for that Operation.

Being a new Improvement lately made in Chronology, since the former Rule was printed.

Or, universally, bef. Chr.

If the Date and 17, you by 19 divide, What remains from 19, will the Prime then decide. † Comp. of 2 to 19.

G. No. or Prime 6 required.

N. B. When the Year before Christ is less than 2, deduct it therefrom, and the Remainder is the Golden Number for that Year, by 1 Rule.

TO find the Cycle of the Sun for any Year before Christ?
FROM the Date, before Christ, deducting 10 Years,
28 next divides—Take from that what is Left, and the Cycle appears.

Yrs.
Example. 397 before Christ.

$$\begin{array}{r} 397 \\ - 10 \\ \hline 387 \\ 28 \overline{) 387} \\ \underline{28} \\ 107 \\ \underline{84} \end{array}$$

N. B. When the Number of the Year before Christ is less than 10, deduct it therefrom, and the Remainder is the Cycle for that Year.

Or, universally, bef. Chr.

Divide by 4 Sevens the Date and 18, Take what's left from 4 Sevens, the Sun's Cycle is seen.

Cycle of ☉ . . 5 required.

† Comp. of 10 to 28.

TO find the Roman Indiction for Years before Christ?
From the Date deduct 4, then divide by Fifteen,
What remains take therefrom, and the Cycle is seen.

Yrs.
789 before Christ

$$\begin{array}{r} 789 \\ - 4 \\ \hline 785 \\ 15 \overline{) 785} \\ \underline{75} \end{array}$$

N. B. When the Number of the Year before Christ is less than 4, deduct it from 4, and the Remainder is the Indiction for that Year.

Or, universally, bef. Chr.

Divide by 15, the Date and 11, Take what's left from 15, the Indiction is given. † Comp. of 4 to 15.

10 Cycle of Indiction, required.

As 1, 9, 3, the Golden Number, Cycle of the Sun, and Roman Indiction, for the Year before Christ's Coming, are added to any Year since Christ, before a Division by 19, 28, & 15 respectively, for determining the respective Cycles for that Year; so likewise 2, 10, 4, the Golden Number, Cycle of the Sun, and Roman Indiction, for the 1 Year of Christ, or in the Year since his Coming are to be subtracted from or added to any Year before Christ, the contrary Way on the Scale of Chronology, before a Division by 19, 28, 15, be made for determining the respective Cycles for that Year. — As herein many Chronologers have been mistaken in their Computations, by subtracting or adding 1, 9, 3 instead of 2, 10, and 4, from or to the Year before Christ, thereby giving their Results for a Year 1 less than the True, on the backward Scale of Chronology.

TO find the Year of the Dionysian, or any other, Period, for any Year before Christ, according to the foregoing Improvement?

GENERAL RULE. Subtract (if subtractable) the Number of any Cycle, in the 1st Year of Christ, from the Date of the Year before Christ, and divide the Remainder by the Number of the whole Cycle, what now remains deduct from the Number of the said whole Cycle, and the last Remainder will be the Year of the Cycle, required.

N. B. When the Date of the Year before Christ is less than the Number of the Period, in the 1 Year of Christ, deduct the Date therefrom, and the Remainder will be the Number of the Cycle, required, for that Year.

Yrs.

Example. For 1000 before Christ.

Subtract 458 the No. of Dion. Per. } it being 457
in 1st Yr. Chr. } for the Year before Chr.

$$\begin{array}{r} 1000 \\ - 458 \\ \hline 542 \end{array}$$

Or, universally, instead of subtracting the Period for 1st Yr. Chr. from the Date, add its Comp. to a whole Period.

522 (not 521, See p. 160.) required.

For any Year since Christ there is no Correction to former Rules; our Improvement in Chronology only taking Place for Years before Christ.

TO find the Year before Christ answerable to any Year of any Period, according to the foregoing Improvement?

GENERAL RULE, the Reverse of the former General Rule.

Deduct the Number of the Period from the whole Period, and add the Remainder to 458, (the Number in the 1st Year since Christ) and the Sum will be the least Date before Christ, when the Number of the Period happened as given; to which Date continually adding the Number of the whole Period, and you have all the Dates when the Number of Cycle or Period so happened.

Whole Period 532

Ex. The Reverse of the Former. The No. of given Per. 522

$$\begin{array}{r} 532 \\ - 522 \\ \hline 10 \end{array}$$

Add 458

The least DATE, before Christ, when the No. of the Period so happened } 468 bef. Chr.

+ 532

Also 1000 bef. Chr.

+ 532

Also 1532, &c.

TO find the Number of the Julian Period, according to the foregoing Correction of Chronology, for any Year before Christ?

Yrs.

Ex. For 585 before Christ.

4714 No. of the Per. in 1st Chr. greater than the Yrs. bef.

[Chr. Therefore the Date deducts from it.

Dif. 4129 Julian Period, reqd. according to the General Rule.

TO find the Number of Years before Christ, answering to any Julian Period for that Year?

Ex. Suppose 5320 Julian Period before Christ.

7980 Number of whole Period.

$$\begin{array}{r} 5320 \\ - 7980 \\ \hline 2660 \end{array}$$

+ 4714

Least Date, in the Yr. 7374 before Christ.

The contrary. Yrs.

By General Rule. 7374 before Christ.

$$\begin{array}{r} 7374 \\ - 4714 \\ \hline 2660 \end{array}$$

Rem. — 2660 will not div. by wh. Period.

Therefore 7980

5320 N. S. Julian Period, required.

The Chronological Rules (before these Corrections) take Place by 1st subtracting 1 from the Date of the Year before Christ, or adding 1 to the Results, to agree with Truth.

CHRONOLOGICAL RULES.

CYCLE of the SUN and DOMINICAL LETTER TABLE, O. S.

1	GF	5	BA	9	DC	13	FE	17	AG	21	CB	25	ED
2	E	6	G	10	B	14	D	18	F	22	A	26	C
3	D	7	F	11	A	15	C	19	E	23	G	27	B
4	C	8	E	12	G	16	B	20	D	24	F	28	A

To find the Dominical Letter for any Year, O. or N. S. arithmetically?

Find it correspondent to the Sun's Cycle, in the annexed Table for O. S. Then reckon the Number of Days between O. and N. S. or the Number of Letters Places forward, that the Difference of Days from O. to N. S. is above 7, 14, 21, &c. the Products of 7.

Example. To find the Dominical Letter for 1759, N. S. ?

Yrs.

1759 since Christ.

+ 9

28)1768(63 Number of Cycles completed.

Rem 4, the Number of the present Cycle @. . . Hence Dom. Letter correspondent by the above Table is C, for O. S. To which adding the 4th Letter's Place forward (the Difference between 11 and 7) G is the Dominical Letter, required, for N. S.

A General Method to find the Epact for O. and N. S. arithmetically?

The Epact is the Difference of Days between the solar Year of 365 and lunar of 364 Days.

For the mean Solar Year = 365 5 48. 54
The mean Lunar Year = 364 8 48 36

The Use of the Epact is to find the Moon's Age nearly.

Dif. Days 10 21 0 18
So that the whole Days are made up in the Reckoning.

To find the Epact for any Year O. S.

RULE. Multiply the Golden Number for the Year by 11, (for the Periods of Hundred Years from 16 to 18,00) divide the Product by 30, the Remainder will be the Epact for that Year, O. S. — Which Epact is reduced to That for N. S. by subtracting from it (or 30 added) the Days from O. to N. S. : The Remainder being the Epact answerable to the New.

Yrs.

Ex. For 1759 since Cbr. O. S. The G. No. is 12

11

12

12

3)0)13)2(4

Rem. 12 Epact O. S.

Deduct 11 Ds. Dif. fr. O. [to N. S.]

Rem. 1 Epact N. S. 1759.

only for this period to 1800
1769 this is not general as the result of the Cy must be ded.

To find the Moon's Age, for any Day of the Month in any Year, for which the Epact is given?

You reckon the Number of the Month from March, which is called 1, April 2, May 3, &c. to which Number, you add the Day of the Month, and the No. of the Epact, rejecting 30, and the Moon's Age remains, within a Day. Or you may use the following Memorial Rule.

Jan. 0, Feb. 2, Mar. 1, Ap. 2, May 3.
Ju. 4, Jul. 5, Au. 6, Sep. 8, agree;
Oc. 8, No. 10, De. 10, — to Epact add—
With Day — bate 30 — Age or Change is had.

Example. For July 25th, 1759.

CHRONOLOGICAL RULES.

MEMORIAL RULE.

To the Epact add 6, and reject 3 Times three,
What remains take from 50, your Limit you'll see,
If 50 or 49's Left—when you've done,
And the Prime above 11—Take, for each, left by 1.

Example. for 1759. . . Epact 1 . . G. No. 12.
Add 6

No 30 to reject . . 7—
50

The Limit, . . Days 43 from March 1, inclusive.
March 31—
April 12.

To find Easter Day, or the Number of Days it falls from the first of March, inclusive, arithmetically, N. S.?

MEMORIAL RULE.

Take the Letter and 4, from the Limit Day's Fall,
The Remain from next Sevens—Add the Limit.—That's all.
Example for 1759. Limit . . Days 43 . . G, Dom. Let. + 7
Sub. 11— Add . . 4

Rem. 32—
Next Sevens 35

Rem. . . 3
Add 12 April, the Limit.

15 April, Easter, 1759,
[N. S. required,

A TABLE for readily finding
the YEAR of the Dionysian
and Julian PERIODS, cor-
respondent to given Cycles of
the Sun, Moon, and Indiction.

No. of D, or Indictn Cycles.	Solar Years.	Lunar Years.	Indic- tion Years.
1	57	476	6916
2	114	420	5852
3	171	364	4788
4	228	308	3724
5	285	252	2660
6	342	196	1596
7	399	140	532
8	456	84	7448
9	513	28	6384
10	570	504	5320
11	627	448	4256
12	684	392	3192
13	741	336	2128
14	798	280	1064
15	855	224	7980
16	912	168	
17	437	112	Conf.
18	494	56	For 2d,
19	19	532	3d, 4th,
20	76		Col.
21	133		multi-
22	190		ply 57, 476 and
23	247		6916 respect-
24	304		ively, by the
25	361		No. of each Cy-
26	418		cle in 1st Col.
27	475		taking the Re-
28	532		mainders after

Products by 532, for 2d and 3d
Cols. but after Division by 7980
for 4th Column.

N. B. 57 is the Yr. Dion.
Period, when solar Cycle was
1, and lunar Cycle 19. And
476, Yr. Dion. Period, when
solar Cycle was 28, and lunar
Cycle 1. Also 6916 is Yr. Ju-
lian Period when 1 was the In-
diction, and 19 the lunar Cycle,
as may be proved.

Ex. To find the Year of the
Dionysian Period answering to 3
Gold. No. and 15 Cycle ☉?

Against 3, Golden Number, . . 364 Lunar Years.
15, Cycle of the Sun, . . 323 Solar Years.

Sum 687
Deduct 532 whole Period.

Rem. Dionysian Period 155 Yr. required.

Example. To find the Year of the Julian Period,
answering to 17, Cycle of the Sun; 12, Cycle of the
Moon; and 13, Cycle of Indiction?

Against 17, Cycle of the Sun . . 437 Solar Years.
12, Cycle of the Moon, . . 392 Lunar Years.

Sum 829
Deduct 532 whole Period.

Rem. Dionysian Period 297 ÷ 15.

Rem. 12—
Indiction 13

Rem. 1 . . Tab. No. 6916 4th Column.

Julian Period 7213 required.

Or, from above, 829 ÷ 15.

Rem. 4—
Indiction 13

Rem. 9 . . Tab. No. 6384

Julian Period 7213 as before.

So for the Rest.

FROM the 3 CYCLES of the SUN, MOON, and IN-
DICTION, given, to find the Year of the Dionysian and
Julian Period arithmetically?

See Rules, Page 160.

ANOTHER GENERAL RULE.

Subtract the Lunar from the Solar Cycle, (or from
28 added, if not otherwise subtractable) and multiply
the Remainder by 56: To this Product add the solar
Cycle, dividing the last Sum by 532 (if greater than
that Number) and the Remainder will be the Dionysian
Period answering to the Solar and Lunar Cycles.

From this Period found, subtract the Number of the
Indiction (or from 532 added if not otherwise subtract-
able) dividing the Remainder by 15; then multiply
what remains by 1064, deducting (if possible) 7980
from that Product, add the Year of the said Dionysian
Period to the Remainder, and the last Sum will be the
Number of the Julian Period, required.

Example. To find the Year of the Julian Period an-
swering to 17, Cycle of the Sun; 12, Cycle of the Moon;
and 13, Cycle of Indiction.

17 Solar Cycle.
—12 Lunar Cycle.

Rem. 5, X 56 = 280
Solar Cycle + 17

Dionysian Period 297
Indiction —13

÷ 15 . . 284

{ Rem. 14 }
X 1064 } 14896

Dionysian Period from above + 297

Sum, 15193
Subtracting 7980

Rem. the Julian Period 7213

USE of the TABLE to the LEFT.

Take out the Number of Solar and Lunar Years
from the 2d and 3d Columns answering to the Number
of the Solar and Lunar Cycles: and add together for
the Dionysian Period, if less than 532; but if greater,
take 532 therefrom, and the Dionysian Period will
remain. — Now divide this Period by 15, and take the
Remainder from the Indiction Cycle, (or from 15
added thereto) and with the last Remainder in the 1st
Column, take out the Number in the 4th Column cor-
respondent: the Sum of which, and the Dionysian Pe-
riod, will be the Julian Period.

Or divide the Sum of the Numbers in 2d and 3d
Columns by 15, and the Remainder taken from the
Indiction will give a Number in the 4th Column,
(answerable to that Remainder in the 1st Column),
which being added to the Sum of the Numbers in the
2d and 3d Cols. will be the Julian Period, required.

If epact
is 24 call it 25
and if
25 & 26
above 11
call it 26

If 50 is left
subt 1
and if 49
& 50 above
11 subt 1

CHRONOLOGICAL RULES.

The ANTIEN *Paschal*
TABLE for finding
EASTER, O. S.

Golden No.	Dom. Letter.	Month Day.
16 5	D	22
13 2	E F G	23 24 25
10 18 7 15	A B C D E F G	26 27 28 29 30 31 Apr. 1
4 12 1 9	A B C D E F G	2 3 4 5 6 7 8
17 6 14 5 11	A B C D E F G	9 10 11 12 13 14 15
19 8	A B C D E F G	16 17 18 19 20 21 22
	A B C	23 24 25

Ex. To find when Easter
happened 1593, when the
G. No. was 17, and Do-
minical Letter G?

Against 17, G. No. in
the 1st Column, stands A,
and against G, the Domi-
nical Let. following, stands
April 15, for the Time of
Easter, required.

N. B. The Day of the
Month against the Golden
Number, April 9, is the
Day of Easter Full Moon,
preceding Easter - Sunday.
If the Dominical Letter
stands against the Golden
Number, or the Easter
Moon be on a Sunday,
then Easter is on the Sun-
day following, for the
Dominical Letter must fol-
low the Golden Number.

So for the Rest.

To find the Letter in the Roman Calendar, correspondent to any Month-Day in the Year, before or since Christ, arith-
metically?

RULE. Divide the Number of Days, counting from the Beginning of the Year, or January 1, to the given
Month-day (see Tab. p. 129.) by 7, and what *remains will be the Number of the Letter, required.

Example. Required the Letter answerable to the 15th of November, in any common or Bissextile Years?

Com. Yr.
7)319(45
39

* Rem. 4 . . Letter D, required,
for a common Year.

Bif. Yr.
7)320(45
40

* Rem. 5 . . Letter E, required,
for a Bissextile Year.
A B C D E F G.
* 1. 2. 3. 4. 5. 6. 7.

RULES FOR FINDING THE WEEK-DAYS.

To find the Week-Day, arithmetically, answerable to a given Month-day, in any Year before Christ, O. S.?

RULE. To the Years, less 1, before Christ, add their Fourth, (being the Number of their Bissextiles) and 3,
and also the Complement of the Number of the Month-day, from Jan. 1. (from Tab. p. 129.) to next Sevens,
or (universally) to 371, divide the last Sum by 7, and deduct the Remainder from 7, and what last *remains will
be the Week-day, required. . . Sun. Mon. Tu. Wed. Th. Fr. Sat.

Example. To find the Week-day answering to May 28, 585 Years before Christ, O. S.

Yrs.
585
—1

From Jan. 1, to May 28, Biff. 149—
371

Comp. 222

584 } add.
146 }
3 }
222 }

Or
7)149(21

Rem. 2 Week-days, 28th May
is forward of Jan. 1.

Otherwise

584
146
3

7)955(136
25
45

7)733(104
7

4 . . Wednesday, O. S. required

2 Monday, Jan. 1, or the 31st of December, preceding Jan. 1.
+ 2 Week-days 28th of May is forward.

4 . . Wednesday, as before.

N. B. The Dominical Letter, before Christ, advances, (in Reckoning back, as the Dom. Let. Tab. shows, p. 148,
for Years before Christ) and consequently the Week-day, on Jan. 1, each Year goes back or retreats.
Hence the Reason of adding the Comp. of the Month-days to Sevens from January.

To find the Week-day for any Month-day and Year before Christ, N. S. arithmetically?

RULE. To the Years, less 1, before Christ, add their Fourth and 3, (but 2 from Feb. 25 to the End of Cen-
turies, not Lp. Yrs. N. S.) and also the Complement of the No. of the Month-day since Jan. 1. (fr. Tab. p. 129.)
to next Sevens, or (universally) to 371, and likewise the Complement of the Day's Difference between O. and N. S.
to next Sevens, then divide the Sum by 7, taking the Remainder from 7, and the last Remainder will shew the
Week-day, N. S. required.

Let the Week-day for the 28th of May, N. S. be required?

585
—1

From Jan. 1, to May 28, Biff. 149
Ds. 6 Dif. bet. O. and N. S. 371

584 } add.
146 }
3 }
222 }

Next 7

1 Comp.

Comp. 222

Otherwise.

4 Wednesday, O. S.

+ 6 Dif. Ds. fr. O. to N. S.

Rem. 4—
7

7)10(1

3 . . Tuesday, N. S. required

Rem. 3 . . Tuesday, N. S. as before. (Adding the Days Difference of Styles to the Week-Day of O. S. dividing
by 7, taking the Remainder for the Week-Day, N. S.)

CHRONOLOGICAL RULES.

N. B. The Week-day for corresponding Month-days of O. and N. S. being the same, when the Dif. of Month-days subtracts from O. for N. S. (before Christ) the Difference of Week-days from O. to N. S. adds or advances, as much; or, when (since Christ) the Difference of Month-days adds to O. for N. S. the Difference of Week-days subtracts or retreats as much; respecting the same Month-Day in both Styles.

To find the Week-Day arithmetically, answerable to a given Month-Day, in any Year since Christ, O. S. ?

RULE. To the complete Years since Christ (viz. 1 less than the Years current) add their Fourth, —1, and also the Number of the Days since the Beginning of the Year, or Jan. o. (from Tab. p. 129.) divide the Remainder by 7, and what last † remains, will be the Number of the Week-Day, required, O. S.

Sund.	Mon.	Tu.	Wed.	Th.	Fr.	Sat.
† 1.	2.	3.	4.	5.	6.	7.

Example. To find the Week-Day answering to May 15, 1759, current since Christ, O. S.

—1

1758

1, —1 . . . 438

Days 135 fr. Jan. o. to May 15, com. Year.

7)2331(333

23

21

† Rem. o. or 7. Saturd. O. S. required.

To find the Week-Day, arithmetically, answerable to a given Month-Day, in any Year since Christ, O. S. ?

RULE. To the complete Years since Christ (viz. 1 less than the Years current) add their Fourth, —1, and the Number of the Days since the Beginning of the Year, or Jan. o. (from Tab. p. 129.) and from the Sum of these subtract the Number of Days Difference between O. and N. Style, then divide the Remainder by 7, and the last Remainder will be the Number of the Week-Day, required, N. S.

Example. To find the Week-day answering to May 15, 1759, current since Christ, N. S.

—1

1758

1, —1 . . . 438

Days 135 from Jan. o. to May 15 com. Yr.

2331 Sum as before.

—11 Dif. bet. O. and N. S.

Otherwise, 7)2320(331

o Saturday, O. S. 22

—11 (adding Sevens to o.) 10

Rem. 3 Tuesday, N. S. *Rem. 3 . . Tuesday, N. S. required. [as before.]

N. B. The Dominical Letter, since Christ, retreats, (in reckoning forward, as the Dom. Letter Table shows, p. 148. for Years since Christ) and consequently the Week-day, on Jan. o. each Year, goes forward or advances. Hence the Reason of adding the Month-days from January, and not their Complement to Sevens, as before Christ.

UNIVERSAL and CORRECT RULES for finding the DOMINICAL LETTERS before CHRIST, according to N. and O. STYLE; deduced from the Principles of the DOMINICAL LETTER TABLES, Pages 148 and 149: being a further Illustration of Rules VII. & VIII. p. 171.

To find the Dominical Letter for any Year before Christ, O. S. arithmetically ?

RULE. To Years current less 1, add their Fourth, and twice 2, Quote by 7, and what's † Left will the Letter's Place shew.

A.	B.	C.	D.	E.	F.	G.
† 1.	2.	3.	4.	5.	6.	7.

O. S. before Christ,

N. B. Before Christ when it is Leap-Year the first Letter is shewn of the Two; and the Second Letter succeeds the Number of the Former.

Example. 1759 before Christ.

—1

1758

1, . . . 439

4

7)2201(314

10

31

Add Sevens . . 3 . . Dom. Let C, for O. S. reqd.

Subtract 15 . . No. Days Dif. from O. S. to N.

Rem. 2 Dom. Let. B. N. S.

To find the Dominical Letter for any Year before Christ, N. S. arithmetically ?

RULE. Divide Hundreds by 4, take what's left from twice 2, One with twice what remains will that Year's Letter shew; Add once and the Fourth, of the plus Years less one, Quote by 7, and the Letter by what's left is known.

Example for } 1759. } Hund. N. S. before Christ.

4)17(4

Rem. 1—

4

3, twice which +1=7, Letr. G for 1700.

58 plus years —1.

N. B. Bef. Christ, when it is Leap Year the first Letter is shewn of the two; and the 2d Letter follows the Number of the former.

14

7)79(11

Twice what rem. +1 making 9, for Leap- 2 rem. B. Dom. Let. year Cent^s, 7 deducts, and the Dom. Let. is [N. S. required. B, answering to 2.

When the Dominical Letter retreats (reckoning for Years forward from 200 since Christ) the Difference of Days between O. and N. Style, advances, or adds to the Month-days of Old, for the Month-days of New Style. And also adds to the Number of the Dominical Letter of the Old, for the Number of the Dominical Letter for the New Style. And when the Dominical Letter advances (reckoning for Years backward before Christ) the Difference of Days between Old and New Style, retreats, or subtracts from the Month-days of Old, for the Month-days of New Style, (from 200 Years since Christ.) And also subtracts from the Number of the Dominical Letter of the Old, for the Number of the Dominical Letter for the New Style, from 200 Years since Christ, when both Styles were exactly alike; as may be seen by Tab. p. 168.

UNIVERSAL PRACTICAL RULE for finding the DAY'S DIFFERENCE from OLD to NEW STYLE, before CHRIST for Years above Hundreds.

Before Christ, take the Fourth from the Hundreds—that done—Then add two: and the Difference of Style will be known.

N. B. The above Rule agrees with the Difference in the Dominical Letter Tables for Old and New Style before Christ, and with Table, p. 168. Or, if you add 1 to the Centuries, in Dates above Centuries, before Christ, the FIRST RULE (at Bottom of the Table, p. 168.) will serve, universally, for finding the Difference of Style for Dates above complete Centuries—It then amounting to no more than the Universal Rule last given: The said first Rule serving only for complete Centuries, that are not Leap Years, from the 25th of February to the End of those Centuries.

The Difference of Days (for the greatest Part of a Century) increased by 1, from Old to New Style, does not take Place till the 25th of February, after the Bissextile-Hundred is begun.

CHRONOLOGICAL RULES.

In Table, p. 168, the Day's Difference of Style between O. S. Leap-year, and those Centuries, not Leap-years, N. S. before Christ, take Place only for those compleat Centuries, and only from February 25, to the End of those Centuries, not Leap-years, N. S. respectively; and the Rest of the Numbers likewise take Place in the same Manner, for the same compleat Centuries. And therefore, any Dates above compleat Centuries before Christ will have the same Days Difference in the 2d Column between Old and New Style, and likewise the same Numbers in the 4th and 5th Columns, as at the preceding compleat Centuries, and so in Succession.

That, to find the Difference of Days between Old and New Style, Retreat of Lunations in Old Style, and Advance and Retreat of Lunations and Moon's Age, from Old to New Style, [in Col. 2, 4, & 5, from Tab. p. 168.] for Years above compleat Centuries before Christ.

You must take out the Numbers for 100 Years preceding, as you do for Years above compleat Centuries since Christ, which will then exactly agree with the foregoing PRACTICAL RULE, for finding the Difference of Styles for Years above compleat Hundreds before Christ.

TO reduce YEARS BEFORE, to YEARS SINCE, CHRIST, and the contrary; Having the SAME DOMINICAL LETTER, or LETTERS, and consequently, beginning with the SAME DAY OF THE WEEK, OLD and NEW STYLE.

Years before Christ reduced to Years since Christ, O. S.

RULE. Before Christ, take the Number of Years less by One, From Hundreds by 7, and the Year since is known.

Example. 585 Years current before Christ. $\begin{array}{r} 584 \\ -1 \\ \hline 584 \\ 1400 \dots 2 \text{ Hundreds by } 7. \end{array}$ *right*

Rem. 816 Years current since Christ, having the same Dominical Letters, [viz. FE for O. S.] and beginning with the same Week-day, [viz. Tuesday] as 585 Years current before Christ, O. S.

Years since Christ reduced to Years before Christ, O. S.

RULE. Years current since Christ take from Hundreds by Seven, Add 1 to what's Left; Years before Christ are given.

Example. 1759 Years current since Christ. $\begin{array}{r} 1759 \\ 2800 \dots 4 \text{ Hundreds by } 7. \\ \hline 1041 \\ +1 \\ \hline 1042 \end{array}$ *right*

1042 . . Years current before Christ, having the same Dominical Letter [viz. C, O. S.] and beginning with the same Week-day, [viz. Friday] as 1759 Years since Christ, O. S.

N. B. The RULES ABOVE serve for N. S. by using Hundreds by 4, instead of Hundreds by 7. Viz. for N. S.

Before Christ take the Number of Years less by 1, From Hundreds by 4, the Year since Christ is done.

Example. 585 Yrs. current bef. Chr.

Yrs. $\begin{array}{r} 585 \\ -1 \\ \hline 584 \\ 1600 \dots 4 \text{ Hundreds by } 4. \\ \hline 1016 \end{array}$ 1016 Years since Christ, Dom. Letters GF, Year begins on Monday, N. S.

$\begin{array}{r} 584 \\ 800 \\ \hline 216 \end{array}$

Since Christ the Years current from 4 Hundreds take, Adding 1 to what's Left, Years before Christ will make.

Yrs. $\begin{array}{r} 1759 \text{ since Christ,} \\ \text{Dominical Letter G,} \\ \text{Year begins on a} \\ \text{Monday, N. S.} \end{array}$

1759 Years current since Christ.
3200 . . 8 Hundreds by 4.

$\begin{array}{r} 1441 \\ +1 \\ \hline 1442 \end{array}$

1442 . . Yrs current bef. Christ, Dominical Letter G, Year begins on a Monday, N. S.

$\begin{array}{r} 1759 \\ 2000 \\ \hline 241 \\ 242 \end{array}$

In the same Manner, the Years since Christ, are found correspondent to the Years before Christ, having the same Golden Number, Cycle of the Sun, or Indiction; by subtracting 1 from current Years before Christ, and taking the Remainder from any Number of Hundreds by 19, 28, and 15, and the Years since Christ will, respectively, remain. And, on the contrary, taking the current Years since Christ, from any Number of Hundreds by 19, 28, and 15, adding 1, to each Remainder, and the Years current before Christ will result, having the same Golden Number, Cycle of the Sun, and Indiction, respectively.

Example I. Of the Golden Number 1612 before Christ.

Yrs. $\begin{array}{r} 1612 \\ -1 \\ \hline 1611 \end{array}$

1900 . . 1 Hundred by 19.

Years current since Christ 289

And 1612 before, and 289 Years since Christ, have the same Golden Number, viz. 5.

Example. II. Of the Golden Number 2189 since Christ.

3800 . . 2 Hundreds by 19.

$\begin{array}{r} 1611 \\ +1 \\ \hline 1612 \end{array}$

Yrs. 2189 since Christ, Golden Number 5. . . By Tab. p. 150. . . Golden Number 5.

The Examples for the Solar Cycle and Indiction are similar to the foregoing.

As the antient Egyptian, Arabic, Grecian, Persian, Jewish, and later Accounts of Time, are reduced, (for astronomical and chronological Purposes) to correspond with the Julian Form of the Year, or Account of Time, long before any such Year had Existence, so, in Conformity to the late Correction of the Julian Account of Time, we have reduced the Julian to the Gregorian Style, as far backward and forward as remotest Antiquity, and Futurity, for the Improvement of Chronology and Astronomy. This is performed at Sight, in two Pages, 148 and 149, foregoing, with the following Rules to those Pages.

For the COMMENCEMENT of the most famous AERAS (since Creation) and the REDUCTION of CHRONOLOGY, see further on.

A TABLE of KALENDS, IDES, and NONES, for the ROMAN YEAR.
Of USE in HISTORY and CHRONOLOGY.

Days	January.	February.	March.	April.	May	June.	July.	August.	Sept.	Octo.	November	December
1	Kal.	Kal.	Kal.	Kal.	Kal.	Kal.	Kal.	Kal.	Kal.	Kal.	Kal.	Kal.
2	4 No.	4 No.	6 No.	4	6 N.	4 N.	6 N.	4 N.	4 N.	6 N.	4 N.	4 N.
3	3 No.	3 No.	5 No.	3 No.	5	3	5	3 N.	5 N.	5	3 N.	3 N.
4	Pr. No.	Pr. No.	4 No.	Prid.	4	Prid.	4	Prid.	Prid.	4	Prid.	Prid.
5	Nonæ.	Nones.	3 No.	Non.	3 N.	Non.	3 N.	Non.	Non.	3 N.	Non.	Non.
6	8 Id.	8 Id.	Prid.	8 Id.	Prid.	8 Id.	Prid.	8 Id.	8 Id.	Prid.	8 Id.	8 Id.
7	7 January.	7 February.	Non.	7	Non.	7 June.	Non.	7 August.	7 Sept.	Non.	7 Nov.	7 December.
8	6	6	8 Id.	6 April.	8 Id.	6	8	6	6	8	6	6
9	5	5	7 March.	5	7 May.	5	7 July.	5	5	7 October.	5	5
10	4	4	6	4	6	4	6	4	4	6	4	4
11	3 Id.	3 Id.	5	3	5	3 Id.	5	3 Id.	3 Id.	5	3 Id.	3 Id.
12	Pr. Id.	Pr. Id.	4	Prid.	4	Prid.	4	Prid.	Prid.	4	Prid.	Prid.
13	Idus.	Idus.	3 Id.	Ides.	3 Id.	Idus.	3 Id.	Idus.	Idus.	3 Id.	Idus.	Idus.
14	19 K.	16 K.	Prid.	18 K.	Prid.	18 K.	Prid.	19 K.	18 K.	Prid.	18 K.	19
15	18	15	Idus.	17	Idus.	17	Idus.	18	17	Idus.	17	18
16	17	14	17 K.	16	17 K.	16	17 K.	17	16	17	16	17
17	16	13	16	15	16	15	16	16	15	16	15	16
18	15	12	15 April.	14	15	14	15	15	14	15	14	15
19	14	11	14	13	14	13	14	14	13	14	13	14
20	13	10	13	12	13	12	13	13	12	13	12	13
21	12	9	12	11	12	11	12	12	11	12	11	12
22	11	8	11	10	11	10	11	11	10	11	10	11
23	10	7	10	9	10	9	10	10	9	10	9	10
24	9	6	9	8	9	8	9	9	8	9	8	9
25	8	5	8	7	8	7	8	8	7	8	7	8
26	7	4	7	6	7	6	7	7	6	7	6	7
27	6	3 K.	6	5	6	5	6	6	5	6	5	6
28	5	Prid.	5	4	5	4	5	5	4	5	4	5
29	4	—	4	3 K.	4	3 K.	4	4	3 K.	4	3 K.	4
30	3 K.	—	3 K.	Prid.	3 K.	Prid.	3 K.	3 K.	Prid.	3 K.	Prid.	3 K.
31	Prid.	—	Prid.	—	Prid.	—	Prid.	Prid.	—	Prid.	—	Prid.

An ACCOUNT of the OLD and NEW Ecclesiastical and Civil Calendar. Of USE in HISTORY and CHRONOLOGY.

CALENDAR, or Account of the Year, borrows it's Name from *Kalend*, signifying the first Day of each Month, reckoning according to the Roman Account by the Moon. *Kalend* is derived from *καλέω*, *voco*, I call: For the *Priest*, whose Business it was to observe the New Moon, by which the first Day of each Month was reckoned, gave Notice thereof to the *President* of the Sacrifices, who called the People together, and ordained how the rest of the Days, in that Month, were to be set apart for religious and other Uses. Hence it was, that an Annual Account of divine and human Affairs, came to be kept throughout all the Months of the Year; which Account (from reckoning by the *Kalends*) was therefore called *The Calendar-Table*, and considered both as *Ecclesiastical* and *Civil*.

The Day preceding the *Kalends*, or first Day of each Month, was called *pridie Kalendas*, the preceding Day to this the 3d of the *Kalends*, the next preceding Day the 4th of the *Kalends*, &c. reckoning backwards to the *Ides* of the preceding Month.

The Day immediately following the *Kalends* of each Month was called the 4th or 6th of the *Nones*; always reckoning backwards in each Month from the first Day of *Nones*. In the Full Months of *March*, *May*, *July* and *October*, wherein are six of the *Nones*, the first of the *Nones* fell upon the 7th Day; but in all the Months called *Hollow* or *Cave*, wherein there are reckoned but four of the *Nones*; the first of those *Nones* fell upon the 5th Day. And there being eight of the *Ides* in each Month, throughout the Year, consequently the 15th Day, in the Full Months, and 13th in the *Hollow*, will be the Days of the *Ides*, from whence the rest are reckoned, like the *Kalends*, by inverted Order. The Day preceding the *Nones*, or *Ides*, is called *pridie Nonas*, or *pridie Idus*, the next preceding Days to these, the 2d, 3d, 4th, &c. of the *Nones*, or of the *Ides*, according to the following Observation of the Schools.

*Prima Dies Mensis cujusque est dicta Kalendæ,
Sex Nonas, Maius, October, Julius, et Mars;
Quatuor at Reliqui: dabit Idus quilibet octo,
Inde Dies reliquos omnes dic esse Kalendas,
Quas retro numerans, dies à Mense sequenti.*

This was the ancient Way of reckoning Time in the Latin Churches, and is used in the *Prayer-Books* of the Church of England at this Day.

The *Ecclesiastical* and *Civil Calendar*, or *Almanack*, also disposes of all the Days of the Year into *Months*, *Week-Days*, *Vigils*, *Fast-Days*, *Holy Days*, *Law-Days*, *University Terms*, &c. in a successive Order of reckoning; which being interrupted cannot constitute, literally, a *Calendar*.

The *Week-Days* are distinguished by seven Letters, *a, b, c, d, e, f, g*. The first Letter *a* is placed against the 1st of *January*, *b* against the 2d, *c* against the 3d, *d* against the 4th, and so on to *g* against the 7th, then *a* against 8th, *b* against 9th, and so on, in Succession of the seven Letters, through the Year: By which Means *a, b, c, d, e, f, g*, will always point out the same *Week Days* in any Year. And *a, d, g, b, e, f, c*, the 1st Day of each of the 12 Months, respectively.

The principal Letter of the seven, pointing out the *Sundays* for any Year, is always made *Capital*, and called the *Dominical Letter* for that Year.

Dividing 365 Days, in a common Year, by 7, the Quotient is 52 Weeks, and one Day over; if there had been no Remainder, all Years would begin and end on the same Day of the Week: But the odd Day occasions each Year to end on the Week-Day with which it began, whereby the next Year begins on the Week-Day following. And a Year of 366 Days divided by 7, having two Days remaining, the next Year begins two Week Days after the Week-Day on which the former Leap-Year begun: Whence, if a common Year begins on a *Sunday*, pointed out by the *Dominical Letter A*, it will also end on a *Sunday*, and the next, or second Year, will begin on a *Monday*, against which stands *a*; against *Tuesday b*; against *Wednesday c*; and so on to *Sunday*, against which stands *G*, the *Dominical Letter* for that Year. Against the Beginning of the third common Year, on a *Tuesday*, stands *a*, against *Wednesday b*, &c. to *F*, the *Dominical Letter* against *Sunday* for that Year.

And therefore E is the Dominical Letter for the fourth Year, beginning on a *Wednesday*, and so on, in a retrograde Order of Dominical Letters for each succeeding Year. But in this fourth Year, containing 366 Days, called *Bissextile*, or Leap-Year, the 24th Day of *February*, answering to the 6th of the Kalends of *March*, is twice reckoned, by placing the Letter *f* against the 24th and 25th of *February*, which interrupts the annual Retrogression of Dominical Letters, from E, the Sunday-Letter to 24th of *February*, to D, the Sunday-Letter following the 24th of *February*, to the Year's End. And the fifth Year (of 365 Days) begins two Week-Days after the Week-Day beginning the Leap-Year, viz. on a *Friday*, when the Dominical Letter is C; and so for the sixth Year it is B, the seventh A, and the eighth G, and F, and so on. Whereby it is observable, that, after seven Years, if each Year consisted but of 365 Days, the Days of the Month would again return on the same Days of the Week: But by the Interruption, every four Years, to each Letter, it is 7 Times 4, or 28 Years, before the Days of the Months return exactly on the same Days of the Week; which Term of Years is called the Cycle of the Sun. But, in *New Style*, this Order is now farther interrupted, by a Day less reckoned, in every odd Hundredth Year, being now made common, instead of *Bissextile*, as in *Old Style*.

The Cycle of the Sun-Table, for any Year of the Cycle, of Use in History and Chronology, O. S.

1	GF	5	BA	9	DC	13	FE	17	AG	21	CB	25	ED
2	E	6	G	10	B	14	D	18	F	22	A	26	C
3	D	7	F	11	A	15	C	19	E	23	G	27	B
4	C	8	E	12	G	16	B	20	D	24	F	28	A

This Table and the following Rule, are here repeated for Use.

N. B. In the first Year of Christ, the Year of the Cycle being 10, therefore adding 9 to the Year since Christ, and dividing the Sum by 28, the Remainder will be the Year of the present Cycle as above, to which the Dominical Letters are determined. See Pages foregoing for finding the Dominical Letters O. S. arithmetically.

The Fixed Feasts in O. S. were regulated by the Sun's Motion, but the Moveable ones were governed by the Moon's Motion, as the Fixed and Moveable Feasts N. S. are now governed; especially *Easter*, on which the rest depend. God ordained that the Feast of the Jewish *Pasover*, should be kept on the 14th Day of the first Lunar Month at Even (see *Levit. Chap. xiii.*) which was at the Time of the first Full Moon in the first Month. But the Christian Church ordained the Celebration of *Easter* on the Sunday following; because our Saviour rose upon that Day. The Primitive Fathers had no Way of determining the Time of the New-Moon, but by observing it, emerging out of the Sun's Rays at *Even*, which was a Day after Conjunction. The Council of *Nice*, Anno 325, settled the Fall of *Easter* by the Metonic Cycle of Moons, (so called from it's Inventor *Meton*) as in the Ecclesiastical *Kalendar*; inserting the Order of their happening there. They supposed that, in every Succession of 19 Julian Years (the Period they allotted to the Lunar-Cycle) the Moons would return on the same Days of the Months, and Hours of the Day, as in the Order of the Golden Number, pointing out the New-Moons, in the Bible-Calendar: Which, besides Anticipation, is incompatible, by the odd Hours not reckoned for the last 3 common Years. Those Ecclesiastical New-Moons (reckoned from and to Leap-Year) were about five Days later than the Time of the true ones, according to O. S. before it's Alteration to the New. Hereby *Easter* came not to be observed according to the Time of it's original Institution. [The next Sunday after the first Full-Moon happening next after the 21st of *March*, or *Vernal Equinox*, tho' at the Time of the *Nicene* Council happening on the 20th, but before the *Stile* was altered and the New *Style* took Place, in 1752, on the 9th of *March*.] If the Celebration of *Easter* had been ordered on the next Sunday after the first true Full-Moon happening next after the Time of the *Vernal Equinox*, it would always have been observed according to it's first Institution.

The Anticipation of the *Vernal Equinox* O. S. (and of the Seasons or any Points of the *Ecliptic*) was 44^m 9^s 21^h every four Julian Years, by reckoning so much more Time than was contained in four Solar Years (each Year, according to *Sir Isaac Newton*, being about 365^d 5^h 48^m 57^s 39th) by which in 33 Leap Years to come, the Time of the *Vernal Equinox* would be 1^d 0^h 17^m 8^s sooner. And, in 356 Leap-Years backward from 1748, since the Leap-Year 324 (preceding the Year 325, the Time of the *Nicene* Council) the Anticipation was 10^d 21^h 59^m 32^s. The *Equinox*, Anno 324, being on *March* 19^d 15^h 29^m, and consequently Anno 325 on *March* 19th, 21^h 18^m. Also, in the Year before Christ, being *Bissextile*, the *Vernal Equinox* was *March* 22^d 3^h 6^m P. M. and therefore Anno Christi 1, it was 22^d 8^h 55^m, nearly agreeing with the Time by the *Caroline* Tables of *March* 22, 7^h 22^m. Three Years after each Leap-Year, the *Equinox* and Seasons went forward, at the Rate of 5^h 48^m 57^s 39th, &c. each Year; the Year of 365 Days, under-reckoning the solar-Year till every fourth Year of 366 Days, which over-reckon'd what Time was lost, by 44^m 9^s, &c. and then brought back, or retrograded the Seasons again.

The Ancients supposed that 235 Lunations exactly completed the Lunar Cycle of 19 Years: But 235 Lunations, allowing 29^d 12^h 44^m 3^s 39th, &c. to each, are equal to 6939^d 16^h 34^m 17^s. And 19 Julian Years, of 365^d 6^h each, are equal to 6939^d 18^h. Therefore the Difference, 1^h 25^m 43^s, was the Lunar-Anticipation in 19 Julian Years: Four Times that Time, 5^h 42^m 52^s, was the proper Lunar-Anticipation for 19 Leap-Years. For it is to be observed, that every 1, 2, 3 common Years after each Leap-Year, the Mean New-Moons anticipated or happened sooner by 10^d 15^h 11^m, 21^d 6^h 22^m, 2^d 8^h 42^m, respectively, than the New-Moons in the same Months in the Leap-Year preceding: The 6 Hours above 365 Days in each Julian Year being counted but once in 4 Years. The Mean New-Moons anticipated 14^d 0^h 1^m in every four Julian Years; therefore in four Leap-Years, or 16 Julian, the Lunar-Anticipation was 56^d 0^h 4^m from whence deducting a Lunation=29^d 12^h 44^m, there remain 26^d 11^h 20^m, the nearest Anticipation; to which adding the third common Year's Anticipation after Leap-Year, viz. 2^d 8^h 49^m, the Sum is 28^d 20^h 9^m, Anticipation at the End of four Leap and three common Years: The Mean New-Moons then happening so much sooner in the Months, than in the preceding 19th Year.

Say 1^h 25^m 43^s: 19th: 24^h. 319 Julian Years, or 79 Leap, three common, and 18 Hours, when the Lunar-Anticipation is a Day; consequently, in 1424 Julian, or 356 Leap-Years, since the Leap-Year, preceding the *Nicene*-Council to 1748, the Lunar-Anticipation was about four Days, eleven Hours; if to which one Day be added, the Time when the New-Moons were observed, later than they really happened, at the *Nicene*-Council, will make almost five Days and a half difference; which accounts for the Difference between the Ecclesiastical and true Moons, before the Alteration of *Stile*, and Correction of the Calendar. N. B. The greatest Difference of Time between a Mean and True Conjunction, or Opposition, is, about 14 Hours.

In the *Victorian*, or *Dyonisian*-Period of 532 Years, the New and Full-Moons were supposed by the Antients to return on the same Days of the Month, and also on the same Week-Days, which Error has been shewn; there being a Day Lunar-Anticipation in that Time: Yet the Period is still retained for it's Use in History and Chronology.

Sir Isaac Newton, in his Account of the Birth and Passion of Christ, printed Anno 1733, observes that the Primitive Christians had little Regard to the Times of the Birth and Passion of our Saviour, as being not material to Religion; though there since has been such a Strictness observed in the Keeping the Times of the Fasts and Festivals. They were first celebrated on the Cardinal Points of the Year. The Annunciation on 25th of *March*, the Time of the *Vernal Equinox*, when *Julius Caesar* corrected the Calendar. The Feast of St. John on June 24th, the Time of the Summer Solstice. St. Michael on September 29, the Time of the Autumnal Equinox; and the Birth of Christ on December 25, then the Day of the Winter Solstice. The Feast of St. Thomas on December 20; St. Matthias on the 21st of September; and the other Saints on the Sun's Entrance into the different Signs; as may be read.

He fixes the first *Pasover* in the 16th Year of *Tiberius*, in the 4728th Year of the Julian-Period, between the Baptism of Christ, and Imprisonment of John. The second, four Months after St. John's Imprisonment, and Christ's beginning to preach in *Galilee*. The third *Pasover* was the Feast following that when the Corn was eared and ripe. And the Fifth, at the Time in which Christ suffered. He thence concludes that the Time of Crucifixion to be in the 34th Year from the Incarnation, or in 33d Year from the Time of the Birth.

PERPETUAL TIME - TABLES: OR

General RULES for CHRONOLOGISTS, HISTORIANS, MEMORANDUM-BOOK-MAKERS, TRAVELLERS, and NAVIGATORS. Determining, *at Sight*, the POINTS of TIME, for any one *Season* or *YEAR*, *past*, *present*, or *to come*. Particularly useful for all Persons writing ALMANACS for the STATIONERS COMPANY.

WHICH TABLES (of themselves) are no more a CALENDAR, or ALMANAC, serving for any one *Season*, or *Time*, than the Timber cut out in a *Dock-Yard* for Building, but not united into Form, is a *SHIP*; or no more so than Timber framed for Building, is a *House*; or than RULES for making ALMANACS is an ALMANAC. And therefore these TABLES or RULES are not subject to *Stamp-duty*; since all *Rules of Art*, for making any *Performance*, are allowable, without Restraint: For, without Rules to make them, there could be no *Almanacs*.

This *Truth* we think ourselves obliged to explain, as the *Stamp-Office* has a Right, by Act of Parliament, to charge a Duty on what is *literary* and *truly* an ALMANAC for any one Time, or Number of Times: or on *what chiefly* serves that Purpose, for any one Time, or Number of Times, to come: Since every printed Figure or Letter, or even the Paper of any Book of Chronology, and especially all *chronological Rules*, may partly serve that Purpose. In which Sense of *chiefly* serving that Purpose of an ALMANAC, for any one Time, or particular Number of Times, for the Year or Season, or Years or Seasons, present, or to come, the *Act* takes Place, and no further.

But, our Tables consist of general RULES, not serving *chiefly*, or made for any one Time, or Number of Times, or Years, present or to come, but *generally* (and not *chiefly*) serve, for all *past* and *future* TIME, and for no Year or Season, or Years and Seasons, in particular: And of which Time, in general, as *infinite Duration*, no particular Year or Years, Season or Seasons, can be understood. So that our *Perpetual Tables* are only subser- vient (like all other *general Rules*) to the *general*, and therein particular, Uses for which they were wanted and designed.

For Time, though in *Eternity* applied
To Motion, measures all Things durable
By present, past, and future. —

MILTON b. V. l. 580.

RULES for CHRONOLOGISTS, HISTORIANS, &c.

RULES for finding the Moveable FEASTS and HOLIDAYS.

EASTER, on which the Rest depends, is always the 1st Sunday after the first F. Moon, happening upon, or next after, March 21; and if the F. M. is on a Sunday, Easter is the Sunday following.

Septuagesima S.	9	} be- fore Ea- ster.	Rogation-Sunday	5 Wks	} af- ter Ea- ster.
Sexagesima	8		Ascension-day	40 Ds	
Quinquagesima	7		Whitsunday	7 Wks	
Quadragesima	5		Trinity Sunday	8 Wks	

ADVENT SUNDAY the nearest Sund. to the Feast of St Andrew.

All the moveable and fixed FEASTS observed in the Church of England.

All Sundays	St. Barnabas	St. Andrew Ap.
Circumcision	Nat. St. J. Bap.	St. Thomas Ap.
Epiphany	St. Peter Ap.	Nat. of Christ
Conv. St. Paul	St. James Ap.	St. Stephen Mar.
Purification V. M.	St. Barth. Ap.	St. John Evan.
St. Matthias	St. Matth. Ap.	Holy Innocents
Annun. V. Mary	St. Mic. & all An.	Mo. & Tu. in East.
St. Mark Evan.	St. Luke Ev.	Mo. & Tu. in Whit.
St. Ph. and Ja. Ap.	St. Sim. & Ju. Ap.	
Ascension J. Chr.	All Saints	

VIGILS and FASTS.

Eve	Nat. Chr.	Pentecost	St. James	St. Andrew
or	Purif.	St. Matthias	St. Bartholom.	St. Thomas
Vigil	Easter Day	St. John Bap.	St. Matthew	All Saints.
before	Ascension	St. Peter	St. Sim. & Jude	

N. B. When a Feast-day falls on a Monday, the Vigil or Fast is kept on the Saturday-Eve preceding, and not on that of the Sunday, which is a Feast-day.

DAYS of FASTING and ABSTINENCE.

- I. Forty Days in Lent.
- II. The Ember Days, at the 1st Sunday in Lent
- 4 SEASONS, on the Wednesday, Feast of Pentecost
- Friday and Saturday after the September the 14th
- III. The 3 Rogation-days, Monday, Tuesday, Wednesday, before Christ's Ascension or Holy Thursday. December the 13th
- IV. All the Fridays in the Year, except Christmas-day.

Solemn Days kept, for which particular Church-Service is appointed.

- I. November 5, in Memory of the Popish Plot.
- II. January 30, in Memory of the Martyrdom of K. Charles I.
- III. May 29, in Memory of the Birth and Return of K. Charles II.

Hil. T. b. Jan. 23. ends Feb. 12	Michaelmas T. beg. Nov. 6, ends
1 Return Jan. 20	[Nov. 29]
2 Return 27	Nov. 3
3 Return Feb. 3	12
4 Return 9	18
Easter Term	25
1 Ret. 14 Days from	Trinity Term.
2 Ret. 21 Easter	1 Ret. Morrow of Hol. Tr.
3 Ret. 28 exclusive.	2 Ret. 7 Days Fr. Holy
4 Ret. 35	3 Ret. 14 Trinity,
5 Ret. on Morrow of Ascen.	4 Ret. 21 exclusive.

PERPETUAL

PERPETUAL TIME - TABLES:

Or General RULES for CHRONOLOGISTS, HISTORIANS, &c.

MOVEABLE FEASTS for ever, according to the Fall of
EASTER, N. S.

Easter Sunday.	Septua- gesima Sunday	Ash- Wednes.	Rogat. Sunday.	Ascen. Sunday.	Whit- Sunday.	Advent Sunday.
Mar. 22	Jan. 18	Feb. 4	Apr. 26	Apr. 30	May 10	Nov. 29
23	19	5	27	May 1	11	30
24	20	6	28	2	12	Dec. 1
25	21	7	29	3	13	2
26	22	8	30	4	14	3
27	23	9	May 1	5	15	Nov. 27
28	24	10	2	6	16	28
29	25	11	3	7	17	29
30	26	12	4	8	18	30
31	27	13	5	9	19	Dec. 1
Apr. 1	28	14	6	10	20	2
2	29	15	7	11	21	3
3	30	16	8	12	22	Nov. 27
4	31	17	9	13	23	28
5	Feb. 1	18	10	14	24	29
6	2	19	11	15	25	30
7	3	20	12	16	26	Dec. 1
8	4	21	13	17	27	2
9	5	22	14	18	28	3
10	6	23	15	19	29	Nov. 27
11	7	24	16	20	30	28
12	8	25	17	21	31	29
13	9	26	18	22	June 1	30
14	10	27	19	23	2	Dec. 1
15	11	28	20	24	3	2
16	12	Mar. 1	21	25	4	3
17	13	2	22	26	5	Nov. 27
18	14	3	23	27	6	28
19	15	4	24	28	7	29
20	16	5	25	29	8	30
21	17	6	26	30	9	Dec. 1
22	18	7	27	31	10	2
23	19	8	28	June 1	11	3
24	20	9	29	2	12	Nov. 27
25	21	10	30	3	13	28

N. B. When any Feasts fall in January or February in Leap-Year, such Feasts must be taken a Day later than by Table.

To find the Golden Number for any Year?

Add 1 to the Date, and divide the Sum by 19, what remains is the Golden Number required.

When 0 remains 19 is the Golden Number.

The above Rule is here repeated for Use.

MOVEABLE TERMS for ever, according
to the Fall of EASTER, N. S.

Easter Term begins.	Ends.	Trinity Term begins.	Ends.
Apr. 8	May 4	May 22	June 10
9	5	23	11
10	6	24	12
11	7	25	13
12	8	26	14
13	9	27	15
14	10	28	16
15	11	29	17
16	12	30	18
17	13	31	19
18	14	June 1	20
19	15	2	21
20	16	3	22
21	17	4	23
22	18	5	24
23	19	6	25
24	20	7	26
25	21	8	27
26	22	9	28
27	23	10	29
28	24	11	30
29	25	12	July 1
30	26	13	2
May 1	27	14	3
2	28	15	4
3	29	16	5
4	30	17	6
5	31	18	7
6	June 1	19	8
7	2	20	9
8	3	21	10
9	4	22	11
10	5	23	12
11	6	24	13
12	7	25	14

N. B. Easter - Term begins 17 Days, and ends 43 Days after E A S T E R. Trinity-Term begins 5 Days, and ends 24 Days after Trinity. Therefore Easter-Term continues 27 and Trinity 20, inclusive.

PERPETUAL

PERPETUAL TIME-TABLES:

Or General RULES for CHRONOLOGISTS, HISTORIANS, &c.

MOVEABLE FEASTS from 1759 to 1800, N. S.

Years of Christ.	Gol. N ^o .	Dom. Letter. N. S.	Dom. Letter. O. S.	Septua- gesima Sunday	Ash- Wednes.	Easter Sunday.	Rogat. Sunday.	Ascens. Day.	Whit- Sunday.	Advent Sunday.	E- pact.
1759	12	G	C	Feb. 11	Feb. 28	Apr. 15	May 20	May 24	Jun. 3	Nov. 2	1
B. 1760	13	FE	BA	3	20	6	11	15	May 25	Nov. 30	12
1761	14	D	G	Jan. 18	4	Mar. 22	Apr. 26	Apr. 30	10	29	23
1762	15	C	F	Feb. 7	24	Apr. 11	May 16	May 2	30	28	4
1763	16	B	E	Jan. 30	16	3	8	12	22	27	15
B. 1764	17	AG	DC	Feb. 19	Mar. 7	22	27	31	Jun. 10	Dec. 2	26
1765	18	F	B	3	Feb. 20	7	12	16	May 26	1	7
1766	19	E	A	Jan. 20	12	Mar. 30	4	8	18	Nov. 30	18
1767	1	D	G	Feb. 15	Mar. 4	Apr. 19	24	28	Jun. 7	29	0
B. 1768	2	CB	FE	Jan. 31	Feb. 17	3	8	12	May 22	27	11
1769	3	A	D	22	8	Mar. 20	Apr. 30	4	14	Dec. 3	22
1770	4	G	C	Feb. 11	28	Apr. 15	May 20	24	Jun. 3	2	3
1771	5	F	B	Jan. 27	13	Mar. 31	5	9	May 19	1	14
B. 1772	6	ED	AG	Feb. 16	Mar. 4	Apr. 19	24	28	Jun. 7	Nov. 29	25
1773	7	C	F	7	Feb. 24	11	16	20	May 30	28	6
1774	8	B	E	Jan. 30	16	3	8	12	22	27	17
1775	9	A	D	Feb. 12	Mar. 1	16	21	25	Jun. 4	Dec. 3	28
B. 1776	10	GF	CB	4	Feb. 21	7	12	16	May 26	1	9
1777	11	E	A	Jan. 20	12	Mar. 30	4	8	18	Nov. 30	20
1778	12	D	G	Feb. 15	Mar. 4	Apr. 19	24	28	Jun. 7	29	1
1779	13	C	F	Jan. 31	Feb. 17	4	9	13	May 23	28	12
B. 1780	14	BA	ED	23	9	Mar. 26	Apr. 30	4	14	Dec. 3	23
1781	15	G	C	Feb. 11	28	Apr. 15	May 20	24	Jun. 5	2	4
1782	16	F	B	Jan. 27	13	Mar. 31	5	9	May 19	1	15
1783	17	E	A	Feb. 16	Mar. 5	Apr. 20	25	29	Jun. 8	Nov. 30	26
B. 1784	18	DC	GF	8	Feb. 25	11	16	20	May 30	28	7
1785	19	B	E	Jan. 23	9	Mar. 27	1	5	15	27	18
1786	1	A	D	Feb. 12	Mar. 1	Apr. 16	21	25	Jun. 4	Dec. 3	0
1787	2	G	C	4	Feb. 21	8	13	17	May 27	2	11
B. 1788	3	FE	BA	Jan. 20	6	Mar. 23	Apr. 27	1	11	Nov. 30	22
1789	4	D	G	Feb. 8	25	Apr. 12	May 17	21	31	29	3
1790	5	C	F	Jan. 31	17	4	9	13	23	28	14
1791	6	B	E	Feb. 20	Mar. 9	24	29	June 2	Jun. 12	27	25
B. 1792	7	AG	DC	5	Feb. 22	8	13	May 17	May 27	Dec. 2	6
1793	8	F	B	Jan. 27	13	Mar. 31	5	9	19	1	17
1794	9	E	A	Feb. 16	Mar. 5	Apr. 20	25	29	Jun. 8	Nov. 30	28
1795	10	D	G	1	Feb. 18	5	10	14	May 24	29	9
B. 1796	11	CB	FE	Jan. 24	10	Mar. 27	1	5	15	27	20
1797	12	A	D	Feb. 12	Mar. 1	Apr. 16	21	25	Jun. 4	Dec. 3	1
1798	13	G	C	Feb. 4	Feb. 21	8	13	17	May 27	2	12
1799	14	F	B	Jan. 20	6	Mar. 24	Apr. 28	2	12	1	23
1800	15	E	AG	Feb. 9	26	Apr. 13	May 18	22	Jun. 1	Nov. 30	4

PERPETUAL

PERPETUAL TIME-TABLES:

Or General RULES for CHRONOLOGISTS, HISTORIANS, &c.

JANUARY xxxi Days.

G	F	E	D	C	B	A	Days	Fixed Feasts, Remarkable Days, &c. according to New Style.
m	t	w	th	f	f	S	1	CIRCUMCISION. Day Br. 5 ^h 59 ^m . T. e.
t	w	th	f	f	S	m	2	Sun rises 8 4, sets 3 56, Lond[6h 1m.
w	th	f	f	S	m	t	3	
th	f	f	S	m	t	w	4	Sir Isaac Newton born, 1643, N. S.
f	f	S	m	t	w	th	5	Old Christmas-day.
f	S	m	t	w	th	f	6	EPIPHANY.
S	m	t	w	th	f	f	7	Sun flow of Clocks 7m 10s.
m	t	w	th	f	f	S	8	Lucian.
t	w	th	f	f	S	m	9	
w	th	f	f	S	m	t	10	Princess Elizabeth born, 1740.
th	f	f	S	m	t	w	11	Day Break 5 22. Tw. ends 6 8.
f	f	S	m	t	w	th	12	Sun rises 7 56, sets 4 5.
f	S	m	t	w	th	f	13	St. Hillary, Bishop of Poitiers.
S	m	t	w	th	f	f	14	Ox. & Cam. Terms begin.
m	t	w	th	f	f	S	15	Exchequer opens.
t	w	th	f	f	S	m	16	
w	th	f	f	S	m	t	17	Old Twelfth-day. ☉ fl. of Cl. 10 ^m 56 ^s .
th	f	f	S	m	t	w	18	Prisca.
f	f	S	m	t	w	th	19	Sun enter'd ☿, Noon, 1756.
f	S	m	t	w	th	f	20	Fabian.
S	m	t	w	th	f	f	21	Agnes. Day Br. 5 43. Tw. e. 6 17.
m	t	w	th	f	f	S	22	Vincent. Sun rises 7 44, sets 4 16.
t	w	th	f	f	S	m	23	Hillary Term begins.
w	th	f	f	S	m	t	24	
th	f	f	S	m	t	w	25	CONVERSION OF ST. PAUL.
f	f	S	m	t	w	th	26	
f	S	m	t	w	th	f	27	Sun flow of Clocks 13m 29s.
S	m	t	w	th	f	f	28	
m	t	w	th	f	f	S	29	
t	w	th	f	f	S	m	30	K. CHA. II. MAR. 1648-9, O. S.
w	th	f	f	S	m	t	31	[12m past One.

The NEW, FULL, and QUARTER MOONS.

N	F	F	L	Gol. No.	N	F	F	L	Gol. No.	N	F	F	L	Gol. No.	N	F	F	L	Gol. No.
M	Q	M	Q		M	Q	M	Q		M	Q	M	Q		M	Q	M	Q	
●	☾	☉	☾		●	☾	☉	☾		●	☾	☉	☾		●	☾	☉	☾	
1	30	7	15	23	7	24	1	31	9	17	13	17	25	3	10				
2	19	26	4	12	8	12	20	27	5	14	6	14	21	29					
3	8	16	23	1	9	1	31	9	16	24	15	25	3	11	18				
4	26	4	11	19	10	20	28	6	13	16	14	21	29	6					
5	16	23	1	31	8	11	10	17	25	3	17	3	11	18	25				
6	5	12	20	27	12	28	6	13	20	18	22	29	7	15					
										19	11	18	26	4					

FEBRUARY xxviii or xxix Days.

G	F	E	D	C	B	A	Days	Fixed Feasts, Remarkable Days, &c. according to New Style.
th	f	f	S	m	t	w	1	Sun rises 7 26, sets 4 34, London.
f	f	S	m	t	w	th	2	PURIF. V. MARY, or Candlemas-day.
f	S	m	t	w	th	f	3	Blaize, Bishop.
S	m	t	w	th	f	f	4	
m	t	w	th	f	f	S	5	Agatha.
t	w	th	f	f	S	m	6	
w	th	f	f	S	m	t	7	Sun flow of Clocks 14m 43s.
th	f	f	S	m	t	w	8	
f	f	S	m	t	w	th	9	
f	S	m	t	w	th	f	10	Dies Scholastica, at Oxford.
S	m	t	w	th	f	f	11	Day Br. 5 12. Tw. ends 6 48.
m	t	w	th	f	f	S	12	Hillary T. ends. Sun r. 7 8. f. 4 52.
t	w	th	f	f	S	m	13	Old Candlemas-day.
w	th	f	f	S	m	t	14	Valentine, a Bishop of Rome & M.
th	f	f	S	m	t	w	15	
f	f	S	m	t	w	th	16	
f	S	m	t	w	th	f	17	Sun flow of Clocks, 14m 29s
S	m	t	w	th	f	f	18	Sun in ♄ 21' Noon, 1756.
m	t	w	th	f	f	S	19	
t	w	th	f	f	S	m	20	
w	th	f	f	S	m	t	21	Day Br. 4 55. Twil. ends 7 5.
th	f	f	S	m	t	w	22	Sun rises 6 49, sets 5 11.
f	f	S	m	t	w	th	23	
f	S	m	t	w	th	f	24	St. MATTHIAS.
S	m	t	w	th	f	f	25	
m	t	w	th	f	f	S	26	
t	w	th	f	f	S	m	27	Sun flow of Clocks 13m 4s
w	th	f	f	S	m	t	28	Hare-Hunting goes out.
th	f	f	S	m	t	w	29	Leap-Year.
								Camb. Com. for B. A. the Day after Ash-Wednesday.

The NEW, FULL, and QUARTER MOONS.

N	F	F	L	Gol. No.	N	F	F	L	Gol. No.	N	F	F	L	Gol. No.	N	F	F	L	Gol. No.
M	Q	M	Q		M	Q	M	Q		M	Q	M	Q		M	Q	M	Q	
●	☾	☉	☾		●	☾	☉	☾		●	☾	☉	☾		●	☾	☉	☾	
1	28	6	13	21	7	22	—	8	15	13	16	23	1	9					
2	17	25	3	10	8	11	18	25	3	14	5	13	20	27					
3	7	14	22	29	9	—	8	15	23	15	24	2	9	17					
4	25	3	10	17	10	19	27	4	12	16	12	20	27	5					
5	14	22	29	7	11	8	16	23	1	17	2	9	16	24					
6	3	11	18	26	12	26	4	11	19	18	20	8	6	13					
										19	10	18	25	3					

☞ Under the Dom. Letter for O. or N. Stile, against the Day of the Month stands the Week-Day Letter correspondent.
 * * * Oxford and Cambridge Terms end Thursday before Whitsunday—Trinity College Election on Trinity Monday.

☞ Oxford and Cambridge Terms begin Wednesday after Trinity Sunday—Trinity College Election, Monday after Trinity Sunday.

PERPETUAL TIME-TABLES:

Or General RULES for CHRONOLOGISTS, HISTORIANS, &c.

MARCH xxxi Days.

G	F	E	D	C	B	A	Days	Fixed Feasts, Remarkable Days, &c. according to New Style.
th	f	f	S	m	t	w	1	S DAVID, Abp. M. b. 441. T. e. 7 20
f	f	S	m	t	w	th	2	Cedde or Chad. B. Sun r. 6 34 f. 5 27
f	S	m	t	w	th	f	3	
S	m	t	w	th	f	f	4	Sun flow of Clocks 12m 1s.
m	t	w	th	f	f	S	5	Princess Mary of Hesse b. 1722.
t	w	th	f	f	S	m	6	
w	th	f	f	S	m	t	7	Perpetua Mauric. Mart.
th	f	f	S	m	t	w	8	Camb. latter Aet Thurs. af. 4 S. in
f	f	S	m	t	w	th	9	Lent.
f	S	m	t	w	th	f	10	Day br. 4 19. Twil. ends 7 42.
S	m	t	w	th	f	f	11	Sun rises 6 14, sets 5 47.
m	t	w	th	f	f	S	12	Gregory.
t	w	th	f	f	S	m	13	
w	th	f	f	S	m	t	14	Sun flow of Clocks 9m 21s.
th	f	f	S	m	t	w	15	Oxf. & Cam. Terms end the Day bef.
f	f	S	m	t	w	th	16	Palm S.
f	S	m	t	w	th	f	17	St. Patrick, a Bishop of Ireland.
S	m	t	w	th	f	f	18	Ed. K. of West Sax.
m	t	w	th	f	f	S	19	Prise Louisa-Ann b. 1749. Joseph.
t	w	th	f	f	S	m	20	Cuthbert. Equal Day and Night.
w	th	f	f	S	m	t	21	Benedict Abbot. D. b. 3 57. T. e. 8 4.
th	f	f	S	m	t	w	22	Sun rises 5 54, sets 6 7.
f	f	S	m	t	w	th	23	Sun in V 20' on 20 D. Noon, 1756.
f	S	m	t	w	th	f	24	Sun flow of Clocks 6m 18s.
S	m	t	w	th	f	f	25	LADY-DAY. ANNUNCIATION B.
m	t	w	th	f	f	S	26	[V. M. Pr. Edward b. 1739.
t	w	th	f	f	S	m	27	N. B. According to Gregorian
w	th	f	f	S	m	t	28	Reckoning, the Golden Numbers to the
th	f	f	S	m	t	w	29	Left of all, shew the Times of the
f	f	S	m	t	w	th	30	Full Moons preceding Easter-Sunday,
f	S	m	t	w	th	f	31	till 1900.

The NEW, FULL, and QUARTER MOONS.

Go. No.	N	F	F	L	Go. No.	N	F	F	L
M	Q	M	Q	M	M	Q	M	Q	M
●	☾	☉	☾	☉	●	☾	☉	☾	☉
1	30	8	15	23	7	23	1	30	8
2	19	27	5	12	8	12	20	27	5
3	7	15	22	30	9	1	31	9	16
4	26	4	11	19	10	21	28	6	14
5	16	23	1	31	11	9	17	24	1
6	5	13	20	28	12	28	6	13	21

N. B. Corpus Christ Th. af. Tr. S. — Cam. Commencement 1st Tuesd. July.

The above is not always correct for Easter

APRIL xxx Days.

G	F	E	D	C	B	A	Days	Fixed Feasts, Remarkable Days, &c. according to New Style.
13	S	m	t	w	th	f	1	All Fools-day. D. b. 3 31. T. e.
2	m	t	w	th	f	f	2	Sun rises 5 32, sets 6 28. [8 29.
t	w	th	f	f	S	m	3	Richard B. Chich.
10	w	th	f	f	S	m	4	S. Ambrose. Sun fl. of Cl. 2m 55s
th	f	f	S	m	t	w	5	Old Lady day.
18	f	f	S	m	t	w	16	Oxf. & Cam. Terms beg. Wed. af.
7	f	S	m	t	w	th	7	1st S. aft. Easter.
S	m	t	w	th	f	f	8	
15	m	t	w	th	f	f	9	
4	t	w	th	f	f	S	10	D. br. 3 34. T. e. 8 57.
w	th	f	f	S	m	t	11	Sun rises 5 13, sets 6 48.
12	th	f	f	S	m	t	12	Easter Ter. beg. 17 Days af. Easter,
f	f	S	m	t	w	th	13	ends in 17 Days.
f	S	m	t	w	th	f	14	Sun flow of Clocks om 16s.
9	S	m	t	w	th	f	15	Sun and Clocks together.
m	t	w	th	f	f	S	16	Westminster Election Day af. 4 S. af.
17	t	w	th	f	f	S	17	Easter.
6	w	th	f	f	S	m	18	[in 20 Days.
th	f	f	S	m	t	w	19	Alphege. Tr. T. beg. 5Ds. af. Ea. ends
f	f	S	m	t	w	th	20	Sun in 8 46' Noon, 1756.
f	S	m	t	w	th	f	21	Day Br. 2 37. Tw. ends 9 24.
S	m	t	w	th	f	f	22	Sun rises 4 54, sets 7 7.
m	t	w	th	f	f	S	23	S. Geo. of Cappadocia, Patr. Eng.
t	w	th	f	f	S	m	24	Sun fast of Clocks 2m 6s.
w	th	f	f	S	m	t	25	St. MARK, Evangelist.
th	f	f	S	m	t	w	26	DUKE OF CUMBERLAND, b. 1721.
f	f	S	m	t	w	th	27	Vict. of Culloden.
f	S	m	t	w	th	f	28	N. B. According to the Greg. Reck-
S	m	t	w	th	f	f	29	ing, the Gold. Nos. to the Left of all,
m	t	w	th	f	f	S	30	shew the Times of the Full Moons pre-
							ceding Easter Sunday, till 1900.	

The NEW, FULL, and QUARTER MOONS.

Go. No.	N	F	F	L	Go. No.	N	F	F	L
M	Q	M	Q	M	M	Q	M	Q	M
●	☾	☉	☾	☉	●	☾	☉	☾	☉
1	28	6	13	21	7	22	29	7	14
2	18	25	3	11	8	11	19	26	3
3	6	14	21	28	9	30	8	15	23
4	25	3	10	18	10	19	27	4	12
5	14	22	29	7	11	7	15	22	30
6	4	11	18	26	12	27	4	12	19

N. B. The Rules for variable Days are in the Body of the Months from January to April, and then in Succession from January to April at the Bottom.

* * Oxf. Aet. 7 Days fr. Cam. Commencement. — Bailed Col. Election Th. bef. St. Andrew.

PERPETUAL TIME-TABLES:

Or General RULES for CHRONOLOGISTS, HISTORIANS, &c.

MAY xxxi Days.

F	E	D	C	B	A	Days	Fixed Feasts, Remarkable Days, &c. according to New Style.
w	th	f	f	S	m	1	St. PHIL. & JAMES. D. br. 2 2. T. E.
th	f	f	S	m	t	2	Sun rises 5 36, sets 7 25. [9 59.
f	f	S	m	t	w	3	Invention of the Cross.
f	S	m	t	w	th	4	Sun fast of Clocks 3m 33s.
S	m	t	w	th	f	5	
m	t	w	th	f	f	6	St. John Evang. ante Port. Lat.
t	w	th	f	f	S	7	
w	th	f	f	S	m	8	
th	f	f	S	m	t	9	
f	f	S	m	t	w	10	
f	S	m	t	w	th	11	Day br. 1 24. Twil. ends 10 38.
S	m	t	w	th	f	12	Old May-day. Sun r. 4 19, f. 7 42.
m	t	w	th	f	f	13	
t	w	th	f	f	S	14	Sun fast of Clocks 4m 2s.
w	th	f	f	S	m	15	
th	f	f	S	m	t	16	
f	f	S	m	t	w	17	
f	S	m	t	w	th	18	
S	m	t	w	th	f	19	Dunstan.
m	t	w	th	f	f	20	Sun in II 44' on 21 D. Noon, 1756.
w	th	f	f	S	21	Day br. 0 12. Twil. ends 11 49.	
th	f	f	S	m	22	Sun rises 4 5, sets 7 55.	
f	f	S	m	t	23		
f	S	m	t	w	24	Pr. Fred. Will. b. 1750. Sun fast of Cl.	
S	m	t	w	th	25	[3m 36s.	
m	t	w	th	f	26	Augustine. No Night.	
t	w	th	f	f	S	27	Venerable Bede.
w	th	f	f	S	28		
th	f	f	S	m	29	K. CHARLES II's NAT. and RESTOR.	
f	f	S	m	t	30		
f	S	m	t	w	31		

JUNE xxx Days.

G	F	E	D	C	B	A	Days	Fixed Feasts, Remarkable Days, &c. according to New Style.
f	f	S	m	t	w	th	1	Nicomedes. No Night.
f	S	m	t	w	th	f	2	Sun rises 3 52, sets 8 8.
S	m	t	w	th	f	f	3	
m	t	w	th	f	f	S	4	George Prince of Wales born, 1738.
t	w	th	f	f	S	m	5	Boniface. Sun fast of Clocks 2m 1s.
w	th	f	f	S	m	t	6	
th	f	f	S	m	t	w	7	
f	f	S	m	t	w	th	8	
f	S	m	t	w	th	f	9	
S	m	t	w	th	f	f	10	Prsse Amelia b. 1711.
m	t	w	th	f	f	S	11	St. BARNABAS.
t	w	th	f	f	S	m	12	Sun rises 3 45, sets 8 15. No Nt.
w	th	f	f	S	m	t	13	
th	f	f	S	m	t	w	14	
f	f	S	m	t	w	th	15	Sun fast of Clocks om 2s.
f	S	m	t	w	th	f	16	Sun and Clocks together.
S	m	t	w	th	f	f	17	St. Alban.
m	t	w	th	f	f	S	18	
t	w	th	f	f	S	m	19	Sun in 22' on 21 D. Noon, 1756.
w	th	f	f	S	m	t	20	Transl. of Edw. K. West Sax.
th	f	f	S	m	t	w	21	Sun rises 3 42, sets 8 17. No Night.
f	f	S	m	t	w	th	22	K GEO. II's Inaug. Longest Day.
f	S	m	t	w	th	f	23	
S	m	t	w	th	f	f	24	St. JOHN BAPT. born.
m	t	w	th	f	f	S	25	Sun fl. of Cl. 2m 5s. St. John's Col. El.
t	w	th	f	f	S	m	26	King GEORGE II. proclaimed.
w	th	f	f	S	m	t	27	
th	f	f	S	m	t	w	28	
f	f	S	m	t	w	th	29	St. PETER.
f	S	m	t	w	th	f	30	Buck hunting comes in and continues till Holy Rood. Exeter Col. Elec.

The NEW, FULL, and QUARTER MOONS.

N	F	F	L	Go. No.	N	F	F	L	Go. No.	N	F	F	L	Go. No.
M	Q	M	Q	Go. No.	M	Q	M	Q	Go. No.	M	Q	M	Q	Go. No.
8	6	13	21	7	21	29	6	14	13	15	23	1	30	8
7	25	3	10	8	10	18	25	3	14	5	12	19	27	
6	13	20	28	9	29	17	15	22	15	23	30	8	15	
4	2	10	17	10	19	27	4	12	16	12	19	26	4	
4	21	28	6	11	7	15	22	29	17	1	31	9	16	24
3	11	18	26	12	26	4	11	19	18	20	18	6	13	
									19	9	16	23	1	31

The NEW, FULL, and QUARTER MOONS.

N	F	F	L	Go. No.	N	F	F	L	Go. No.	N	F	F	L	Go. No.
M	Q	M	Q	Go. No.	M	Q	M	Q	Go. No.	M	Q	M	Q	Go. No.
1	26	4	11	19	7	20	27	5	12	13	14	21	28	6
2	14	23	1	30	8	9	17	24	2	14	3	11	18	26
3	4	12	19	26	9	28	6	13	21	15	21	29	6	14
4	23	1	30	8	10	27	25	3	10	16	11	18	25	3
5	12	20	27	5	11	6	13	20	28	17	29	7	14	22
6	1	30	9	17	12	24	2	10	17	18	19	26	4	12
										19	7	15	22	29

REMARKS. P. 172, Right Hand Col. l. 18. fr. Top. for *since Christ*, read *before Christ*.
P. 177, Right Hand Col. l. 3. fr. Bot. for *Rules*, read *Rules*.
P. 170, Left Hand Col. l. 16. fr. Top. for *required*, read *require*.

PERPETUAL TIME-TABLES:

Or General RULES for CHRONOLOGISTS, HISTORIANS, &c.

JULY xxxi Days.

G	F	E	D	C	B	A	Days	Fixed Feasts, Remarkable Days, &c. according to New Style.
S	m	t	w	th	f	f	1	No Night.
m	t	w	th	f	f	S	2	Visit. of B. V. M. Sun r. 3 45, f. 8 15.
t	w	th	f	f	S	m	3	Dies Comitiar.
w	th	f	f	S	m	t	4	Transf. of St. Mart. B.
th	f	f	S	m	t	w	5	Old Midsummer-day.
f	f	S	m	t	w	th	6	
f	S	m	t	w	th	f	7	Thomas à Becket.
S	m	t	w	th	f	f	8	Sun flow of Clocks 4m 30s.
m	t	w	th	f	f	S	9	
t	w	th	f	f	S	m	10	
w	th	f	f	S	m	t	11	No Night. Sun rises 8 52, sets 8 7.
th	f	f	S	m	t	w	12	
f	f	S	m	t	w	th	13	
f	S	m	t	w	th	f	14	
S	m	t	w	th	f	f	15	St. Swithin, or Swithun, B. of Winch.
m	t	w	th	f	f	S	16	[transf.]
t	w	th	f	f	S	m	17	
w	th	f	f	S	m	t	18	Sun flow of Clocks 5m 39s.
f	f	S	m	t	w	th	19	
f	S	m	t	w	th	f	20	Margaret V. and M.
f	S	m	t	w	th	f	21	No Night. Sun rises 4 3, sets 7 56.
S	m	t	w	th	f	f	22	Mary Magdalen. <i>Prise Car. Matilda</i>
m	t	w	th	f	f	S	23	Sun in Ω 53' Noon, 1756. [b. 1751.
t	w	th	f	f	S	m	24	Magdalen Coll. Election.
w	th	f	f	S	m	t	25	St. JAMES.
th	f	f	S	m	t	w	26	St. Ann, Mother B. V. M.
f	f	S	m	t	w	th	27	
f	S	m	t	w	th	f	28	Sun flow of Clocks 5m 56s.
S	m	t	w	th	f	f	29	
m	t	w	th	f	f	S	30	Dog-days beg. Sirius rises with the
t	w	th	f	f	S	m	31	[Sun.]

The NEW, FULL, and QUARTER MOONS.

Go. No.	N	F	F	L	Go. No.	N	F	F	L
	M	Q	M	Q		M	Q	M	Q
1	26	4	11	19	7	19	27	4	12
2	15	23	10	8	8	8	16	23	31
3	3	11	18	26	9	27	5	12	30
4	22	30	8	15	10	17	24	2	10
5	12	19	26	4	11	5	13	20	27
6	1	31	9	15	12	24	1	31	9

AUGUST xxxi Days.

G	F	E	D	C	B	A	Days	Fixed Feasts, Remarkable Days, &c. according to New Style.
w	th	f	f	S	m	t	1	Lammas-day. Day br. 1 27.
th	f	f	S	m	t	w	2	Sun r. 4 19, sets 7 40. T. ends 10 31.
f	f	S	m	t	w	th	3	
f	S	m	t	w	th	f	4	
S	m	t	w	th	f	f	5	Sun flow of Clocks 5m 31s.
m	t	w	th	f	f	S	6	Transfiguration.
t	w	th	f	f	S	m	7	Name of Jesus.
w	th	f	f	S	m	t	8	
th	f	f	S	m	t	w	9	
f	f	S	m	t	w	th	10	St. Laurence, Arch-deac. Rome, & M.
f	S	m	t	w	th	f	11	Princess Augusta born, 1737.
S	m	t	w	th	f	f	12	D. br. 2 5. T. e. 9 53. Sun r. 4 36.
m	t	w	th	f	f	S	13	[f. 7 23.]
t	w	th	f	f	S	m	14	
w	th	f	f	S	m	t	15	Assumption of V. M. Sun fl. of Cl.
th	f	f	S	m	t	w	16	[3m 56s.]
f	f	S	m	t	w	th	17	
f	S	m	t	w	th	f	18	
S	m	t	w	th	f	f	19	
m	t	w	th	f	f	S	20	
t	w	th	f	f	S	m	21	Athanasius. D. br. 2 38. T. e. 9 21.
w	th	f	f	S	m	t	22	Sun rises 4 54, sets 7 5.
th	f	f	S	m	t	w	23	Sun in Ω 38' Noon, 1756.
f	f	S	m	t	w	th	24	St. BARTHOLOMEW.
f	S	m	t	w	th	f	25	Sun flow of Clocks 1m 35s.
S	m	t	w	th	f	f	26	
m	t	w	th	f	f	S	27	
t	w	th	f	f	S	m	28	St. Austin B. of Hip.
w	th	f	f	S	m	t	29	Beheading St. John Bapt.
th	f	f	S	m	t	w	30	Sun and Clocks together.
f	f	S	m	t	w	th	31	

The NEW, FULL, and QUARTER MOONS.

Go. No.	N	F	F	L	Go. No.	N	F	F	L
	M	Q	M	Q		M	Q	M	Q
1	24	2	9	17	7	18	25	3	11
2	14	21	28	6	8	7	15	22	29
3	2	10	17	25	9	26	4	11	19
4	21	28	6	14	10	15	23	1	30
5	10	18	25	3	11	4	11	19	26
6	29	7	14	22	12	22	30	7	15

PERPETUAL

PERPETUAL TIME-TABLES:

Or General RULES for CHRONOLOGISTS, HISTORIANS, &c.

SEPTEMBER xxx Days.

G	F	E	D	C	B	A	Days	Fixed Feasts, Remarkable Days, &c. according to New Style.
f	S	m	t	w	th	f	1	Giles, Abbot. D. br. 3 9. T. e. 8 50.
S	m	t	w	th	f	f	2	LONDON burnt 1666, O.S. Sun r. 5 14.
m	t	w	th	f	f	S	3	[f. 6 44.
t	w	th	f	f	S	m	4	
w	th	f	f	S	m	t	5	Sun fast of Clocks 1m 45s.
th	f	f	S	m	t	w	6	
f	f	S	m	t	w	th	7	Eunurchus.
f	S	m	t	w	th	f	8	Nativity B. V. M.
S	m	t	w	th	f	f	9	Dog-days end. <i>Canis Major</i> r. 3 M.
m	t	w	th	f	f	S	10	
t	w	th	f	f	S	m	11	Day Br. 3 35. Twil. ends 8 24.
w	th	f	f	S	m	t	12	Sun rises 5 34, sets 6 25.
th	f	f	S	m	t	w	13	
f	f	S	m	t	w	th	14	Holy Cross day, or Exalt. of the Cross.
f	S	m	t	w	th	f	15	Sun fast of Clocks 5m 11s.
S	m	t	w	th	f	f	16	
m	t	w	th	f	f	S	17	Lambert B.
t	w	th	f	f	S	m	18	
w	th	f	f	S	m	t	19	
th	f	f	S	m	t	w	20	D. br. 3 58. Twil. ends 8 1.
f	f	S	m	t	w	th	21	St. MATTHEW. Sun r. 5 53, f. 6 5.
f	S	m	t	w	th	f	22	Equal Day and Nt. in all the World.
S	m	t	w	th	f	f	23	Sun in $\triangle$ 48' Noon, 1756.
m	t	w	th	f	f	S	24	
t	w	th	f	f	S	m	25	Sun fast of Clocks 8m 37s.
w	th	f	f	S	m	t	26	St. Cyprian.
th	f	f	S	m	t	w	27	
f	f	S	m	t	w	th	28	Sheriffs of London sworn.
f	S	m	t	w	th	f	29	St. MICHAEL. Hare-hunting comes
S	m	t	w	th	f	f	30	St. Jerom. [in and lasts till the End [of Feb.

The NEW, FULL, and QUARTER MOONS.

Go. No.	N	F	F	L	Go. No.	N	F	F	L	Go. No.	N	F	F	L
M	Q	M	Q	M	M	Q	M	Q	M	M	Q	M	Q	M
●	☾	☉	☾	☉	●	☾	☉	☾	☉	●	☾	☉	☾	☉
1	23	1	31	8	16	7	16	24	1	9	13	10	18	25
2	12	20	27	5	8	8	5	14	20	28	14	29	7	14
3	30	8	15	23	9	9	24	2	10	17	15	18	25	3
4	20	27	5	12	10	10	14	21	28	6	16	7	15	22
5	9	17	24	2	11	11	2	10	17	25	17	26	4	11
6	28	5	13	21	12	12	21	29	6	14	18	15	23	1
											19	4	11	18

OCTOBER xxxi Days.

G	F	E	D	C	B	A	Days	Fixed Feasts, Remarkable Days, &c. according to New Style.
m	t	w	th	f	f	S	1	Remegius, B. of Rhemes.
t	w	th	f	f	S	m	2	D. br. 4 20. T. e. 7 39. Sun r. 6 13.
w	th	f	f	S	m	t	3	[f. 5 46.
th	f	f	S	m	t	w	4	Sun fast of Clocks, 11m 30s.
f	f	S	m	t	w	th	5	
f	S	m	t	w	th	f	6	St. Faith, V. & M.
S	m	t	w	th	f	f	7	
m	t	w	th	f	f	S	8	
t	w	th	f	f	S	m	9	St. Denis.
w	th	f	f	S	m	t	10	Old Michaelmas-day. Ox. & Cam. T. b.
th	f	f	S	m	t	w	11	D. br. 4 40. T. e. 7 19. Sun r. 6 33.
f	f	S	m	t	w	th	12	[f. 5 22.
f	S	m	t	w	th	f	13	Transf. K. Ed. Conf.
S	m	t	w	th	f	f	14	Sun fast of Clocks 14m 5s.
m	t	w	th	f	f	S	15	
t	w	th	f	f	S	m	16	
w	th	f	f	S	m	t	17	Etheldred Virg.
th	f	f	S	m	t	w	18	St. LUKE the Evangelist.
f	f	S	m	t	w	th	19	
f	S	m	t	w	th	f	20	
S	m	t	w	th	f	f	21	D. br. 5 0. T. e. 6 59. Sun r. 6 52.
m	t	w	th	f	f	S	22	K. Geo. II. crowned, 1727. [f. 5 7.
t	w	th	f	f	S	m	23	Sun in η 31' Noon, 1756.
w	th	f	f	S	m	t	24	Sun fast of Clocks 15m 45s.
th	f	f	S	m	t	w	25	<i>Crispin</i> , Patron of Shoe-makers.
f	f	S	m	t	w	th	26	
f	S	m	t	w	th	f	27	
S	m	t	w	th	f	f	28	St. SIMON and JUDE.
m	t	w	th	f	f	S	29	
t	w	th	f	f	S	m	30	
w	th	f	f	S	m	t	31	

The NEW, FULL, and QUARTER MOONS.

Go. No.	N	F	F	L	Go. No.	N	F	F	L	Go. No.	N	F	F	L
M	Q	M	Q	M	M	Q	M	Q	M	M	Q	M	Q	M
●	☾	☉	☾	☉	●	☾	☉	☾	☉	●	☾	☉	☾	☉
1	22	30	8	15	7	16	23	1	31	8	13	10	18	25
2	12	19	26	4	8	5	13	19	27	14	29	7	14	24
3	30	8	15	23	9	24	2	9	17	15	17	25	3	10
4	19	27	4	12	10	13	21	28	7	16	6	14	21	29
5	8	16	23	1	11	1	31	19	16	24	17	25	3	11
6	27	5	12	20	12	20	28	6	13	18	15	23	29	7
										19	3	11	18	26

RULES for the STARS.

RIGHT ASCENSION and DECLINATION of STARS.

To determine the right Ascension of any Star from the given right Ascension of any other Star.

SAY, as the Time marked by a Clock, going uniformly, whether regulated or not, during the Interval of a Star's Revolution, is to 360°, so is the Time marked by the same Clock, between the Passage of that and any other Star, through the Meridian, to their Difference of Right Ascension.

To determine the Right Ascension of any particular Star?

RULE. When the Sun is near the Equinox, where its Change in Declination is swiftest, observe its Meridian Height or Declination some Day at Noon. And by the Method of corresponding Altitudes, or otherwise, observe the Difference in R. A. between the chosen Star and the Sun, at the same Instant of Noon. When the Sun has passed the following Solstice, and is returned nearly to the same Parallel, observe, for three or four Days together, its Meridian Altitudes, and its Difference in R. A. with the same Star, for determining, from these Observations, when the Sun comes into the same Parallel, as in the first Observation, and the Difference of R. A. for the same Instant.

This Method gives two Instants when the Sun was at equal Distances from the same Tropic; because, at equal Distances, on either Side a Tropic, the Declinations are equal, as are likewise the corresponding Arcs of the Equator.

The Difference of R. A. answering to these two Instants, will give (the Star being fixed) the Arc of the Equator, or Sun's Motion in R. A. in the Interval of these two Instants.

The Solstitial Colure therefore bisects that Arc, the Complement of Half which will be the Sun's true R. A. at the first Observation.

The Sun's R. A. being thus determined, the R. A. of the Star is likewise determined by the observed Difference.

	Merid. Alt. ☉ at Noon.		Dif. R. A. bet. ☉ & * Procyon, at Noon.
EXAMPLE. 1745, April 4. . . .	46 58 41	97 52 10 East	
Sept. 6. . . .	47 29 32	53 39 29 West	
7. . . .	47 7 1	54 33 36 West	
8. . . .	46 44 24	55 27 43 West	

Interpolating these Observations, it appears that if the Sun had been in the Meridian Sept. 7^d 8^h 50^m P. M. he would have had equal Altitude, with his Altitude on April 4th at Noon, preceeding; viz. . . . 46° 58' 41"

The Difference between his R. A. and that of the Star, had been . . . 54 53 39 West

Therefore, from April 4, to September 7^d 8^h 50^m Evening, the Sun had run through, in R. A. . . . 152 45 49

Whence, on April 4th Noon, Sun's Distance in R. A. from the Tropic of $\overline{26}$ was . . . 76 22 54½

And had R. A. . . .	{ 13 37 5½
The Star being to the East . . .	{ 97 52 10

The Star Procyon had R. A. required . . . Sum . . . 111° 29' 15½"

USES of the RIGHT ASCENSION and DECLINATION of the STARS.

1. To find the Longitude and Latitude of those Stars.
2. To shew the Order in diurnal Revolution, and the Intervals of Time they take in succeeding each other in their Passage through the Meridian.
3. To compute at what Time each Star passes the Meridian. Thus, Take the Difference between the Star's R. A. and the R. A. of the Sun, for the Noon of the given Day, reduce this Difference to Time, by Tab. p. 28, which will be nearly the Interval of Time from Noon to Noon of the Star's Passage through the Meridian.

The above Computation gives the Time of the Passage but nearly; because neither the Sun or Star are supposed to have any Motion in R. A.

But, to find the correct Time, compute the Sun and Star's R. A. for the Time already found, and their Difference, reduced to Time, will give the correct Instant of the Star's passing the Meridian.

EXAMPLE. Suppose the R. A. of Mars, on a given Day at Noon be 112° 18', and the R. A. of the Sun 183° 42', the Difference 71° 24' reduced to Time is 4^h 45^m 36^s. Now Mars being eastward of the Sun, must pass the Meridian about 4^h 45^m 36^s before Noon, that is, 7^h 14^m 24^s in the Morning.

	☉	1	"
The R. A. of Mars for that Time is . . .	112	13	12
The R. A. of the Sun for that Time is . . .	183	30	10

The Difference . . . 71 16 58

Reduced to Time, is the Instant of the Passage of Mars through the Meridian . . . 7^h 14^m 52^s required.

This Computation is plainly the Reverse of the former, in finding the R. A. of the Stars by observing their Passages over the Meridian.

By the above Computation, the Times marked by a Clock may be proved.

For, observing at what Instant any Star, whose R. A. is known, passes the Meridian, that Instant being compared with the other, found by Computation, will shew whether the Clock agrees with true Time, or what it differs from it.

4. Another Use of the R. A. and Declination of Stars is, To find the Distance of any Star from the Meridian of a Place at a given Time. Or which is the same,

To find the Angle at the Pole, formed by the Meridian of a Place and the Circle of Declination passing through the Star.

METHOD. Reduce the Interval of Time between Noon and the given Instant, into Degrees, by Tab. p. 28, add them to the Sun's R. A. at that Instant; and from the Sum subtract the Star's R. A.

N. B. When the said Sum is less than the Star's R. A. add to it 360°.

RULES for SUN and STARS.

FROM Two of these THREE THINGS, the POLES HEIGHT, SUN or STAR'S DECLINATION, and Meridian ALTITUDE (signified by P, D, M. and their Complements by p, d, and m, respectively) being given to find a THIRD, by Addition and Subtraction, only.

EXAMPLE

RULES for SUN and STARS.

EXAMPLE I.

The Meridian Altitude of the SUN or a Star, and the Pole's Elevation being given, to find the SUN or Star's Declination?

The elevated Pole, and Meridian Altitude, ON THE SAME SIDE OF THE MERIDIAN.

Pole's Height greater than the Meridian Altitude.

CASE 1. The Sum of the Complement of the Pole's Height and Meridian Altitude will be the SUN or Star's Declination, sought, of the same Name with that of the Pole.

Or, $p + M = D$. Let the Pole's Height be 50° N. and Meridian Altitude 30° . Then, $40^\circ + 30^\circ = 70^\circ$, the Star's Declination N. required.

N. B. The Difference of the Pole's Height and Meridian Altitude in this Case being added to the Pole's Height will give the Meridian Altitude above the Pole, as in 2d Case, viz. $50^\circ - 30^\circ = 20^\circ$; $+ 50^\circ = 70^\circ$ Altitude above the Pole.

Pole's Height less than the Meridian Altitude.

CASE 2. The Sum of the Complement of the Meridian Altitude and the Pole's Height (being the Reverse of the former Case) will be the SUN or Star's Declination of the same Name with that of the Pole.

Or, $m + P = D$. Let the Pole's Height be 20° N. and Meridian Altitude 40° . Then $50^\circ + 20^\circ = 70^\circ$, the Star's Declination N. required.

N. B. The Difference of the Pole's Height and Meridian Altitude in this Case being subtracted from the Pole's Height gives the Meridian Altitude below the Pole, as in 1st Case, viz. $40^\circ - 20^\circ = 20^\circ$ And $20^\circ - 20^\circ = 0^\circ$ Altitude beneath the Pole.

The elevated Pole and Meridian Altitude, ON CONTRARY SIDES OF THE MERIDIAN.

The Height of the Equator greater than the Meridian Altitude.

CASE 3. Subtract the Meridian Altitude from the Equator's Height, and the Remainder will be the SUN or Star's Declination, with a contrary Name to that of the Pole.

Or, $p - M = D$. Let the Pole's Height be 70° N. and Meridian Alt. 15° . Then, $20^\circ - 15^\circ = 5^\circ$, the SUN or Star's Declination S. required.

The Height of the Equator less than the Meridian Altitude.

CASE 4. Subtract the Equator's Height from the Meridian Altitude (being the Reverse of the former Case) and the Remainder will be the Declination of like Name with that of the Pole.

Or, $M - p = D$. Let the Pole's Height be 50° N. and Meridian Altitude 50° . Then, $50^\circ - 40^\circ = 10^\circ$, the Sun or Star's Declination, N. required.

EXAMPLE II.

The Pole's Height and SUN or Star's Declination being given to find the SUN or Star's Meridian Altitude?

The Pole's Elevation and Star's Declination, OF THE SAME NAME.

CASE 1. Add the Declination to the Equator's Height, and if the Sum be less than 90° , it will be the Meridian Altitude sought, on the contrary Side of the Meridian to that of the elevated Pole. For by Transposition of Case 4. Exam. I.

$D + p = M$. Let the Pole's Height be 50° N. and Declination $* 10^\circ$ N. Then, $10^\circ + 40^\circ = 50^\circ$, the Meridian Altitude, S. required.

CASE 2. When the said Sum exceeds 90° , its Supplement to 180° will be the Meridian Altitude, sought, between the Zenith and Pole. Or by Transposition of Case 2. Exam. I.

$D - P = m$. Let the Pole's Height be 20° N. and Star's Declination 70° N. $70^\circ - 20^\circ = 50^\circ$ Comp. Merid. Alt. and 40° Meridian Altitude N. sought, above the Pole.

Also, 70° Equator's Height $+ 70^\circ$ *'s Declination $= 140^\circ -$
180

Supplement, the Merid. Alt. 40° N. as before.

CASE 3. The Height of the Equator subtracted from the Declination of the SUN or Star will give the Meridian Altitude on the same Side of the Meridian with the Pole, by Transposition of Case 1. Exam. I.

That is, $D - p = M$. Let the Pole's Height be 60° N. and Declination 40° . Then, $40^\circ - 30^\circ = 10^\circ$, The Meridian Altitude N. required, below the Pole.

Universally, when the Declination is of the same Name with the Pole. The Com. of Declination (or Distance from Pole) being subtracted (if possible) from the Pole's Height will give the Meridian Altitude below the Pole; and being added to the Pole's Altitude will give the Meridian Altitude above the Pole if under 90° .

When the Sum of the Comp. of Declination and Pole's Height exceed 90° , then the Supplement to 180° will be the Meridian Altitude on the contrary Side of the Meridian to that of the Pole.

Thus, in Case 1, foregoing, 80° *'s Distance from Pole $+ 50^\circ$ N. the Pole's Height $= 130^\circ$, whose Supplement is 50° S. the Meridian Altitude, as before.

In Case 2. 20° *'s Distance from Pole $+ 20^\circ$ Pole's Height N. $= 40^\circ$, Meridian Altitude N. above the Pole, as before.

In Case 3. 50° *'s Distance from Pole from 60° Pole's Height N. remains 10° , Meridian Altitude, below the Pole, as before.

The Pole's Elevation and SUN or Star's Declination, HAVING DIFFERENT NAMES.

CASE 4. Subtract the Declination from the Height of the Equator and the Remainder will be the Meridian Altitude, of a contrary Name to that of the Pole, by Transposition of Case 3. Exam. I.

$p - D = M$. Let the Pole's Height be 70° N. and *'s Declination 5° . Then, $20^\circ - 5^\circ = 15^\circ$ Meridian Altitude S. required.

EXAMPLE III. PARTICULARLY USEFUL IN NAVIGATION.

THE STAR's or SUN's Declination and Meridian Altitude being given for some one Place of the Earth, to find the Elevation of the Pole at that Place.

When the Meridian Altitude is between the Pole and Horizon, and DECLINATION OF THE SAME NAME WITH THE POLE.

CASE 1. Add the Complement of Declination to the Meridian Altitude, and the Sum will be the Height of the Pole.

Or, $d + M = P$. Let the Meridian Altitude 30° and Declination 70° N. Then, $20^\circ + 30^\circ = 50^\circ$, the Height of the Pole required, or $D - M = p$, by Case 1. Exam. I. Or $70^\circ - 30^\circ = 40^\circ$ Comp. Pole, whence 50° Pole's Height, N. as before.

When the Meridian Altitude is between the Pole and Zenith.

CASE 2. Subtract the Comp. of Declination (or Distance from Pole) from the Meridian Altitude, and the Remainder will be the Pole's Height.

Or,

RULES for SUN and STARS.

Or, $M - d = P$. Let the Meridian Altitude be 40° and $\star$'s Declination 70° N. Then, $40^\circ - 20^\circ = 20^\circ$, the Pole's Height N. required. Or $D - m = P$, by Transposition of Case 2. Exam. I. Or $70^\circ - 50^\circ = 20^\circ$, Pole's Height N. as before.

When the Meridian Altitude and the Pole, are on contrary Sides of the Meridian.

CASE 3. Subtract the Declination from the Altitude, and the Remainder is the Height of the Equator, or Comp. of the Pole's Altitude.

Or, $M - D = p$. Let Meridian Altitude 50° and $\star$'s Declination 10° . Then, $50^\circ - 10^\circ = 40^\circ$ Comp. Latitude. Whence 50° Height Pole, N. required.

For, $M - D = p$, by Transposition of Case 4. Exam. I.

When the Declination and Pole are of DIFFERENT NAMES.

CASE 4. Add the Meridian Altitude and $\star$'s Declination, and the Sum will be the Height of the Equator, whose Com. is the Poles Height.

$M + D = p$. Let the Meridian Altitude be 15° and Star's Declination 5° S.

Then, $15^\circ + 5^\circ = 20^\circ$, the Equator's Altitude, whence 70° Pole's Height, N. required.

For $M + D = p$, by Transposition of Case 3. Exam. I.

*** The foregoing are all the Cases that can happen from the different Data of any two of the three foregoing Arcs of great Circles of the Sphere, viz. Pole's Heights, Star's or SUN's Declination, and Meridian Altitude.

The other Circumstances of the Position and Phenomena of the Stars, in respect of the great Circles of the Sphere, for any particular Place of the Earth, are computed by the Rules of Trigonometry.

ANY THREE of these FIVE THINGS being given; the Obliquity of the ECLIPTIC, a Star's Right ASCENSION, DECLINATION, LONGITUDE, LATITUDE, the other Two may be determined.

These are practically reduced to the Obliquity of the Ecliptic, the Measure of Time, and right Ascension and Declination of each Star; from whence the Longitude and Latitude of a Star are easily known.

TABLE of the Apparent SEMI-DIAMETER of the SUN, TIME of his Passing the MERIDIAN, and his DISTANCE from the EARTH, for every 10th Day of the Leap-Year, 1756, N. S. By *Connoissance des Temps*.

Days	JANUARY.				FEBRUARY.				MARCH.				APRIL.			
	Sem. Diam. app. $\odot$	T. of passing Meridian.		Dist. from $\odot$	Sem. Diam. app. $\odot$	Time passing Meridian.		Dist. from $\odot$	Sem. Diam. app. $\odot$	Time passing Meridian.		Dist. from $\odot$	Sem. Diam. app. $\odot$	Time passing Meridian.		Dist. from $\odot$
	" "	M. S.	S. Ds. $\ominus$	" "	" "	M. S.	S. Ds. $\ominus$	" "	" "	M. S.	S. Ds. $\ominus$	" "	" "	M. S.	S. Ds. $\ominus$	" "
10	16 20	2 21	21638	16 16	2 14	21730	16 9	2 10	21886	16 0	2 9	22083	16 0	2 9	22083	
20	16 19	2 19	21659	16 14	2 11	21780	16 7	2 9	21951	15 58	2 10	22141	15 58	2 10	22141	
30	16 18	2 16	21690	16 12	2 10	21836	16 3	2 8	22016	15 56	2 12	22197	15 56	2 12	22197	
Days	MAY.				JUNE.				JULY.				AUGUST.			
	Sem. Diam. app. $\odot$	T. of passing Meridian.		Dist. from $\odot$	Sem. Diam. app. $\odot$	Time passing Meridian.		Dist. from $\odot$	Sem. Diam. app. $\odot$	Time passing Meridian.		Dist. from $\odot$	Sem. Diam. app. $\odot$	Time passing Meridian.		Dist. from $\odot$
	" "	M. S.	S. Ds. $\ominus$	" "	" "	M. S.	S. Ds. $\ominus$	" "	" "	M. S.	S. Ds. $\ominus$	" "	" "	M. S.	S. Ds. $\ominus$	" "
10	15 54	2 14	22253	15 49	2 17	22349	15 48	2 18	22365	15 48	2 18	22370	15 52	2 12	22285	
20	15 52	2 15	22286	15 48	2 18	22365	15 49	2 16	22349	15 54	2 11	22252	15 54	2 11	22252	
30	15 50	2 16	22320	15 48	2 18	22370	15 50	2 14	22321	15 56	2 10	22195	15 56	2 10	22195	
Days	SEPTEMBER.				OCTOBER.				NOVEMBER.				DECEMBER.			
	Sem. Diam. app. $\odot$	T. of passing Meridian.		Dist. from $\odot$	Sem. Diam. app. $\odot$	Time passing Meridian.		Dist. from $\odot$	Sem. Diam. app. $\odot$	Time passing Meridian.		Dist. from $\odot$	Sem. Diam. app. $\odot$	Time passing Meridian.		Dist. from $\odot$
	" "	M. S.	S. Ds. $\ominus$	" "	" "	M. S.	S. Ds. $\ominus$	" "	" "	M. S.	S. Ds. $\ominus$	" "	" "	M. S.	S. Ds. $\ominus$	" "
10	15 59	2 9	22137	16 7	2 10	21949	16 14	2 17	21767	16 19	2 22	21652	16 19	2 22	21652	
20	16 1	2 9	22076	16 9	2 12	21885	16 16	2 19	21718	16 20	2 22	21635	16 20	2 22	21635	
30	16 3	2 9	22014	16 12	2 14	21829	16 18	2 21	21685	16 20	2 22	21630	16 20	2 22	21630	

EQUATION of the SUN'S PLACE, for every 4 Julian YEARS. According to Halley's Motions.

Days	January.	February.	March.	April.	May.	June.	July.	August.	Septemb.	October.	November.	December.
	+	+	+	+	+	+	+	+	+	+	+	+
	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "
1	1 44	1 46	1 47	1 49	1 51	1 52	1 53	1 52	1 51	1 49	1 47	1 45
11	1 45	1 46	1 48	1 50	1 52	1 53	1 53	1 52	1 50	1 48	1 46	1 45
21	1 45	1 47	1 48	1 51	1 52	1 53	1 53	1 51	1 50	1 48	1 46	1 44

EQUATION of the SUN'S PLACE for every 4 Julian YEARS. According to De la Caille.

Days	January.	February.	March.	April.	May.	June.	July.	August.	Septemb.	October.	November.	December.
	+	+	+	+	+	+	+	+	+	+	+	+
	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "
1	1 44	1 47	1 47	1 48	1 48	1 51	1 52	1 52	1 47	1 45	1 44	1 46
11	1 45	1 47	1 47	1 47	1 50	1 52	1 50	1 53	1 45	1 46	1 45	1 47
21	1 46	1 47	1 50	1 48	1 50	1 52	1 51	1 49	1 45	1 44	1 47	1 45

A Solar EPHEMERIS, commencing 1756, and serving by Means of EQUATIONS to the END of the present CENTURY.

A TABLE of the TIME of the DAY by a regulated CLOCK when it is NOON by the SUN. Shewing the EQUATION of natural DAYS, and Advance and Retreat of CLOCK-TIME for the LEAP-YEAR.

J A N.				F E B.				M A R C H				A P R I L				M A Y				J U N E			
Days.		Cl. Eq.		Days.		Cl. Eq.		Days.		Cl. Eq.		Days.		Cl. Eq.		Days.		Cl. Eq.		Days.		Cl. Eq.	
h. m. s.		+ -		h. m. s.		+ -		h. m. s.		+ -		h. m. s.		+ -		h. m. s.		+ -		h. m. s.		+ -	
1	12 4 28	ad	20 37	12	14 12	ad	30 21	12	12 36	re	28 45	12	3 48	re	19 57	11	56 47	re	12 56	11	57 20	ad	13 29
2	12 4 56		21 5	12	14 19		30 28	12	12 23		28 32	12	3 30		19 39	11	56 40		12 49	11	57 29		13 38
3	12 5 23		21 22	12	14 25		30 34	12	12 10		28 19	12	3 11		19 20	11	56 33		12 42	11	57 39		13 48
4	12 5 50		21 59	12	14 30		30 39	12	11 56		28 5	12	1 53		19 2	11	56 27		12 36	11	57 49		13 58
5	12 6 17		22 26	12	14 34		30 43	12	11 42		27 51	12	2 35		18 44	11	56 21		12 30	11	57 59		14 8
6	12 6 43		22 52	12	14 38		30 47	12	11 28		27 37	12	2 17		18 26	11	56 16		12 25	11	58 9		14 18
7	12 7 9		23 18	12	14 40		30 49	12	11 13		27 22	12	2 0		18 9	11	56 12		12 21	11	58 20		14 29
8	12 7 34		23 43	12	14 42		30 51	12	10 58		27 7	12	1 43		17 52	11	56 8		12 17	11	58 31		14 40
9	12 7 59		24 8	12	14 43		30 52	12	10 42		26 51	12	1 26		17 35	11	56 4		12 13	11	58 42		14 51
10	12 8 23		24 32	12	14 44		30 53	12	10 26		26 35	12	1 9		17 18	11	56 1		12 10	11	58 54		15 3
11	12 8 46		24 55	12	14 43		30 52	12	10 10		26 19	12	0 53		17 2	11	55 59		12 8	11	59 6		15 15
12	12 9 9		25 18	12	14 42		30 51	12	9 53		26 2	12	0 37		16 46	11	55 58		12 7	11	59 18		15 27
13	12 9 31		25 40	12	14 40		30 49	12	9 36		25 45	12	0 21		16 30	11	55 57	ad	12 6	11	59 30		15 39
14	12 9 53		26 2	12	14 37		30 46	12	9 19		25 28	12	0 6		16 15	11	55 56		12 5	11	59 42		15 51
15	12 10 14		26 23	12	14 33		30 42	12	9 2		25 11	11	59 51		16 0	11	55 56		12 5	11	59 55		16 4
16	12 10 34		26 43	12	14 29		30 38	12	8 44		24 53	11	59 36		15 45	11	55 57		12 6	12	0 8		16 17
17	12 10 53		27 2	12	14 24		30 33	12	8 26		24 35	11	59 21		15 30	11	55 58		12 7	12	0 21		16 30
18	12 11 12		27 21	12	14 19		30 28	12	8 8		24 17	11	59 7		15 16	11	56 0		12 9	12	0 34		16 43
19	12 11 30		27 39	12	14 19		30 22	12	7 50		23 59	11	58 54		15 3	11	56 2		12 11	12	0 47		16 56
20	12 11 47		27 56	12	14 6		30 15	12	7 32		23 41	11	58 41		14 50	11	56 5		12 14	12	1 0		17 9
21	12 12 4		28 13	12	13 58		30 7	12	7 14		23 23	11	58 28		14 37	11	56 9		12 18	12	1 13		17 22
22	12 12 20		28 29	12	13 50		29 59	12	6 55		23 4	11	58 16		14 25	11	56 13		12 22	12	1 26		17 35
23	12 12 35		28 44	12	13 41		29 50	12	6 36		22 45	11	58 4		14 13	11	56 17		12 26	12	1 39		17 48
24	12 12 49		28 58	12	13 32		29 41	12	6 17		22 26	11	57 52		14 1	11	56 22		12 31	12	1 52		18 1
25	12 13 2		29 11	12	13 22		29 31	12	5 58		22 7	11	57 41		13 50	11	56 28		12 37	12	2 5		18 14
26	12 13 15		29 24	12	13 11		29 20	12	5 40		21 49	11	57 31		13 40	11	56 34		12 43	12	2 17		18 26
27	12 13 26		29 35	12	13 0		29 9	12	5 21		21 30	11	57 21		13 30	11	56 41		12 50	12	2 29		18 38
28	12 13 37		29 46	12	12 48		28 57	12	5 2		21 11	11	57 12		13 21	11	56 48		12 57	12	2 41		18 50
29	12 13 47		29 56					12	4 44		20 53	11	57 3		13 12	11	56 55		13 4	12	2 53		19 2
30	12 13 56		30 5					12	4 25		20 34	11	56 55		13 4	11	57 3		13 12	12	3 5		19 14
31	12 14 5		30 14					12	4 6		20 15		* fast.			11	57 11		13 20		* flow.		
Clock-Time advances 9 ^m 45 ^s in this Month.				Cl. T. advances 0 ^m 33 ^s retreats 2 ^m 6 ^s in this Month.				Clock-Time retreats 8 ^m 49 ^s in this Month.				Clock-Time retreats 7 ^m 1 ^s in this Month.				Clock-Time retreats 0 ^m 50 ^s , advances 1 ^m 24 ^s in this Month.				Clock-Time advances 5 ^m 55 ^s in this Month.			

IN taking the Time of the DAY, and Clock-Time-Equation, out of the above Table, you must take out for 1 Day sooner in the Months of January, and February, for Leap-Year: For other Months in that Year take out for the Days as they stand. And, for the 1st, 2d, and 3d Years after any Leap-Year, you must take out for $\frac{1}{4}$, $\frac{1}{2}$, and $\frac{3}{4}$ of a Day sooner, in all the Months, respectively.

EXAMPLES. REQUIRED the Times by the Clock, when it is Noon by the Sun, January 10th, 1760, February 23, 1761, June 29th, 1762, and December 15th, 1763, respectively?

Leap-Year.	h	m	s	Com. Year.	h	m	s	Com. Year.	h	m	s	Com. Year.	h	m	s
Jan. 9, 1760. . .	12	7	59	Feb. 23. . .	12	13	41	June 29. . .	12	2	53	December 15. . . .	11	55	58
For 10, 1 Day sooner				$\frac{1}{4}$ Day sooner		+	2	$\frac{1}{2}$ Day sooner		—	6	$\frac{1}{4}$ Day sooner . . .		—	22
Answers, required				1761. . .	12	13	43	1762. . .	12	2	47	1763. . . .	11	55	36

* Compare this short Method with Ferguson's EQUATION-TABLES, in his ASTRONOMY repeated for three Years after Leap-Year.

The Days, on which the Sun's and Clock's Time meet, or when the Time by a Sun-Dial, and that by a regulated Clock is the same, once in 24 Hours, are observed to be on April 15th, June 16th, August 31st, and December 24th, one Year with another. N. S.

A Solar EPHEMERIS, commencing 1756, and serving by Means of EQUATIONS to the END of the present CENTURY.

A TABLE of the TIME of the DAY by a regulated CLOCK when it is NOON by the SUN. Shewing the EQUATION of natural DAYS, and Advance and Retreat of CLOCK-TIME for the LEAP-YEAR.

JULY.				AUG.				SEPT.				OCT.				NOV.				DEC.										
Cl. Eq.				Cl. Eq.				Cl. Eq.				Cl. Eq.				Cl. Eq.				Cl. Eq.										
+				+				+				+				+				+										
h m s				h m s				h m s				h m s				h m s				h m s										
1	12	3	16	ad	19	25	12	5	44	re	21	53	11	59	33	19	15	42	11	43	51	o	0	0	11	49	43	24	5	52
2	12	3	27	11	19	36	12	5	40	4	21	49	11	59	14	19	15	23	11	49	11	1	0	0	11	50	7	24	6	16
3	12	3	38	11	19	47	12	5	36	4	21	45	11	58	55	19	15	4	11	48	53	1	0	1	11	50	31	25	6	40
4	12	3	49	11	19	58	12	5	31	5	21	40	11	58	36	19	14	45	11	48	35	2	0	2	11	50	56	25	7	5
5	12	4	0	10	20	9	12	5	25	6	21	34	11	58	17	20	14	26	11	48	17	3	0	4	11	51	21	25	7	30
6	12	4	10	9	20	19	12	5	19	6	21	28	11	57	57	20	14	6	11	48	0	4	0	7	11	51	47	26	7	56
7	12	4	19	9	20	28	12	5	12	7	21	21	11	57	35	20	13	46	11	47	43	5	0	11	11	52	13	27	8	22
8	12	4	28	9	20	37	12	5	5	8	21	14	11	57	17	20	13	26	11	47	27	6	0	16	11	52	40	27	8	40
9	12	4	37	9	20	46	12	4	57	9	20	57	11	56	57	20	13	6	11	47	12	7	0	22	11	53	7	28	9	16
10	12	4	46	8	20	55	12	4	48	9	20	57	11	56	37	21	12	46	11	46	57	15	0	29	11	53	35	28	9	44
11	12	4	54	8	21	3	12	4	39	10	20	48	11	56	16	21	12	25	11	46	42	15	0	36	11	54	3	28	10	12
12	12	5	2	7	21	11	12	4	29	10	20	38	11	55	55	21	12	4	11	46	27	15	0	44	11	54	31	29	10	40
13	12	5	9	7	21	13	12	4	19	11	20	28	11	55	34	21	11	43	11	46	13	13	0	53	11	55	0	29	11	9
14	12	5	16	6	21	25	12	4	8	12	20	17	11	53	13	21	11	22	11	46	0	13	1	3	11	55	29	29	11	38
15	12	5	22	6	21	36	12	3	56	12	20	5	11	54	52	21	11	1	11	45	57	12	1	56	11	55	58	29	12	7
16	12	5	28	5	21	37	12	3	44	12	19	53	11	54	31	20	10	40	11	45	35	11	1	44	11	56	27	29	12	36
17	12	5	33	5	21	42	12	3	32	13	19	41	11	54	11	21	10	20	11	45	24	11	1	33	11	56	56	30	13	5
18	12	5	38	4	21	41	12	3	19	13	19	28	11	53	50	21	9	59	11	45	13	10	1	22	11	57	26	30	13	35
19	12	5	42	4	21	31	12	3	6	14	19	15	11	53	29	21	9	38	11	45	3	10	1	12	11	57	56	30	14	5
20	12	5	46	3	21	55	12	2	52	14	19	1	11	53	8	21	9	17	11	44	53	9	1	2	11	58	26	30	14	35
21	12	5	49	2	21	38	12	2	38	15	18	47	11	52	47	21	8	56	11	44	44	9	0	53	11	58	56	30	15	1
22	12	5	51	2	22	0	12	2	23	15	18	32	11	52	26	20	8	35	11	44	35	8	0	44	11	59	26	30	15	35
23	12	5	53	2	22	2	12	2	8	16	18	17	11	52	6	20	8	15	11	44	27	7	0	36	11	59	56	30	16	5
24	12	5	55	1	22	4	12	1	52	16	18	1	11	51	46	20	7	55	11	44	20	6	0	29	11	47	18	30	16	35
25	12	5	56	0	22	5	12	1	36	17	17	45	11	51	26	20	7	35	11	44	14	5	0	23	11	47	37	30	17	5
26	12	5	56	1	22	5	12	1	19	17	17	28	11	51	6	20	7	15	11	44	9	5	0	18	11	47	56	20	17	35
27	12	5	55	1	22	4	12	1	2	17	17	11	11	50	46	20	6	55	11	44	4	5	0	13	11	47	16	21	18	5
28	12	5	54	2	22	3	12	0	45	17	16	54	11	50	26	19	6	35	11	43	59	4	0	8	11	48	37	21	18	34
29	12	5	53	3	22	0	12	3	28	18	16	37	11	50	7	19	6	16	11	43	55	2	0	4	11	48	58	22	19	3
30	12	5	51	4	22	0	12	0	10	18	16	19	11	49	48	19	5	57	11	43	53	1	0	2	11	49	20	23	19	32
31	12	5	48	4	22	57	11	59	52	18	16	1				19			11	43	52		0	1				29	20	1

Cl. T. adv. 2^m 40^s. re. 6^m 11^s in this Month. * re. Clock-Time retreats 10^m 4^s in this Month. Clock-Time retreats 5^m 38^s in this Month. Clock-Time advances 14^m 38^s in this Month. * flow

N. B. The CLOCK EQUATION for each Month of the Year, shews the Advance or Retreat of the Clock-Time, *correcting*, or beginning its *Era*, or *Epocha*, from Nov. 2d in any Year, one of the Days in the Year (the other being Feb. 10, May 15, and July 26, N. S.) when the Sun's mean and true Motion is very nearly alike for 24 Hours. And the Advance or Retreat of Clock Time, from any one Day to any other Day in the same, or following Month, is immediately determined by taking the Difference of the Clock-Equations for those Days, from the above Table.

EXAMPLE I. To find the Retreat of Clock-Time, from Sept. 1, to Oct. 1, (a Leap-Year,) following ?

For Sept. 1.	15	42
Oct. 1.	5	38
Dif.	10	4

Dif. 10 4 Retreat, as at the Bottom of September required.

EXAMPLE II. To find the Advance of Clock-Time, from 4th July, to 29th of the same Month following.

July 4th, $\frac{1}{4}$ D. sooner	12	3	40
July 29th, $\frac{1}{4}$ D. sooner	12	5	54
Dif.	2	14	advance, as by the Clock Equation.

Dif. 2 14 advance, as by the Clock Equation.

Dif. 2 14 advance, required.

N. B. Against Nov. 2d. . . stands 11 43 51 Clock-Time, when 12 by the Sun.

Suppose for Aug. 6. a Leap-Year. . . Dif. -16 9; With which connecting the Clock Equation, and you will have the Equation of [Time for that natural Day.

+5 19 Equation of Time, which added to 12 Hours.
Makes 12^h 5^m 19^s the same Clock-Time, as by Tab. when it is Noon by the Sun.
So for the Rest.

A Solar EPHEMERIS, commencing 1756, and serving by Means of EQUATIONS to the END of the present CENTURY.

JANUARY.							FEBRUARY.						
Days	☉ Place by.	Dif.	☉'s Dec. S.	Dif.	☉'s R. Afc.	Dif.	Days	☉'s Pl. by.	Dif.	☉'s Dec. S.	Dif.	☉'s R. Afc.	Dif.
1	11 38 7	61 10	22 58 22	4 34	282 39 12	66 9	1	13 10 41	60 49	16 53 35	17 29	315 39 11	60 49
2	12 39 17	61 11	22 52 48	6 2	283 45 21	66 3	2	14 11 30	60 48	16 36 6	17 45	316 40 0	60 37
3	13 40 28	61 10	22 46 46	6 28	284 51 24	65 55	3	15 12 18	60 46	16 18 21	18 3	317 40 37	60 25
4	14 41 38	61 10	22 40 18	6 56	285 57 19	65 50	4	16 13 4	60 44	16 0 18	18 19	318 41 2	60 13
5	15 42 48	61 10	22 33 22	7 22	287 3 9	65 44	5	17 13 48	60 43	15 41 59	18 36	319 41 15	60 1
6	16 43 58	61 9	22 26 0	7 48	288 8 53	65 36	6	18 14 31	60 42	15 23 23	18 51	320 41 16	59 49
7	17 45 7	61 10	22 18 12	8 15	289 14 29	65 29	7	19 15 13	60 41	15 4 32	19 6	321 41 5	59 36
8	18 46 17	61 9	22 9 57	8 42	290 19 58	65 20	8	20 15 54	60 40	14 45 26	19 21	322 40 41	59 24
9	19 47 26	61 9	22 1 15	9 6	291 25 18	65 9	9	21 16 34	60 38	14 26 5	19 35	323 40 5	59 13
10	20 48 35	61 8	21 52 9	9 32	292 30 27	65 3	10	22 17 12	60 37	14 6 30	19 49	324 39 18	59 2
11	21 49 43	61 8	21 42 37	9 58	293 35 30	64 52	11	23 17 49	60 35	13 46 41	20 2	325 38 20	58 47
12	22 50 51	61 7	21 32 39	10 23	294 40 22	64 42	12	24 18 24	60 33	13 26 39	20 16	326 37 7	58 37
13	23 51 58	61 6	21 22 16	10 46	295 45 4	64 33	13	25 18 57	60 32	13 6 23	20 29	327 35 44	58 26
14	24 53 4	61 5	21 11 30	11 12	296 49 37	64 23	14	26 19 29	60 30	12 45 54	20 41	328 34 10	58 18
15	24 54 9	61 5	21 0 18	11 36	297 54 0	64 13	15	27 19 59	60 29	12 25 13	20 52	329 32 28	58 4
16	26 55 14	61 4	20 48 4	11 58	298 58 13	64 1	16	28 20 28	60 27	12 4 21	21 4	330 30 32	57 55
17	27 56 18	61 3	20 36 44	12 22	300 2 14	63 51	17	29 20 55	60 25	11 43 17	21 15	331 28 27	57 44
18	28 57 21	61 3	20 24 22	12 46	301 6 5	63 40	18	30 21 20	60 23	11 22 2	21 25	332 26 11	57 34
19	29 58 24	61 2	20 11 36	13 57	302 9 45	63 28	19	1 21 43	60 22	11 0 37	21 37	333 23 45	57 23
20	29 59 26	61 1	19 58 29	13 32	303 13 13	63 15	20	2 22 5	60 20	10 39 0	21 45	334 21 8	57 13
21	2 0 27	61 0	19 44 57	13 52	304 16 28	63 4	21	3 22 25	60 18	10 17 15	21 55	335 18 21	57 5
22	3 1 27	60 59	19 31 5	14 15	305 19 32	62 52	22	4 22 43	60 17	9 55 20	22 4	336 15 26	56 55
23	4 2 26	60 58	19 16 50	14 35	306 22 24	62 41	23	5 23 0	60 15	9 33 16	22 13	337 12 21	56 46
24	5 3 24	60 58	19 2 15	14 59	307 25 5	62 30	24	6 23 15	60 13	9 11 3	22 21	338 9 7	56 37
25	6 4 22	60 57	18 47 16	15 13	308 27 35	62 16	25	7 23 28	60 12	8 48 42	22 28	339 5 44	56 28
26	7 5 19	60 56	18 32 3	15 37	309 29 51	62 4	26	8 23 40	60 10	8 20 14	22 35	340 2 12	56 21
27	8 6 15	60 55	18 16 26	15 58	310 31 55	61 52	27	9 23 50	60 8	8 3 39	22 34	340 58 33	56 13
28	9 7 10	60 55	18 0 28	16 16	311 33 47	61 40	28	10 23 58	60 7	7 40 45	22 34	341 54 46	
29	10 8 5	60 53	17 44 12	16 33	312 35 27	61 28							
30	11 8 58	60 52	17 27 39	16 53	313 36 55	61 14							
31	12 9 50	60 52	17 10 46		314 38 9								

In Leap-Year, take out for a Day sooner, for Jan. and Feb. only. —

1	Years after	5 ^h 43 55	respectively	2423	6158
For 2	Lp-Yr. take	11 37 50	for all	4846	3145
3	out for	17 26 45	Months.	7269	1385

for Dif. for Dif.

MARCH.							APRIL.						
Days	☉	Dif.	☉'s Dec. S.	Dif.	☉'s R. Afc.	Dif.	Days	☉	Dif.	☉'s Dec. S.	Dif.	☉'s R. Afc.	Dif.
1	11 24 4	60 5	7 18 6	22 46	342 50 50	55 57	1	12 10 40	59 2	4 49 19	23 2	11 11 45	54 32
2	12 24 9	60 2	6 55 20	23 13	343 46 47	55 49	2	13 9 42	59 0	5 12 21	22 55	12 6 17	54 35
3	13 24 11	60 0	6 32 7	23 5	344 42 36	55 44	3	14 8 42	58 58	5 35 16	22 53	13 0 52	54 35
4	14 24 11	59 59	6 9 2	23 13	345 38 20	55 36	4	15 7 40	58 55	5 58 9	22 45	13 55 31	54 40
5	15 24 10	59 56	5 45 49	23 19	346 33 56	55 31	5	16 6 35	58 52	6 20 54	22 38	14 50 11	54 43
6	16 24 6	59 54	5 22 30	23 17	347 29 27	55 23	6	17 5 27	58 51	6 43 32	22 30	15 44 54	54 46
7	17 24 0	59 52	4 59 13	23 25	348 24 50	55 18	7	18 4 18	58 49	7 6 2	22 24	16 39 40	54 51
8	18 23 52	59 50	4 35 48	23 29	349 20 8	55 14	8	19 3 7	58 47	7 28 26	22 17	17 34 31	54 54
9	19 23 42	59 48	4 12 19	23 31	350 15 22	55 8	9	20 1 54	58 45	7 50 43	22 9	18 29 25	54 52
10	20 23 30	59 46	3 48 48	23 33	351 10 30	55 1	10	21 0 39	58 42	8 12 52	21 59	19 24 21	55 1
11	21 23 16	59 45	3 25 15	23 37	352 5 31	54 57	11	21 59 21	58 41	8 34 51	21 52	20 19 22	55 6
12	22 23 1	59 43	3 1 38	23 38	353 0 28	54 54	12	22 58 2	58 39	8 56 43	21 43	21 14 28	55 11
13	23 22 44	59 41	2 38 0	23 39	353 55 22	54 48	13	23 56 41	58 37	9 18 26	21 34	22 9 39	55 16
14	24 22 25	59 38	2 14 21	23 42	354 50 10	54 48	14	24 55 18	58 36	9 40 0	21 24	23 4 55	55 21
15	25 22 3	59 36	1 50 39	23 42	355 44 58	54 43	15	25 53 54	58 33	10 1 24	21 14	24 0 16	55 27
16	26 21 39	59 34	1 26 57	23 43	356 39 41	54 41	16	26 52 27	58 32	10 22 38	21 4	24 55 43	55 32
17	27 21 13	59 33	1 3 14	23 41	357 34 22	54 37	17	27 50 59	58 30	10 43 42	20 53	25 51 15	55 36
18	28 20 46	59 30	0 39 33	23 42	358 28 59	54 34	18	28 49 29	58 28	11 4 35	20 44	26 46 51	55 47
19	29 20 16	59 28	0 15 51	23 43	359 23 33	54 33	19	29 47 57	58 25	11 25 19	20 30	27 42 38	55 52
20	29 19 44	59 26	N 7 52	23 41	0 18 6	54 30	20	30 46 22	58 20	11 45 49	20 21	28 38 30	55 57
21	1 19 10	59 24	0 31 33	23 39	1 12 36	54 28	21	1 44 45	58 22	12 6 10	20 8	29 34 27	56 2
22	2 18 34	59 22	0 55 12	23 37	2 7 4	54 28	22	2 43 7	58 20	12 26 18	19 55	30 30 29	56 11
23	3 17 56	59 20	1 18 49	23 36	3 1 32	54 28	23	3 41 27	58 18	12 46 13	19 43	31 16 40	56 18
24	4 17 16	59 18	1 42 25	23 34	3 56 0	54 27	24	4 39 45	58 16	13 5 56	19 30	32 22 58	56 27
25	5 16 34	59 15	2 5 59	23 32	4 50 27	54 27	25	5 38 1	58 14	13 25 26	19 18	33 19 25	56 31
26	6 15 49	59 14	2 29 31	23 28	5 54 54	54 27	26	6 36 15	58 13	13 44 44	19 4	34 15 56	56 41
27	7 15 3	59 12	2 52 59	23 23	6 39 21	54 27	27	7 34 28	58 11	14 3 48	18 49	35 12 37	56 47
28	8 14 15	59 9	3 16 22	23 21	7 33 48	54 28	28	8 32 39	58 9	14 22 37	18 35	36 9 24	56 54
29	9 13 24	59 7	3 39 43	23 16	8 28 16	54 28	29	9 30 48	58 8	14 41 12	18 23	37 6 18	57 2
30	10 12 31	59 5	4 2 59	23 12	9 22 44	54 30	30	10 28 56		14 59 35		38 3 10	
31	11 11 36		4 26 11		10 17 14								

N. B. The Difs. for a Day preceding any Month-day of Leap-year being Xd by 2423; 4846; 7269, will give the Quantity to be subtracted from or added to the Leap-year Quantity, for That to the same Month-day, 1, 2, 3 Yrs. after Leap-year, respectively. Or 6158, 3145, 1385, being respectively added to the Lo. Logs. of Difs. for a Day preceding the Month-day of Leap-year will give the Lo. Log. of a Quantity to be subd. fr. or added to the Leap-year Quantity for the Quant. for the same Month-day 1, 2, 3 Years after Leap-year, respectively. See Examples farther on, and Equations for 4 Years forward of the Quantities, or Places, for any given Year.

A Solar EPHEMERIS, commencing 1756, and serving by Means of EQUATIONS to the END of the present CENTURY.

M A Y.										J U N E.									
Days	☉'s Pl.	☉'s Pl.	☉'s Pl.	Dif.	☉'s Dec. N.	Dif.	☉'s R. Afc.	Dif.	Days	☉'s Pl.	☉'s Pl.	☉'s Pl.	Dif.	☉'s Dec. N.	Dif.	☉'s R. Afc.	Dif.		
1	11 27 2	58 4	15 17 42	17 51	39 0 33	57 19	2	11 15 56	57 23	22 10 15	8 45	69 42 25	61 26						
2	12 25 6	58 2	15 35 33	17 35	39 57 52	57 28	3	12 13 19	57 23	22 18 0	7 21	70 43 51	61 32						
3	13 23 8	58 1	15 53 8	17 20	40 55 20	57 35	4	13 10 42	57 22	22 25 21	6 58	71 45 23	61 38						
4	14 21 9	57 59	16 10 28	17 5	41 52 55	57 44	5	14 8 4	57 22	22 32 19	6 34	72 47 1	61 42						
5	15 19 8	57 58	16 27 33	16 48	42 50 39	57 53	6	15 5 26	57 20	22 38 53	6 12	73 48 43	61 48						
6	16 17 6	57 56	16 44 21	16 31	43 48 32	58 3	7	16 2 46	57 20	22 45 5	5 45	74 50 31	61 54						
7	17 15 2	57 55	17 0 52	16 16	44 46 35	58 10	8	17 0 6	57 19	22 50 50	5 22	75 52 25	61 55						
8	18 12 57	57 53	17 17 8	15 57	45 44 45	58 18	9	17 57 25	57 18	22 56 12	4 58	76 54 20	62 1						
9	19 10 50	57 51	17 33 5	15 40	46 43 3	58 29	10	18 54 43	57 17	23 1 10	4 31	77 56 21	62 4						
10	20 8 41	57 49	17 48 45	15 21	47 41 32	58 35	11	19 52 0	57 16	23 5 44	4 10	78 58 25	62 6						
11	21 6 30	57 48	18 4 6	15 4	48 40 7	58 45	12	20 49 16	57 16	23 9 54	3 45	80 0 31	62 10						
12	22 4 18	57 47	18 19 10	14 45	49 38 52	58 53	13	21 46 32	57 15	23 13 39	3 20	81 2 41	62 12						
13	23 2 5	57 46	18 33 55	14 28	50 37 45	59 2	14	22 43 47	57 15	23 16 59	2 57	82 4 53	62 16						
14	23 59 51	57 45	18 48 23	14 37	51 36 47	59 10	15	23 41 2	57 15	23 19 56	2 32	83 7 9	62 17						
15	24 57 36	57 43	19 23 0	13 51	52 35 57	59 21	16	24 38 17	57 14	23 22 28	2 7	84 9 26	62 19						
16	25 55 19	57 42	19 16 21	13 27	53 35 18	59 26	17	25 35 31	57 14	23 24 35	1 42	85 11 45	62 18						
17	26 53 1	57 41	19 29 48	13 9	54 34 44	59 37	18	26 32 45	57 14	23 26 17	1 18	86 14 3	62 21						
18	27 50 42	57 39	19 42 57	12 50	55 34 21	59 44	19	27 29 59	57 13	23 27 35	0 53	87 16 24	62 23						
19	28 48 21	57 37	19 55 47	12 29	56 34 5	59 54	20	28 27 12	57 12	23 28 28	0 27	88 18 47	62 23						
20	29 45 58	57 35	20 8 16	12 8	57 33 59	60 1	21	29 24 24	57 11	23 28 55	0 3	89 21 10	62 22						
21	30 43 33	57 34	20 20 24	11 48	58 34 0	60 10	22	30 21 35	57 11	23 28 58	0 22	90 23 32	62 22						
22	1 41 7	57 33	20 32 12	11 27	59 34 10	60 17	23	1 18 47	57 11	23 28 36	0 46	91 25 54	62 21						
23	2 38 40	57 33	20 43 39	11 5	60 34 27	60 23	24	2 15 58	57 11	23 27 50	1 11	92 28 15	62 20						
24	3 36 13	57 32	20 54 44	10 44	61 34 50	60 32	25	3 13 9	57 12	23 26 39	1 37	93 30 35	62 18						
25	4 33 45	57 30	21 5 28	10 22	62 35 22	60 41	26	4 10 21	57 11	23 25 2	2 0	94 32 53	62 16						
26	5 31 15	57 29	21 15 50	10 1	63 36 3	60 48	27	5 7 32	57 11	23 23 2	2 25	95 35 9	62 14						
27	6 28 44	57 28	21 25 51	9 37	64 36 51	60 53	28	6 4 43	57 11	23 20 37	2 51	96 37 23	62 12						
28	7 26 12	57 28	21 35 28	9 16	65 37 44	61 0	29	7 1 54	57 11	23 17 46	3 15	97 39 35	62 8						
29	8 23 40	57 27	21 44 44	8 53	66 38 44	61 7	30	7 59 5	57 10	23 14 31	3 38	98 41 43	62 5						
30	9 21 7	57 25	21 53 37	8 31	67 39 51	61 14	31	8 56 15	57 10	23 10 53		99 43 48							
31	10 18 32	57 25	22 2 8		68 41 5														

J U L Y.										A U G U S T.									
Days	☉'s Pl.	☉'s Pl.	☉'s Pl.	Dif.	☉'s Dec. N.	Dif.	☉'s R. Afc.	Dif.	Days	☉'s Pl.	☉'s Pl.	☉'s Pl.	Dif.	☉'s Dec. N.	Dif.	☉'s R. Afc.	Dif.		
1	9 53 26	57 11	23 6 50	4 25	100 45 49	61 57	1	9 29 2	57 28	17 54 44	15 29	131 55 55	58 8						
2	10 50 37	57 11	23 2 25	4 54	101 47 46	61 55	2	10 26 30	57 29	17 39 15	15 44	132 54 3	58 8						
3	11 47 48	57 12	22 57 31	5 16	102 49 41	61 49	3	11 23 59	57 30	17 23 31	16 1	133 52 2	57 59						
4	12 45 0	57 11	22 52 15	5 39	103 51 30	61 44	4	12 21 29	57 31	17 7 30	16 18	134 49 51	57 49						
5	13 42 11	57 11	22 46 36	6 3	104 53 14	61 39	5	13 19 0	57 33	16 51 12	16 34	135 47 32	57 32						
6	14 39 22	57 11	22 40 33	6 27	105 54 53	61 34	6	14 16 33	57 33	16 34 38	16 50	136 45 4	57 23						
7	15 36 33	57 12	22 34 6	6 50	106 56 27	61 27	7	15 14 6	57 35	16 17 48	17 4	137 42 27	57 15						
8	16 33 45	57 12	22 27 16	7 4	107 57 54	61 23	8	16 11 41	57 36	16 0 44	17 23	138 39 42	57 7						
9	17 30 57	57 11	22 20 2	7 36	108 59 17	61 16	9	17 9 17	57 37	15 43 21	17 34	139 36 49	56 57						
10	18 28 8	57 13	22 12 26	7 59	110 0 33	61 9	10	18 6 54	57 38	15 25 47	17 55	140 33 46	56 49						
11	19 25 21	57 13	22 4 27	8 22	111 1 42	61 2	11	19 4 32	57 40	15 7 52	18 7	141 30 35	56 41						
12	20 22 34	57 14	21 56 5	8 45	112 2 44	60 56	12	20 2 12	57 41	14 49 45	18 19	142 27 16	56 33						
13	21 19 48	57 14	21 47 20	9 7	113 3 40	60 49	13	20 59 53	57 43	14 31 26	18 34	143 23 49	56 25						
14	22 17 2	57 15	21 38 13	9 30	114 4 29	60 42	14	21 57 36	57 44	14 12 52	18 49	144 20 14	56 15						
15	23 14 17	57 15	21 28 43	9 51	115 5 11	60 36	15	22 55 20	57 45	13 54 3	19 1	145 16 29	56 7						
16	24 11 32	57 16	21 18 52	10 13	116 5 47	60 27	16	23 53 5	57 47	13 35 2	19 15	146 12 36	55 59						
17	25 8 48	57 15	21 8 39	10 35	117 6 14	60 19	17	24 50 52	57 48	13 15 47	19 26	147 8 35	55 53						
18	26 6 3	57 16	20 58 4	10 55	118 6 33	60 11	18	25 48 40	57 49	12 56 21	19 35	148 4 28	55 47						
19	27 3 19	57 16	20 47 9	11 17	119 6 44	60 3	19	26 46 29	57 51	12 36 42	19 52	149 0 15	55 38						
20	28 0 35	57 13	20 35 52	11 47	120 6 47	59 53	20	27 44 20	57 52	12 16 50	20 2	149 55 53	55 31						
21	28 57 53	57 19	20 24 15	11 59	121 6 40	59 45	21	28 42 12	57 54	11 56 48	20 15	150 51 24	55 25						
22	29 55 12	57 19	20 12 16	12 16	122 6 25	59 36	22	29 40 6	57 55	11 36 33	20 26	151 46 49	55 18						
23	☉ 52 31	57 20	20 0 0	12 40	123 6 1	59 27	23	☉ 38 1	57 57	11 16 7	20 36	152 42 7	55 12						
24	1 49 51	57 20	19 47 20	12 59	124 5 28	59 19	24	1 35 58	57 59	10 55 31	20 46	153 37 19	55 6						
25	2 47 11	57 22	19 34 21	13 16	125 4 47	59 12	25	2 33 57	57 59	10 34 45	20 58	154 32 25	55 0						
26	3 44 33	57 23	19 21 5	13 38	126 3 59	59 3	26	3 31 58	58 0	10 13 47	21 7	155 27 25	54 54						
27	4 41 56	57 24	19 7 27	13 57	127 3 2	58 52	27	4 30 0	58 4	9 52 40	21 15	156 22 19	54 48						
28	5 39 20	57 24	18 53 30	14 14	128 1 54	58 43	28	5 28 4	58 5	9 31 25	21 25	157 17 7	54 42						
29	6 36 44	57 25	18 39 16	14 33	129 0 37	58 34	29	6 26 9	58 7	9 10 0	21 36	158 11 40	54 38						
30	7 34 9	57 26	18 24 43	14 52	129 59 11	58 25	30	7 24 16	58 9	8 48 24	21 42	159 6 27	54 35						
31	8 31 35		18 9 51		130 57 36		31	8 22 25		8 26 42		160 1 2							

SEPTEMBER.										OCTOBER.											
Days	o	i	u	Dif.	o	i	u	Dif.		Days	o	i	u	Dif.	o	i	u	Dif.			
1	9	20	36	58 12	8	4	52	21 59	160 55 34	54 22	1	8	40	34	59 11	3	26	47	23 20	187 58 3	54 30
2	10	18	48	58 14	7	42	53	22 7	161 49 56	54 21	2	9	39	45	59 13	3	50	7	23 17	188 52 33	54 34
3	11	17	2	58 16	7	20	46	22 13	162 44 17	54 17	3	10	38	58	59 15	4	13	24	23 14	189 47 7	54 41
4	12	15	18	58 18	6	58	33	22 21	163 38 34	54 13	4	11	38	13	59 17	4	36	38	23 10	190 41 48	54 46
5	13	13	36	58 17	6	36	12	22 26	164 32 47	54 11	5	12	37	30	59 19	4	59	48	23 6	191 36 34	54 50
6	14	11	55	58 21	6	13	46	22 34	165 26 58	54 8	6	13	36	49	59 21	5	22	54	23 4	192 31 24	54 57
7	15	10	16	58 23	5	51	12	22 39	166 21 6	54 5	7	14	36	10	59 23	5	45	58	22 59	192 26 21	55 5
8	16	8	39	58 25	5	28	33	22 44	167 15 11	54 3	8	15	35	33	59 26	6	8	57	22 53	194 21 26	55 12
9	17	7	4	58 28	5	5	49	22 51	168 9 14	53 59	9	16	34	59	59 28	6	31	50	22 43	195 16 38	55 17
10	18	5	32	58 29	4	42	58	22 55	169 3 13	53 59	10	17	34	27	59 31	6	54	33	22 47	196 11 55	55 25
11	19	4	1	58 30	4	20	3	22 59	169 57 12	53 57	11	18	33	58	59 32	7	17	20	22 37	197 7 20	55 33
12	20	2	31	58 32	3	57	4	23 3	170 51 9	53 55	12	19	33	30	59 34	7	39	57	22 32	198 2 53	55 39
13	21	1	3	58 34	3	34	1	23 8	171 45 4	53 53	13	20	33	4	59 36	8	2	29	22 23	198 58 32	55 49
14	21	59	37	58 37	3	10	53	23 10	172 38 57	53 54	14	21	32	40	59 38	8	24	52	22 16	199 54 21	55 57
15	22	58	14	58 39	2	47	43	23 15	173 32 51	53 54	15	22	32	18	59 39	8	47	8	22 9	200 50 18	56 5
16	23	56	53	58 41	2	24	28	23 18	174 26 45	53 55	16	23	31	57	59 41	9	9	17	22 2	201 46 23	56 14
17	24	55	34	58 43	2	1	10	23 20	175 20 40	53 54	17	24	31	38	59 44	9	31	19	21 52	202 42 37	56 23
18	25	54	17	58 45	1	37	50	23 22	176 14 34	53 55	18	25	31	21	59 46	9	53	11	21 45	203 39 0	56 33
19	26	53	2	58 47	1	14	28	23 24	177 8 29	53 56	19	26	31	8	59 49	10	14	56	21 35	204 35 33	56 42
20	27	51	49	58 48	0	51	4	23 26	178 2 25	53 58	20	27	30	57	59 50	10	36	31	21 25	205 32 15	56 53
21	28	50	37	58 50	0	27	38	23 26	178 56 23	53 58	21	28	30	47	59 52	10	57	56	21 16	206 29 8	57 1
22</																					

I. CATALOGUE of remarkable FIXED STARS, their Longitudes and Latitudes, for the Beginning of the Year 1720, O. S. according to HALLEY. For comparing with the CHANGES of their observed PLACES, in remote YEARS, and thereby to find the exact Quantity of their MOTION.

N. B. Sparkling and lucid Stars are generally distinguished by brightest and bright.

Names of Stars.	Mag.	Longitude.	Latitude.	N or S	Names of Stars.	Mag.	Longitude.	Latitude.	N or S
		° ' "	° ' "				° ' "	° ' "	
T H E					T H E				
Extreme of Pegasus Wing	2	5 14 50	12 35 12	N	Northern among the same	3	27 5 40	51 39 36	N
Head of Andromeda	2	10 23 53	25 41 1	N	Last but 2 in Dragon's Tail	3	3 28 11	66 21 43	N
Northern in the Whale's Belly	3	18 1 47	20 21 19	S	Beginning Gr. Bear's Tail	2	4 56 25	54 20 16	N
Knot of the Fishes	3	25 27 33	9 5 10	S	Section Ship's Deck	2	7 17 0	55 52 30	S
Brightest in Andromeda's Girdle	2	26 27 44	25 56 19	N	Bright * in Lyon's Loins	2	7 22 21	14 19 4	N
Southern in Horn of ♈. * ♈	4	29 16 0	7 8 58	N	Southern in Lyon's Buttock	3	9 30 31	9 39 50	N
Northern in same Horn	3	0 3 0	8 28 16	N	Last but 1 Gr. Bear's Tail	2	11 44 0	56 23 15	N
Bright * Cassiop. Chair	2	1 13 7	51 13 50	N	Southern in Lyon's Thigh	3	13 38 0	6 5 10	N
Bright * ♈, above the Neck	2	3 44 18	9 57 12	N	Section Ship's Keel	2	15 2 18	57 11 11	S
* in Cassiopea's Breast, Schedir	2	3 55 21	46 35 54	N	Lyon's Tail	1	17 44 14	12 16 51	N
Flexure of Cassiop. Hip.	2	10 3 44	48 47 35	N	Bright * in Ship's Keel	2	19 17 10	72 39 32	S
Southern Foot of Andromeda	2	10 20 44	27 46 7	N	Brightest bet. Bear's and Lyon's Tails	2	20 39 22	40 7 53	N
Bright * Whale's Jaw	2	10 24 15	12 37 0	S	Last Gr. Bear's Tail	2	22 59 24	54 24 0	N
— in the Knee of Cassiop.	3	14 2 15	46 23 26	N	Bent of Southern Wing, Virgin	3	23 11 14	0 40 47	N
Head of Medusa, Algol	2	22 15 42	22 23 47	N	North. following Section Ship's Keel	3	24 59 30	63 42 23	S
Brightest of Pleiades	3	26 5 8	4 0 37	N	Preceding in Southern Wing of Virgin	3	0 55 52	1 22 1	N
Brightest in the Side of Perseus	2	28 11 4	30 5 20	N	South. following Section Ship's Keel	2	1 28 30	57 5 20	S
First of Hyades, in Bull's Nose	3	1 52 34	5 46 22	S	North. in Virg. Wing, Vindematrix	2	6 2 40	16 12 54	N
Northern Eye of the Bull	3	4 32 11	2 35 58	S	2d of Virgin's Southern Wing	3	6 17 11	2 48 53	N
Bull's Southern Eye, Aldebaran	1	5 52 0	5 29 50	S	Side, under Virgin's Girdle	3	7 34 54	8 38 27	N
Bright * Orion's Foot, Rigel	1	12 55 0	31 10 11	S	Bootes preceding Shoulder	3	13 43 18	49 33 0	N
Orion's Right Shoulder	2	17 2 33	16 51 30	S	Spike of the Virgin	1	19 56 22	2 2 0	S
Goat, Hircus	1	17 56 41	22 51 47	N	Brightest between Thighs of Bootes	1	20 18 52	30 57 0	N
Preceding of brightest in the Dove	2	18 16 38	57 24 15	S	Root of Caroline Oak [Arcturus]	2	28 5 57	13 6	S
First of Orion's Girdle	2	18 26 38	23 36 7	S	Bright * in Centaur's Loins	2	28 26 56	40 7 20	S
Northern Horn of the Bull	2	18 38 56	5 21 34	N	Pre. or hind Knee of Cent. Bor. Crucis	2	2 51 9	47 45 51	S
Middle * in Orion's Girdle	2	19 32 44	24 33 23	S	Ankle of same Foot, Seq. Crucis	2	7 46 12	48 35 3	S
Following in the same Girdle	2	20 46 45	25 20 17	S	Following in lower Part Cent's Foot	2	8 0 5	52 49 15	S
Tip of Bull's Southern Horn	3	20 52 55	2 14 24	S	Bright * North. Crown [Pes Crucis]	2	8 20 56	44 21 17	N
Following the brightest in the Dove	2	22 31 5	59 15 8	S	Southern Scale of Libra	2	11 11 40	0 22 51	N
Last of the Tail of little Bear, Polar	2	24 39 41	66 4 11	N	Brightest in Centaur's Breast	2	11 38 20	39 32 0	S
Following Shoulder Orion	1	24 50 0	16 4 26	S	Northern Scale of Libra	2	15 27 40	8 31 45	N
Following Shoulder Auriga, or Driver	2	26 0 32	21 28 20	N	Bright * Serpent's Neck, Ophiuchus	2	18 8 22	25 31 56	N
Fore Foot of preceding Twin	3	29 31 43	0 56 0	S	Left Knee, Centaur	2	19 54 30	14 4 47	S
Following in same Foot, Calx	3	25 1 23	0 51 22	S	Right Foot same	1	26 1 16	42 27 48	S
Extremity G. Dog's Fore Foot	2	3 17 58	41 17 47	S	Brightest in left Hand Ophiuchus, Yed	3	28 23 15	17 17 15	N
Foot of the following Twin	2	5 11 18	6 47 19	S	Middle of 3 in Forehead of Scorpion	2	28 40 50	1 56 31	S
Great Dog's Mouth, Sirius	1	10 14 0	39 32 8	S	Southern among same	3	29 2 25	5 25 46	S
Knee of following Twin	3	11 4 40	2 5 27	S	Northern in Forehead of Scorp.	2	29 17 56	1 3 9	N
Rudder of Ship Argus, Canopus	1	11 9 0	75 51 0	S	Preceding Knee of Ophiuchus	3	5 18 55	11 25 27	N
Groin of the following Twin	3	14 36 20	0 13 7	S	Scorpion's Heart, Antares	1	5 51 4	4 31 26	S
Head of preceding Twin, Castor	2	16 20 20	10 3 48	N	Head of Hercules	3	12 13 47	37 18 55	N
* between Thighs Gr. Dog	2	16 55 0	51 22 48	S	Following Knee, Ophiuchus	3	14 4 28	7 14 12	N
Head of following Twin, Pollux	2	19 21 9	6 39 27	N	Point of Southern Triangle	2	16 56 27	46 5 33	S
* in G. Dog's Belly	2	19 32 5	48 27 33	S	Sole of the Foot, Ophiuchus	3	17 30 2	1 47 38	S
Little Dog, Procyon	1	21 55 21	15 57 55	S	Head of Ophiuchus	3	18 30 32	35 53 16	N
Tail of Gr. Dog	2	25 40 27	50 37 41	S	Sting of Scorpion	2	20 40 0	13 43 25	S
Northern Afs	4	3 38 0	3 9 41	N	Following Shoulder, Ophiuchus	3	21 25 44	27 58 0	N
Southern Afs	4	4 48 40	0 3 46	N	Fifth Joint Scorpion's Tail	2	21 40 16	19 36 15	S
Northern of preceding in □ Gr. Bear	2	11 15 0	49 40 5	N	Bright * in Dragon's Head	2	24 0 35	74 58 26	N
Brightest in Deck of Ship Argus	2	14 41 24	58 21 6	S	Southern Part Bow, Sagittary	2	1 10 33	10 59 54	S
Northern of preceding in □ Gr. Bear	2	15 29 12	45 6 16	N	Left Shoulder Sagittary	3	8 28 12	3 23 32	S
The Lyon's Head, more Southern	3	16 47 16	9 41 4	N	Armpit of Sagittary	3	9 42 22	7 7 55	S
Hydra's Head	2	23 23 0	22 24 32	S	Bright * in Harp	1	11 22 18	61 45 31	N
* under Ship's Deck	2	23 29 0	64 27 32	S	Following in Head, Sagittary	3	12 21 44	1 29 0	N
Northern of 3, in Lyon's Neck	3	23 38 41	11 50 13	N	Peacock's Eye	2	19 53 56	36 11 0	S
Southern in the same Neck	3	23 59 24	4 50 20	N	Beak of the Swan.	2	27 20 37	49 0 31	N
Middle and bright * in same Neck	2	25 40 5	8 47 27	N	Bright * of the Eagle	1	27 48 24	29 19 11	N
Lyon's Heart, Regulus	1	25 56 20	0 26 38	N	Following of several in Goat's Horn	3	29 57 21	6 58 6	N
Lower of the following in □ Gr. Bear	2	26 31 35	47 7 26	N	Southern in same Horn	3	0 8 57	4 37 27	N

N. B. Following and preceding Shoulder, Thigh, Leg, &c.
Signify the Right and Left Shoulder, Thigh, Leg, &c.

I. CATALOGUE of remarkable STARS, continued, according to HALLEY.

Names of Stars.	Mag.	Longitude.			Latitude.	S	N	Names of Stars.	Mag.	Longitude.			Latitude.	S	N
		°	'	"						°	'	"			
Preceding Wing of Crane	2	11	56	35	32	50	22	In Mouth of the Fish, Fomalhaut	1	29	54	0	21	4	54
Northern Wing of Swan	3	12	22	17	64	27	14	Tail of the Swan	2	1	26	32	59	56	37
Preceding of 2 in Tail of the Goat	3	17	52	42	2	31	18	In Leg of Aquarius, Scephat	3	4	58	49	8	11	17
At Setting on of Crane's Tail	2	18	18	53	35	22	46	Head of Hydra	2	7	59	0	64	11	0
Preceding Shoulder Aquarius	3	19	29	23	8	38	43	At the End of the River, Achernar	1	11	14	6	59	19	40
Following in Tail of Goat	3	19	38	14	2	32	19	Brightest in Head of Phoenix	2	11	30	27	40	34	10
Breast of the Swan	3	20	57	51	57	9	20	In the Shou. of Pegasus Wing, Markab.	2	19	34	13	19	24	37
Southern Wing of the Swan	13	23	47	52	49	26	21	In Thigh of Pegasus, Scephat	2	25	27	13	31	8	6
Mouth of Pegasus	3	27	58	32	22	7	16	Northern of the Whale's Tail.	3	27	0	0	10	0	41
Following Shoulder of Aquarius	3	29	27	16	10	40	38	Southern of the Whale's Tail	2	28	38	2	20	46	52

II. CATALOGUE of remarkable FIXED STARS, their R. ASCENSIONS and DECLINATIONS, for the Beginning of the YEAR 1756, of immediate Use in ASTRONOMY and NAVIGATION. From the *Connoissance des Temps*, published, in France, by Order of the Academy of Sciences there.

Names of STARS and MAGNITUDE, accord- ing to Bayer.	Bayer. Mag.	R. Ascen. Time. h m s	R. Ascen. Equator. o ' "	Declin. o ' "	N S	Names of STARS and MAGNITUDE, accord- ing to Bayer.	Bayer. Mag.	R. Ascen. Time. h m s	R. Ascen. Equator. o ' "	Declin. o ' "	N S
Algenib, Pegasus	γ 2	0 0 42	0 10 31	13 49 37	N	Wing of the Virgin	ε 3	12 47 55	192 30 22	12 16 37	N
Cassiopea's Breast	α 3	0 26 44	6 42 2	55 11 38	N	Spike of the Virgin	α 1	13 19 12	198 5 35	9 52 45	S
Whale's Tail	β 2	0 31 14	7 49 59	19 19 42	S	2d of the Breast of G. Bear	ζ 2	13 11 52	198 30 34	56 12 18	N
Cassiopea's Girdle	γ 3	4 42 3	10 32 28	59 23 18	N	3d of the Br. of Gr. Bear	η 2	13 35 40	204 28 36	50 32 16	N
Polar Star	α 2	0 43 34	10 55 27	87 59 59	N	Arcturus	α 1	14 2 14	211 8 15	20 27 56	N
Andromeda's Girdle	β 2	0 55 51	13 59 59	34 19 12	N	Southern Scale of Ballance	α 2	14 35 1	219 21 18	15 0 43	S
Ram's Ear	γ 4	1 39 46	25 0 36	18 5 23	N	Shoulder of little Bear	β 2	14 49 14	222 55 10	75 9 7	N
Preceding Horn of Ram.	β 3	1 40 47	25 16 0	19 36 23	N	Northern Scale of Ballance	β 2	15 1 27	225 58 47	8 27 54	S
Foot of Andromeda. Alam	γ 2	1 48 43	27 15 21	41 8 47	N	Brightest of the Crown	α 2	15 21 47	231 4 45	27 33 3	N
Following Horn of the Ram	α 3	1 53 10	28 22 5	22 17 55	N	Brightest in Serpent's Neck	α 2	15 29 48	233 4 11	7 12 37	N
Jaw of the Whale	α 2	2 49 6	42 23 15	3 7 1	N	Northern in Front Scorpion	β 2	15 48 39	237 48 52	19 6 59	S
Head of Medusa, Algol	β 2	2 51 15	43 5 50	39 59 46	N	Scorpion's Heart, Antares	α 1	16 11 50	243 37 22	25 52 2	S
Brightest of Perseus	α 2	3 6 32	46 45 40	48 58 9	N	Knee of Serpentarius	ζ 3	16 21 4	245 56 16	10 3 5	S
Bull's Eye, Aldebaran	α 1	4 21 14	65 29 12	15 59 53	N	Head of Hercules	α 3	17 0 44	255 52 56	14 41 15	N
The Goat, Capella	α 1	4 57 53	74 40 32	45 43 9	N	Head of Serpentarius	α 2	17 20 46	260 54 16	12 45 29	N
Orion's Foot, Rigel	β 1	5 2 0	75 42 32	8 30 8	S	Preceding in H. of Dragon	β 3	16 22 5	261 14 9	52 29 26	N
Northern Horn of Bull	β 2	5 10 1	77 43 7	28 22 32	N	Shoulder of Serpentarius	β 3	17 28 33	262 51 22	4 41 25	N
West Shoulder of Orion	γ 2	5 11 12	78 0 52	6 6 21	N	Bright. of the H. of Dragon	γ 3	17 48 6	267 45 22	51 32 33	N
1st of Orion's Belt	α 2	5 18 42	79 53 31	0 30 2	S	Southern Bow of Sagittary	ε 3	11 5 0	271 59 40	34 28 13	S
2d	ε 2	5 22 58	80 57 46	1 22 42	S	Brightest of the Harp, Wega	α 1	18 25 39	277 10 12	38 34 15	N
3d	ζ 2	5 27 35	82 7 7	2 5 32	S	Fol. Shoulder of Sagittary	σ 3	18 37 14	280 1 50	26 34 30	S
Auriga's Shoulder	β 2	5 40 42	85 24 35	44 53 26	N	Brightest in the Eagle	α 2	19 35 39	294 43 9	8 14 34	N
Eastern Shoulder of Orion	α 1	5 41 2	85 29 39	7 20 20	N	Following Horn of Goat	β 2	20 13 59	301 49 8	15 31 59	S
Brightest Foot of Twins	γ 2	6 22 33	95 54 8	16 35 5	N	Breast of the Swan	α 2	20 29 45	308 16 43	44 25 9	N
Great Dog, Sirius	α 1	6 33 19	98 36 2	16 23 54	S	Pre. Shoulder of Aquarius	β 3	21 15 13	319 40 40	6 37 50	S
Northern Head of Twins	α 2	7 17 47	109 44 51	32 23 54	N	Mouth of Pegasus	ε 3	21 28 39	323 2 47	8 46 6	N
Little Dog, Procyon	α 2	7 25 18	111 37 48	5 49 58	N	Following in Breast of Goat	δ 3	21 30 1	323 23 8	17 13 12	S
Southern Head of Twins	β 2	7 29 8	112 35 23	28 35 37	N	Scephat	δ 3	22 37 57	340 25 5	17 6 47	S
Hydra's Heart	α 2	9 14 5	138 54 6	7 36 44	S	Southern Fish, Fomalhaut	α 1	22 40 23	341 1 42	30 54 31	S
Lyon's Heart, Regulus	α 1	9 53 44	148 50 18	13 9 1	N	Scephat, Pegasus	β 2	22 48 13	342 59 35	26 45 48	N
The Great Bear	β 2	10 45 10	161 44 4	57 40 57	N	Markab, Pegasus	α 1	22 48 53	343 9 26	13 53 51	N
Ditto	α 2	10 46 40	162 6 35	63 3 40	N	Head of Andromeda	α 2	23 51 54	358 20 27	27 44 32	N
Breast of the Lion	α 2	11 34 42	174 8 58	15 56 8	N	Fol. the Chair of Cassiopea	β 2	23 52 21	359 4 15	57 48 7	N
Thigh of the Gr. Bear	γ 2	11 38 57	175 12 59	55 3 1	N						
Of the Gr. Bear	δ 3	12 1 16	180 48 42	58 23 19	N						
1 of the Breast of G. Bear	γ 3	12 41 7	190 48 8	57 17 17	N						

Enquire at Jonathan's Coffee-House, near the Royal Exchange, London, for the Secret Astronomical Tables from Greenwich Observations, (2 printed Sheets) by Mr. Noel Morris.

Enquire at Jonathan's Coffee-House, near the Royal Exchange, London, for the Secret Astronomical Tables from Greenwich Observations, (3 printed Sheets) by Mr. Gael Morris.

REMARK. We are obliged to insert the foregoing and other foreign astronomical Observations and Improvements, because our present astronomical Professors do not publish their Observations for the Improvement of Astronomy, and Benefit of this Nation: though they are paid, with the public Money, that we may see and taste the Fruit of their Industry. — Our late astronomical Professor, Dr. Edm. Halley, published an excellent Ephemerides, we are told, for the Use of our Royal Society, whereby our Astronomy must be then greatly improved. But (by our present astronomical Professors making an impenetrable Secret of their Observations and Improvements) this Nation is obliged to send to France for Ephemerides, published by M. De la Caille, Miraldi, and others there, for their Accuracy and Improvement; because our able astronomical Professors suffer themselves to be excelled herein by Foreigners, and these Sort of curious and useful Productions are now not encouraged among us. — Whereas if proper Encouragement was given us by our NOBILITY, or the ROYAL SOCIETY of this Nation, we would undertake to compute A ROYAL EPHEMERIDES (for many Years to come) to improve the Theory of the celestial Motions, and to excell in Usefulness, — Observations being, at the same Time, allowed us. Then we need not send our Money abroad for foreign Manufacture; having sufficient of our own.

EQUATION of MIDDLE TIME, between the Times of Forenoon and Afternoon Observations of the same Height of the SUN, (marked by the Clock or Watch) to the correct Noon by the SUN: Correspondent to the Increase or Decrease of Day, or of Sun's Declination, in that INTERVAL: Being the METHOD of corresponding ALTITUDES.

From Dec. 21, to June 22, N. S. Days increase; therefore Mid-Time, — Equation = Time of correct Noon } by the SUN.
From June 22, to Dec. 21, N. S. Days decrease; therefore Mid-Time, + Equation = Time of correct Noon }

N. S.	Lat. N. or S. 20°	Lat. N. or S. 30°	Lat. N. or S. 40°	Lat. N. or S. 50°	Lat. N. or S. 60° of the ☉
Dec.	Hrs. betw. Obser.	Hours between Observ.	Hours between Observ.	Hours between Observ.	Hours between Observ.
☉	9 8 7 6 5 4	9 8 7 6 5 4	10 9 8 7 6 5 4	10 9 8 7 6 5 4	10 9 8 7 6 5 4
N. S.	Sec Sec Sec Sec Sec Sec	Sec Sec Sec Sec Sec Sec	Sec Sec Sec Sec Sec Sec	Sec Sec Sec Sec Sec Sec	Sec Sec Sec Sec Sec Sec
23°	1 0 0 0 0 0	2 1 1 0 0 0	3 3 2 2 2 1	4 4 3 3 3 2	7 6 6 5 5 4
22	1 1 0 0 0 0	3 2 2 1 1 0	5 4 4 4 3 2	8 7 7 6 5 5	12 11 10 9 8 8
21	2 1 1 1 0 0	4 3 3 2 2 1	7 6 5 5 4 3	10 9 8 8 7 6	15 14 13 12 11 10
20	2 2 1 1 1 0	4 4 3 2 2 2	8 7 6 5 4 3	11 10 9 8 7 6	18 16 15 14 13 12
19	3 2 2 1 1 1	5 4 4 3 3 2	9 8 7 6 5 4	13 12 11 10 9 8	20 18 17 16 15 14
18	3 2 2 1 1 1	5 5 4 4 3 3	10 9 8 8 6 5	14 13 12 11 10 9	21 20 18 17 16 15
17	4 3 2 2 2 1	6 5 5 4 4 3	11 10 9 8 7 6	15 14 13 12 11 10	23 21 20 19 17 16
16	4 3 3 2 2 1	6 5 5 5 4 4	11 10 9 9 8 7	16 15 14 13 12 11	24 23 21 20 18 17
15	4 3 3 3 2 2	7 6 5 5 5 4	12 11 10 9 8 7	17 16 15 14 13 12	26 24 22 21 19 18
14	4 4 3 3 2 2	7 6 6 5 5 5	12 11 10 10 9 8	18 17 16 15 14 13	27 26 23 22 20 19
13	5 4 3 3 3 2	7 7 6 6 5 5	13 12 11 10 9 8	19 18 17 16 15 14	27 26 24 23 21 20
12	5 4 4 3 3 2	7 7 6 6 6 5	13 12 11 11 10 9	20 19 17 16 15 14	28 25 23 22 21 21
11	5 4 4 4 3 3	8 7 7 6 6 5	14 13 12 11 10 9	20 19 18 16 15 14	30 26 24 23 22 22
10	5 5 4 4 3 3	8 8 7 7 6 5	14 13 12 11 11 10	21 20 18 17 16 15	31 27 25 24 23 22
9	5 5 4 4 4 3	8 8 7 7 6 4	15 14 12 12 11 10	21 20 19 17 16 15	— 28 26 25 23 23
8	5 5 4 4 4 3	9 8 8 7 7 4	15 14 13 12 11 11	21 21 19 18 17 16	— 28 27 26 24 23
7	6 5 5 4 4 3	9 8 8 7 7 4	16 14 13 12 11 11	22 21 20 18 17 16	— 29 27 26 25 24
6	6 5 5 4 4 3	9 9 8 8 7 4	16 15 13 13 12 11	22 22 20 19 17 16	— 29 28 27 25 24
5	6 6 5 5 4 4	9 9 8 8 7 7	16 15 14 13 12 11	23 22 20 19 18 17	— 29 28 27 26 25
4	6 6 5 5 5 4	10 9 9 8 8 7	17 15 14 13 13 12	23 22 20 19 18 17	— 30 29 27 26 25
3	6 6 6 5 5 5	10 9 9 8 8 7	17 15 14 14 13 12	23 22 21 20 18 17	— 30 29 28 27 26
2	6 6 6 5 5 5	10 9 9 9 8 8	17 16 14 14 13 12	23 22 21 20 19 18	— 30 29 28 27 26
1	7 6 6 5 5 5	10 10 9 9 8 8	17 16 15 14 13 12	23 22 21 20 19 18	— 31 29 28 27 27
0	7 6 6 6 5 5	10 10 9 9 8 8	17 16 15 14 13 13	23 22 21 21 20 19	— 31 29 28 28 27
S. N.	7 6 6 6 5 5	10 10 10 9 9 8	16 15 14 14 13 13	22 21 21 20 19 18	— 29 28 28 27
0°	7 6 6 6 6 5	10 10 10 9 9 9	16 15 14 14 13 13	22 21 21 20 19 18	— 29 29 28 27
1	7 6 6 6 6 6	11 10 10 9 9 9	16 15 15 14 14 13	22 22 21 20 19 18	— 30 29 28 27
2	7 7 6 6 6 6	11 10 10 10 9 9	16 15 15 14 14 13	22 21 20 19 18	— 30 29 28 27
3	7 7 6 6 6 6	11 10 10 10 10 9	16 15 15 14 14 13	22 21 20 19 19	— 30 29 29 28
4	7 7 6 6 6 6	11 10 10 10 10 10	16 16 15 15 14 14	22 21 20 19 19	— 30 29 29 28
5	7 7 7 6 6 6	11 10 10 10 10 10	16 16 15 15 14 14	22 21 20 19 19	— 30 29 29 28
6	7 7 7 7 6 6	10 10 10 10 10 10	16 16 15 15 14 14	21 20 20 19 19	— 29 29 28 27
7	7 7 7 7 7 6	10 10 10 10 10 10	16 16 15 15 14 14	21 20 20 19 19	— 29 28 27 26
8	7 7 7 7 7 7	10 10 10 10 10 10	16 16 15 15 14 14	21 20 20 19 19	— 29 28 27 26
9	7 7 7 7 7 7	10 10 10 10 10 10	16 15 15 15 14 14	21 20 20 19 19	— 28 28 27 26
10	8 8 7 7 7 7	11 10 10 10 10 10	16 15 15 15 14 14	20 19 19 19 18	— 28 28 27 26
11	7 7 8 7 7 7	11 11 10 10 10 10	15 15 15 14 14 14	19 19 19 18 18	— 27 27 26 26
12	7 7 7 7 7 7	10 11 10 10 10 10	15 15 14 14 14 14	19 19 18 18 18	— 26 26 25 25
13	7 7 7 7 7 7	10 10 10 10 9 9	15 14 14 14 14 14	18 18 17 17 17	— 25 25 24 24
14	7 7 7 7 7 7	10 10 10 9 9 9	14 14 14 14 13 13	18 18 17 17 17	— 24 24 23 23
15	7 7 7 7 7 7	10 10 9 9 9 9	14 14 14 13 13 13	17 17 17 16 16	— 23 23 22 22
16	6 6 6 6 6 6	9 9 9 9 8 8	13 13 13 12 12 12	16 16 16 15 15	— 21 21 20 20
17	6 6 6 6 6 6	9 9 9 8 8 8	13 12 12 12 12 12	15 15 15 14 14	— 19 19 18 18
18	6 6 6 6 6 6	9 9 8 8 8 8	12 12 12 12 12 12	14 14 14 14 14	— 17 17 16 16
19	6 6 6 6 6 6	8 8 8 8 8 8	11 11 11 11 11 11	13 13 13 13 13	— 14 14 13 13
20	5 5 5 5 5 5	8 8 7 7 7 7	10 10 10 10 10 10	11 11 11 11 11	— 11 11 10 10
21	5 5 5 5 5 5	7 7 7 7 7 7	9 9 9 9 9 9	10 10 10 10 10	— 9 9 8 8
22	4 4 4 4 4 4	6 6 6 6 6 6	7 7 7 7 7 7	9 9 9 9 9 9	— 7 7 6 6
23	2 2 2 2 2 2	3 3 3 3 3 3	4 4 4 4 4 4	5 5 5 5 5 5	— 5 5 4 4

EXAMPLE. To find the Clock-Time of the correct Noon, by the SUN, for Latitude of Greenwich $51^{\circ} 30' 28''$, on 20th Jan. ($\odot$ 20° Declination South) from $\odot$'s Altitude observed $10^{\circ} 40'$, at $9^h 28^m 19^s$ Forenoon, and again at $2^h 38^m 45^s$ Afternoon, by the same Clock or Watch, by which a Clock or Watch is regulated to the Sun's Time?

	h	m	s
1st Observ. . .	9	28	19
2d Observ. . .	2	38	45
Sum . . .	5	10	26
Time between Observations.			
Time, 1st Observ. 12	3	32	
Hence the Clock or Watch may be kept going with the Sun, at 12.	12	3	19

Middle-Time found . . . 12 3 32
Lat. 50° . Under 5^h between Observations against 20° South Declination, Equation at Noon. }
— 13

For finding the Time of RISING and SETTING of the SUN and STARS.

SEMIDIURNAL or DURATION ARCS.

N.S.	North or South LATITUDE of the PLACE, correspondent to N. S. Declination of SUN or STAR.																
or *s Dec.	1°	2°	3°	4°	5°	6°	7°	8°	9°	10°	11°	12°	13°	14°	15°	16°	17°
N.S.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.
1	6 2	6 2	6 2	6 2	6 2	6 3	6 3	6 3	6 3	6 3	6 3	6 3	6 3	6 3	6 3	6 3	6 3
2	6 2	6 2	6 3	6 3	6 3	6 3	6 3	6 3	6 3	6 4	6 4	6 4	6 4	6 4	6 4	6 4	6 5
3	6 2	6 3	6 3	6 3	6 3	6 4	6 4	6 4	6 4	6 5	6 5	6 5	6 5	6 5	6 5	6 6	6 6
4	6 2	6 3	6 3	6 3	6 4	6 4	6 4	6 4	6 5	6 5	6 5	6 6	6 6	6 6	6 6	6 7	6 7
5	6 2	6 3	6 3	6 4	6 4	6 4	6 5	6 5	6 5	6 6	6 6	6 6	6 7	6 7	6 8	6 8	6 8
6	6 3	6 3	6 3	6 4	6 4	6 4	6 5	6 6	6 6	6 6	6 7	6 7	6 8	6 8	6 9	6 9	6 10
7	6 3	6 3	6 4	6 4	6 5	6 5	6 6	6 6	6 7	6 7	6 8	6 8	6 9	6 9	6 10	6 10	6 11
8	6 3	6 3	6 4	6 4	6 5	6 5	6 6	6 7	6 7	6 8	6 8	6 9	6 10	6 10	6 11	6 11	6 12
9	6 3	6 3	6 4	6 4	6 5	6 6	6 6	6 7	6 8	6 8	6 9	6 9	6 10	6 11	6 11	6 12	6 13
10	6 3	6 4	6 4	6 5	6 6	6 6	6 7	6 8	6 9	6 9	6 10	6 10	6 11	6 12	6 12	6 13	6 14
11	6 3	6 4	6 4	6 5	6 6	6 7	6 8	6 8	6 9	6 10	6 11	6 12	6 13	6 13	6 14	6 15	6 16
12	6 3	6 4	6 5	6 6	6 6	6 7	6 8	6 9	6 10	6 11	6 12	6 13	6 14	6 14	6 15	6 16	6 17
13	6 3	6 4	6 5	6 6	6 7	6 8	6 9	6 10	6 11	6 12	6 12	6 13	6 15	6 15	6 16	6 17	6 18
14	6 3	6 4	6 5	6 6	6 7	6 8	6 9	6 10	6 11	6 12	6 13	6 14	6 15	6 16	6 18	6 19	6 20
15	6 3	6 4	6 5	6 6	6 7	6 8	6 9	6 10	6 11	6 12	6 13	6 14	6 15	6 16	6 18	6 20	6 21
16	6 3	6 4	6 6	6 7	6 8	6 9	6 10	6 11	6 13	6 14	6 15	6 16	6 17	6 19	6 20	6 21	6 22
17	6 3	6 5	6 6	6 7	6 8	6 10	6 11	6 12	6 13	6 15	6 16	6 17	6 18	6 20	6 21	6 22	6 24
18	6 4	6 5	6 6	6 7	6 9	6 10	6 11	6 13	6 14	6 15	6 17	6 18	6 19	6 21	6 22	6 24	6 25
19	6 4	6 5	6 6	6 8	6 9	6 11	6 12	6 13	6 15	6 16	6 18	6 19	6 21	6 22	6 23	6 25	6 27
20	6 4	6 5	6 7	6 8	6 9	6 11	6 12	6 14	6 15	6 17	6 19	6 20	6 22	6 23	6 25	6 26	6 28
21	6 4	6 5	6 7	6 8	6 10	6 12	6 13	6 15	6 16	6 18	6 19	6 21	6 23	6 24	6 26	6 28	6 29
22	6 4	6 6	6 7	6 8	6 10	6 12	6 14	6 15	6 17	6 19	6 20	6 22	6 24	6 25	6 27	6 29	6 31
23	6 4	6 6	6 7	6 9	6 11	6 13	6 14	6 16	6 18	6 19	6 21	6 23	6 25	6 27	6 28	6 30	6 32
24	6 4	6 6	6 8	6 9	6 11	6 13	6 15	6 17	6 19	6 20	6 22	6 24	6 26	6 28	6 30	6 32	6 34
25	6 4	6 6	6 8	6 10	6 12	6 14	6 15	6 17	6 19	6 21	6 23	6 25	6 27	6 29	6 31	6 33	6 35
26	6 4	6 6	6 8	6 10	6 12	6 14	6 16	6 18	6 20	6 22	6 24	6 26	6 28	6 30	6 32	6 35	6 37
27	6 4	6 6	6 8	6 11	6 13	6 15	6 17	6 19	6 21	6 23	6 25	6 27	6 29	6 32	6 34	6 36	6 38
28	6 5	6 7	6 9	6 11	6 13	6 15	6 17	6 20	6 22	6 24	6 26	6 28	6 31	6 33	6 35	6 38	6 40
29	6 5	6 7	6 9	6 11	6 14	6 16	6 18	6 20	6 23	6 25	6 27	6 30	6 32	6 34	6 37	6 39	6 42
30	6 5	6 7	6 9	6 12	6 14	6 16	6 17	6 21	6 23	6 26	6 28	6 31	6 33	6 36	6 38	6 41	6 45
31	6 5	6 7	6 10	6 12	6 15	6 17	6 19	6 22	6 24	6 27	6 29	6 32	6 34	6 37	6 40	6 42	6 45
32	6 5	6 7	6 10	6 12	6 15	6 18	6 20	6 23	6 25	6 28	6 30	6 33	6 36	6 38	6 41	6 44	6 47
S. N.	1°	2°	3°	4°	5°	6°	7°	8°	9°	10°	11°	12°	13°	14°	15°	16°	17°
1	6 2	6 2	6 2	6 2	6 2	6 2	6 3	6 1	6 1	6 1	6 1	6 1	6 1	6 1	6 1	6 1	6 1
2	6 2	6 1	6 1	6 1	6 1	6 1	6 3	6 1	6 1	6 1	6 1	6 0	6 0	6 0	6 0	6 0	6 0
3	6 2	6 1	6 1	6 1	6 1	6 1	6 4	6 0	6 0	6 0	6 0	6 0	5 59	5 59	5 59	5 59	5 59
4	6 2	6 1	6 1	6 1	6 1	6 0	6 4	6 0	6 0	5 59	5 59	5 59	5 59	5 58	5 58	5 58	5 57
5	6 2	6 1	6 1	6 1	6 0	6 0	6 5	5 59	5 59	5 59	5 58	5 58	5 57	5 57	5 57	5 56	5 56
6	6 2	6 1	6 1	6 0	6 0	6 0	6 5	5 59	5 58	5 58	5 57	5 57	5 57	5 56	5 56	5 55	5 55
7	6 2	6 1	6 1	6 0	6 0	5 59	6 6	5 58	5 58	5 57	5 57	5 56	5 56	5 55	5 55	5 54	5 54
8	6 1	6 1	6 0	6 0	5 59	5 59	6 6	5 57	5 57	5 56	5 56	5 55	5 55	5 54	5 54	5 53	5 52
9	6 1	6 1	6 0	6 0	5 59	5 58	6 7	5 56	5 57	5 56	5 55	5 55	5 54	5 54	5 53	5 52	5 51
10	6 1	6 1	6 0	5 59	5 59	5 58	6 7	5 56	5 56	5 55	5 54	5 54	5 53	5 53	5 52	5 51	5 50
11	6 1	6 1	6 0	5 59	5 58	5 57	6 8	5 56	5 55	5 54	5 54	5 53	5 53	5 52	5 51	5 50	5 49
12	6 1	6 1	6 0	5 59	5 58	5 57	6 8	5 55	5 54	5 54	5 53	5 53	5 52	5 51	5 50	5 49	5 48
13	6 1	6 0	5 59	5 58	5 57	5 57	6 9	5 55	5 54	5 53	5 53	5 52	5 51	5 50	5 49	5 48	5 47
14	6 1	6 0	5 59	5 58	5 57	5 56	6 9	5 54	5 53	5 52	5 51	5 50	5 49	5 48	5 47	5 46	5 45
15	6 1	6 0	5 59	5 58	5 57	5 56	6 10	5 54	5 52	5 51	5 50	5 49	5 48	5 47	5 46	5 45	5 44
16	6 1	6 0	5 59	5 58	5 56	5 55	6 10	5 53	5 52	5 51	5 49	5 48	5 47	5 46	5 45	5 43	5 42
17	6 1	6 0	5 59	5 57	5 56	5 55	6 11	5 52	5 51	5 50	5 49	5 47	5 46	5 45	5 43	5 42	5 41
18	6 1	6 0	5 58	5 57	5 56	5 54	6 11	5 52	5 50	5 49	5 48	5 46	5 45	5 44	5 42	5 41	5 40
19	6 1	5 59	5 58	5 57	5 55	5 54	6 12	5 51	5 50	5 48	5 47	5 45	5 44	5 43	5 41	5 40	5 38
20	6 1	5 59	5 58	5 56	5 55	5 53	6 12	5 51	5 49	5 48	5 46	5 45	5 43	5 41	5 40	5 38	5 37
21	6 1	5 59	5 58	5 56	5 55	5 53	6 13	5 50	5 48	5 47	5 45	5 44	5 42	5 40	5 39	5 37	5 35
22	6 1	5 59	5 57	5 56	5 54	5 53	6 14	5 49	5 48	5 46	5 44	5 43	5 41	5 39	5 38	5 36	5 34
23	6 1	5 59	5 57	5 55	5 54	5 52	6 14	5 49	5 47	5 45	5 43	5 42	5 40	5 38	5 36	5 34	5 33
24	6 1	5 59	5 57	5 55	5 53	5 52	6 15	5 48	5 46	5 44	5 42	5 41	5 39	5 37	5 35	5 33	5 31
25	6 1	5 59	5 57	5 55	5 53	5 51	6 15	5 47	5 45	5 43	5 42	5 40	5 38	5 36	5 34	5 32	5 30
26	6 0	5 58	5 56	5 55	5 53	5 51	6 16	5 47	5 45	5 43	5 41	5 39	5 37	5 34	5 32	5 30	5 28
27	6 0	5 58	5 56	5 54	5 52	5 50	6 17	5 46	5 44	5 42	5 40	5 38	5 36	5 33	5 31	5 29	5 26
28	6 0	5 58	5 56	5 54	5 52	5 50	6 17	5 45	5 43	5 41	5 39	5 37	5 34	5 32	5 30	5 27	5 25
29	6 0	5 58	5 56	5 54	5 51	5 49	6 18	5 45	5 42	5 40	5 38	5 36	5 33	5 31	5 28	5 26	5 24
30	6 0	5 58	5 55	5 53	5 51	5 49	6 19	5 44	5 41	5 39	5 37	5 35	5 32	5 29	5 27	5 24	5 22
31	6 0	5 58	5 55	5 53	5 50	5 48	6 19	5 43	5 41	5 38	5 36	5 34	5 31	5 28	5 26	5 23	5 20
32	6 0	5 57	5 55	5 52	5 50	5 47	6 20	5 43	5 40	5 37	5 35	5 32	5 29	5 27	5 24	5 21	5 18

For

SEMIDIURNAL or DURATION ARCS.

North or South LATITUDE of the PLACE, correspondent to N. S. Declination of SUN or STAR.

For

FOR finding the RISING and SETTING of the SUN and STARS.

SEMIDIURNAL or DURATION ARCS.

North or South LATITUDE of the PLACE, correspondent to N. S. Declination of SUN or STAR.

N.S.	35°	36°	37°	38°	39°	40°	41°	42°	43°	44°	45°	46°	47°	48°	49°	50°	51°
Dec.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.
1	6 5	6 6	6 6	6 6	6 6	6 6	6 6	6 6	6 7	6 7	6 7	6 7	6 7	6 8	6 8	6 8	6 8
2	6 8	6 8	6 9	6 9	6 9	6 9	6 10	6 10	6 10	6 11	6 11	6 11	6 12	6 12	6 12	6 13	6 13
3	6 11	6 11	6 12	6 12	6 12	6 13	6 13	6 14	6 14	6 15	6 15	6 15	6 16	6 17	6 17	6 18	6 18
4	6 14	6 14	6 15	6 15	6 16	6 16	6 17	6 17	6 18	6 18	6 19	6 20	6 20	6 21	6 22	6 22	6 22
5	6 17	6 17	6 18	6 18	6 19	6 20	6 20	6 21	6 22	6 22	6 23	6 24	6 25	6 25	6 26	6 27	6 27
6	6 19	6 20	6 21	6 22	6 22	6 23	6 24	6 25	6 25	6 26	6 27	6 28	6 29	6 30	6 31	6 32	6 33
7	6 22	6 24	6 24	6 25	6 26	6 26	6 27	6 28	6 29	6 30	6 31	6 32	6 33	6 34	6 36	6 37	6 38
8	6 25	6 26	6 27	6 28	6 29	6 30	6 31	6 32	6 33	6 34	6 35	6 37	6 38	6 39	6 41	6 42	6 43
9	6 28	6 29	6 30	6 31	6 32	6 33	6 35	6 36	6 37	6 38	6 40	6 41	6 42	6 44	6 45	6 47	6 48
10	6 31	6 32	6 33	6 34	6 36	6 37	6 38	6 39	6 41	6 42	6 44	6 45	6 47	6 48	6 50	6 52	6 54
11	6 34	6 35	6 36	6 38	6 39	6 40	6 42	6 43	6 45	6 46	6 48	6 50	6 51	6 53	6 55	6 57	6 59
12	6 37	6 38	6 40	6 41	6 42	6 44	6 45	6 47	6 49	6 50	6 52	6 55	6 56	6 58	7 0	7 2	7 4
13	6 40	6 41	6 43	6 44	6 46	6 48	6 49	6 51	6 53	6 55	6 57	6 59	7 1	7 3	7 5	7 7	7 10
14	6 43	6 44	6 46	6 48	6 49	6 51	6 53	6 55	6 57	6 59	7 1	7 3	7 5	7 8	7 10	7 13	7 15
15	6 46	6 48	6 49	6 51	6 53	6 55	6 57	6 59	7 1	7 3	7 5	7 8	7 10	7 13	7 15	7 18	7 21
16	6 49	6 51	6 53	6 55	6 57	6 59	7 1	7 3	7 5	7 7	7 10	7 12	7 15	7 18	7 21	7 24	7 27
17	6 52	6 54	6 56	6 58	7 0	7 2	7 5	7 7	7 9	7 12	7 14	7 17	7 20	7 23	7 26	7 29	7 33
18	6 55	6 57	7 0	7 2	7 4	7 6	7 9	7 11	7 14	7 16	7 19	7 22	7 25	7 28	7 31	7 35	7 38
19	6 59	7 1	7 3	7 5	7 8	7 10	7 13	7 15	7 18	7 21	7 24	7 27	7 30	7 34	7 37	7 41	7 45
20	7 2	7 4	7 7	7 9	7 12	7 14	7 17	7 20	7 23	7 26	7 29	7 32	7 35	7 39	7 43	7 47	7 51
21	7 5	7 8	7 10	7 13	7 15	7 18	7 21	7 24	7 27	7 30	7 34	7 37	7 41	7 45	7 49	7 53	7 57
22	7 9	7 11	7 13	7 17	7 19	7 22	7 25	7 29	7 32	7 35	7 39	7 43	7 46	7 50	7 55	7 59	8 4
23	7 12	7 15	7 18	7 21	7 24	7 27	7 30	7 33	7 37	7 40	7 44	7 48	7 52	7 56	8 1	8 6	8 11
24	7 16	7 19	7 21	7 25	7 28	7 31	7 34	7 38	7 42	7 45	7 49	7 54	7 58	8 3	8 7	8 12	8 18
25	7 19	7 22	7 25	7 29	7 32	7 35	7 39	7 43	7 47	7 51	7 55	7 59	8 4	8 9	8 14	8 19	8 25
26	7 23	7 26	7 29	7 33	7 36	7 40	7 44	7 48	7 52	7 56	8 1	8 5	8 10	8 15	8 21	8 27	8 33
27	7 27	7 30	7 34	7 37	7 41	7 45	7 49	7 53	7 57	8 2	8 6	8 12	8 17	8 22	8 28	8 34	8 41
28	7 31	7 34	7 38	7 42	7 45	7 49	7 54	7 58	8 3	8 7	8 12	8 18	8 23	8 29	8 35	8 42	8 49
29	7 35	7 38	7 42	7 46	7 50	7 54	7 59	8 4	8 9	8 13	8 19	8 24	8 30	8 37	8 43	8 50	8 58
30	7 39	7 43	7 47	7 51	7 55	8 0	8 5	8 9	8 14	8 20	8 25	8 31	8 38	8 44	8 52	8 59	9 8
31	7 43	7 47	7 51	7 56	8 0	8 5	8 10	8 15	8 20	8 26	8 32	8 38	8 45	8 52	9 0	9 9	9 18
32	7 47	7 51	7 56	8 1	8 5	8 10	8 16	8 21	8 27	8 33	8 39	8 46	8 53	9 1	9 9	9 19	9 28
S.N.	6 0	6 0	6 0	6 0	5 59	5 59	5 59	5 59	5 59	5 59	5 59	5 59	5 59	5 59	5 59	5 59	5 58
1	5 57	5 57	5 57	5 56	5 56	5 56	5 56	5 56	5 55	5 55	5 55	5 55	5 55	5 54	5 54	5 54	5 53
2	5 54	5 54	5 54	5 53	5 53	5 53	5 52	5 52	5 52	5 51	5 51	5 51	5 50	5 50	5 49	5 49	5 49
3	5 51	5 51	5 51	5 50	5 50	5 49	5 49	5 48	5 48	5 47	5 47	5 46	5 46	5 45	5 45	5 44	5 44
4	5 49	5 48	5 48	5 47	5 46	5 46	5 45	5 45	5 44	5 44	5 43	5 42	5 42	5 41	5 40	5 39	5 39
5	5 46	5 45	5 44	5 44	5 43	5 43	5 42	5 41	5 40	5 39	5 38	5 37	5 36	5 35	5 35	5 34	5 34
6	5 43	5 42	5 41	5 41	5 40	5 39	5 38	5 37	5 36	5 35	5 34	5 33	5 32	5 31	5 30	5 29	5 29
7	5 40	5 39	5 38	5 37	5 37	5 36	5 35	5 34	5 33	5 32	5 31	5 30	5 28	5 27	5 26	5 25	5 25
8	5 37	5 36	5 35	5 34	5 33	5 32	5 31	5 30	5 29	5 28	5 27	5 25	5 24	5 23	5 21	5 20	5 18
9	5 34	5 33	5 32	5 31	5 30	5 29	5 28	5 26	5 25	5 24	5 22	5 21	5 20	5 18	5 17	5 15	5 13
10	5 31	5 30	5 29	5 28	5 27	5 25	5 24	5 23	5 21	5 20	5 18	5 17	5 15	5 13	5 12	5 10	5 8
11	5 28	5 27	5 26	5 25	5 23	5 22	5 20	5 19	5 17	5 16	5 14	5 12	5 11	5 9	5 7	5 5	5 3
12	5 25	5 24	5 23	5 21	5 20	5 18	5 17	5 15	5 13	5 12	5 10	5 8	5 6	5 4	5 2	5 0	4 57
13	5 22	5 21	5 19	5 18	5 16	5 15	5 13	5 11	5 9	5 7	5 5	5 3	5 1	4 59	4 57	4 54	4 52
14	5 19	5 18	5 16	5 14	5 13	5 11	5 9	5 7	5 5	5 3	5 1	4 59	4 57	4 54	4 52	4 49	4 46
15	5 16	5 15	5 13	5 11	5 9	5 7	5 5	5 3	5 1	4 59	4 57	4 54	4 52	4 49	4 46	4 45	4 41
16	5 13	5 11	5 10	5 8	5 6	5 4	5 2	4 59	4 57	4 55	4 53	4 50	4 47	4 44	4 41	4 38	4 35
17	5 10	5 8	5 6	5 4	5 2	4 59	4 57	4 55	4 53	4 50	4 47	4 45	4 42	4 39	4 36	4 33	4 29
18	5 7	5 5	5 3	5 0	4 58	4 56	4 53	4 51	4 48	4 46	4 43	4 40	4 37	4 34	4 30	4 27	4 23
19	5 4	5 2	4 59	4 57	4 54	4 52	4 49	4 47	4 44	4 41	4 38	4 35	4 32	4 28	4 25	4 21	4 17
20	5 1	4 58	4 56	4 53	4 51	4 48	4 45	4 42	4 39	4 36	4 33	4 30	4 26	4 23	4 19	4 15	4 11
21	4 57	4 55	4 52	4 49	4 47	4 44	4 41	4 38	4 35	4 32	4 28	4 25	4 21	4 17	4 13	4 9	4 4
22	4 54	4 51	4 49	4 46	4 43	4 40	4 37	4 33	4 30	4 27	4 23	4 19	4 15	4 11	4 7	4 3	3 58
23	4 50	4 48	4 45	4 42	4 39	4 35	4 32	4 29	4 25	4 22	4 18	4 14	4 10	4 5	4 1	3 56	3 51
24	4 47	4 44	4 41	4 38	4 34	4 31	4 28	4 24	4 20	4 17	4 13	4 8	4 4	3 59	3 54	3 49	3 44
25	4 43	4 40	4 36	4 34	4 30	4 27	4 23	4 19	4 15	4 11	4 7	4 3	3 58	3 53	3 48	3 42	3 37
26	4 39	4 36	4 33	4 29	4 26	4 22	4 18	4 14	4 10	4 6	4 1	3 57	3 52	3 46	3 41	3 35	3 29
27	4 36	4 32	4 29	4 25	4 21	4 17	4 13	4 9	4 5	4 0	3 55	3 50	3 45	3 40	3 34	3 28	3 21
28	4 32	4 28	4 25	4 21	4 17	4 13	4 8	4 4	3 59	3 54	3 49	3 44	3 38	3 33	3 26	3 20	3 12
29	4 28	4 24	4 20	4 16	4 12	4 8	4 3	3 59	3 54	3 48	3 43	3 37	3 31	3 25	3 18	3 11	3 4
30	4 24	4 20	4 15	4 12	4 7	4 8	3 58	3 53	3 48	3 42	3 37	3 31	3 24	3 17	3 10	3 3	2 54
31	4 19	4 15	4 11	4 7	4 2	3 57	3 52	3 47	3 42	3 36	3 30	3 23	3 17	3 9	3 2	2 53	2 44
32																	

FOR

FOR finding the RISING and SETTING of the SUN and STARS.

SEMI-DIURNAL or DURATION ARCS.

North or South LATITUDE of the PLACE, correspondent to N. S. Declination of SUN or STAR.

N.S. Dec. or Star.	52°	53°	54°	55°	56°	57°	58°	59°	60°	61°	62°	63°	64°	65°	66°
N.S.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.	H. M.
1	6 9	6 9	6 9	6 9	6 10	6 10	6 10	6 11	6 11	6 12	6 13	6 13	6 13	6 14	6 14
2	6 14	6 14	6 15	6 15	6 16	6 16	6 17	6 17	6 18	6 19	6 20	6 20	6 21	6 22	6 23
3	6 19	6 19	6 20	6 21	6 22	6 22	6 23	6 24	6 25	6 26	6 27	6 28	6 30	6 31	6 32
4	6 24	6 25	6 26	6 27	6 28	6 29	6 30	6 31	6 32	6 33	6 35	6 36	6 38	6 40	6 41
5	6 29	6 30	6 31	6 32	6 34	6 35	6 36	6 38	6 39	6 41	6 42	6 44	6 46	6 48	6 51
6	6 34	6 36	6 37	6 38	6 40	6 41	6 43	6 44	6 46	6 48	6 50	6 52	6 55	6 57	7 0
7	6 40	6 41	6 43	6 44	6 46	6 48	6 49	6 51	6 53	9 55	6 58	7 1	7 3	7 6	7 10
8	6 45	6 47	6 48	6 50	6 52	6 54	6 56	6 58	7 1	7 3	7 6	7 9	7 12	7 15	7 19
9	6 50	6 52	6 54	6 56	6 58	7 1	7 3	7 5	7 8	7 11	7 14	7 17	7 21	7 25	7 29
10	6 56	6 58	7 0	7 2	7 5	7 7	7 10	7 13	7 16	7 19	7 22	7 26	7 30	7 34	7 39
11	7 1	7 3	7 6	7 8	7 11	7 14	7 17	7 20	7 23	7 27	7 31	7 35	7 39	7 44	7 49
12	7 7	7 9	7 12	7 15	7 18	7 21	7 24	7 27	7 31	7 35	7 39	7 44	7 49	7 54	8 0
13	7 12	7 15	7 18	7 21	7 24	7 28	7 31	7 35	7 39	7 43	7 48	7 53	7 59	8 5	8 11
14	7 18	7 21	7 24	7 28	7 31	7 35	7 39	7 43	7 47	7 52	7 57	8 3	8 9	8 15	8 23
15	7 24	7 27	7 31	7 34	7 39	7 42	7 46	7 51	7 56	8 1	8 6	8 13	8 19	8 27	8 35
16	7 30	7 33	7 37	7 41	7 45	7 49	7 54	7 59	8 4	8 10	8 16	8 23	8 30	8 38	8 48
17	7 36	7 40	7 44	7 48	7 52	7 57	8 2	8 7	8 13	8 19	8 26	8 34	8 42	8 51	9 1
18	7 42	7 46	7 51	7 55	8 0	8 5	8 10	8 16	8 22	8 29	8 37	8 45	8 54	9 4	9 16
19	7 49	7 53	7 58	8 2	8 7	8 13	8 19	8 25	8 32	8 40	8 48	8 57	9 7	9 18	9 32
20	7 55	8 0	8 5	8 10	8 15	8 21	2 28	8 35	8 42	8 50	8 59	9 10	9 21	9 34	9 49
21	8 2	8 7	8 12	8 18	8 24	8 30	8 37	8 45	8 53	9 2	9 12	9 23	9 37	9 51	10 10
22	8 9	8 14	8 20	8 26	8 32	8 39	8 47	8 55	9 4	9 14	9 25	9 38	9 53	10 12	10 35
23	8 16	8 22	8 28	8 34	8 41	8 49	8 57	9 6	9 16	9 27	9 40	9 55	10 13	10 36	11 12
24	8 24	8 30	8 36	8 43	8 51	8 59	9 8	9 18	9 29	9 42	9 57	10 15	10 38	11 13	
25	8 31	8 38	8 45	8 53	9 1	9 10	9 20	9 31	9 44	9 58	10 16	10 39	11 14		
26	8 39	8 47	8 54	9 2	9 11	9 21	9 33	9 45	10 0	10 17	10 40	11 14			
27	8 48	8 56	9 4	9 13	9 23	9 34	9 46	10 1	10 18	10 41	11 14				
28	8 57	9 5	9 14	9 24	9 35	9 48	10 2	10 19	10 42	11 15					
29	9 6	9 14	9 25	9 36	9 49	10 3	10 20	10 42	11 16						
30	9 17	9 26	9 38	9 50	10 4	10 21	10 43	11 16							
31	9 28	9 38	9 51	10 5	10 22	10 44	11 17								
32	9 39	9 52	10 6	10 23	10 44	11 17									

S.N.	52°	53°	54°	55°	56°	57°	58°	59°	60°	61°	62°	63°	64°	65°	66°
1	5 58	5 58	5 58	5 58	5 58	5 58	5 58	5 57	5 57	5 57	5 57	5 57	5 57	5 56	5 56
2	5 53	5 53	5 53	5 52	5 52	5 52	5 51	5 51	5 50	5 50	5 49	5 49	5 48	5 48	5 47
3	5 48	5 48	5 47	5 47	5 46	5 45	5 45	5 44	5 43	5 43	5 42	5 41	5 40	5 39	5 38
4	5 43	5 42	5 42	5 41	5 40	5 39	5 38	5 37	5 36	5 35	5 34	5 33	5 32	5 31	5 29
5	5 38	5 37	5 36	5 35	5 34	5 33	5 32	5 31	5 29	5 28	5 27	5 25	5 24	5 22	5 20
6	5 33	5 31	5 30	5 29	5 28	5 27	5 25	5 24	5 22	5 21	5 19	5 17	5 15	5 13	5 11
7	5 27	5 26	5 25	5 23	5 22	5 20	5 19	5 17	5 15	5 13	5 11	5 9	5 7	5 4	5 1
8	5 22	5 21	5 19	5 17	5 16	5 14	5 12	5 10	5 8	5 6	5 3	5 1	4 58	4 55	4 52
9	5 17	5 16	5 13	5 12	5 10	5 8	5 5	5 3	5 1	4 58	4 55	4 53	4 49	4 46	4 42
10	5 11	5 10	5 8	5 5	5 3	5 1	4 59	4 56	4 53	4 50	4 47	4 44	4 40	4 37	4 32
11	5 6	5 4	5 2	4 59	4 57	4 54	4 52	4 49	4 46	4 43	4 39	4 35	4 31	4 27	4 22
12	5 0	4 58	4 56	4 53	4 51	4 48	4 45	4 42	4 38	4 35	4 31	4 27	4 22	4 17	4 12
13	4 55	4 52	4 50	4 47	4 44	4 41	4 38	4 34	4 30	4 26	4 22	4 18	4 13	4 7	4 1
14	4 49	4 47	4 44	4 41	4 37	4 34	4 30	4 27	4 23	4 18	4 13	4 8	4 3	3 56	3 50
15	4 44	4 41	4 37	4 34	4 31	4 27	4 23	4 19	4 14	4 9	4 4	3 59	3 53	3 46	3 39
16	4 38	4 34	4 31	4 27	4 24	4 20	4 15	4 11	4 6	4 1	3 55	3 49	3 42	3 35	3 27
17	4 32	4 28	4 23	4 21	4 17	4 12	4 8	4 3	3 57	3 52	3 45	3 39	3 31	3 23	3 14
18	4 26	4 22	4 18	4 14	4 9	4 5	4 0	3 54	3 48	3 42	3 35	3 28	3 20	3 11	3 0
19	4 19	4 15	4 11	4 7	4 2	3 56	3 51	3 45	3 39	3 32	3 25	3 17	3 8	2 58	2 46
20	4 13	4 9	4 4	3 59	3 54	3 49	3 43	3 36	3 29	3 22	3 14	3 5	2 55	2 43	2 30
21	4 6	4 2	3 57	3 52	3 46	3 40	3 34	3 27	3 19	3 11	3 2	2 52	2 41	2 28	2 12
22	4 0	3 55	3 50	3 44	3 38	3 31	3 24	3 17	3 9	3 0	2 50	2 38	2 25	2 10	1 52
23	3 53	3 47	3 42	3 36	3 29	3 23	3 15	3 6	2 57	2 47	2 36	2 23	2 8	1 50	1 27
24	3 46	3 40	3 34	3 27	3 20	3 13	3 5	2 55	2 45	2 34	2 21	2 7	1 49	1 26	
25	3 38	3 32	3 25	3 18	3 11	3 3	2 53	2 43	2 32	2 20	2 5	1 47	1 25		
26	3 30	3 24	3 17	3 9	3 1	2 52	2 42	2 31	2 18	2 3	1 46	1 23			
27	3 22	3 15	3 8	2 59	2 50	2 40	2 29	2 16	2 2	1 45	1 22				
28	3 14	3 6	2 58	2 49	2 38	2 28	2 15	2 1	1 43	1 21					
29	3 5	2 56	2 47	2 37	2 26	2 14	2 0	1 42							
30	2 55	2 46	2 36	2 25	2 13	1 58	1 41	1 20							
31	2 45	2 35	2 24	2 12	1 57	1 41	1 19								
32	2 34	2 23	2 11	1 57	1 40	1 18									

For finding the VARIATION of the COMPASS.

AMPLITUDE of RISING and SETTING, from EAST and WEST.

N. S.		North or South LATITUDE of the PLACE, correspondent to N. S. Declination of SUN or STAR.														
Declin. Sun or Star.	N. S.	1°	2°	3°	4°	5°	6°	7°	8°	9°	10°	11°	12°	13°	14°	15°
1	N. S.	0 1	0 1	0 1	0 1	0 1	0 1	0 1	0 1	0 1	0 1	0 1	0 1	0 1	0 1	0 1
2		1 0	1 1	1 2	1 2	1 3	1 4	1 4	1 5	1 6	1 7	1 7	1 8	1 9	1 10	1 11
3		2 0	2 1	2 2	2 2	2 3	2 4	2 5	2 6	2 6	2 7	2 8	2 9	2 10	2 12	2 13
4		3 0	3 1	3 2	3 2	3 3	3 4	3 5	3 6	3 7	3 8	3 9	3 10	3 12	3 13	3 15
5		4 0	4 1	4 2	4 3	4 4	4 5	4 6	4 7	4 8	4 9	4 11	4 12	4 14	4 15	4 17
6		5 0	5 1	5 2	5 3	5 4	5 5	5 6	5 8	5 9	5 10	5 12	5 14	5 15	5 17	5 19
7		6 1	6 2	6 3	6 4	6 5	6 6	6 7	6 8	6 10	6 11	6 13	6 15	6 17	6 19	6 21
8		7 1	7 2	7 3	7 4	7 5	7 6	7 7	7 9	7 10	7 12	7 14	7 16	7 19	7 21	7 24
9		8 1	8 2	8 3	8 4	8 5	8 6	8 8	8 9	8 11	8 13	8 15	8 18	8 20	8 23	8 26
10		9 1	9 2	9 3	9 4	9 5	9 6	9 8	9 10	9 12	9 14	9 16	9 19	9 22	9 25	9 28
11		10 1	10 2	10 3	10 4	10 5	10 7	10 8	10 10	10 13	10 15	10 18	10 20	10 23	10 27	10 30
12		11 1	11 2	11 3	11 4	11 5	11 7	11 9	11 11	11 13	11 16	11 19	11 22	11 25	11 28	11 32
13		12 1	12 2	12 3	12 4	12 5	12 7	12 9	12 12	12 14	12 17	12 20	12 23	12 27	12 30	12 34
14		13 1	13 2	13 3	13 4	13 5	13 7	13 10	13 12	13 15	13 18	13 21	13 25	13 29	13 32	13 37
15		14 1	14 2	14 3	14 4	14 5	14 7	14 10	14 13	14 16	14 19	14 22	14 26	14 30	14 34	14 39
16		15 1	15 2	15 3	15 4	15 5	15 7	15 11	15 14	15 17	15 20	15 24	15 28	15 32	15 36	15 42
17		16 1	16 2	16 3	16 4	16 5	16 7	16 11	16 14	16 17	16 21	16 25	16 29	16 34	16 39	16 44
18		17 1	17 2	17 3	17 4	17 5	17 7	17 12	17 15	17 18	17 22	17 26	17 30	17 35	17 41	17 46
19		18 1	18 2	18 3	18 4	18 5	18 7	18 13	18 16	18 19	18 23	18 27	18 32	18 37	18 43	18 48
20		19 1	19 2	19 3	19 4	19 5	19 7	19 13	19 16	19 20	19 24	19 28	19 34	19 39	19 45	19 51
21		20 1	20 2	20 3	20 4	20 5	20 7	20 14	20 17	20 21	20 25	20 30	20 35	20 41	20 47	20 53
22		21 1	21 2	21 3	21 4	21 5	21 7	21 14	21 18	21 22	21 26	21 31	21 37	21 43	21 49	21 56
23		22 1	22 2	22 3	22 4	22 5	22 7	22 15	22 19	22 23	22 28	22 33	22 38	22 45	22 52	22 59
24		23 1	23 2	23 3	23 4	23 5	23 7	23 15	23 19	23 24	23 29	23 34	23 40	23 47	23 54	24 1
25		24 1	24 2	24 3	24 4	24 5	24 7	24 16	24 20	24 25	24 30	24 36	24 42	24 49	24 56	25 4
26		25 1	25 2	25 3	25 4	25 5	25 7	25 16	25 21	25 26	25 31	25 37	25 43	25 51	25 58	26 6
27		26 1	26 2	26 3	26 4	26 5	26 7	26 17	26 22	26 27	26 32	26 38	26 45	26 53	27 0	27 9
28		27 1	27 2	27 3	27 4	27 5	27 7	27 17	27 22	27 28	27 34	27 40	27 47	27 55	28 3	28 12
29		28 1	28 2	28 3	28 4	28 5	28 7	28 18	28 23	28 29	28 35	28 41	28 49	28 57	29 5	29 15
30		29 1	29 2	29 3	29 4	29 5	29 7	29 19	29 24	29 30	29 36	29 43	29 51	29 59	30 8	30 18
31		30 1	30 2	30 3	30 4	30 5	30 7	30 19	30 25	30 31	30 38	30 44	30 52	31 1	31 10	31 20
32		31 1	31 2	31 3	31 4	31 5	31 7	31 20	31 26	31 32	31 39	31 46	31 54	32 3	32 13	32 23
33		32 1	32 2	32 3	32 4	32 5	32 7	32 21	32 26	32 33	32 40	32 48	32 56	33 6	33 16	33 27
1	S. N.	0 59	0 59	0 58	0 58	0 57	0 57	0 56	0 56	0 56	0 55	0 55	0 54	0 54	0 53	0 53
2		1 59	1 59	1 58	1 58	1 57	1 57	1 57	1 56	1 56	1 56	1 56	1 56	1 56	1 56	1 56
3		2 59	2 59	2 58	2 58	2 58	2 57	2 57	2 57	2 57	2 57	2 57	2 57	2 57	2 57	2 58
4		3 59	3 59	3 58	3 58	3 58	3 58	3 58	3 58	3 58	3 58	3 58	3 58	3 59	3 59	4 0
5		4 59	4 59	4 58	4 58	4 58	4 58	4 58	4 58	4 59	4 59	4 59	5 0	5 1	5 1	5 2
6		5 59	5 59	5 59	5 59	5 59	5 59	5 59	5 59	5 59	6 0	6 1	6 1	6 2	6 3	6 4
7		6 59	6 59	6 59	6 59	6 59	6 59	6 59	6 59	6 59	7 0	7 1	7 2	7 3	7 4	7 6
8		7 59	7 59	7 59	7 59	7 59	8 0	8 0	8 0	8 1	8 2	8 3	8 4	8 5	8 7	8 8
9		8 59	8 59	8 59	8 59	8 59	9 0	9 0	9 1	9 2	9 3	9 4	9 5	9 7	9 8	9 10
10		9 59	9 59	9 59	9 59	9 59	10 0	10 0	10 1	10 2	10 4	10 5	10 7	10 8	10 10	10 13
11		10 59	10 59	10 59	10 59	11 0	11 0	11 1	11 2	11 3	11 4	11 6	11 8	11 10	11 12	11 15
12		11 59	11 59	11 59	11 59	12 0	12 1	12 1	12 3	12 4	12 5	12 7	12 9	12 12	12 14	12 17
13		12 59	12 59	12 59	12 59	13 0	13 1	13 2	13 3	13 5	13 6	13 8	13 11	13 13	13 16	13 19
14		13 59	13 59	13 59	14 0	14 1	14 1	14 2	14 4	14 6	14 7	14 10	14 12	14 15	14 18	14 21
15		14 59	14 59	14 59	15 0	15 1	15 2	15 3	15 4	15 6	15 8	15 11	15 13	15 17	15 20	15 24
16		15 59	15 59	15 59	16 0	16 1	16 2	16 3	16 5	16 7	16 9	16 12	16 15	16 18	16 22	16 26
17		16 59	16 59	16 59	17 0	17 1	17 2	17 4	17 6	17 8	17 10	17 13	17 16	17 20	17 24	17 28
18		17 59	17 59	17 59	18 0	18 1	18 2	18 3	18 4	18 6	18 9	18 11	18 14	18 18	18 22	18 30
19		18 59	18 59	18 59	19 0	19 1	19 2	19 3	19 5	19 7	19 9	19 12	19 15	19 19	19 23	19 33
20		19 59	19 59	19 59	20 0	20 1	20 2	20 3	20 5	20 8	20 10	20 13	20 17	20 21	20 25	20 35
21		20 59	20 59	20 59	21 0	21 1	21 2	21 4	21 6	21 8	21 11	21 14	21 18	21 22	21 27	21 38
22		21 59	21 59	21 59	22 0	22 1	22 2	22 4	22 6	22 9	22 12	22 15	22 19	22 24	22 29	22 40
23		22 59	22 59	22 59	23 0	23 1	23 2	23 4	23 7	23 9	23 13	23 16	23 21	23 25	23 30	23 42
24		23 59	23 59	23 59	24 0	24 1	24 3	24 5	24 7	24 10	24 13	24 17	24 22	24 27	24 32	24 45
25		24 59	24 59	24 59	25 0	25 1	25 3	25 5	25 8	25 11	25 14	25 18	25 23	25 28	25 34	25 47
26		25 59	25 59	25 59	26 0	26 1	26 3	26 5	26 8	26 11	26 15	26 20	26 24	26 30	26 36	26 50
27		26 59	26 59	26 59	27 0	27 1	27 2	27 4	27 7	27 10	27 14	27 19	27 23	27 29	27 35	27 50
28		27 59	27 59	27 59	28 0	28 1	28 2	28 4	28 7	28 10	28 14	28 19	28 23	28 30	28 37	28 52
29		28 59	28 59	28 59	29 0	29 1	29 2	29 4	29 7	29 10	29 14	29 19	29 23	29 30	29 38	29 53
30		29 59	29 59	29 59	30 0	30 1	30 2	30 4	30 7	30 10	30 14	30 19	30 23	30 30	30 38	30 53
31		30 59	30 59	30 59	31 0	31 1	31 2	31 4	31 7	31 10	31 14	31 19	31 23	31 30	31 38	31 53
32		31 59	31 59	31 59	32 0	32 1	32 2	32 4	32 7	32 10	32 14	32 19	32 23	32 30	32 38	32 53

FOR

FOR finding the VARIATION of the COMPASS.

AMPLITUDE of RISING and SETTING, from EAST and WEST.

N. S.		North or South LATITUDE of the PLACE, correspondent to N. S. Declination of SUN or STAR.														
Declin. Sun or Star.	N. S.	16°	17°	18°	19°	20°	21°	22°	23°	24°	25°	26°	27°	28°	29°	30°
1°		1 12	1 13	1 14	1 15	1 16	1 17	1 18	1 19	1 20	1 21	1 22	1 24	1 25	1 26	1 28
2		2 14	2 15	2 17	2 18	2 19	2 21	2 22	2 24	2 26	2 27	2 29	2 31	2 33	2 35	3 37
3		3 16	3 18	3 20	3 21	3 23	3 25	3 27	3 29	3 31	3 34	3 36	3 38	3 41	3 44	3 46
4		4 19	4 21	4 23	4 25	4 27	4 29	4 32	4 34	4 37	4 40	4 43	4 46	4 49	4 52	4 56
5		5 21	5 23	5 26	5 28	5 31	5 34	5 37	5 40	5 43	5 46	5 50	5 54	5 57	6 1	6 5
6		6 24	6 26	6 29	6 32	6 35	6 38	6 42	6 45	6 48	6 52	6 56	7 1	7 5	7 10	7 15
7		7 26	7 29	7 32	7 36	7 39	7 42	7 46	7 50	7 54	7 59	8 3	8 8	8 13	8 18	8 24
8		8 29	8 32	8 35	8 39	8 43	8 47	8 51	8 56	9 0	9 5	9 10	9 16	9 22	9 27	9 34
9		9 31	9 35	9 38	9 43	9 47	9 51	9 56	10 1	10 6	10 12	10 17	10 23	10 30	10 36	10 43
10		10 34	10 38	10 42	10 46	10 51	10 56	11 1	11 6	11 12	11 18	11 24	11 31	11 38	11 45	11 53
11		11 36	11 40	11 45	11 50	11 55	12 0	12 6	12 12	12 18	12 24	12 31	12 38	12 46	12 54	13 2
12		12 38	12 43	12 48	12 53	12 59	13 5	13 11	13 17	13 24	13 31	13 38	13 46	13 55	14 3	14 12
13		13 41	13 46	13 52	13 57	14 3	14 9	14 16	14 23	14 30	14 38	14 46	14 54	15 3	15 12	15 22
14		14 44	14 49	14 55	15 1	15 7	15 14	15 21	15 28	15 36	15 44	15 53	16 2	16 12	16 22	16 32
15		15 47	15 52	15 58	16 5	16 11	16 18	16 26	16 34	16 42	16 51	17 0	17 10	17 20	17 31	17 43
16		16 49	16 55	17 2	17 8	17 15	17 23	17 32	17 40	17 49	17 58	18 8	18 18	18 29	18 41	18 53
17		17 52	17 58	18 5	18 12	18 20	18 28	18 36	18 45	18 55	19 5	19 16	19 27	19 38	19 50	20 4
18		18 55	19 2	19 9	19 16	19 24	19 33	19 42	19 51	20 1	20 12	20 23	20 35	20 47	21 0	21 14
19		19 58	20 5	20 12	20 20	20 29	20 38	20 47	20 57	21 8	21 19	21 31	21 44	21 57	22 11	22 25
20		21 0	21 8	21 16	21 24	21 33	21 43	21 53	22 3	22 15	22 26	22 39	22 52	23 6	23 21	23 36
21		22 3	22 11	22 19	22 28	22 38	22 48	22 58	23 9	23 21	23 34	23 47	24 1	24 16	24 31	24 47
22		23 6	23 14	23 23	23 32	23 42	23 53	24 4	24 16	24 29	24 42	24 55	25 10	25 25	25 41	25 58
23		24 9	24 18	24 27	24 37	24 47	24 58	25 10	25 22	25 35	25 49	26 3	26 19	26 35	26 52	27 10
24		25 12	25 21	25 31	25 41	25 52	26 3	26 16	26 29	26 43	26 56	27 12	27 28	27 45	28 3	28 22
25		26 15	26 25	26 35	26 46	26 57	27 9	27 22	27 35	27 49	28 5	28 21	28 37	28 55	29 14	29 34
26		27 18	27 28	27 39	27 50	28 2	28 14	28 28	28 42	28 57	29 13	29 29	29 47	30 6	30 25	30 46
27		28 21	28 32	28 43	28 54	29 7	29 20	29 34	29 49	30 4	30 21	30 38	30 57	31 16	31 37	31 59
28		29 25	29 35	29 47	29 59	30 12	30 26	30 40	30 56	31 12	31 29	31 48	32 7	32 27	32 49	33 12
29		30 29	30 39	30 51	31 4	31 17	31 32	31 47	32 3	32 20	32 38	32 57	33 17	33 39	34 1	34 25
30		31 32	31 43	31 55	32 9	32 23	32 38	32 53	33 10	33 28	33 47	34 7	34 28	34 50	35 14	35 39
31		32 35	32 47	33 0	33 14	33 28	33 44	34 0	34 18	34 37	34 56	35 17	35 39	36 2	36 27	36 52
32		33 38	33 51	34 4	34 18	34 34	34 50	35 7	35 26	35 45	36 5	36 27	36 50	37 14	37 40	38 7
S. N.																
1°		0 53	0 53	0 53	0 53	0 52	0 52	0 52	0 52	0 52	0 51	0 51	0 51	0 51	0 51	0 51
2		1 56	1 56	1 56	1 56	1 56	1 57	1 57	1 57	1 57	1 58	1 58	1 58	1 59	1 59	2 0
3		2 58	2 59	2 59	3 0	3 0	3 1	3 1	3 2	3 3	3 4	3 5	3 6	3 7	3 8	3 9
4		4 0	4 1	4 2	4 3	4 4	4 5	4 6	4 7	4 8	4 10	4 11	4 13	4 13	4 15	4 17
5		5 3	5 4	5 5	5 6	5 8	5 9	5 11	5 12	5 14	5 16	5 18	5 21	5 23	5 25	5 28
6		6 5	6 7	6 8	6 10	6 11	6 13	6 15	6 17	6 19	6 21	6 23	6 26	6 28	6 31	6 34
7		7 8	7 9	7 11	7 13	7 15	7 18	7 20	7 23	7 26	7 29	7 32	7 35	7 39	7 43	7 47
8		8 10	8 12	8 14	8 17	8 19	8 22	8 25	8 28	8 31	8 35	8 39	8 43	8 47	8 51	8 55
9		9 13	9 15	9 17	9 20	9 23	9 26	9 30	9 33	9 37	9 41	9 45	9 50	9 55	10 0	10 6
10		10 15	10 18	10 21	10 24	10 27	10 31	10 35	10 39	10 43	10 48	10 53	10 58	11 3	11 9	11 15
11		11 18	11 21	11 24	11 27	11 31	11 35	11 39	11 44	11 49	11 54	11 59	12 5	12 12	12 18	12 25
12		12 20	12 24	12 27	12 31	12 35	12 39	12 44	12 49	12 55	13 0	13 6	13 13	13 20	13 27	13 34
13		13 23	13 26	13 30	13 34	13 39	13 44	13 49	13 55	14 1	14 7	14 14	14 21	14 28	14 36	14 44
14		14 25	14 29	14 33	14 38	14 43	14 48	14 54	15 0	15 6	15 14	15 21	15 28	15 36	15 45	15 54
15		15 27	15 32	15 37	15 42	15 47	15 53	15 59	16 6	16 13	16 20	16 28	16 36	16 45	16 54	17 4
16		16 30	16 35	16 40	16 45	16 51	16 57	17 4	17 11	17 19	17 27	17 35	17 44	17 53	18 3	18 14
17		17 33	17 38	17 43	17 49	17 55	18 2	18 9	18 17	18 25	18 33	18 42	18 52	19 2	19 13	19 24
18		18 35	18 41	18 47	18 53	19 0	19 7	19 14	19 25	19 31	19 40	19 50	20 0	20 11	20 23	20 35
19		19 38	19 44	19 50	19 57	20 4	20 12	20 20	20 28	20 37	20 47	20 57	21 8	21 20	21 32	21 45
20		20 41	20 47	20 54	21 1	21 8	21 16	21 25	21 34	21 44	21 54	22 5	22 17	22 29	22 42	22 56
21		21 44	21 50	21 57	22 4	22 12	22 21	22 30	22 40	22 50	23 1	23 13	23 25	23 38	23 52	24 6
22		22 46	22 53	23 1	23 8	23 17	23 26	23 36	23 46	23 57	24 9	24 21	24 34	24 48	25 2	25 17
23		23 49	23 56	24 4	24 12	24 21	24 31	24 41	24 52	25 4	25 16	25 29	25 42	25 57	26 12	26 28
24		24 52	25 0	25 8	25 17	25 26	25 36	25 47	25 58	26 10	26 23	26 37	26 52	27 7	27 23	27 40
25		25 55	26 3	26 12	26 21	26 31	26 41	26 52	27 4	27 17	27 31	27 45	28 0	28 16	28 33	28 51
26		26 58	27 6	27 15	27 25	27 36	27 47	27 58	28 11	28 25	28 39	28 54	29 10	29 26	29 44	30 3
27		28 1	28 10	28 19	28 29	28 40	28 52	29 4	29 17	29 32	29 46	30 2	30 19	30 37	30 56	31 16
28		29 4	29 13	29 23	29 34	29 45	29 57	30 10	30 24	30 39	30 54	31 11	31 29	31 47	32 7	32 28
29		30 7	30 17	30 27	30 38	30 50	31 3	31 17	31 31	31 46	32 2	32 21	32 39	32 58	33 19	33 40
30		31 10	31 20	31 31	31 43	31 55	32 8	32 23	32 38	32 54	33 11	33 30	33 49	34 9	34 31	34 54
31		32 13	32 24	32 35	32 47	33 0	33 14	33 29	33 45	34 2	34 20	34 39	34 59	35 20	35 43	36 7
32		33 16	33 27	33 39	33 52	34 6	34 20	34 36	34 52	35 10	35 29	35 49	36 10	36 32	36 55	37 20

FOR

For finding the VARIATION of the COMPASS.

AMPLITUDE of RISING and SETTING, from EAST and WEST.

N. S.		North or South LATITUDE of the PLACE, correspondent to N. S. Declination of SUN or STAR.														
Declin. Sun or Star.		31°	32°	33°	34°	35°	36°	37°	38°	39°	40°	41°	42°	43°	44°	45°
N. S.		o /	o /	o /	o /	o /	o /	o /	o /	o /	o /	o /	o /	o /	o /	o /
1°		1 29	1 31	1 32	1 34	1 36	1 38	1 39	1 41	1 43	1 45	1 47	1 50	1 52	1 54	1 57
2		2 39	2 42	2 44	2 46	2 49	2 52	2 54	2 57	3 0	3 4	3 7	3 10	3 14	3 18	3 22
3		3 49	3 52	3 55	3 59	4 2	4 6	4 10	4 14	4 18	4 22	4 26	4 31	4 36	4 41	4 47
4		4 59	5 3	5 7	5 11	5 16	5 20	5 25	5 30	5 35	5 40	5 46	5 52	5 58	6 5	6 12
5		6 9	6 14	6 19	6 24	6 29	6 34	6 40	6 46	6 52	6 59	7 6	7 13	7 21	7 29	7 37
6		7 20	7 25	7 30	7 36	7 42	7 49	7 56	8 3	8 10	8 18	8 26	8 34	8 43	8 53	9 3
7		8 30	8 36	8 42	8 49	8 56	9 3	9 11	9 19	9 28	9 37	9 46	9 56	10 6	10 16	10 28
8		9 40	9 47	9 54	10 2	10 10	10 18	10 27	10 36	10 45	10 55	11 6	11 17	11 29	11 41	11 54
9		10 50	10 58	11 6	11 15	11 24	11 33	11 42	11 53	12 3	12 14	12 26	12 39	12 52	13 5	13 20
10		12 1	12 9	12 18	12 28	12 37	12 48	12 58	13 9	13 21	13 34	13 47	14 0	14 15	14 30	14 46
11		13 11	13 21	13 31	13 41	13 51	14 2	14 14	14 27	14 40	14 53	15 8	15 23	15 38	15 55	16 13
12		14 22	14 32	14 43	14 54	15 5	15 18	15 30	15 44	15 58	16 13	16 28	16 45	17 2	17 20	17 40
13		15 33	15 44	15 55	16 7	16 20	16 33	16 47	17 1	17 17	17 33	17 50	18 8	18 26	18 46	19 7
14		16 44	16 55	17 8	17 21	17 34	17 48	18 3	18 19	18 36	18 53	19 11	19 31	19 51	20 12	20 34
15		17 55	18 7	18 20	18 34	18 49	19 4	19 20	19 37	19 55	20 14	20 33	20 54	21 16	21 38	22 3
16		19 6	19 19	19 33	19 48	20 4	20 20	20 37	20 55	21 14	21 34	21 55	22 17	22 41	23 5	23 31
17		20 17	20 31	20 46	21 2	21 19	21 36	21 54	22 14	22 34	22 55	23 18	23 42	24 6	24 33	25 1
18		21 28	21 44	22 0	22 16	22 34	22 53	23 12	23 32	23 54	24 17	24 41	25 6	25 32	26 1	26 31
19		22 40	22 56	23 13	23 31	23 50	24 9	24 30	24 52	25 15	25 39	26 4	26 31	26 59	27 29	28 1
20		23 52	24 9	24 27	24 46	25 6	25 26	25 48	26 11	26 36	27 1	27 28	27 57	28 27	28 59	29 32
21		25 4	25 22	25 41	26 1	26 22	26 44	27 7	27 31	27 57	28 24	28 53	29 23	29 55	30 29	31 4
22		26 16	26 35	26 55	27 16	27 38	28 1	28 26	28 52	29 19	29 47	30 18	30 50	31 24	31 59	32 37
23		27 29	27 49	28 10	28 32	28 55	29 19	29 45	30 12	30 41	31 11	31 43	32 17	32 53	33 31	34 11
24		28 42	29 3	29 25	29 48	30 12	30 38	31 5	31 34	32 4	32 36	33 10	33 46	34 24	35 3	35 46
25		29 55	30 17	30 40	31 4	31 30	31 57	32 25	32 56	33 28	34 1	34 37	35 15	35 55	36 37	37 22
26		31 8	31 31	31 55	32 21	32 48	33 16	33 40	34 18	34 52	35 27	36 5	36 45	37 28	38 12	39 0
27		32 22	32 46	33 11	33 38	34 6	34 36	35 8	35 41	36 17	36 54	37 34	38 16	39 1	39 48	40 39
28		33 36	34 1	34 28	34 56	35 26	35 57	36 30	37 5	37 42	38 22	39 4	39 48	40 35	41 26	42 19
29		34 50	35 16	35 44	36 14	36 45	37 18	37 53	38 30	39 9	39 51	40 35	41 22	42 11	43 5	44 2
30		36 5	36 32	37 2	37 33	38 6	38 40	39 17	39 56	40 37	41 20	42 7	42 57	43 49	44 45	45 46
31		37 20	37 49	38 20	38 52	39 26	40 3	40 41	41 22	42 5	42 51	43 40	44 33	45 28	46 28	47 32
32		38 36	39 6	39 38	40 12	40 48	41 26	42 7	42 50	43 35	44 24	45 16	46 11	47 10	48 13	49 21
N. S.																
1°		0 51	0 51	0 51	0 51	0 51	0 51	0 51	0 51	0 51	0 52	0 52	0 52	0 52	0 52	0 53
2		2 1	2 2	2 2	2 3	2 4	2 5	2 6	2 7	2 8	2 10	2 11	2 13	2 14	2 16	2 18
3		3 11	3 12	3 14	3 15	3 17	3 19	3 21	3 23	3 26	3 28	3 31	3 34	3 37	3 40	3 43
4		4 21	4 23	4 25	4 28	4 30	4 33	4 36	4 39	4 43	4 46	4 50	4 54	4 58	5 3	5 7
5		5 31	5 34	5 37	5 40	5 44	5 48	5 52	5 56	6 0	6 5	6 10	6 15	6 21	6 27	6 33
6		6 41	6 45	6 49	6 53	6 57	7 2	7 7	7 12	7 18	7 24	7 30	7 36	7 43	7 50	7 58
7		7 51	7 56	8 0	8 5	8 11	8 16	8 22	8 28	8 35	8 42	8 49	8 57	9 5	9 14	9 23
8		9 1	9 7	9 12	9 18	9 24	9 31	9 38	9 45	9 53	10 1	10 9	10 18	10 28	10 38	10 48
9		10 12	10 18	10 24	10 31	10 38	10 45	10 53	11 1	11 10	11 20	11 29	11 40	11 50	12 2	12 14
10		11 22	11 29	11 36	11 43	11 51	12 0	12 9	12 18	12 28	12 39	12 49	13 1	13 13	13 26	13 40
11		12 32	12 40	12 48	12 56	13 5	13 14	13 24	13 35	13 46	13 58	14 10	14 23	14 36	14 51	15 6
12		13 42	13 51	14 0	14 9	14 19	14 29	14 41	14 52	15 4	15 17	15 30	15 45	16 0	16 16	16 33
13		14 53	15 2	15 12	15 22	15 33	15 44	15 56	16 9	16 22	16 36	16 51	17 7	17 23	17 41	17 59
14		16 4	16 14	16 24	16 35	16 47	17 0	17 13	17 27	17 41	17 56	18 12	18 29	18 47	19 6	19 26
15		17 14	17 25	17 37	17 49	18 2	18 15	18 29	18 44	19 0	19 16	19 34	19 52	20 12	20 32	20 54
16		18 25	18 37	18 49	19 2	19 16	19 30	19 46	20 2	20 19	20 36	20 55	21 15	21 36	21 58	22 22
17		19 36	19 49	20 2	20 16	20 31	20 46	21 2	21 20	21 38	21 57	22 17	22 39	23 1	23 25	23 50
18		20 47	21 1	21 15	21 30	21 46	22 2	22 20	22 38	22 57	23 18	23 41	24 3	24 27	24 52	25 19
19		21 59	22 13	22 28	22 44	23 1	23 18	23 37	23 57	24 17	24 39	25 3	25 27	25 53	26 20	26 49
20		23 10	23 25	23 41	23 58	24 16	24 35	24 55	25 16	25 38	26 1	26 26	26 52	27 19	27 48	28 19
21		24 22	24 38	24 55	25 13	25 32	25 52	26 13	26 35	26 58	27 23	27 49	28 17	28 46	29 17	29 50
22		25 34	25 51	26 9	26 28	26 48	27 9	27 31	27 55	28 20	28 46	29 14	29 43	30 14	30 47	31 22
23		26 46	27 4	27 23	27 43	28 4	28 26	28 50	29 15	29 41	30 9	30 38	31 9	31 42	32 17	32 54
24		27 58	28 17	28 37	28 58	29 20	29 44	30 9	30 35	31 3	31 33	32 3	32 37	33 12	33 49	34 28
25		29 10	29 30	29 52	30 14	30 38	31 3	31 29	31 56	32 26	32 57	33 30	34 5	34 42	35 21	36 3
26		30 23	30 44	31 7	31 30	31 55	32 21	32 49	33 18	33 49	34 22	34 57	35 33	36 13	36 54	37 38
27		31 36	31 58	32 22	32 47	33 13	33 40	34 9	34 40	35 13	35 47	36 24	37 3	37 44	38 28	39 15
28		32 50	33 13	33 37	34 3	34 31	35 0	35 31	36 3	36 37	37 14	37 53	38 34	39 17	40 4	40 53
29		34 3	34 28	34 54	35 21	35 50	36 20	36 52	37 26	38 3	38 41	39 22	40 5	40 52	41 41	42 33
30		35 18	35 43	36 10	36 39	37 8	37 41	38 15	38 51	39 29	40 10	40 53	41 38	42 27	43 19	44 15
31		36 32	36 59	37 27	37 57	38 29	39 2	39 38	40 16	40 56	41 39	42 24	43 13	44 4	45 0	45 59
32		37 47	38 15	38 45	39 16	39 49	40 25	41 2	41 42	42 24	43 9	43 57	44 48	45 43	46 42	47 45
Amplitude Northerly from E. or. W. Lat. N. but Southerly Lat. S.																
1°		1 29	1 31	1 32	1 34	1 36	1 38	1 39	1 41	1 43	1 45	1 47	1 50	1 52	1 54	1 57
2		2 39	2 42	2 44	2 46	2 49	2 52	2 54	2 57	3 0	3 4	3 7	3 10	3 14	3 18	3 22
3		3 49	3 52	3 55	3 59	4 2	4 6	4 10	4 1							

For

FOR finding the VARIATION of the COMPASS.

AMPLITUDE of RISING and SETTING, from EAST and WEST.

North or South LATITUDE of the PLACE, correspondent to N. S. Declination of SUN or STAR.

Dec. or * N.S.	46°	47°	48°	49°	50°	51°	52°	53°	54°	55°	56°	57°	58°	59°	60°
1°	2 0	2 2	2 5	2 8	2 12	2 15	2 18	2 22	2 26	2 30	2 35	2 40	2 45	2 50	2 56
2°	3 26	3 30	3 35	3 40	3 45	3 50	3 56	4 2	4 8	4 15	4 22	4 30	4 38	4 47	4 56
3°	4 53	4 59	5 5	5 11	5 18	5 26	5 34	5 42	5 51	6 0	6 10	6 20	6 32	6 44	6 56
4°	6 20	6 27	6 35	6 43	6 52	7 2	7 12	7 22	7 33	7 45	7 58	8 11	8 26	8 41	8 57
5°	7 46	7 55	8 5	8 15	8 26	8 38	8 50	9 3	9 16	9 31	9 46	10 2	10 20	10 39	10 59
6°	9 13	9 24	9 35	9 47	10 0	10 14	10 28	10 43	10 59	11 16	11 33	11 54	12 15	12 37	13 1
7°	10 40	10 52	11 6	11 20	11 35	11 50	12 7	12 24	12 43	13 3	13 24	13 46	14 11	14 36	15 4
8°	12 7	12 22	12 37	12 52	13 9	13 27	13 46	14 6	14 27	14 50	15 14	15 39	16 7	16 36	17 8
9°	13 35	13 51	14 8	14 26	14 45	15 5	15 26	15 48	16 12	16 37	17 4	17 33	18 4	18 37	19 12
10°	15 3	15 21	15 39	15 59	16 20	16 42	17 6	17 31	17 57	18 26	18 56	19 28	20 2	20 39	21 19
11°	16 31	16 51	17 11	17 33	17 56	18 21	18 47	19 14	19 43	20 14	20 48	21 22	22 1	22 42	23 26
12°	18 0	18 21	18 44	19 8	19 33	19 59	20 28	20 58	21 30	22 4	22 41	23 20	24 2	24 47	25 36
13°	19 29	19 52	20 16	20 42	21 10	21 39	22 10	22 43	23 18	23 55	24 35	25 18	26 4	26 53	27 47
14°	20 58	21 23	21 50	22 18	22 48	23 19	23 53	24 29	25 7	25 47	26 31	27 18	28 8	29 2	30 0
15°	22 28	22 55	23 24	23 54	24 26	25 1	25 37	26 16	26 57	27 41	28 28	29 19	30 13	31 12	32 16
16°	23 59	24 28	24 58	25 31	26 6	26 43	27 22	28 4	28 47	29 36	30 27	31 22	32 21	33 25	34 34
17°	25 30	26 1	26 34	27 9	27 46	28 26	29 8	29 53	30 41	31 32	32 27	33 27	34 31	35 41	36 56
18°	27 2	27 35	28 10	28 48	29 28	30 10	30 55	31 43	32 35	33 31	34 30	35 34	36 34	37 59	39 22
19°	28 35	29 10	29 48	30 28	31 10	31 56	32 44	33 36	34 31	35 31	36 35	37 45	39 0	40 22	41 51
20°	30 8	30 46	31 26	32 8	32 54	33 43	34 34	35 30	36 29	37 34	38 43	39 58	41 19	42 49	44 27
21°	31 42	32 22	33 5	33 51	34 39	35 31	36 27	37 26	38 30	39 39	40 54	42 15	43 43	45 20	47 8
22°	33 18	34 0	34 46	35 34	36 26	37 21	38 21	39 24	40 33	41 48	43 8	44 36	46 12	47 58	49 56
23°	34 54	35 39	36 28	37 19	38 14	39 14	40 17	41 25	42 39	43 59	45 27	47 2	48 47	50 44	52 54
24°	36 31	37 20	38 11	39 6	40 5	41 8	42 16	43 29	44 49	46 16	47 50	49 34	51 29	53 38	56 4
25°	38 10	39 2	39 56	40 55	41 57	43 5	44 18	45 37	47 3	48 36	50 19	52 13	54 20	56 44	59 28
26°	39 51	40 45	41 43	42 46	43 53	45 5	46 23	47 48	49 21	51 2	52 55	55 1	57 22	60 4	63 14
27°	41 33	42 30	43 32	44 39	45 50	47 8	48 32	50 4	51 44	53 36	55 40	57 59	60 39	63 46	67 32
28°	43 16	44 18	45 24	46 35	47 52	49 15	50 46	52 25	54 15	56 17	58 35	61 12	64 16	67 59	72 46
29°	45 2	46 8	47 18	48 34	49 56	51 26	53 4	54 53	56 53	59 9	61 44	64 45	68 24	73 6	80 20
30°	46 50	48 0	49 15	50 37	52 5	53 43	55 30	57 29	59 43	62 16	65 14	68 49	73 26	80 31	
31°	48 41	49 56	51 16	52 44	54 20	56 5	58 3	60 15	62 45	65 41	69 12	73 45	80 42		
32°	50 35	51 54	53 21	54 56	56 40	58 36	60 46	63 14	66 7	69 35	74 2	80 53			
S. N.	0 53	0 53	0 54	0 55	0 55	0 56	0 56	0 57	0 58	0 59	1 0	1 1	1 2	1 3	1 4
1°	2 20	2 22	2 24	2 26	2 28	2 31	2 34	2 37	2 40	2 43	2 47	2 51	2 55	3 0	3 5
2°	3 46	3 50	3 54	3 58	4 2	4 6	4 11	4 17	4 22	4 28	4 34	4 41	4 49	4 56	5 4
3°	5 12	5 18	5 23	5 29	5 35	5 42	5 49	5 57	6 5	6 13	6 22	6 32	6 42	6 53	7 5
4°	6 39	6 46	6 53	7 1	7 9	7 18	7 27	7 37	7 47	7 58	8 10	8 22	8 36	8 51	9 6
5°	8 6	8 14	8 23	8 33	8 43	8 54	9 5	9 17	9 30	9 44	9 58	10 14	10 30	10 48	11 7
6°	9 32	9 43	9 53	10 5	10 17	10 30	10 43	10 58	11 13	11 29	11 47	12 5	12 25	12 46	13 9
7°	10 59	11 11	11 24	11 37	11 51	12 6	12 22	12 39	12 56	13 15	13 36	13 58	14 21	14 46	15 12
8°	12 27	12 40	12 55	13 10	13 26	13 43	14 1	14 20	14 40	15 2	15 25	15 50	16 17	16 45	17 16
9°	13 54	14 9	14 26	14 43	15 1	15 20	15 40	16 2	16 25	16 49	17 16	17 44	18 14	18 46	19 20
10°	15 22	15 39	15 57	16 16	16 36	16 58	17 20	17 44	18 10	18 37	19 7	19 38	20 12	20 47	21 26
11°	16 50	17 9	17 29	17 50	18 12	18 36	19 1	19 27	19 56	20 27	20 59	21 33	22 11	22 51	23 34
12°	18 19	18 39	19 1	19 24	19 48	20 14	20 42	21 11	21 43	22 16	22 52	23 30	24 11	24 55	25 42
13°	19 48	20 10	20 34	20 59	21 25	21 54	22 24	22 56	23 30	24 8	24 46	25 28	26 13	27 1	27 53
14°	21 17	21 41	22 7	22 34	23 3	23 34	24 7	24 42	25 19	25 59	26 41	27 27	28 16	29 9	30 6
15°	22 47	23 13	23 41	24 10	24 42	25 15	25 50	26 28	27 8	27 52	28 38	29 28	30 21	31 19	32 21
16°	24 17	24 45	25 15	25 47	26 21	26 57	27 35	28 16	28 59	29 46	30 36	31 30	32 28	33 31	34 39
17°	25 48	26 18	26 50	27 24	28 1	28 39	28 20	30 5	30 52	31 42	32 36	33 35	34 38	35 46	37 0
18°	27 19	27 52	28 26	29 3	29 42	30 22	31 7	31 55	32 45	33 40	34 38	35 42	36 50	38 4	39 25
19°	28 52	29 26	30 3	30 42	31 24	32 8	32 56	33 47	34 41	35 40	36 43	37 51	39 5	40 26	41 55
20°	30 25	31 2	31 41	32 23	33 7	33 55	34 46	35 40	36 39	37 42	38 50	40 4	41 24	42 52	44 29
21°	31 59	32 38	33 20	34 5	34 52	35 43	36 38	37 36	38 39	39 47	41 0	42 20	43 47	45 23	47 9
22°	33 34	34 16	35 0	35 48	36 39	37 33	38 31	39 34	40 42	41 55	43 14	44 41	46 16	48 0	49 57
23°	35 10	35 54	36 42	37 32	38 26	39 24	40 27	41 34	42 47	44 6	45 32	47 6	48 49	50 44	52 53
24°	36 47	37 34	38 25	39 18	40 16	41 18	42 25	43 37	44 56	46 21	47 54	49 37	51 30	53 37	56 1
25°	38 25	39 15	40 9	41 7	42 8	43 15	44 26	45 44	47 8	48 41	50 22	52 14	54 20	56 41	59 23
26°	40 5	40 58	41 55	42 57	44 3	45 14	46 31	47 57	49 26	51 6	52 57	55 1	57 20	60 0	63 7
27°	41 46	42 43	43 44	44 49	46 0	47 16	48 39	50 9	51 48	53 38	55 40	57 58	60 35	63 29	67 21
28°	43 30	44 30	45 35	46 45	48 0	49 22	50 51	52 30	54 18	56 18	58 34	61 9	64 10	67 48	72 28
29°	45 15	46 19	47 28	48 43	50 4	51 32	53 9	54 56	56 55	59 9	61 41	64 39	68 14	72 49	
30°	47 2	48 11	49 25	50 45	52 2	53 48	55 33	57 31	59 42	62 13	65 8	68 39	73 9		
31°	48 52	50 5	51 25	52 51	54 25	56 9	58 5	60 15	62 43	65 35	69 3	73 28			
32°															

III. CATALOGUE of Eminent FIXED STARS: With their Number, Right Ascension, Declination for January 1, 1750, and annual Variation from thence. According to M. DE LA CAILLE.

No.	NORTHERN STARS NAMES, AND Places in their CONSTELLATIONS.	Bayer's Letters.	Magnitude	Right Ascen.	R. Ascen.	Ann.	Declination	Ann.
				in Degrees.	in Time.	Var.	North.	Var.
				° ' "	H. M. S.	+	° ' "	+
THE								
1	Extreme of the Wing of <i>Pegasus</i>	γ	2	0 5 53,5	0 0 24	3,08	13 47 36,3	20,04
2	Shoulder of <i>Andromeda</i>	δ	3	6 30 15,0	0 26 1	3,16	29 29 18,3	20,01
3	Breast of <i>Cassiopea</i>	α	3	6 37 4,8	0 26 28	3,30	55 9 37,9	20,00
4	Waist of <i>Cassiopea</i>	γ	3	10 27 14,2	0 41 49	3,49	59 21 19,2	19,70
5	Polar Star	α	2	10 40 23,0	0 42 42	10,05		19,69
6	Waist of <i>Andromeda</i>	β	2	13 57 3,6	0 55 48	3,29	34 17 15,7	19,43
7	Knee of <i>Cassiopea</i>	δ	3	17 25 14,5	1 9 41	3,71	58 55 27,1	19,07
8	Leg of <i>Cassiopea</i>	ϵ	3	24 10 22,9	1 36 42	4,13	62 25 20,6	18,27
9	Preceding of the Northern Triangle	α	4	24 43 28,0	1 38 54	3,49	28 20 59,3	18,19
10	Middle of Two, at the Ear of the Ram	γ	4	24 57 46,3	1 39 51	3,27	18 3 34,0	18,16
11	Preceding Horn of the Ram	β	3	25 13 4,7	1 40 52	3,28	19 34 34,1	18,12
12	Foot of <i>Andromeda</i>	γ	2	27 9 56,5	1 48 40	3,61	41 7 0,5	17,82
13	Node of the Fishes	α	3	27 17 7,0	1 49 8	3,09	1 32 50,2	17,80
14	Following Horn of the Ram	α	3	28 17 5,0	1 53 8	3,34	22 16 9,6	17,63
15	Northern of the Triangle	β	4	28 41 15,6	1 54 45	3,48	33 47 28,2	17,56
16	Southern of the Triangle	γ	4	30 38 0,7	2 2 32	3,52	32 40 37,7	17,22
17	Following in the Whale's Cheek	γ	3	37 35 34,0	2 30 22	3,12	2 10 10,5	15,87
18	Northern in the Flower de Luce		4	38 16 0,0	2 33 4	3,53	28 11 26,3	15,73
19	Southern in the Flower de Luce		4	38 49 39,5	2 35 19	3,50	26 12 44,4	15,61
20	Shoulder of <i>Perseus</i>	γ	3	41 42 55,6	2 46 52	4,24	52 30 13,8	14,96
21	Jaw of the Whale	α	2	42 18 33,6	2 49 14	3,13	3 5 32,6	14,81
22	Head of <i>Medusa</i>	β	2	43 0 4,6	2 52 0	3,84	39 58 17,9	14,64
23	Bright Star of <i>Perseus</i>	α	2	46 39 22,5	3 6 37	4,19	48 56 46,6	13,74
24	Thigh of <i>Perseus</i>	δ	3	51 18 33,3	3 25 14	4,17	46 57 40,0	12,52
25	Brightest of the <i>Pleiades</i>	η	3	53 10 1,6	3 32 40	3,54	23 18 35,8	12,01
26	Extreme of the Foot of <i>Perseus</i>	ζ	3	54 37 4,6	3 38 28	3,72	31 7 2,6	11,60
27	Knee of <i>Perseus</i>	ϵ	3	55 17 24,1	3 41 10	3,94	38 15 41,6	11,41
28	First of the <i>Hyades</i>	γ	3	61 23 56,2	4 5 36	3,39	15 0 6,8	9,59
29	Second of the <i>Hyades</i>	δ	4	62 8 17,3	4 8 33	3,44	16 56 2,8	9,35
30	Northern Eye of the Bull	ϵ	3	63 30 42,4	4 14 3	3,48	18 36 13,7	8,93
31	Southern Eye of the Bull. Aldebaran	α	1	65 24 4,0	4 21 36	3,43	15 59 3,2	8,33
32	Goat	α	1	74 33 56,5	4 58 16	4,40	45 42 36,7	5,33
33	Northern Horn of the Bull	β	2	77 37 27,2	5 10 30	3,78	28 22 6,2	4,29
34	Western Shoulder of <i>Orion</i>	γ	2	77 56 2,2	5 11 44	3,22	6 5 55,7	4,19
35	Southern Horn of the Bull	ζ	3	80 40 42,9	5 22 43	3,58	20 57 52,7	3,24
36	Shoulder of the Coachman	β	3	85 17 57,8	5 41 12	4,41	44 53 16,5	1,64
37	Eastern Shoulder of <i>Orion</i>	α	1	85 24 46,2	5 41 39	3,25	7 20 10,6	1,60
38	Hand of the Coachman	θ	3	85 40 8,5	5 42 41	4,08	37 9 51,6	1,51
39	Foot of <i>Castor</i>	η	4	89 56 36,3	5 59 48	3,63	21 33 15,5	0,02
40	Foot of <i>Pollux</i> , the brightest	μ	4	91 58 25,0	6 7 54	3,63	22 30 59,2	0,07
41	Bright Star in the Foot of <i>Pollux</i>	γ	3	95 48 54,7	6 23 16	3,48	16 35 16,9	2,02
42	Knee of <i>Castor</i>	ϵ	3	97 8 9,0	6 28 33	3,71	25 21 9,4	2,49
43	Knee of <i>Pollux</i>	ζ	3	102 18 49,4	6 49 15	3,58	20 54 45,1	4,27
44	Thigh of <i>Pollux</i>	δ	3	106 17 26,4	7 5 10	3,61	22 25 8,5	5,61
45	Neck of the Little Dog	β	3	108 23 46,5	7 13 35	3,27	8 46 20,2	6,32
46	Bright Star in the Head of <i>Castor</i>	α	2	109 39 2,7	7 18 36	3,87	32 24 34,5	6,73
47	Brightest in the Little Dog. Procyon	α	1	111 32 59,4	7 26 12	3,21	5 50 42,5	7,35
48	Head of <i>Pollux</i>	β	2	112 29 45,6	7 29 59	3,75	28 36 22,5	7,66
49	Southern Foot of the Crab	β	4	120 44 4,3	8 2 56	3,15	9 56 7,0	10,23
50	Northern <i>Afs</i>	γ	4	127 11 42,2	8 27 47	3,52	22 20 59,3	12,11
51	Southern <i>Afs</i>	δ	4	127 36 43,4	8 30 27	3,44	19 3 22,0	12,22

III. A CATALOGUE of Eminent FIXED STARS: With their Number, Right Ascension, Declination for January 1, 1750, and annual Variation from thence.

No.	NORTHERN STARS NAMES, AND Places in their CONSTELLATIONS.	Bayer's Letters.	Magnitude	Right Ascen. in Degrees.			R. Ascen. in Time.			Ann. Var.		Declination North.			Ann. Var.		
				°	'	"	H.	M.	S.	+	S.	°	'	"	—	"	
THE																	
52	Northern in the Foot of the <i>Great Bear</i>	α	3	130	29	7,9	8	41	57	4,25		48	59	58,7	13,00		
53	Claw of the <i>Crab</i>	α	3	130	32	21,3	8	42	9	3,31		12	48	33,5	13,01		
54	Southern in the Foot of the <i>Great Bear</i>	α	4	131	36	28,0	8	46	26	4,20		48	7	16,6	13,30		
55	Preceding Knee of the <i>Great Bear</i>	θ	3	138	59	39,2	9	15	59	4,22		52	47	42,3	15,11		
56	Foot of the <i>Lion</i>	ο	4	141	56	46,0	9	27	47	3,24		10	40	56,5	15,79		
57	Eye of the <i>Lion</i>	α	3	142	54	6,0	9	31	36	3,45		24	24	35,2	15,99		
58	Northern of the Head of the <i>Lion</i>	μ	3	144	37	21,0	9	38	29	3,47		27	10	8,5	16,33		
59	Southern in the Neck of the <i>Lion</i>	η	3	148	24	53,8	9	53	40	3,31		17	58	18,4	17,07		
60	Heart of the <i>Lion</i> . <i>Regulus</i>	α	1	148	45	26,0	9	55	2	3,24		13	10	43,7	17,13		
61	Northern in the Neck of the <i>Lion</i>	ζ	3	150	40	55,0	10	2	44	3,34		24	39	4,0	17,47		
62	Following in the Neck of the <i>Lion</i>	γ	3	151	32	6,1	10	6	0	3,31		21	5	49,3	17,61		
63	In the <i>Belly</i> of the <i>Lion</i>	ε	4	154	54	20,0	10	19	37	3,18		10	35	6,4	18,14		
64	Southern in the □ of the <i>Great Bear</i>	β	2	161	38	26,5	10	46	34	3,75		57	42	51,0	19,01		
65	Northern following in the <i>Great Bear</i>	α	2	162	0	45,1	10	48	3	3,89		63	5	33,9	19,05		
66	Thigh of the <i>Lion</i>	δ	3	165	11	27,2	11	0	46	3,22		21	53	28,6	19,37		
67	Back of the <i>Lion</i>	θ	3	165	16	13,1	11	1	5	3,18		16	47	32,5	19,38		
68	Tail of the <i>Lion</i>	β	2	174	4	17,4	11	36	17	3,12		15	58	7,2	19,93		
69	Northern Wing of the <i>Virgin</i>	β	3	174	25	5,0	11	37	40	3,08		3	10	20,5	19,94		
70	Southern following in the □ of the <i>Gr. Bear</i>	γ	2	175	8	7,1	11	40	32	3,24		55	5	0,7	19,98		
71	Last in the □ of the <i>Great Bear</i>	δ	2	180	43	47,3	12	2	55	3,05		58	25	19,3	20,04		
72	Southern Wing of the <i>Virgin</i>	η	3	181	46	48,1	12	7	7	3,07		0	43	31,1	20,02		
73	First of the Tail of the <i>Great Bear</i>	ε	2	190	44	5,6	12	42	56	2,69		57	19	15,1	19,69		
74	Foregoing in the Waist of the <i>Virgin</i>	δ	3	190	45	18,1	12	43	1	3,06		4	45	43,2	19,69		
75	The brightest under the <i>Great Bear</i>	3	3	191	4	20,0	12	44	17	2,86		39	40	25,8	19,66		
76	Northern Wing of the <i>Virgin</i>	ε	3	192	25	51,8	12	49	43	3,01		12	18	34,3	19,57		
77	Middle of the Tail of the <i>Great Bear</i>	ζ	2	198	26	53,8	13	13	48	2,45		56	14	12,2	19,01		
78	Following in the Waist of the <i>Virgin</i>	ζ	3	200	29	37,0	13	21	58	3,08		0	41	27,5	18,77		
79	Extremity of the Tail of the <i>Great Bear</i>	η	2	204	24	58,6	13	37	40	2,41		50	34	5,4	18,24		
80	Preceding Thigh of the <i>Cowherd</i> , or <i>Clown</i>	η	3	205	41	34,1	13	42	46	2,88		19	39	45,0	18,06		
81	Extremity of the <i>Dragon's Tail</i>	α	3	209	24	28,7	13	57	38	1,50		65	34	32,5	17,45		
82	Brightest of the <i>Clown</i> . <i>Arcturus</i>	α	1	211	4	1,0	14	4	16	2,82		20	29	39,2	17,16		
83	Preceding Shoulder of the <i>Clown</i>	γ	3	215	29	55,5	14	22	0	2,43		38	24	19,0	16,31		
84	Following Foot of the <i>Clown</i>	ζ	3	217	18	14,6	14	29	13	2,86		14	48	52,2	15,93		
85	Following Thigh of the <i>Clown</i>	ε	3	218	32	15,8	14	34	9	2,63		28	8	27,0	15,67		
86	Preceding in □ of the <i>Little Bear</i>	β	3	222	55	42,3	14	51	43	0,36	—				14,66		
87	Head of the <i>Clown</i> , or <i>Cowherd</i>	β	3	223	8	0,0	14	52	32	2,28		41	23	15,7	14,62		
88	Following Shoulder of the <i>Clown</i>	δ	3	226	23	12,5	15	5	33	2,44		34	15	40,8	13,82		
89	Preceding in the last Joint of the <i>Dragon</i>	ι	4	229	51	2,1	15	19	24	1,31		59	50	54,3	12,91		
90	Following in □ of the <i>Little Bear</i>	γ	3	230	19	58,5	15	21	20	0,22	—				12,79		
91	Preceding in Neck of the <i>Serpent</i>	δ	3	230	43	9,0	15	22	53	2,87		11	23	32,0	12,68		
92	Brightest in the <i>Crown</i>	α	2	231	1	40,2	15	24	7	2,05		27	34	18,2	12,60		
93	Brightest in the Neck of the <i>Serpent</i>	α	3	232	59	39,8	15	31	59	3,01		7	13	50,5	12,06		
94	Southern in the Neck of the <i>Serpent</i>	β	3	233	39	52,0	15	34	39	2,76		16	13	16,9	11,87		
95	Following the brightest of the <i>Serpent</i>	ε	4	234	35	30,2	15	38	22	2,98		5	14	56,2	11,62		
96	Following of Southern in <i>Serpent's Neck</i>	γ	3	236	13	52,5	15	44	55	2,75		14	29	41,4	11,14		
97	Following in the last Joint of the <i>Dragon</i>	θ	4	239	18	37,2	15	57	14	1,14		59	14	17,5	10,23		
98	Preceding Arm of <i>Hercules</i>	γ	3	242	43	30,8	16	10	54	2,66		19	45	29,6	9,18		
99	Preceding Shoulder of <i>Hercules</i>	β	3	244	52	29,7	16	19	30	2,59		22	3	7,1	8,51		
100	Last Joint of the <i>Dragon</i>	η	4	245	9	46,9	16	20	39	0,78		62	5	2,5	8,41		
101	Foregoing in the Side of <i>Hercules</i>	ζ	3	247	58	6,2	16	31	52	2,30		32	4	19,4	7,51		
102	Loins of <i>Hercules</i>	η	4	248	34	59,6	16	34	20	2,07		39	24	46,9	7,31		

III. A CATALOGUE of Eminent FIXED STARS: With their Number, Right Ascension, Declination for
January 1, 1750, and annual Variation from thence.

N ^o .	NORTHERN STARS NAMES, AND Places in their CONSTELLATIONS.	Bayer's Letters.	Magnitude.	Right Ascen. in Degrees.			R. Ascen. in Time.			Ann. Var. +	Declination North.			Ann. Var. -
				°	'	"	H.	M.	S.		°	'	"	
THE														
103	Following in the Side of <i>Hercules</i>	α	3	252	40	52,2	16	50	43	2,29	31	18	41,6	5,96
104	Head of <i>Hercules</i>	α	2	255	48	49,2	17	3	15	2,74	14	41	44,5	4,91
105	Following Shoulder of <i>Hercules</i>	δ	3	256	26	36,4	17	5	46	2,47	25	9	4,7	4,69
106	Head of <i>Ophiuchus</i>	α	2	260	50	5,3	17	23	20	2,78	12	45	47,9	3,17
107	Eye of the <i>Dragon</i>	β	3	261	12	6,8	17	24	48	1,36	52	29	44,5	3,06
108	Northern in Shoulder of <i>Ophiuchus</i>	β	3	262	46	55,0	17	31	8	2,97	4	41	40,6	2,52
109	Southern in Shoulder of <i>Ophiuchus</i>	γ	3	263	50	35,2	17	35	22	3,01	2	49	28,3	2,15
110	Elbow of <i>Hercules</i>	μ	4	264	10	9,5	17	36	41	2,38	27	53	3,9	2,03
111	Knee of <i>Hercules</i>	θ	3	266	55	25,6	17	47	42	2,06	37	17	56,4	1,08
112	Head of the <i>Dragon</i>	γ	3	267	42	3,0	17	50	48	2,21	51	31	37,9	0,80
+														
113	Brightest in the <i>Harp</i>	α	1	277	7	7,0	18	28	28	2,06	38	34	0,0	2,48
114	Preceding in the Lozange of the <i>Harp</i>	β	3	280	12	47,3	18	40	51	2,22	33	5	23,4	3,55
115	Preceding in the Extremity of <i>Serpent's Tail</i>	θ	4	280	56	51,2	18	43	47	3,00	3	54	6,0	3,80
116	Following in the Lozange of the <i>Harp</i>	δ	3	281	26	36,3	18	45	46	2,11	36	35	48,6	3,97
117	Preceding in the Tail of the <i>Eagle</i>	ϵ	4	282	4	17,5	18	48	17	2,73	14	44	56,1	4,18
118	Following in the Lozange of the <i>Harp</i>	γ	3	282	23	54,3	18	49	36	2,26	32	21	49,0	4,29
119	Following in the Tail of the <i>Eagle</i>	ζ	4	283	28	50,4	18	53	55	2,69	13	30	42,5	4,67
120	Second Joint of the <i>Dragon</i>	δ	3	288	6	23,0	19	12	26	0,06	67	13	17,1	6,22
121	Preceding Wing of the <i>Eagle</i>	δ	3	288	13	24,7	19	12	54	3,02	2	38	18,2	6,26
122	Beak of the <i>Swan</i>	β	3	290	9	25,3	19	20	38	2,43	27	27	4,0	6,90
123	Brightest of the <i>Arrow</i>	α	4	292	14	1,0	19	28	56	2,69	17	27	28,9	7,58
124	Preceding in the Neck of the <i>Eagle</i>	γ	3	293	35	30,2	19	34	22	2,28	10	1	22,0	8,02
125	Northern Wing of the <i>Swan</i>	δ	3	294	17	28,4	19	37	10	1,86	44	31	58,5	8,24
126	Brightest of the <i>Eagle</i>	α	2	294	38	46,7	19	38	35	2,91	8	13	44,3	8,36
127	Following Shoulder of <i>Antinous</i>	η	4	294	55	55,0	19	39	44	3,07	0	23	5,4	8,45
128	Following in Neck of the <i>Eagle</i>	β	3	295	45	31,0	19	43	2	3,02	5	48	10,6	8,71
129	Breast of the <i>Swan</i>	γ	3	303	18	50,0	20	13	15	2,16	39	8	12,8	11,00
130	Preceding in Tail of the <i>Dolphin</i>	ϵ	4	305	18	56,4	20	21	16	2,88	10	28	20,0	11,58
131	Preceding in the Lozange of the <i>Dolphin</i>	ζ	4	305	54	17,8	20	23	37	2,82	13	49	52,2	11,74
132	Southerly in the Lozange of the <i>Dolphin</i>	β	3	306	27	34,3	20	25	50	2,82	13	44	32,6	11,91
133	Northerly in the Lozange of the <i>Dolphin</i>	α	3	307	0	22,2	20	28	1	2,79	15	2	51,5	12,05
134	Southerly following in the Loz. of the <i>Dolphin</i>	δ	4	307	56	44,9	20	31	47	2,81	14	11	36,6	12,32
135	Tail of the <i>Swan</i>	α	2	308	13	38,5	20	32	55	2,05	44	23	54,5	12,39
136	Last in Lozange of <i>Dolphin</i>	γ	4	308	46	3,0	20	35	4	2,79	15	14	22,5	12,54
137	Brightest in Southern Wing of the <i>Swan</i>	ϵ	3	309	1	19,6	20	36	5	2,40	33	2	50,4	12,62
138	Extremity of Southern Wing of the <i>Swan</i>	ζ	4	315	34	26,4	21	2	18	2,55	29	12	50,7	14,31
139	Head of the little <i>Horse</i>	α	4	315	49	36,0	21	3	18	3,01	4	14	8,1	14,37
140	First of <i>Pegasus</i>	ϵ	4	317	37	25,3	21	10	30	2,79	18	44	53,7	14,80
141	Preceding Shoulder of <i>Cepheus</i>	α	3	318	8	41,8	21	12	34	1,43	61	31	55,7	14,92
142	Girdle of <i>Cepheus</i>	β	4	321	19	34,4	21	25	18	0,81	69	27	54,8	15,64
143	Mouth of <i>Pegasus</i>	ϵ	3	322	58	21,6	21	31	53	2,95	8	44	29,8	16,00
144	Lower Part of Southern Wing of the <i>Swan</i>	μ	4	323	14	41,1	21	32	59	2,64	27	37	24,1	16,05
145	Neck of <i>Pegasus</i>	ζ	3	337	14	43,9	22	28	59	2,99	9	32	4,3	18,46
146	Northern in the Knee of <i>Pegasus</i>	η	3	337	49	37,7	22	31	18	2,80	28	55	14,5	18,55
147	Ring of the Chain of <i>Andromeda</i>	θ	4	342	36	49,0	22	50	27	2,72	40	59	22,0	19,12
148	Thigh of <i>Pegasus</i>	β	2	342	55	14,5	22	51	41	2,89	26	43	52,7	19,15
149	Wing of <i>Pegasus</i>	α	2	343	4	57,3	22	52	20	2,99	13	51	56,0	19,17
150	Following Foot of <i>Cepheus</i>	γ	4	352	19	22,1	23	29	17	2,31				19,86
151	Head of <i>Andromeda</i>	α	2	358	52	44,1	23	55	31	3,06	27	42	32,1	20,04
152	Chair of <i>Cassiopea</i>	β	3	358	59	41,4	23	55	59	3,04	57	46	7,0	20,04

III. A CATALOGUE of Eminent FIXED STARS: With their Number, Right Ascension, Declination for January 1, 1750, and annual Variation from thence.

No.	SOUTHERN STARS NAMES, AND Places in their CONSTELLATIONS.	Bayer's Letters.	Magnitude	Right Ascen. in Degrees.			R. Ascen. in Time.			Ann. Var.		Declination South.			Ann. Var.	
				°	'	"	H. M. S.			+	S.	°	'	"	—	
THE																
1	Tail of the Male <i>Hydra</i>	β	3	3	2	58,3	0	12	12	2,72		78	39	50,3	20,01	
2	Head of the <i>Phoenix</i>	α	2	3	28	0,9	0	13	52	3,01		43	39	43,4	20,00	
3	Brightest in the <i>Whale's Tail</i>	β	2	7	45	28,6	0	31	2	3,01		19	21	51,1	19,85	
4	Thigh of the <i>Phoenix</i>	β	3	13	43	21,7	0	54	53	2,73		48	4	42,1	19,46	
5	Preceding towards the <i>Whale's Tail</i>	η	4	14	0	16,1	0	56	1	3,00		11	30	48,8	19,44	
6	Following towards the <i>Whale's Tail</i>	θ	4	17	53	9,1	1	11	33	3,01		9	28	49,5	19,07	
7	Following Wing of the <i>Phoenix</i>	γ	3	19	22	10,7	1	17	29	2,67		44	36	18,8	18,90	
8	Source of the River. <i>Achernar</i>	α	1	22	5	42,2	1	28	23	2,26		58	30	52,1	18,56	
9	Head of the Male <i>Hydra</i>	α	3	27	43	22,5	1	50	53	1,87		62	47	36,3	17,73	
10	Variable of the <i>Whale</i>	$\circ$	4	31	40	53,8	2	6	44	3,03		4	7	28,0	17,05	
11	Preceding in the Cheek of the <i>Whale</i>	δ	3	36	40	35,4	2	26	42	3,08		0	45	50,2	16,07	
12	Breast of the <i>Whale</i>	ϵ	3	36	52	22,9	2	27	30	2,90		12	56	51,9	16,03	
13	First Turning of the River	θ	3	42	11	52,7	2	48	48	2,30		41	19	4,0	14,85	
14	Third Turning of the River	ζ	4	45	21	58,8	3	1	28	2,54		29	59	13,7	14,07	
15	River, before the <i>Whale</i>	ζ	3	45	55	39,1	3	3	43	2,91		0	45	51,2	13,93	
16	Following, in the River	ϵ	3	50	17	32,0	3	21	10	2,89		10	19	12,0	12,79	
17	Following in the River	δ	3	52	49	30,1	3	31	18	2,88		10	37	32,0	12,10	
18	Fourth Turn of the River	γ	3	56	35	48,3	3	46	23	2,79		14	14	21,6	11,02	
19	Middle of the Male <i>Hydra</i>	γ	4	57	51	51,2	3	51	27	1,14		74	59	55,9	10,66	
20	Brightest in the rhomboidal <i>Reticula</i>	α	3	62	49	11,8	4	11	17	0,74		63	6	16,3	9,15	
21	Tail of the Gold-Fish, or <i>Dorado</i>	α	3	67	9	19,7	4	28	37	1,28		55	34	12,9	7,78	
22	Last of the River	β	3	73	53	44,0	4	55	35	2,95		5	25	49,2	5,56	
23	Bright Foot of <i>Orion</i> . <i>Rigel</i>	β	1	75	38	12,2	5	2	33	2,89		8	30	37,4	4,97	
24	Northern in the Sword of <i>Orion</i>	η	3	77	58	48,0	5	11	55	3,02		2	38	56,9	4,17	
25	Belly of the <i>Hare</i>	β	4	79	23	11,3	5	17	33	2,58		20	58	43,1	3,69	
26	Preceding of <i>Orion's Girdle</i>	δ	2	79	48	54,4	5	19	16	3,07		0	30	22,8	3,54	
27	Brightest of the <i>Hare</i>	α	3	80	25	49,7	5	21	43	2,65		18	1	19,0	3,34	
28	Middle of <i>Orion's Girdle</i>	ϵ	2	80	53	11,1	5	23	33	3,05		1	23	1,2	3,17	
29	Last of <i>Orion's Girdle</i>	ζ	2	82	2	33,5	5	28	10	3,04		2	5	48,9	2,77	
30	Preceding of the brightest of the <i>Dove</i>	α	2	82	39	12,2	5	30	37	2,20		34	13	22,7	2,56	
31	Belly of the Gold-Fish, or <i>Dorado</i>	β	4	82	52	17,5	5	31	19	0,50		62	39	29,7	2,48	
32	Southern in the <i>Hare's Foot</i>	γ	4	83	30	43,2	5	34	3	2,53		22	32	58,7	2,26	
33	Knee of <i>Orion</i>	κ	3	83	58	48,2	5	35	55	2,85		9	46	38,3	2,10	
34	Northern in the <i>Hare's Foot</i>	δ	4	85	8	46,0	5	40	35	2,57		20	55	16,1	1,68	
35	Following of the brightest of the <i>Dove</i>	β	3	85	32	30,4	5	42	10	2,11		35	52	40,3	1,55	
+																
36	Preceding Foot of the <i>Great Dog</i>	ζ	3	92	40	58,9	6	10	44	2,31		29	58	11,8	0,94	
37	Knee of the <i>Great Dog</i>	β	3	92	55	25,3	6	11	42	2,63		17	51	11,4	1,02	
38	Brightest of the Ship. <i>Canopus</i>	α	1	94	36	7,6	6	18	25	1,34		52	34	8,9	1,60	
39	Summit of the Ship's Rudder	ν	3	97	31	44,4	6	30	7	1,84		42	59	25,2	2,62	
40	Brightest of the <i>Great Dog</i> . <i>Syrus</i>	α	1	98	32	0,3	6	34	8	2,69		16	23	36,6	2,97	
41	Thigh of the <i>Great Dog</i>	ϵ	3	102	12	7,2	6	48	48	2,36		28	38	59,6	4,23	
42	Back of the <i>Great Dog</i>	δ	2	104	33	26,8	6	58	14	2,45		26	0	56,1	5,03	
43	Poop or Stern of the Ship	π	3	107	4	44,0	7	8	19	2,13		36	39	44,4	5,88	
44	Tail of the <i>Great Dog</i>	η	2	108	33	3,6	7	14	12	2,39		28	49	59,7	6,37	
45	Poop of the Ship	σ	3	110	19	40,5	7	21	19	1,94		42	48	25,6	6,96	
46	Poop of the Ship	ζ	2	118	42	4,5	7	54	48	2,12		39	18	39,5	9,62	
47	Preceding, in Body of the Ship	γ	2	120	27	48,0	8	1	51	1,85		46	36	34,9	10,16	
48	Following in Body of the Ship	ϵ	3	124	20	10,7	8	17	21	1,26		58	42	52,5	11,30	
49	Brightest in Middle of the Ship	δ	2	129	27	5,0	8	37	48	1,61		53	48	2,1	12,73	
50	Sail of the Ship	λ	3	134	42	17,7	8	57	49	2,21		42	26	4,1	14,09	
51	Brightest about the Oars of the Ship	β	1	137	35	26,0	9	10	22	0,75		68	41	26,5	14,79	
52	Mast of the Ship	ϵ	3	137	36	3,3	9	10	24	1,65		58	14	8,3	14,79	

III. A CATALOGUE of Eminent FIXED STARS: With their Number, Right Ascension, Declination for
January 1, 1750, and annual Variation from thence.

No.	SOUTHERN STARS NAMES, AND Places in their CONSTELLATIONS.	Bayer's Letters.	Magnitude	Right Ascen. in Degrees.			R. Ascen. in Time.			Ann. Var. +	Declination South.			Ann. Var. +
				°	'	"	H. M. S.				°	'	"	
							S.							
THE														
53	Brightest more <i>Northerly</i>	α	3	138	35	52,9	9	14	24	1,86	53	57	0,1	15,03
54	Heart of the female <i>Hydra</i>	α	2	138	49	40,0	9	15	19	2,96	7	35	13,1	15,08
55	Oars of the <i>Ship</i>	υ	3	145	12	41,2	9	40	51	1,50	63	55	6,7	16,45
56	Southern in Section of the <i>Ship</i>	θ	3	153	31	34,9	10	34	6	2,12	63	3	24,9	18,64
57	Northern in Section of the <i>Ship</i>	η	2	158	51	29,3	10	35	26	2,27	58	22	39,2	18,68
58	Top of the <i>Ship's</i> Section	μ	3	159	1	11,4	10	36	5	2,55	48	6	15,0	18,81
59	Foot of the <i>Cup</i>	α	4	161	54	13,7	10	47	37	2,95	16	58	29,0	19,05
60	Preceding in the Crupper of the <i>Centaur</i>	δ	3	178	51	51,4	11	55	27	3,05	49	19	39,8	20,03
61	Beak of the <i>Raven</i>	α	4	178	53	31,4	11	55	34	3,07	23	20	6,0	20,03
62	Head of the <i>Raven</i>	ε	4	179	19	46,1	11	57	19	3,06	21	13	44,6	20,04
63	Preceding Arm of the <i>Cross</i>	δ	3	180	30	34,7	12	2	2	3,10	57	21	30,0	20,04
64	Preceding Wing of the <i>Raven</i>	γ	3	180	44	44,5	12	2	59	3,09	16	9	32,5	20,04
65	Foot of the <i>Cross</i>	α	1	183	13	53,7	12	12	56	3,22	61	42	47,4	20,01
66	Preceding in following Wing of the <i>Raven</i>	δ	4	184	14	29,6	12	16	58	3,09	15	7	18,2	19,98
67	Head of the <i>Cross</i>	γ	2	184	28	35,0	12	17	54	3,36	70	44	54,3	19,97
68	Foot of the <i>Raven</i>	β	3	185	19	30,9	12	21	18	3,14	22	0	40,0	19,95
69	Preceding of the <i>Fly</i>	α	4	185	38	41,6	12	22	35	3,42	67	46	17,2	19,94
70	Top of the <i>Centaur's</i> Crupper	γ	2	186	57	53,8	12	27	52	3,27	47	34	52,9	19,89
71	Southern in the <i>Virgin's</i> Girdle	γ	3	187	15	7,2	12	29	0	3,08	0	4	22,3	19,88
72	Head of the <i>Fly</i>	β	4	187	48	53,4	12	31	16	3,52	66	44	6,1	19,84
73	Following Arm of the <i>Cross</i>	β	2	188	19	40,1	12	33	19	3,42	58	19	8,3	19,83
74	Southern Wing of the <i>Virgin</i>	θ	4	194	15	26,2	12	57	2	3,09	4	11	47,2	19,41
75	Tail of the female <i>Hydra</i>	γ	3	196	20	46,6	13	5	23	3,22	21	50	43,8	19,22
76	Preceding Shoulder of <i>Centaur</i>	ε	3	196	39	27,6	13	6	38	3,34	35	23	6,6	19,20
77	Spike of the <i>Virgin</i>	α	1	198	0	51,4	13	12	3	3,15	9	50	51,1	19,06
78	Belly of the <i>Centaur</i>	ε	3	201	3	30,4	13	24	14	3,69	52	10	55,2	18,70
79	Waist of the <i>Centaur</i>	ζ	3	205	1	14,6	13	40	5	3,66	46	2	38,3	18,15
80	Preceding Leg of the <i>Centaur</i>	β	2	206	36	41,8	13	46	27	4,09	59	9	1,8	17,91
81	Following Shoulder of the <i>Centaur</i>	θ	3	207	59	57,5	13	52	0	3,52	35	7	34,9	17,69
82	Robe of the <i>Virgin</i>	κ	4	209	53	57,8	13	59	36	3,19	9	5	43,3	17,37
83	Southern Foot of the <i>Virgin</i>	λ	4	211	24	18,4	14	5	37	3,22	12	12	29,0	17,20
84	Southern in the Shield of the <i>Centaur</i>	η	3	214	56	9,1	14	19	45	3,75	41	2	28,8	16,43
85	Head of the <i>Compasses</i>	α	3	215	39	45,4	14	22	39	4,68	63	51	48,2	16,26
86	Brightest in the Foot of the <i>Centaur</i>	α	1	215	42	31,1	14	22	50	4,41	59	47	9,4	16,26
87	Foot of the <i>Wolf</i>	α	3	216	21	48,0	14	25	27	3,89	46	17	40,1	16,14
88	Southern Scale of the <i>Ballance</i>	α	2	219	16	20,9	14	37	5	3,30	14	59	9,7	15,50
89	Extremity of the Foot of the <i>Wolf</i>	β	3	220	34	6,6	14	42	16	3,86	42	6	13,9	15,22
90	Following Foot of the <i>Centaur</i>	κ	3	220	45	12,7	14	43	1	3,84	41	4	45,6	15,17
91	Preceding Claw of the <i>Scorpion</i>	γ	4	222	22	28,5	14	49	30	3,48	24	16	53,5	14,70
92	Preceding of Southern Triangle	γ	3	224	0	2,8	14	56	0	5,29	67	43	32,3	14,40
93	The Northern Scale of the <i>Ballance</i>	β	2	225	53	50,2	15	3	35	3,30	8	26	30,5	13,94
94	Following in Shoulder of the <i>Wolf</i>	γ	3	229	38	36,2	15	18	34	3,94	40	18	4,6	12,97
95	Following in Northern Scale of the <i>Ballance</i>	γ	4	230	23	42,6	15	21	35	3,34	13	56	10,4	12,77
96	Top of the Southern Triangle	β	3	233	20	42,1	15	33	23	5,06	62	37	34,2	11,96
97	Southern Claw of the <i>Scorpion</i>	ρ	4	235	22	40,5	15	41	31	3,68	28	27	36,0	11,38
98	Southern in the Forehead of the <i>Scorpion</i>	π	4	235	56	38,8	15	43	47	3,60	25	22	18,7	11,32
99	Middle of the Forehead of the <i>Scorpion</i>	δ	3	236	23	52,2	15	45	35	3,52	21	53	17,6	11,09
100	Northern in the Forehead of the <i>Scorpion</i>	β	2	237	44	7,0	15	50	56	3,50	19	5	55,2	10,69
101	Top of the Forehead of the <i>Scorpion</i>	ν	4	239	15	5,4	15	57	0	3,47	18	47	19,6	10,24
102	Preceding Foot of <i>Ophiuchus</i>	δ	3	240	18	58,8	16	1	16	3,14	3	1	46,3	9,92
103	Following in the same Foot of <i>Ophiuchus</i>	ε	3	241	16	51,2	16	5	7	3,16	4	3	36,0	9,63

III. A CATALOGUE of Eminent FIXED STARS: With their Number, Right Ascension, Declination for January 1, 1750, and annual Variation from thence.

No.	SOUTHERN STARS NAMES, AND Places in their CONSTELLATIONS.	Bayer's Letters.	Magnitude	Right Ascen. in Degrees.			R. Ascen. in Time.			Ann. Var.		Declination South.			Ann. Var.	
				°	'	"	H.	M.	S.	+	S.	°	'	"	+	"
104	Preceding of the <i>Scorpion's Heart</i>	σ	4	241	30	40,7	16	6	3	3,62		24	58	2,9	9,56	
105	Brightest in the <i>Scorpion's Heart</i> . <i>Antares</i>	α	1	243	31	52,8	16	14	8	3,66		25	51	8,4	8,93	
106	Following of the <i>Scorpion's Heart</i>	τ	4	245	5	32,3	16	20	22	3,70		27	40	14,7	8,44	
107	Following of the Southern <i>Triangle</i>	α	3	245	37	5,0	16	22	28	6,16		68	31	24,3	8,26	
108	Southern in the <i>Knee of Ophiuchus</i>	ζ	3	245	51	19,0	16	23	25	3,30		10	2	16,0	8,20	
109	First Joint of the <i>Scorpion's Tail</i>	ϵ	3	248	30	14,8	16	34	1	3,90		33	48	41,1	7,34	
110	Second Joint of the <i>Scorpion's Tail</i>	μ	3	248	44	54,6	16	35	0	4,04		37	35	18,6	7,26	
111	Brightest in the third Joint	ζ	3	249	15	51,0	16	37	3	4,20		41	54	3,6	7,09	
112	Fourth Joint	η	3	253	34	34,1	16	54	18	4,28		42	52	39,2	5,66	
113	Following Knee of <i>Ophiuchus</i>	η	2	254	0	53,9	16	56	4	3,44		15	23	28,4	5,52	
114	Following Foot of <i>Ophiuchus</i>	θ	3	256	40	11,2	17	6	21	3,67		24	43	20,6	4,61	
115	Middle of the <i>Altar</i>	α	3	258	8	27,0	17	12	34	4,61		49	38	31,8	4,12	
116	Little Star in Extremity of <i>Scorpion's Tail</i>	ν	4	258	27	0,8	17	13	48	4,09		37	3	55,9	4,01	
117	Brightest in Extremity of the same Tail	λ	3	259	9	59,4	17	16	40	4,08		36	53	28,6	3,77	
118	5th Joint of the <i>Scorpion's Tail</i>	θ	3	259	50	56,1	17	19	24	4,30		42	48	23,2	3,53	
119	7th Joint of the same Tail	κ	3	261	18	19,3	17	25	13	4,13		38	52	14,6	3,03	
120	6th Joint of the same Tail	ι	3	262	31	53,5	17	30	8	4,18		39	59	49,0	2,60	
121	In the last Joint of the <i>Serpent</i>	ζ	4	266	49	26,9	17	47	18	3,16		3	38	56,3	1,11	
122	Follows at the Point of the Arrow of <i>Sagittary</i>	γ	4	267	26	21,5	17	49	45	3,87		30	23	53,7	0,89	
123	Top of the Bow of <i>Sagittary</i>	μ	4	269	42	12,7	17	58	49	3,60		21	5	56,9	0,10	
124	Foot of <i>Sagittary</i>	δ	3	271	14	31,8	18	4	58	3,85		29	54	18,4	0,43	
125	Southern Extreme of the Bow of <i>Sagittary</i>	ϵ	3	271	53	39,7	18	7	35	4,00		34	28	17,4	0,67	
126	Northern Extreme of the Bow of <i>Sagittary</i>	λ	3	273	8	5,2	18	12	32	3,72		25	31	57,4	1,09	
127	Arrow of <i>Sagittary</i>	ϕ	4	277	30	26,7	18	30	2	3,77		27	13	13,0	2,61	
128	Preceding Shoulder of <i>Sagittary</i>	τ	3	279	56	13,4	18	39	45	3,74		26	34	51,0	3,45	
129	Arm of <i>Sagittary</i>	ζ	3	281	40	16,6	18	46	41	3,84		30	12	36,0	4,06	
130	Head of <i>Sagittary</i>	ι	4	282	25	19,1	18	49	41	3,60		22	5	0,8	4,31	
131	Preceding Shoulder of <i>Sagittary</i>	τ	4	282	49	38,3	18	51	19	3,77		28	0	30,6	4,45	
132	Following in the Head of <i>Sagittary</i>	π	3	283	35	45,4	18	54	23	3,59		21	23	48,8	5,70	
133	Preceding of 2 in Foot of <i>Sagittary</i>	β	4	286	9	8,7	19	4	37	4,34		44	53	42,4	5,57	
134	Leg of <i>Sagittary</i>	α	4	286	37	43,3	19	6	31	4,20		41	3	21,5	5,73	
135	Side of <i>Antinous</i>	ι	4	290	58	30,6	19	23	54	4,01		1	49	12,2	7,17	
136	Back of the <i>Peacock</i>	δ	4	295	58	55,0	19	43	56	5,92		66	46	50,8	8,78	
137	Brightest preceding in the Head of <i>Capricorn</i>	α	3	300	56	28,9	20	3	46	3,35		13	18	2,8	10,30	
138	Eye of the <i>Peacock</i>	α	2	301	25	20,0	20	5	41	4,89		57	30	27,4	10,44	
139	Following in the Head of <i>Capricorn</i>	β	3	301	44	3,0	20	6	56	3,39		15	33	2,7	10,54	
140	Zagaye of the <i>Indian</i>	α	3	304	57	56,9	20	19	52	4,29		48	8	14,0	11,48	
141	Wing of the <i>Peacock</i>	β	3	305	31	18,6	20	22	5	5,65		67	4	8,5	11,64	
142	Breast of the <i>Peacock</i>	γ	4	316	21	5,2	21	5	24	5,28		66	28	19,8	14,50	
143	Preceding Shoulder of <i>Aquarius</i>	β	3	319	35	53,7	21	18	24	3,18		6	39	22,4	15,35	
144	Preceding in the Tail of the <i>Goat</i>	γ	3	321	27	12,5	21	25	49	3,35		17	46	42,0	15,66	
145	Following in the Tail of the <i>Goat</i>	δ	3	323	18	8,6	21	23	13	3,33		17	14	48,3	16,06	
146	Head of the <i>Crane</i>	γ	3	324	40	33,5	21	38	42	3,70		38	31	28,7	16,35	
147	Preceding Wing of the <i>Crane</i>	α	2	328	5	7,7	21	52	21	3,98		48	9	21,4	17,00	
148	Following Shoulder of <i>Aquarius</i>	α	3	328	14	0,6	21	52	56	3,10		1	31	25,9	17,03	
149	Beak of the <i>Toucan</i>	α	3	330	16	46,0	22	1	7	4,31		61	29	34,0	17,40	
150	Following Arm of <i>Aquarius</i>	γ	3	332	11	5,6	22	8	44	3,09		2	38	13,5	17,71	
151	Thigh of the <i>Crane</i>	β	3	336	54	3,2	22	27	36	3,68		48	10	51,0	18,43	
152	Water of the <i>Water-Bearer</i>	λ	4	339	53	27,8	22	39	34	3,16		8	54	8,2	18,82	
153	Leg of the <i>Water-Bearer</i>	δ	3	340	20	14,6	22	41	21	3,22		17	8	39,9	18,87	
154	Brightest of the Southern <i>Fishes</i> . <i>Fomalhaut</i>	α	1	340	56	41,7	22	43	47	3,34		30	56	24,8	18,94	
155	Water of the <i>Water-Bearer</i>	ϕ	4	354	20	23,1	23	1	22	3,13		7	23	31,2	19,38	

SMALL

SMALL EQUATIONS OF CLOCK-TIME, and of the SUN'S Place and Declination, for every 4th Year from 1750, (or from any common, or Leap Year after) to 1800, N. S. Which will serve, with contrary Sign, back to 1700, N. or O. Style. According to M. De la Caille's Motions.

Days.	January.			February.			March.			April.			May.			June.		
	Eqn.	Eq. °	Eq. °	Eqn.	Eq. °	Eq. °	Eqn.	Eq. °	Eq. °	Eqn.	Eq. °	Eq. °	Eqn.	Eq. °	Eq. °	Eqn.	Eq. °	Eq. °
	Time.	Place.	Decl.	Time.	Place.	Decl.	Time.	Place.	Decl.	Time.	Place.	Decl.	Time.	Place.	Decl.	Time.	Place.	Decl.
	+	+	-	-	+	-	-	+	-	-	+	+	-	+	+	+	+	+
	th.	"	"	th.	"	"	th.	"	"	th.	"	"	th.	"	"	th.	"	"
1	19	1 44	0 10	2	1 47	0 33	41	1 47	0 43	39	1 48	0 41	2	1 48	0 29	47	1 51	0 14
2	17	1 44	0 12	4	1 47	0 32	41	1 47	0 43	38	1 48	0 41	1	1 48	0 29	47	1 52	0 13
3	16	1 44	0 13	6	1 47	0 34	43	1 47	0 43	37	1 48	0 41	-0	1 48	0 29	46	1 52	0 12
4	17	1 44	0 14	7	1 47	0 34	44	1 47	0 44	36	1 47	0 41	+1	1 48	0 28	47	1 52	0 11
5	16	1 44	0 14	9	1 47	0 34	45	1 47	0 44	35	1 48	0 41	2	1 48	0 28	47	1 52	0 11
6	18	1 45	0 15	11	1 47	0 35	45	1 47	0 43	34	1 47	0 41	3	1 48	0 28	48	1 52	0 9
7	17	1 45	0 17	12	1 47	0 36	45	1 47	0 44	34	1 48	0 40	4	1 48	0 27	48	1 52	0 9
8	17	1 45	0 17	13	1 47	0 36	46	1 47	0 45	33	1 47	0 40	6	1 49	0 27	46	1 52	0 9
9	17	1 45	0 18	14	1 47	0 37	48	1 47	0 44	33	1 47	0 39	7	1 50	0 27	50	1 52	0 8
10	17	1 45	0 18	15	1 47	0 37	47	1 47	0 45	32	1 47	0 40	7	1 50	0 27	49	1 52	0 8
11	18	1 45	0 18	17	1 47	0 38	48	1 47	0 45	32	1 47	0 39	8	1 50	0 27	50	1 52	0 6
12	17	1 45	0 20	18	1 47	0 40	47	1 48	0 45	30	1 47	0 39	9	1 50	0 26	51	1 52	0 6
13	18	1 45	0 21	17	1 47	0 40	47	1 47	0 45	29	1 47	0 39	10	1 50	0 25	52	1 52	0 5
14	17	1 45	0 21	19	1 47	0 40	48	1 48	0 45	27	1 47	0 39	12	1 50	0 26	52	1 52	0 4
15	16	1 45	0 22	20	1 47	0 41	46	1 47	0 47	26	1 47	0 39	13	1 49	0 25	52	1 52	0 3
16	16	1 46	0 22	22	1 47	0 41	45	1 48	0 45	24	1 47	0 38	16	1 50	0 25	52	1 52	0 3
17	13	1 46	0 23	24	1 47	0 41	44	1 48	0 45	23	1 47	0 38	18	1 50	0 24	53	1 52	0 2
18	12	1 46	0 24	25	1 47	0 41	42	1 48	0 44	22	1 48	0 37	20	1 50	0 23	53	1 52	0 1
19	12	1 46	0 24	27	1 47	0 41	39	1 48	0 44	21	1 47	0 36	22	1 50	0 23	53	1 52	0 1
20	11	1 46	0 25	29	1 47	0 41	36	1 48	0 44	19	1 48	0 36	24	1 50	0 21	52	1 52	0 0
21	9	1 46	0 26	29	1 47	0 42	35	1 49	0 44	17	1 48	0 35	25	1 50	0 20	51	1 52	-
22	10	1 46	0 27	30	1 47	0 42	35	1 48	0 43	15	1 48	0 34	28	1 50	0 20	52	1 52	0 1
23	10	1 46	0 28	31	1 47	0 42	35	1 48	0 43	13	1 48	0 34	30	1 50	0 18	52	1 53	0 1
24	8	1 46	0 27	31	1 47	0 42	35	1 48	0 43	11	1 48	0 32	32	1 50	0 18	52	1 53	0 1
25	7	1 46	0 28	33	1 47	0 43	36	1 48	0 43	9	1 48	0 31	33	1 50	0 17	52	1 53	0 1
26	6	1 46	0 29	38	1 47	0 43	36	1 48	0 43	8	1 48	0 30	36	1 50	0 17	53	1 53	0 2
27	4	1 46	0 29	37	1 47	0 43	37	1 48	0 42	6	1 48	0 30	39	1 51	0 16	52	1 52	0 2
28	2	1 47	0 29	39	1 47	0 44	38	1 48	0 42	5	1 48	0 29	41	1 51	0 17	54	1 52	0 2
29	1	1 47	0 30	41	1 47	0 43	39	1 48	0 42	3	1 48	0 29	43	1 52	0 16	53	1 52	0 2
30	+1	1 47	0 31				39	1 48	0 42	2	1 48	0 29	45	1 52	0 15	53	1 52	0 3
31	-0	1 47	0 31				41	1 48	0 42				47	1 52	0 14			
	July.			August.			September.			October.			November.			December.		
	+	+	-	+	+	-	-	+	-	-	+	+	-	+	+	+	+	+
	Time.	Place.	Decl.	Time.	Place.	Decl.	Time.	Place.	Decl.	Time.	Place.	Decl.	Time.	Place.	Decl.	Time.	Place.	Decl.
	th.	"	"	th.	"	"	th.	"	"	th.	"	"	th.	"	"	th.	"	"
1	51	1 52	0 4	15	1 52	0 26	31	1 47	0 39	45	1 45	0 41	43	1 44	0 32	7	1 46	0 17
2	53	1 51	0 4	15	1 52	0 25	34	1 47	0 39	43	1 45	0 41	42	1 44	0 31	9	1 46	0 16
3	46	1 51	0 5	16	1 52	0 28	36	1 47	0 40	44	1 45	0 41	41	1 44	0 31	12	1 46	0 16
4	45	1 51	0 6	16	1 52	0 28	38	1 47	0 39	42	1 45	0 41	39	1 44	0 31	14	1 46	0 17
5	41	1 51	0 8	16	1 53	0 29	40	1 47	0 40	41	1 45	0 41	39	1 45	0 31	16	1 46	0 15
6	39	1 51	0 8	17	1 53	0 29	42	1 47	0 40	38	1 45	0 40	36	1 44	0 32	18	1 46	0 14
7	36	1 51	0 10	17	1 53	0 30	44	1 46	0 40	38	1 45	0 40	33	1 44	0 32	18	1 46	0 13
8	34	1 51	0 11	17	1 53	0 30	46	1 46	0 40	38	1 45	0 41	31	1 45	0 32	19	1 46	0 13
9	34	1 50	0 11	18	1 53	0 31	47	1 46	0 40	37	1 46	0 41	28	1 44	0 31	19	1 46	0 11
10	35	1 50	0 13	16	1 53	0 32	48	1 46	0 40	39	1 45	0 41	26	1 45	0 30	22	1 47	0 11
11	30	1 50	0 14	14	1 53	0 32	49	1 45	0 41	39	1 46	0 41	22	1 45	0 30	21	1 47	0 9
12	29	1 50	0 15	13	1 52	0 32	50	1 45	0 41	39	1 45	0 40	22	1 46	0 29	21	1 46	0 9
13	28	1 50	0 15	11	1 52	0 34	50	1 45	0 42	40	1 46	0 40	18	1 45	0 29	21	1 46	0 7
14	27	1 51	0 17	8	1 52	0 34	51	1 45	0 42	40	1 45	0 40	16	1 46	0 28	22	1 46	0 6
15	26	1 51	0 16	5	1 52	0 35	51	1 46	0 42	41	1 45	0 40	15	1 46	0 27	22	1 46	0 5
16	25	1 51	0 19	+2	1 51	0 35	51	1 45	0 42	41	1 45	0 39	14	1 47	0 27	22	1 46	0 4
17	25	1 51	0 19	-1	1 51	0 35	52	1 45	0 43	41	1 44	0 39	13	1 46	0 26	22	1 46	0 2
18	25	1 51	0 20	4	1 50	0 36	51	1 45	0 43	41	1 44	0 39	13	1 46	0 26	22	1 46	0 2
19	26	1 51	0 20	8	1 49	0 36	51	1 45	0 43	42	1 44	0 39	12	1 46	0 25	22	1 46	0 1
20	27	1 51	0 21	11	1 49	0 36	51	1 45	0 43	42	1 44	0 38	10	1 46	0 25	23	1 45	0 0
21	26	1 51	0 21	13	1 49	0 37	52	1 45	0 43	43	1 44	0 38	8	1 47	0 24	23	1 45	-
22	26	1 51	0 22	15	1 49	0 37	53	1 45	0 43	43	1 44	0 37	9	1 46	0 23	24	1 45	0 2
23	24	1 51	0 22	17	1 48	0 37	53	1 45	0 43	44	1 43	0 36	4	1 46	0 22	24	1 45	0 3
24	23	1 51	0 23	19	1 49	0 38	53	1 45	0 43	45	1 43	0 35	-1	1 46	0 22	24	1 45	0 3
25	19	1 52	0 24	19	1 48	0 38	52	1 45	0 43	45	1 43	0 35	+1	1 45	0 21	24	1 45	0 4
26	18	1 51	0 25	21	1 48	0 39	51	1 45	0 43	45	1 43	0 35	3	1 46	0 20	24	1 45	0 5
27	17	1 51	0 25	22	1 48	0 38	49	1 45	0 42	45	1 43	0 35	3	1 46	0 19	23	1 44	0 6
28	16	1 51	0 25	24	1 48	0 39	48	1 46	0 43	46	1 43	0 35	4	1 45	0 19	22	1 44	0 7
29	14	1 51	0 25	26	1 47	0 38	46	1 46	0 42	47	1 43	0 34	6	1 46	0 19	21	1 44	0 8
30	14	1 52	0 26	28	1 47	0 39	45	1 46	0 42	45	1 43	0 34	8	1 46	0 17	19	1 44	0 9
31	14	1 52	0 27	30	1 47	0 39	44	1 43	0 35	44	1 43	0 35				17	1 44	0 10

N. B. The Equation of Declination for every 4 Julian Years adds in the Sun's Passage from the Equator to the Solstices; but subtracts in the Passage from the Solstices to the Equator. —The Equation of the Sun's Pl. for every 4 Years continually adds. —The Equation of Clock-Time, in Thirds, for every 4 Years, adds and subtracts, alternately. From Jan. 1st, it adds to Jan. 30; —subt. fr. thence to May 3. adds fr. thence to 16 Aug. sub. fr. thence to 24 Nov. adds fr. thence to 30 Jan. —The Signs printed are for Years forward; and the contrary Signs serve for Yrs. back.

SMALL

SMALL EQUATIONS continued, &c. according to M. De la Caille.

EQUATION of the SUN's Right Ascension in Time for every four Years forward, serving with a contrary Sign and Years forward, for the Equation of the Distance in Time, of the *Section of Aries*, from the Meridian, for finding the Time of Stars passing the Meridian by Addition; being the Complement of the SUN's Right Ascension in Time.

N. B. For Years back use the contrary Sign.

Days	Jan.	Feb.	Mar.	Apr.	May.	June	July	August	Sept.	Oct.	Nov.	Dec.
	+	+	+	+	+	+	+	+	+	+	+	+
	s.	s.	s.	s.	s.	s.	s.	s.	s.	s.	s.	s.
1	7	7	6	6	7	8	9	7	7	7	7	7
11	7	7	6	6	7	8	9	7	7	7	7	7
21	7	6	6	6	7	9	8	7	7	7	7	7

The SUN's apparent Semi-Diameter for every fifth Day, in Degrees and Time, for 1760, or Leap Year.

Days	January.	February.	March.	April.	May.	June.
	m. s. t.	m. s. t.	m. s. t.	m. s. t.	m. s. t.	m. s. t.
1	16 20 20	1 10 50	16 17 50	1 8 2	16 11 55	1 5 8
6	16 20 15	1 10 33	16 17 0	1 7 30	16 10 15	1 4 50
11	16 20 0	1 10 12	16 50 55	1 6 55	16 8 55	1 4 34
16	16 19 40	1 9 45	16 14 55	1 6 23	16 7 35	1 4 21
21	16 19 10	1 9 15	16 13 55	1 5 54	16 6 5	1 4 14
26	16 18 40	1 8 42	16 12 50	1 5 26	16 4 50	1 4 12
Days	July.	August.	September.	October.	November.	December.
	m. s. t.	m. s. t.	m. s. t.	m. s. t.	m. s. t.	m. s. t.
1	15 47 50	1 8 31	15 50 10	1 6 24	15 56 10	1 4 15
6	15 48 0	1 8 17	15 51 9	1 5 34	15 57 20	1 4 5
11	15 48 15	1 8 0	15 52 50	1 5 10	15 58 40	1 3 57
16	15 48 35	1 7 41	15 52 40	1 4 48	16 0 0	1 3 54
21	15 48 55	1 7 20	15 53 35	1 4 31	16 1 20	1 3 56
26	15 49 25	1 6 55	15 54 45	1 4 17	16 2 50	1 4 1

EQUATION of the Horizontal SEMI-DIAMETER of the MOON, to the apparent Semi-diameter of the MOON, in ALTITUDE.
The Horizontal Semi-Diameter to the Horizontal Parallax of the Moon being considered as 15' to 55' 40".

Argument. The Moon's Altitude, and Horizontal Semi-Diameter.

Horizontal Semi-Diameter of the MOON.

D	14' 40"	15' 0"	15' 20"	15' 40"	16' 0"	16' 20"	16' 40"	17' 0"
Alt.	+	Dif.	+	Dif.	+	Dif.	+	Dif.
0	0 0	73	0 0	76	0 0	80	0 0	83
5	1 13	72	1 16	76	1 23	82	1 27	86
10	2 25	71	2 32	74	2 45	82	2 53	84
15	3 36	70	3 46	73	4 7	79	4 17	83
20	4 46	67	4 59	70	5 26	77	5 40	81
25	5 53	65	6 9	68	6 43	74	7 1	76
30	6 58	61	7 17	64	7 57	70	8 17	73
35	7 59	58	8 21	61	8 44	63	9 7	66
40	8 57	54	9 22	56	9 47	59	10 13	61
45	9 51	49	10 18	52	10 46	54	11 14	57
50	10 40	44	11 10	46	11 40	48	12 11	52
55	11 24	40	11 56	41	12 28	43	13 0	48
60	12 4	33	12 37	35	13 11	37	13 48	36
65	12 37	28	13 12	29	13 48	30	14 24	32
70	13 5	22	13 41	24	14 18	24	14 56	25
75	13 27	16	14 5	16	14 42	18	15 21	18
80	13 43	10	14 21	10	15 0	10	15 39	12
85	13 53	3	14 31	3	15 10	3	15 51	5
90	13 56	3	14 34	3	15 13	3	15 54	3

Apparent OBLIQUITIES of the Ecliptic and Equator, for January 1, in the Years following.

1748 23° 28' 20"	1757 23° 28' 21"	1760 23° 28' 24"	1763 23° 28' 18"
1755 23 28 16	1758 23 28 23	1761 23 28 23	1764 23 28 15
1756 23 28 19	1759 23 28 24	1762 23 28 21	1765 23 28 11

Still decreasing (See p. 33) 23 28 20 being the Mean Obliquity, according to the French Authority.

EXAMPLE. To find the Moon's apparent Semi-Diameter in Altitude, and apparent Altitude of the Moon's Center, from the Horizontal Semi-Diameter 16' 15", and apparent Altitude of the Moon's upper Limb, 49° 30', observed?

Alt. D 49° 30' 0" observed.

— 16 28

49 13 32 Ap. Alt. D's Center, reqd.

Horizontal or true Semi-Diameter 16' 15"
Under 16' 20", and against 49° 30' Altitude, you find + 13 Equation

Apparent Semi-Diameter D in Altitude 16 28 required.

TABLE of CLOCK-TIME at NOON by the SUN, the SUN's Right Ascension, and his Complement of Right Ascension, or Distance of Aries from the Meridian, in Time, according to M. De la Caille, reduced to Greenwich Observatory for 1760.

JANUARY.												FEBRUARY.												MARCH.															
Days	Clock Time at Noon by the ☉			Dif.	☉ Right Ascension in Time.			Dif.	Co. ☉ R. A. in Time, or ☉ fr. Mer.			Days	Clock Time at Noon by the ☉			Dif.	☉ Right Ascension in Time.			Dif.	Co. ☉ R. A. in Time, or ☉ fr. Mer.			Days	Clock Time at Noon by the ☉			Dif.	☉ Right Ascension in Time.			Dif.	Co. ☉ R. A. in Time, or ☉ fr. Mer.						
	h.	m.	s.		h.	m.	s.		h.	m.	s.		h.	m.	s.		h.	m.	s.		h.	m.	s.		h.	m.	s.		h.	m.	s.		h.	m.	s.	h.	m.	s.	
1	12	4	27		18	50	46		5	9	14		12	14	10		21	2	45		2	57	15		12	12	33		22	51	30		1	8	30				
2	12	4	55	28	18	55	10	4	5	4	50		12	14	17	6	21	6	49	4	4	2	53	11		12	12	21	14	22	55	14	3	44	1	4	46		
3	12	5	22	27	18	59	35	4	5	0	25		12	14	23	5	21	10	52	4	1	2	49	8		12	12	7	13	22	58	58	3	44	1	1	2		
4	12	5	49	27	18	3	59	4	4	56	1		12	14	28	4	21	14	53	4	1	2	45	7		12	11	54	13	23	2	41	0	57	19				
5	12	6	16	26	19	8	22	4	4	51	38		12	14	32	4	21	18	54	4	0	2	41	6		12	11	40	14	23	6	22	3	41	0	53	38		
6	12	6	42		19	12	44		4	47	16		12	14	36		21	22	54	3	59	2	37	6		12	11	25	15	23	10	4	42	0	49	56			
7	12	7	8	26	19	17	7	4	4	42	53		12	14	38	2	21	26	53	3	58	2	33	7		12	11	10	15	23	13	46	3	41	0	46	14		
8	12	7	33	25	19	21	29	4	4	38	31		12	14	40	1	21	30	51	3	58	2	29	9		12	10	55	16	23	17	27	3	41	0	42	33		
9	12	7	58	25	19	25	49	4	4	34	11		12	14	41	0	21	34	49	3	57	2	25	11		12	10	39	16	23	21	8	3	41	0	38	52		
10	12	8	22	24	19	30	10	4	4	29	50		12	14	41	0	21	38	46	3	55	2	21	14		12	10	23	16	23	24	49	3	40	0	35	11		
11	12	8	45	23	19	34	30	4	4	25	30		12	14	41	1	21	42	41	3	55	2	17	19		12	10	7	17	23	28	29	3	40	0	31	31		
12	12	9	8	22	19	38	49	4	4	21	11		12	14	40	2	21	46	36	3	55	2	13	24		12	9	50	17	23	32	9	3	39	0	27	51		
13	12	9	30	22	19	43	9	4	4	16	51		12	14	37	3	21	50	31	3	55	2	9	29		12	9	33	17	23	35	48	3	39	0	24	12		
14	12	9	52	21	19	47	27	4	4	12	33		12	14	35	4	21	54	25	3	54	2	5	35		12	9	16	17	23	39	28	3	39	0	20	32		
15	12	10	13	20	19	51	45	4	4	8	15		12	14	31	4	21	58	18	3	53	2	1	42		12	8	59	18	23	43	7	3	38	0	16	53		
16	12	10	33	19	19	56	2	4	4	3	58		12	14	27	5	22	2	11	3	51	1	57	49		12	8	41	17	23	46	45	3	39	0	13	15		
17	12	10	52	19	20	0	17	4	3	59	43		12	14	22	6	22	6	2	3	51	1	53	58		12	8	24	18	23	50	24	3	38	0	9	36		
18	12	11	11	18	20	4	33	4	3	55	27		12	14	16	6	22	9	53	3	49	1	50	7		12	8	6	18	23	54	2	3	39	0	5	58		
19	12	11	29	17	20	8	48	4	3	51	12		12	14	10	7	22	13	42	3	50	1	46	18		12	7	48	19	23	57	41	3	38	0	2	19		
20	12	11	46	17	20	13	2	4	3	46	58		12	13	3	7	22	17	32	3	49	1	42	28		12	7	29	18	23	0	1	19	3	38	23	58	41	
21	12	12	3	15	20	17	15	4	3	42	45		12	13	56	9	22	21	21	3	48	1	38	39		12	7	11	19	23	0	4	57	3	38	23	55	3	
22	12	12	18	15	20	21	27	4	3	38	33		12	13	47	9	22	25	9	3	47	1	34	51		12	6	52	18	23	0	8	35	3	38	23	51	25	
23	12	12	33	14	20	25	38	4	3	34	22		12	13	38	9	22	28	56	3	48	1	31	4		12	6	34	19	23	0	12	13	3	36	23	47	47	
24	12	12	47	14	20	29	49	4	3	30	11		12	13	29	10	22	32	44	3	46	1	27	16		12	6	15	19	23	0	15	49	3	39	23	44	11	
25	12	13	1	12	20	33	59	4	3	26	1		12	13	19	10	22	36	30	3	46	1	23	30		12	5	57	18	23	0	19	28	3	38	23	40	32	
26	12	13	13	13	20	38	8	4	3	21	52		12	13	8	11	22	40	16	3	45	1	19	44		12	5	38	19	23	0	23	6	3	38	23	36	54	
27	12	13	26	13	20	42	16	4	3	17	44		12	12	57	12	22	44	1	3	45	1	15	59		12	5	19	19	23	0	26	44	3	38	23	33	16	
28	12	13	35	10	20	46	24	4	3	13	36		12	12	45	12	22	47	46	3	45	1	12	14		12	5	0	18	23	0	30	21	3	37	23	29	39	
29	12	13	45	10	20	50	31	4	3	9	29		In Leap Year take out for a Day sooner in January and February only.—For 1, 2, 3, Years after Leap Year take out for $\frac{1}{4}$, $\frac{1}{2}$, $\frac{3}{4}$ of a Day sooner in all the Months (S.p. 194).														12	4	42	19	23	0	33	58	3	37	23	26	1
30	12	13	55	8	20	54	37	4	3	5	23		12	4	23	18	23	0	37	37	3	39	23	22	23		12	4	23	18	23	0	37	37	3	39	23	22	23
31	12	14	3		20	58	41	4	3	1	19		12	14	3		12	4	5							12	4	5	18	23	0	41	16	3	39	23	18	44	

APRIL.												MAY.												JUNE.												
Days	h.	m.	s.	Dif.	h.	m.	s.	Dif.	h.	m.	s.	Days	h.	m.	s.	Dif.	h.	m.	s.	Dif.	h.	m.	s.	Days	h.	m.	s.	Dif.	h.	m.	s.	Dif.	h.	m.	s.	Dif.
1	12	3	46		0	44	54		23	15	6		2	36	10		21	23	56		11	57	24		4	39	0		19	21	0		19	21	0	
2	12	3	28	18	0	48	32	3	38	23	11	28		2	39	59	3	49	21	20	1	11	57	32		4	43	5	6	19	16	55		19	16	55
3	12	3	10	18	8	52	10	3	38	23	7	50		2	43	49	3	50	21	16	11	11	57	42		4	47	11	7	19	12	49		19	12	49
4	12	2	52	18	0	55	49	3	38	23	4	11		2	47	40	3	51	21	12	20	11	57	52		4	51	18	7	19	8	42		19	8	42
5	12	2	34	18	0	59	27	3	39	23	0	33		2	51	31	3	52	21	8	29	11	58	3		4	55	25	7	19	4	35		19	4	35
6	12	2	16		1	3	6	3	39	22	56	54		2	55	23	3	52	21	4	37	11	58	13		4	59	32	7	19	0	28		19	0	28
7	12	1	59	17	1	6	45	3	39	22	53	15		2	59	15	3	52	21	0	45	11	58	24		5	3	39	7	18	56	21		18	56	21
8	12	1	42	17	1	10	24	3	39	22	49	36		3	3	7	3	54	20	56	53	11	58	35		5	7	47	8	18	52	13		18	52	13
9	12	1	25	16	1	14	4	3	40	22	45	56		3	7	1	3	53	20	52	2	11														

TABLE of Clock-Time at NOON by the SUN, the SUN's Right Ascension, and his Complement of Right Ascension, or Distance of Aries from the Meridian, in Time, according to M. De la Caille, reduced to Greenwich Observatory for 1760.

JULY.										AUGUST.										SEPTEMBER.																
Days	Clock Time at Noon by the ☉			Dif.	☉ Right Ascension in Time.			Dif.	Co. ☉ R.A. in Time, or ☿ fr. Mer.			Days	Clock Time at Noon by the ☉			Dif.	☉ Right Ascension in Time.			Dif.	Co. ☉ R.A. in Time, or ☿ fr. Mer.			Days	Clock Time at Noon by the ☉			Dif.	☉ Right Ascension in Time.			Dif.	Co. ☉ R.A. in Time, or ☿ fr. Mer.			
	h.	m.	s.		h.	m.	s.		h.	m.	s.		h.	m.	s.		h.	m.	s.		h.	m.	s.		h.	m.	s.		h.	m.	s.		h.	m.	s.	h.
1	12	3	20	12	6	43	16	4	7	17	16	44	12	5	47	4	8	47	55	3	53	15	12	5	11	59	33	19	10	43	53	3	37	13	16	7
2	12	3	32	10	6	47	23	4	8	17	12	37	12	5	43	5	8	51	48	3	52	15	8	12	11	59	14	19	10	47	30	3	37	13	12	30
3	12	3	42	11	6	51	31	1	8	17	8	29	12	5	38	5	8	55	40	3	51	15	4	20	11	58	55	20	10	51	7	3	38	13	8	53
4	12	3	53	11	6	55	39	4	7	17	4	21	12	5	33	5	8	59	31	3	51	15	0	29	11	58	35	19	10	54	45	3	37	13	5	15
5	12	4	4	9	6	59	46	4	7	17	0	14	12	5	27	6	9	3	22	3	50	14	56	38	11	58	16	20	10	58	22	3	35	13	1	38
6	12	4	13	10	7	3	53	4	6	16	56	7	12	5	21	7	9	7	12	3	49	14	52	48	11	57	56	20	11	1	58	3	37	12	58	2
7	12	4	23	9	7	7	59	4	6	16	52	1	12	5	14	8	9	11	1	3	49	14	48	59	11	57	36	20	11	5	35	3	36	12	54	25
8	12	4	32	9	7	12	5	4	5	16	47	55	12	5	6	9	9	14	50	3	49	14	45	10	11	57	16	21	11	9	11	3	36	12	50	49
9	12	4	41	9	7	16	10	4	5	16	43	50	12	4	58	9	9	18	39	3	48	14	41	21	11	56	55	20	11	12	47	3	35	12	47	13
10	12	4	50	8	7	20	15	4	5	16	39	45	12	4	49	9	9	22	27	3	47	14	37	33	11	56	35	21	11	16	22	3	36	12	43	38
11	12	4	58	7	7	24	20	4	4	16	35	40	12	4	40	10	9	26	14	3	46	14	33	46	11	56	14	21	11	19	58	3	36	12	40	2
12	12	5	5	8	7	28	24	4	4	16	31	36	12	4	30	10	9	30	0	3	46	14	30	0	11	55	53	20	11	23	34	3	35	12	36	26
13	12	5	13	6	7	32	28	4	4	16	27	32	12	4	20	10	9	33	47	3	47	14	26	13	11	55	33	20	11	27	9	3	35	12	32	51
14	12	5	19	7	7	36	31	4	3	16	23	29	12	4	9	11	9	37	32	3	45	14	22	28	11	55	12	21	11	30	45	3	36	12	29	15
15	12	5	26	5	7	40	34	4	2	16	19	26	12	3	58	13	9	41	17	3	44	14	18	43	11	54	51	21	11	34	21	3	35	12	25	39
16	12	5	31	5	7	44	36	4	2	16	15	24	12	3	45	12	9	45	1	3	44	14	14	59	11	54	30	21	11	37	56	3	35	12	22	4
17	12	5	36	5	7	48	38	4	1	16	11	22	12	3	33	13	9	48	45	3	44	14	11	15	11	54	9	21	11	41	31	3	36	12	18	29
18	12	5	41	4	7	52	39	4	0	16	7	21	12	3	20	14	9	52	29	3	43	14	7	31	11	53	48	21	11	45	7	3	35	12	14	53
19	12	5	45	4	7	56	39	4	0	16	3	21	12	3	6	14	9	56	12	3	43	14	3	48	11	53	27	20	11	48	42	3	36	12	11	18
20	12	5	49	4	8	0	39	3	59	15	59	21	12	2	52	14	9	59	55	3	42	14	0	5	11	53	7	21	11	52	18	3	34	12	7	42
21	12	5	52	3	8	4	38	4	0	15	55	22	12	2	38	15	10	3	37	3	42	13	56	23	11	52	46	21	11	55	54	3	36	12	4	6
22	12	5	54	2	8	8	38	3	58	15	51	22	12	2	23	15	10	7	19	3	41	13	52	41	11	52	25	20	11	59	30	3	35	12	0	30
23	12	5	56	2	8	12	36	3	58	15	47	24	12	2	8	16	10	11	0	3	40	13	49	0	11	52	5	21	12	3	5	3	37	11	56	55
24	12	5	58	2	8	16	34	3	58	15	43	26	12	1	52	19	10	14	40	3	40	13	45	20	11	51	44	20	12	6	42	3	36	11	53	18
25	12	5	58	0	8	20	32	3	57	15	39	28	12	1	36	17	10	18	20	3	40	13	41	40	11	51	24	20	12	10	18	3	36	11	49	42
26	12	5	58	0	8	24	29	3	56	15	35	31	12	1	19	17	10	22	0	3	40	13	38	0	11	51	4	20	12	13	55	3	36	11	46	5
27	12	5	58	0	8	28	25	3	55	15	31	35	12	1	2	17	10	25	39	3	39	13	34	21	11	50	44	20	12	17	31	3	36	11	42	29
28	12	5	57	1	8	32	20	3	55	15	27	40	12	0	45	17	10	29	19	3	39	13	30	41	11	50	25	19	12	21	8	3	37	11	38	52
29	12	5	55	2	8	36	15	3	54	15	23	45	12	0	28	18	10	32	58	3	39	13	27	2	11	50	6	19	12	24	45	3	38	11	35	15
30	12	5	53	3	8	40	9	3	53	15	19	51	12	0	10	19	10	36	37	3	37	13	23	23	11	49	47	19	12	28	23	3	38	11	31	37
31	12	5	50	3	8	44	2			15	15	58	11	59	51			10	40	14																
OCTOBER.										NOVEMBER.										DECEMBER.																
1	11	49	28	19	12	32	0	3	38	11	28	0	11	43	49	0	14	28	33	3	57	9	31	27	11	49	43	23	16	32	46	4	20	7	27	14
2	11	49	9	18	12	35	38	3	38	11	24	22	11	43	49	0	14	32	30	3	57	9	27	30	11	50	6	23	16	37	6	4	21	7	22	54
3	11	48	51	18	12	39	17	3	38	11	20	43	11	43	50	1	14	36	27	3	57	9	23	33	11	50	31	24	16	41	27	4	21	7	18	33
4	11	48	33	18	12	42	55	3	38	11	17	5	11	43	51	1	14	40	26	3	59	9	19	34	11	50	55	24	16	45	48	4	22	7	14	12
5	11	48	15	17	12	46	34	3	39	11	13	26	11	43	53	2	14	44	25	3	59	9	15	35	11	51	21	25	16	50	10	4	23	7	9	50
6	11	47	58	16	12	50	13	3	40	11	9	47	11	43	57	4	14	48	24	4	1	9	11	36	11	51	46	26	16	54	33	4	23	7	5	27
7	11	47	42	16	12	53	53	3	41	11	6	7	11	44	1	4	14	52	25	4	1	9	7	35	11	52	12	27	16	58	56	4	24	7	1	4
8	11	47	26	16	12	57	34	3	40	11	2	26	11	44	5	4	14	56	26	4	2	9	3	34	11	52	39	27	17	3	20	4	24	6	56	40
9	11	47	10	15	13	1	14	3	41	10	58	46	11	44	11	6	15	0	28	4	2	8														

Astronomical QUESTIONS (applied to NAVIGATION) answered by the Tables.

QUEST. I. To find the Instant of a Star passing the Meridian on a given Day?

RULE. Add the Complement of the Sun's R. A. in Time, for a given Place, and Day at Noon (being the Distance in Time of the Section of *Aries* from the Meridian, found in Tables, p. 216, 217.) to the R. A. of the Star, in Time, found in the Catalogue of Stars) and the Sum thereof (rejecting 24 Hours when above that Number) will be the near Time, from Noon, of the Star's Passage through the Meridian, required.

To find the Time accurately. Subtract from the near Time, already found, the proportional Decrease, in Time, of the Distance of the Section of *Aries* from the Meridian, to that near Time (the Dist. of the said Section, or Complement of the R. A. of the Sun, having hitherto been only reckoned from the preceding Noon) and the Remainder will be the exact Time of the Star's Passage, required.

EXAMPLE. Required the correct Time of Aldebaran, southing, or passing the Meridian of Greenwich Observatory, on December 30, 1760?

R. A. of Aldebaran, Jan. 1, 1750, by Tab.	h m s
p. 208.	4 21 36
11 Yrs. Equat. (fr. Jan. 1750 to Dec. 1760)	
at 3 ^d .43 yearly	+ 38

R. A. of Aldebaran, 1760, Dec. 30 ^d	4 22 14
By Tab. p. 217. Comp. R. A. of the Sun, at Noon, on that Day	5 19 11

Near Time of Aldebaran's Southing, fr. Noon 9 41 25 Sum.

NOW, as 24 ^h is to 4 ^m 25 ^s (diurnal Dec. fr. <i>Aries</i> , or Comp. R. A. Sun fr. the Merid. fr. 30 to 31 Dec. being the same as the diurnal Incr. of R. A. Sun in Col. of Diff. R. A. Sun) so is 9 ^h 41 ^m 25 ^s , near Time of Southing fr. Noon, to	Decr. of Comp. R.A. Sun.
	- 1 47

Hence, the correct Time of Aldebaran's Southing, at Greenwich Observatory, 1760, } 9 39 38 P. M.
December 30 required.

If the Time of the same Star's Southing was required, at Antigua, in the West Indies, or at any Place at Sea 60° West Longitude from Greenwich Observatory, the following Rule must be regarded.

FIRST, Any fixed Star is observed, apparently, to accelerate, or return to the Meridian of the same Place sooner, each Night, than on the former Night, by the Increase of the Sun's R. A. in Time, eastward, less the annual or progressive Motion of the Star, in each diurnal Revolution of the Earth, eastward. That is, each Star returns nearly 3^m 56^s sooner to the same Meridian, in about 23^h 56^m 4^s mean solar Time, (see p. 29.) while 360° of the Earth's Equator is revolved to the eastward; causing thereby an apparent Motion or Revolution of the whole Heavens, to the Westward. And, while 360°, + the Increase R. A. Sun for 24 Hours, is really revolved Eastward by the

Earth's Equator, and $\frac{50''}{365}$ is progressively moved over, easterly, by

the Star, in 24 Hours, when the Sun returns to the same Meridian, then the Star, that was the Day before on the Meridian with the Sun, will be got the whole easterly Increase of the Sun's right Ascension, less the easterly Motion of the Star, in that Time, to the westward of the Sun; then due South at Noon. Consequently, that Star must pass the Meridian, each successive Day, sooner and sooner than the Sun (passing it at Noon) according to the uniform diurnal, and variable annual Motion of the Earth, easterly; till, a little after the Year's End, the Sun and Star again meet.

This Interval of Time will be proportional to the Difference of the Sun and Star's annual Motions easterly, when the Sun, that departed from the Star, following him at a slow Rate, again overtakes that Star in his annual Circuit, being similar in this Instance, to the Minute Hand departing from, and overtaking the Hour Hand of a Watch; both moving the same Way, in a Circle, at the same Time, and always meeting after the same Interval.

NOW, As the Longitudes of the Earth are measured, in Time, by the Sun's Passage from Noon to Noon, from and to the Meridian of the same Place, on the Earth's Surface, which Place is daily advanced to the eastward, the Difference of the Sun's R. A. in Time, all Places to the westward thereof are advanced to the eastward in that Proportion. And since the Motion of a Star, the same Way, or eastwardly, is inconsiderable for a Day, and as all Stars accelerate with the Increase of R. A. of the Sun, in 24 Hours, (while 360° West Longitude is advanced by the Earth's easterly Rotation, more easterly, by the Dif. Sun's R. A.) therefore, to find the Acceleration of any Star, or its sooner southing, in any Place of West Longitude from First Meridian, we have this

GENERAL RULE. As 24^h West Longitude, or a Revolution of the Earth, is to 4^m 25^s Sun's R. A. in Time for a Day more Motion advanced to the eastward, or that Time by which the same Star sooner returns to the same Meridian, so is any other West Longitude, or 4^h West, the Longitude of Antigua, to the proportional Time, or 44^s, by which that Star is advanced to the Westward, or come sooner to the Meridian of the Second, than to that of the First, Place. And must be subtracted from the Time of the Southing in the first, for the Time of Southing in the second, Place.

The Southing of Aldebaran, at Greenwich, 1760, } h m s	
Dec. 30.	9 39 38
At 60° or 4 ^h West Longitude, proportional Increase R.A. of Sun, or Acceleration, sooner	- 44

Aldebaran souths at Antigua, 60° West of Greenwich, } h m s	
1760, Dec. 30 ^d	9 38 54

N. B. If the Longitude of a Second is Eastward of a first Place, then any Star souths, for that Longitude, proportionally later than at the first Place, as 360° East Longitude is to the Sun's Increase of R. A. for a Day, in Time, being the Southing later for 360° East Longitude, or a Day. This being the Reverse of the former Rule, the proportional Quantity must be added to the Southing in the first Place.

QUEST. II. To find what Star souths nearly at a given Time on a given Day?

RULE. Find (by Tables, p. 216 and 217) the R. A. Sun in Time, for the Day given; to which add the given Time, reckoned forward from Noon, rejecting 24 Hours, when the Sum exceeds that Number. — Seek for the Sum, or nearest Time to it, in the Catalogue of R. A. of Stars, and against it you will find the Name of the Star, nearly southing required?

EXAMPLE. Required what Star nearly souths, at 9^h 41^m 25^s at Night, in December 30, 1760?

By Tab. p. 217. R. A. Sun, at Noon on that Day	18 40 49
	Sum 28 22 14
	- 24
By Tab. p. 208. you find Aldebaran against . . }	4 22 14
R. A. 4 ^h 21 ^m 36 ^s : being the Star required. }	remain.

QUEST. III. To find the Time of Rising and Setting of a given Star, for any Day and Year, at a given Place; the Time of Southing of that Star for the same Place being first found, or given?

RULE. Subtract and add the Star's Semi-Duration Arc, answerable to its Declination, and the Place's Latitude (found by the preceding Tables) from and to the Time of the Star's Southing given, and the Remainder and Sum will be the respective Times of that Star's Rising and Setting, required.

EXAMPLE. Let it be required to find the Time of Rising and Setting of Aldebaran, on December 30, 1760, at Greenwich Observatory, Latitude 51° 28' N. and likewise at Antigua, Latitude 17° 5' N.?

Aldebaran

Astronomical QUESTIONS (applied to NAVIGATION) answered by the Tables.

Aldebaran souths at Greenwich-Observatory, Dec. } h m s
 30, 1760, found by Quest. I. } 9 39 38 P. M.
 Declination of Aldebaran, at that Time, 16°
 $0' 35''$ N. (by Tab. p. 208.) answering to which
 and Lat. $51^{\circ} \frac{1}{2}$ N. The Semi-Duration Arc
 (in Pages 202 and 203) is found } 7 28 30 $\mp$

Aldebaran rises, at Greenwich, Dec. 30^d 2 11 8 Aftern.
 Sets } 17 8 8
 1760, Dec. 31^d 5 8 8 Morn.

Aldebaran souths at Antigua, found by Quest. I. } h m s
 Dec. 30^d } 9 38 54 P. M.
 Sem. Duration Arc answering to Lat. 17° N. }
 and Decl. 16° N. (in p. 200) is found . . . } 6 22 0 $\mp$

Aldebaran rises at Antigua, Dec. 30^d 3 16 54 Aft.
 Sets } 16 0 54
 1760, Dec. 31^d } 4 0 54 Morn.
 required.

QUEST. IV. To find the Time of Sun Rising and Setting
 in any Place, for any Day of the Month, and Year given?

RULE. Subtract and add the Semi-diurnal Arc, answerable to the
 Latitude of the Place and Sun's Declination, for the Day and Year
 given, from and to 12 Hours, or Noon, and the Remainder and Sum
 will be the respective Times of Sun Rising and Setting, required.

EXAMPLE. To find the Time of Sun-Rising and Setting, on April
 15, 1759, at Greenwich-Observatory, and likewise at Antigua, in the
 West-Indies?

Sun souths, in all Places . . . h m s
 12 0 0
 Apr. 15, 1759, 3d Yr. after Lp. (in p. 194)
 ☉'s Decl. $\frac{1}{2}$ Day sooner, is found, accordg
 to Xr. 7269, to be $9^{\circ} 45' 51''$ N. answer-
 able to which, and Lat. $51^{\circ} \frac{1}{2}$ N. (in Pages
 202 and 203) the Semi-diurnal Arc, is found } 6 53 30 $\mp$
 Sun rises at Greenwich, 1759, Apr. 15^d 5 6 30 } allowg for
 Sets } 6 53 30 } Refractn.
 N. B. The Semi-diurnal Arc in Hours, &c. is always the Time of Sun-
 Setting.

Noon . . . h m s
 12 0 0
 Answerable to $9^{\circ} 45' 51''$ N. Decl. Sun and Lat.
 17° N. (in p. 200) Semi-diurnal Arc is }
 found } 6 14 30—Setting }
 At Antigua, 1759, April 15^d 5 45 30 Rising. }
 required.

QUEST. V. To find the Sun's Amplitude of Rising and
 Setting from the East or West Points of the Compass, according
 to any Latitude and Declination of the Sun given; for deter-
 mining the magnetical Variation of the Compass?

RULE. Under the given Latitude N. or S. from Page 204 to 207,
 at the Top of the Pages, and against the given Declination of the Sun,
 N. or S. on the Side of a Page, at the Angle of Meeting, you have,
 at Sight, the Number of Degrees and Minutes of Amplitude Northerly
 from E. or W. Lat. N. but Southerly Lat. South, in the first Half
 Page. Also the Degrees and Minutes of Amplitude Southerly, from E.
 and W. Lat. N. but Northerly Lat. South, in the second half Page.
 By which true Amplitude of the Tables compared with the magnetical
 Amplitude, found by the Azimuth-Compass, at the Sun's Rising or
 Setting, (Refraction being allowed for in the Tables) the Difference,
 or magnetical Variation will be known: the Excess or Defect of Mag-
 netical Amplitude being the Way of Variation.

EXAMPLE I. Required the Sun's Amplitude of Rising and Setting for
 Latitude 20° N. and Declination 20° South?

In the second half Page 205, under 20° N. Lat. and against 20° S.
 Declination, (the inferior Decl. respecting the superior Lat. right up
 and down in the same Line) stands $21^{\circ} 8'$ Amplitude Southerly from
 E. or W. required. This true Amplitude being compared with the
 magnetical Amplitude, the Difference will be the magnetical Variation
 of the Compass, required.

EXAMPLE II. Required the Sun's Amplitude of Rising and Setting
 for Latitude 44° S. and Declination 12° South?

In the first half Page 206, under S. Lat. 44° and against 12° S.
 Declination, stands $17^{\circ} 20'$ Amplitude Southerly from E. or W. re-
 quired; which being compared with the magnetical Amplitude, ob-
 served, the magnetical Variation of the Compass (being the Difference)
 will be known.

N. B. The same Method is to be observed in determining the Amplitudes
 of Rising and Setting of the fixed Stars, of given Declination, when the
 Latitude of the Place is given. By which Stars the magnetical Amplitude
 may be observed, and thence, the magnetical Variation of the Compass is
 found; being the Difference of these two Amplitudes.

QUEST. VI. To find the Length of the longest Day and
 Night in any Latitude, whose Complement is less than the Sun's
 greatest N. or S. Declination?

WHEN the Complement of the Latitude N. or S. is respectively
 equal to the Sun's greatest Declination N. or S. viz. $23^{\circ} 28' 30''$ (the
 Lat. being then $66^{\circ} 31' 30''$) and the Meridian Altitude always equal
 to the Sum of the Comp. Lat. and Sun's Declination to the Pole
 of the same Name, which Altitude is always equal the Depression at
 Midnight) it is evident, that on that Day, the Sun does not set; but
 touches the Edge of the Horizon at the North Part of the Meridian,
 opposite to Noon; and then ascends to the South Part of the Meridian
 at Noon; whose Altitude, on that Day, is Comp. Lat. $+ 23^{\circ} 28' 30''$
 North Declination, and the longest Day there, is just 24 Hours.—
 Therefore, in all Places, where the Complement of Latitude is less than
 the Sun's greatest Declination, on that very Day when the Sun's Declin.
 is equal to the Complement of the Latitude, the Sun will not set, but touch
 the Horizon in the opposite Part of the Meridian to 12 at Noon, (as
 on the longest Day in Latitude $66^{\circ} 31' 30''$) and the Day will be there
 just 24 Hours. In the Sun ascending from which Declination (the
 same with the Complement of Latitude) to the greatest Declination, and
 returning back to the same Declination again, the longest Day in that
 Latitude will be measured; and may be determined at Sight (universally)
 by our Solar Ephemeris from p. 194 to 196, according to what the
 Sun's Declination is; being of the same Name with, and equal to the
 Complement of Latitude.

EXAMPLE. Suppose the Latitude to be the North Cape $71^{\circ} 25'$, in
 which the longest Day is required.

Here the Complement of Latitude $= 18^{\circ} 35' =$ Sun's Declination
 North, answering to May 13, and July 29, N. Style, which Inter-
 val is the longest Day in that Latitude required.

Astronomical QUESTIONS (applied to NAVIGATION) answered by Spherical PROPORTION and LOGARITHMS.

I. QUESTION. To find the Sun's present Declination from his Place in the Ecliptic (Π $2^{\circ} 20' 19'' 40''$) and his greatest Declination ($23^{\circ} 28' 30''$) given?

PROPORTION.		Logarithms.
As Radius		10.0000000
To Sine of the Sun's Long. from Υ $80^{\circ} 10' 40''$		9.9937822
So Sine gr. Declination	$23^{\circ} 28' 30''$	9.6002636

To Sine present Decl. N. required . . . $23^{\circ} 7' 18''$ 9.5940458

N. B. In the above Proportion, or Rule, the Sun's Longitude is always taken from the next equinoctial Point, *Aries*, or *Libra*. Consequently, if the Sun's Place is under 180° , in any of the first six Signs, (*Aries*, *Taurus*, *Gemini*, *Cancer*, *Leo*, *Virgo*, *Libra*) the Declination will be North. If above 180° , in the last six Signs (*Scorpio*, *Sagittary*, *Capricorn*, *Aquarius*, and *Pisces*) the Declination will be South.—This Question is very useful in Navigation.

II. QUESTION. To find the Sun's Longitude from *Aries*, or *Libra*, his present Declination N. or S. being given?

LET the given Declination be $23^{\circ} 7' 18''$ N.

Converse of the former PROPORTION.

As Sine greatest Declination . . . $23^{\circ} 28' 30''$ co.	0.3997364
To Sine present Declination N. or S. $23^{\circ} 7' 18''$	9.5940458
So Radius . . . $90^{\circ} 0' 0''$	10.0000000

To Sine $\odot$'s Lon. fr. Υ , or Δ , reqd. $80^{\circ} 19' 40''$ 9.9937822

N. B. This Question is of Use in Astronomical Observations.

III. QUESTION. To find the Sun's Right Ascension from his Place in the Ecliptic, Π $20^{\circ} 19' 40''$, given?

PROPORTION.		Logarithms.
As Radius		10.0000000
To Tang. Sun's Long. fr. Υ . . . $80^{\circ} 19' 40''$		10.7684433
So Cot. greatest Declination . . . $23^{\circ} 28' 30''$		9.9624801

To Tan. Sun's R. A. from Υ , required $79^{\circ} 28' 25''$ 10.7309234
 When the Sun's Place is in the 1st Quadrant of the Ecliptic, in Υ , δ , Π , then the 4th proportional Arch, is the Sun's right Ascension from *Aries*. If his Place is in the 2d Quadrant, δ , δ , Π , then the 4th proportional Arc of right Ascension subtracts from 180° , for the right Ascension from *Aries*. When in the 3d Quadrant, δ , δ , Π , the 4th proportional Arc is the right Ascension from Δ , and adds to 180° for right Ascension from *Aries*. And when the Sun is in the 4th or last Quadrant, δ , δ , Π , the 4th proportional Arc, subtracts from 360° , for the right Ascension from *Aries*.

IV. QUESTION. To find the Sun's Place in the Ecliptic from the Sun's right Ascension $79^{\circ} 28' 25''$ from *Aries*, given?

Converse of the former PROPORTION.

As Cotine gr. Declination . . . $23^{\circ} 28' 30''$ co.	0.0375199
To Tan. Sun's R. A. from <i>Aries</i> . . . $79^{\circ} 28' 25''$	10.7309234
So Radius . . . $90^{\circ} 0' 0''$	10.0000000

To Tang. Sun's Long. fr. Υ , reqd. $80^{\circ} 19' 40''$ 10.7684433

N. B. In the above Proportion, the Tangent of right Ascension must be taken for the Degrees of right Ascension from *Aries*, or *Libra*, or nearest equinoctial Point, by which the correspondent Longitude from either of those Points may be easily determined, as above.

V. QUESTION. To find the Sun's true Amplitude of Rising or Setting, the Latitude of the Place, Greenwich, $51^{\circ} 28' 30''$, N. and the Sun's Declination, at Rising or Setting, $23^{\circ} 5' 18''$, N. being given?

PROPORTION.		Logarithms.
As Cotine Lat. $51^{\circ} 28' 30''$		9.7943877
To Radius $90^{\circ} 0' 0''$		10.0000000
So Sine Sun's Dec. at Rising or Setting, N. $23^{\circ} 5' 18''$		9.5940458

To Sine Sun's Amplitude N. required . . . $39^{\circ} 5' 3''$ 9.7996581

N. B. The Amplitude is always of the same Name with the Sun's Declination in any Latitude N. or S.

VI. QUESTION. To find the Time of true Sun Rising and Setting before and after Six, by the ascensional Difference, with the Length of Day and Night, the Latitude of the Place $51^{\circ} 28' 30''$ N. and Sun's Declination, (at Rising or Setting) $23^{\circ} 7' 18''$ N. being given?

PROPORTION.		Logarithms.
As Radius $90^{\circ} 0' 0''$		10.0000000
To Tan. Latitude $51^{\circ} 28' 30''$		10.0990059
So Tan. Declination N. $23^{\circ} 7' 18''$		9.6304107

To Sine Ascensional Dif. required . . . $32^{\circ} 25' 58''$ 9.7294166

N. B. The Sun rises before Six, when the Declination is of the same Name with the elevated Pole; but after Six, when of the contrary Name, by this Ascensional Difference turned into Time; allowing 15° in an Hour.

$$\begin{array}{r} 0^{\circ} \quad ' \quad '' \quad h \quad m \quad s \\ 32 \quad 25 \quad 58 = 2 \quad 9 \quad 44 \text{ by Rule, p. 167.} \\ + 6 \end{array}$$

Semi-diurnal Arc $8 \quad 9 \quad 44$ Sun sets.

Comp. to 12^h . . . $3 \quad 50 \quad 16$ Sun rises.

Doub. T. Setting . . . $16 \quad 19 \quad 28$ Length of Day;

Comp. to 24, Doub. Rising . . . $7 \quad 40 \quad 32$ Length of Night.

As double the Time of Sun Setting gives the Length of the Day, so double the Time of Sun Rising gives the Length of the Night.

VII. QUESTION. To find the oblique Ascension or Descension of the Sun, his Declination (at Rising or Setting) $23^{\circ} 7' 18''$ N. and Latitude of the Place, $51^{\circ} 28' 30''$ N. being given?

Find the R. A. Sun from Υ , by III. Question . . . $80^{\circ} 19' 40''$

The Ascensional Dif. by VI. Question $32 \quad 25 \quad 58''$

Oblique Ascension . . . $47 \quad 53 \quad 42''$ reqd.

Oblique Descension . . . $112 \quad 45 \quad 38''$

N. B. When the Latitude and Sun's Declination are both of the same Name, N. or S. the Ascensional Difference subtracts from the R. A. for the oblique Ascension, and adds thereto for the oblique Descension. But,

When the Latitude and Sun's Declination are of a different Name, (N. and S.) the Ascensional Difference adds to the right Ascension for the oblique Ascension; and subtracts therefrom for the oblique Descension.

If the Ascensional Difference exceeds the right Ascension, you must add 360° to the right Ascension, and then subtract the Ascensional Difference therefrom.

Or, if right Ascension, and Ascensional Difference added, exceed 360° , the Excess will be the oblique Ascension, or Descension, according to the above Rules, or the following Scheme.

Astronomical QUESTIONS (applied to NAVIGATION) answered by Spherical PROPORTION and LOGARITHMS.

Decl. and Lat. } R. A. } — } Ascens. Dif. } oblique Ascension.
 Same Name } + } for the } oblique Descension.
 Decl. and Lat. } R. A. } + } Ascens. Dif. } oblique Ascension.
 diff. Name } — } for the } oblique Descension.

VIII. QUESTION. To find the Semi-diurnal or Semi-nocturnal Arc of the Sun, or Star, the Latitude of the Place $51^{\circ} 31' 30''$ N. and Declination $23^{\circ} 7' 18''$ N. being given?

PROPORTION.

As Radius : : : : : 10.0000000
 To Tang. Lat. $51^{\circ} 31' 30''$: : : : : 10.0990059
 So Tang. Decl. same Name with Lat. $23^{\circ} 7' 18''$: : : : : 9.6304197
 To Cof. Semi-nocturnal Arc required $57^{\circ} 34' 2''$: : : : : 9.7294166
 Turned into Time = $3^h 50^m 16^s$ as before;
 being the Time of Sun-Setting; whose Complement to 12^h , or $8^h 9^m 44^s$ is the Time of true Sun Rising.

Hence the ascensional Difference and the Semi-nocturnal Arc are always the Complement of one another to 90° , when the Latitude and Declination are of the same Name. When they are of a different Name, the foregoing Proportion finds the Semi-diurnal Arc of Sun or Star, which Arc, and the ascensional Difference are then Complements of one another to 90° .

IX. QUESTION. To find the Time of apparent Rising and Setting of the Sun, or of the Moon, a Star, or Planet, from the Meridian, the Declination of either, at that Time, and Height of the Pole, being given?

Example. Suppose the Latitude London, $51^{\circ} 32' N$.
 Sun's Declination at Rising 17 $32' N$.

RULE. Add $90^{\circ} 33' 0''$, Sun or *'s Zenith Dist. at Rif. or Setting, Compl. Pole's Height,

And Dist. of Sun or Star from the Pole.

(which last Distance is equal 90° . . . more, or less, the Declination, according as the Pole is of . . . dif. or same . . . Name with the Declination N. and S.)

Take $\frac{1}{2}$ Sum of these 3 Sides of a spherical Triangle. From which half take the Comp. Pole's Height; calling what's left the 1st Remainder.

From the same half take the Sun or Star's Distance from the Pole; calling what's left the 2d Remainder.

Add the Log. Sines of these 1st and 2d Remainders, and double the Log. of Radius.

From which Sum, subtract the Sum of the Log. Sine of the Comp. of Pole's Height, and Log. Sine of the Distance of the Sun or Star from the Pole: — Half of the last Remainder will be the Log. Sine of half the Sun or Star's angular Distance from Noon or the Meridian, at Rising or Setting; which doubled, and reduced into Time, will therefore be the true Interval between the apparent Rising and Setting of the Sun, or Star, and the Time of their passing the Meridian, when the Declination and Latitude of the Place are of the same Denomination. But, if they are of different Denomination, the Difference between the said doubled Arc, reduced into Time, and 12 Hours, will be the Interval required.

Sun's Zenith Distance . . . $90^{\circ} 33'$ Side op. $\angle$ req.

Comp. Pole's Height $38^{\circ} 29'$ Sides including $\angle$ required.

Dist. Sun from Pole $72^{\circ} 28'$
 Sum 201 30
 From Half 100 45

Take Comp. Pole's Height . . 1. Rem. 62 16 Log. S. | 9.9470036
 Take Dist. Sun fr. Pole . . 2. Rem. 28 17 Log. S. | 9.6756245
 Doub. Log. Rad. 20.0000000

From . . Sum 39.6226281

N. B. If a Star, or Planet, had the given Declination, at Rising, then the Semi-diurnal Arc, $114^{\circ} 25' 54''$ is turned into Time, or Semi-Duration Arc, thus.

As 360° is to the Interval of Time, betwixt two next Passages, thro' the same Meridian, so is $114^{\circ} 25' 54''$ to the Semi-Duration Arc, required. Which subtr. from, and added to, the Time of the Star or Planet's Southing, that Day, will give the Time of Rising and Setting, required.

X. QUESTION. To find the Time of the Day, from the Sun's Declination, his true Altitude, and Latitude of the Place given?

RULE. Is exactly the same as for answering the preceding Question, in finding the Time of apparent Sun-Rise, except, that instead of $90^{\circ} 33'$, (the Sun's Zenith Distance apparent) you must use the Compl. of the Sun's Altitude observed, after Refraction is allowed for by Tab. p. 21.

Example. The Sun's Altitude was observed, at Sea, on July 18, 1757, just $146^{\circ} 9'$ (or $9^h 44^m$ Diff. of Time) eastward of Greenwich Observatory, at about 9 in the Forenoon, to be $47^{\circ} 48' 47''$, the Refraction then being $47''$, reduces it to $47^{\circ} 48'$ true Altitude. The Sun's Declination, about that Time, was $21^{\circ} 6' N$. by an Ephemerides (it being then about July 17, $11^h 16^m$ Night, at Greenwich) and the Latitude of the Place of Observation $40^{\circ} 18' N$. required the exact Time, at the Place of Observation, and likewise at Greenwich?

Comp. Sun's Alt. $42^{\circ} 12'$ Side op. $\angle$ req.
 Comp. Lat. . . . $49^{\circ} 42'$ Sides including the
 Sun from the Pole . . . $68^{\circ} 54'$ $\angle$ req.

Sum 160 48
 From . . . Half 80 24

Take Com. Lat. 1. Rem. 30 42 Log. S. 9.7080323
 Take Sun's Dist. fr. Pole. 2. Rem. 11 30 Log. S. 9.2996553
 Doub. Log. Rad. 20.0000000

Otherwise.

Logarithms.
 1. Rem. L. S. 9.7080323
 2. Rem. L. S. 9.2996553
 L. S. $49^{\circ} 42'$ co. 0.1176643
 L. S. $68^{\circ} 54'$ co. 0.0301400

Sum . . 19.1554919
 $\frac{1}{2}$. . 9.5777459
 Double $44^{\circ} 26' 50'' = 2^h 57^m 47^s$ T. fr. Noon.

At Place of Observation, July 18^d 9 2 13 Forenoon.
 Dif. Meridians — 9 44 0 sooner at Greenwich.

At Greenwich, July 17^d 11 18 13, Night
 Times required.

N. B. The Time at Greenwich Observatory, or any Place for which Meridian an Ephemerides is made, is sooner than the Time of all Places to the Eastward, and later than the Time of all Places to the Westward of that first Meridian, by the Difference of these Meridians in Time, and the contrary.

Astronomical QUESTIONS (applied to NAVIGATION) answered by Spherical PROPORTION and LOGARITHMS.

And therefore the Declination of the Sun, Moon, or a Planet, at the Time and Place of Observation, E or W, of the first Meridian, must be found to the correspondent Time, at the first Meridian by an Ephemerides, proportioned to that Interval of Time, betwixt the two next Noons, for which those Declinations, are in the said Ephemerides respectively given.

XI. QUESTION. To find the Sun or Star's true Altitude at a given Time of the Day, 2^h 57' 47" from Noon, or the Meridian, an Arc of 44° 26' 50"; the Sun's Declination at that Time being 21° 6' N. and the Latitude of the Place 40° 18' N.

PROPORTION.			Logarithms.
As Radius			10.0000000
To Sine Complement Arc from Noon	44° 26' 50"		9.8536348
So Tangent Complement Latitude	40 18 0		10.0715726
			9.9252074
To Tangent of an Arc X	40 5 26		
○ from the Pole	68 54 0		
			9.9426168
To Sine Complement Arc Y	Dif. 28 48 34		
As Sine Complement Arc X	40 5 26 co.		0.1163229
So Sine Pole's Height	40 18 0		9.8107631

To Sine Sun's Altitude required 47 48 0 9.8697028
 N. B. Arc X subtracts from the Sun's Distance from the Pole for the Arc Y, when the Time from Noon, or the Meridian, is less than 6^h or 90°, but adds to the Sun's Distance from the Pole, for the Arc Y, when the Time from the Meridian is greater than 6^h or 90°.

The above Proportions find the Side of any Spherical Triangle opposite to a given Angle, included between two given Sides.

Thus, As Radius to Sine Complement $\angle$ included, So Tangent left including Side to Tangent Arc X.

Subtract this Arc X from the other including Side, when the included $\angle$ is less than 90°, but add it thereto, when the included $\angle$ is greater than 90°, calling the Arc of Difference or of the Sum, Y.

Then say, As Sine Complement X is to Sine Complement Y, So is Sine Complement left including Side to the Sine Complement of the Side opposite the included Angle.

XII. QUESTION. To find the apparent Amplitude of Rising or Setting of the Sun, Moon, a Star, or Planet, the Declination of either at that Time, and the Latitude of the Place, being given?

Example. Sun's Declination at that Time is 23° 7' 18" N. and Latitude 51° 28' 30" at Greenwich.

Here are given ○ fr. Pole 66° 52' 42" Side op. $\angle$ required.
 ○ Zenith Distance . . . 90 33 0 } Sides including
 Complement Latitude . . . 38 31 30 } $\angle$ reqd. viz. Azim. first.

	Sum	195 57 12	
From . . .	Half	97 58 36	
Take ○ Zenith D. 1 rem.	7 25 36	Log. S.	9.1114545
Take Comp. Lat. 2 rem.	59 27 6	Log. S.	9.9351046
		Log. S. 90° 33' 0 co.	0.0000200
		Log. S. 38 31 30 co.	0.2056123

Sum 19.2521914
 Log. S. 25° 0' 33" Half 9.6260957
 Doubled 50° 1' 6" $\angle$ of Azimuth fr. N.
 90 0 0

Apparent Amplitude 39 59 44 fr. the East Northerly.
 True Ampl. by V. Qu. 39 5 3 Northerly.
 Dif. bet. true and app. 54 41 in this Case.
 The apparent Amplitude being so much the more Northerly.

N. B. When the Latitude and Declination are of the same Name, the foregoing Proportion finds the Azimuth from the North Point of the Horizon; but if the Latitude and Declination are of contrary Names, the foregoing Proportion finds the Azimuth from the South Point of the Horizon, which Azimuth-Angle being taken from 90°, will leave the Amplitude to the Northward or Southward of the East, or West Point of the Horizon accordingly, required.

XIII. QUESTION. To find the Azimuth of the Sun, Moon, or a Planet, from the Declination 20° 30' N. Altitude 47° 30', and Latitude 51° 32' given; the Declination being for the Time of the Altitude?

Here are given Sun	69° 30'	Side op. $\angle$ required.	Logarithms.
fr. Pole			
Complement Lat.	38 28	Sides includg. } L.S. co.	0.2061683
Complement Alt.	42 30	$\angle$ required. } L.S. co.	0.1703167
			Sum 150 28
From . . .	Half	75 14	
Take Co. Lat. 1 rem.	36 46	Log. Sine	9.7771060
Take Co. Alt. 2 rem.	32 44	Log. Sine	9.7329303

Sum 19.8865713
 Log. S. 61° 21' 6" Half 9.9432856
 Doubled 122 42 12 } Sun's Azimuth fr.
 North required.

N. B. When the Declination and Latitude are of different Names this Rule finds the Azimuth from the South, required.

XIV. QUESTION. To find the Latitude from the Declination 20° 30' N. Altitude 47° 30', and Azimuth of the Sun or any Star 112° 42' 12" from the N. given?

PROPORTION.			Logarithms.
As Radius, or Sine	90° 0' 0"		10.0000000
To Tan. Zenith Distance	42 30 0		9.9620525
So Cos. Azimuth from N.	112 42 12		9.7326264
			9.6946789
To Tan. Arc X	26 20 21		
Now, As Sine Altitude	47 30 10 co.		0.1323691
To Cos. Arc X	26 20 21		9.9523969
So Sine Declination	20 30 0		9.5443253
			9.6290913
To Cos. Arc Y	64 48 21		
Arc X	26 20 21		

Complement Latitude . . . 38 28 0 = Y - X.
 Latitude required . . . 51 32 0 as before.

N. B. If the Observation be made between 6 in the Morning and 6 in the Afternoon, on the same Day, at the same Time the Azimuth is less than 90° (which generally happens when the Declination and Latitude are of different Name, and the Azimuth is taken from the South) then the Sum of the Arcs X and Y, will be the Complement of the Latitude, whose Complement will be the Latitude required. But if the Azimuth from North is less than 90°, and Declination and Latitude of the same Name, the Sum of X and Y will be the Complement of Latitude.

N. B. The foregoing Proportions, answering the XIV Question, find the third Side of any Spherical Triangle, when the other two Sides and an Angle opposite to one of those Sides are given.

Thus, As Radius is to Tangent of the less Side, so is Cosine of the given Angle, (opposite to the greater Side) to Tangent of an Arc X, Part of the Base without or within the Triangle, where a Perpendicular falls, from the End of the longer Side upon the required Side continued, or upon it.

Astronomical QUESTIONS (applied to NAVIGATION) answered by Spherical PROPORTION and LOGARITHMS.

Again, As Cosine less Side is to the Cosine of the Arc X, (without or within the Triangle, where a Perpendicular falls, according as the given Angle opposite to the greater Side, is more or less than 90°) So is the Cosine of the longer Side, To the Cosine of the whole Base, or greater Segment.

Where it must be noted that Y is the whole Base found, when the given $\angle$ exceeds 90° , and then X, the external Segment thereof, subtracts from it, for the Length of the required Side, or Segment. But if the Angle given is less than 90° , then Y is the greater and X the less Segment of the required Base, whose Sum is therefore the whole Base or Side required.

These Things are demonstrable from Spherical Trigonometry, to which we refer the Reader if he requires further Satisfaction, the Truth of the Rules, as here applied to Practice, being sufficient for our Purpose.

XV. QUESTION. To find the Distance, in Time, of the Sun, or a Star, from the Meridian, the Azimuth from the North $122^\circ 42' 12''$, Declination $20^\circ 30' N.$ and Altitude $47^\circ 30'$ at that Time, being given?

PROPORTION.

As Cosine Declination	$20^\circ 30' 0''$	co.	Logarithms.
To Sine Azimuth from North	$122^\circ 42' 12''$		9.9250435
So the Cosine Altitude	$47^\circ 30' 0''$		9.8296833
To Sine angular Dist. from the Merid.	$37^\circ 22' 5''$		9.7831392

XVI. QUESTION. To find the Azimuth of the Sun or a Star, from the Latitude $51^\circ 32' N.$ Declination $20^\circ 30' N.$ and Distance at the same Time of the Sun or Star from the Meridian, $37^\circ 22' 5''$, being given?

RULE. Take the Sum and Difference of the Complement of the Latitude and Distance of the Sun, or Star, from the Pole.

$28^\circ 28'$ Complement Latitude

$69^\circ 30'$ Sun or Star's Distance from the Pole.

Sum $107^\circ 58'$

$\frac{1}{2}$ $53^\circ 59'$

Dif. $31^\circ 22'$

$\frac{1}{2}$ $15^\circ 41'$

Then say by PROPORTION.

As Cosine $\frac{1}{2}$ Sum	$53^\circ 59' 0''$	co.	0.2306075
To Cosine $\frac{1}{2}$ Diff.	$15^\circ 41' 0''$		9.9838755
So Cotang. $\frac{1}{2}$ Angular Dist. from Merid.	$18^\circ 41' 2''$		10.4708675
To Tang. Arc. X	$78^\circ 20' 22''$		10.6853505

Again, As Sine $\frac{1}{2}$ Sum	$53^\circ 59' 0''$	co.	0.0921342
To Sine $\frac{1}{2}$ Diff.	$15^\circ 41' 0''$		9.4273541
So Cotang. $\frac{1}{2}$ $\angle$ Dist. fr. Mer.	$18^\circ 41' 2''$		10.4708675

To Tang. Arc Y.	$44^\circ 21' 50''$		9.9903558
X = $78^\circ 20' 22''$			

Y + X = $122^\circ 42' 12''$ Azim. fr. N. reqd.

X - Y = $33^\circ 58' 32''$ $\angle$ op. co. Lat.

N. B. In the above Proportions, when $\frac{1}{2}$ Sum of the Complement of Latitude and Sun or Star's Distance from the Pole exceed, or is less than 90° , then X is greater or less than 90° .

The Sum of the Arcs X and Y is the Angle of Azimuth from the North, and the Difference of X and Y is the other or less Angle, included by the Distance of the Sun or Star, from the Zenith and Distance of the same, from the Pole, when the Sun or Star's Zenith Distance is less than their Polar Distance, the other Side of the Spherical Triangle being the Complement of Latitude. When the Polar is less than the Zenith Distance from the Sun or Star, then the Difference of X and Y is the Azimuth, and the Sum of X and Y the other alternate $\angle$, opposite the Pole's Distance from the Zenith.

The foregoing Proportions find the $\angle$ s of any Spherical Triangle opposite to two given Sides, including a given Angle, and therefore find the Angle of Position in Sailing by the Arch of a great Circle, continually varying that Course, cross the Meridians, betwixt two Places of given Latitudes, and given Difference of Longitude: The Polar Distances of those Places, and Angle betwixt them, being also given. See Mr. EMERSON'S Navigation, p. 28.

XVII. QUESTION. To find the Latitude of the Place from the Sun's Declination $23^\circ 7' 18'' N.$ and the Time of the Sun's Rising then $3^h 50^m 16^s$, being given.

RULE. Turn the Time of Sun-Rising before or after Six into Degrees (by p. 28) which is called the Ascensional Difference.

Thus, $6^h 00^m 0^s$

$3^\circ 50' 16''$

$2^\circ 9' 44'' = 32^\circ 25' 58''$

Then say, by this PROPORTION.

As Radius		Logarithms.	10.0000000
To Cotang. Declination N.	$23^\circ 7' 18''$		10.3695893
So Sine Ascensional Difference	$32^\circ 25' 58''$		9.7294166
To Tangent of Latitude required	$51^\circ 28' 30''$		10.0990059

XVIII. QUESTION. To find the Latitude of the Place from the Sun's Declination, $23^\circ 7' 18'' N.$ and Amplitude $39^\circ 5' 3''$, being given?

PROPORTION.

As Sine Amplitude from E. or W.	$39^\circ 5' 3''$		9.7996581
To Radius			10.0000000
So Sine Declination N.	$23^\circ 7' 18''$		9.5940458
To Cosine Latitude required	$51^\circ 28' 30''$		9.7943877

N. B. These Questions of Data at the Horizon, are easiest for finding the Latitude of Places; because they come under the Form of a Right angled Spherical Triangle: whereas Questions concerning Altitude of the Sun or Star, betwixt Rising or Setting, and the Meridian, can't be answered without resolving an Oblique angled Spherical Triangle, by two or more Proportions; except where there is no Declination, or the Sun or Star be due East or West.

XIX. QUESTION. To find the Latitude of the Place N. from the Declination of the Sun $20^\circ 30' N.$ Altitude of the Sun $47^\circ 30'$, and Time of the Day $9^h 30^m 32^s$ in the Morning being given?

$12^h 00^m 0^s$

$9^h 30^m 32^s$

Time fr. Noon $2^\circ 29' 28'' = 37^\circ 22' 5''$ Angular Dist. fr. Merid.

First to find the Azimuth.

Say, As Sine Zen. Dist. $\odot$	$42^\circ 30' 0''$	co.	0.1703167
To Sun's Dist. fr. Merid.	$37^\circ 22' 5''$		9.7831396
So Sine $\odot$ Distance from Pole	$69^\circ 30' 0''$		9.9715876

To Sine Azimuth from South	$57^\circ 17' 48''$		9.9250439
	$180^\circ 0' 0''$		

Azimuth from N. required $122^\circ 42' 12''$

Proceed now as in the XIVth Question to which this XIXth Question is reduced.

Astronomical QUESTIONS (applied to NAVIGATION) answered by Spherical PROPORTION and LOGARITHMS.

But, a small Error in the Time of Observation will make a large one in the Azimuth, and also in the Result of the Computation, or Latitude.

And since an *Azimuth* by Observation can scarcely ever be trusted to, for finding the Latitude, except it is derived from the observed Altitude of the Sun or Star, the Method of determining the Latitude from taking an Altitude, or repeated Altitudes, is always the best; as being the most accurate. And the Method by Meridian Altitudes of the Sun or a Star is not only the most easy for determining the Latitudes, but always the most certain, and expeditious.

XX. QUESTION. To find the Time when the Sun will be due East or West, from the Sun's Declination $23^{\circ} 28' 30''$ N. and Latitude $51^{\circ} 28' 30''$, being given?

PROPORTION.		Logarithms.
As Radius	$90^{\circ} 0'$	10.0000000
To Cotan. Lat.	$51^{\circ} 28' 30''$	9.9009941
So Tang. Sun's Declination	$23^{\circ} 28' 30''$	9.6377835
To Sine Sun's Dist. from 6h	$20^{\circ} 13' 42''$	9.5387776
In Time	$\pm 1^h 20^m 55^s$	

Sun due East . . . 7 20 55 Morn. ?

Sun due West . . . 4 39 5 After. } reqd.

N. B. In the Summer the Sun is over the East and West Points at the above Times, and Declination North. But under them in the Winter at the same Declination South.

XXI. QUESTION. To find the Altitude of the Sun, when due East and West, the Sun's Declination, $23^{\circ} 28' 30''$, and Latitude $51^{\circ} 28' 30''$ being given?

PROPORTION.		
As Sine Latitude	$51^{\circ} 28' 30''$ co.	0.1066064
To Radius	$90^{\circ} 0' 0''$	10.0000000
So Sine Declinat. N.	$23^{\circ} 28' 30''$	9.6002636
To Sine Sun's Altitude	$30^{\circ} 36' 33''$	9.7068700

in Summer or N. Latitude; but Depression in Winter or S. Latitude.

XXII. QUESTION. To find the Sun's Altitude at Six, from the Declination $23^{\circ} 28' 30''$ N. and Latitude $51^{\circ} 28' 30''$ N. being given?

PROPORTION.		
As Radius	$90^{\circ} 0' 0''$	10.0000000
To Sine Sun's Declination	$23^{\circ} 28' 30''$	9.6002636
So Sine Latitude	$51^{\circ} 28' 30''$	9.8933936
To Sine Altitude at 6, required	$18^{\circ} 9' 30''$	9.4936572

XXIII. QUESTION. To find the Sun's Azimuth at 6, from the Sun's Declination $23^{\circ} 28' 30''$ N. and Latitude $51^{\circ} 28' 30''$ N. being given?

PROPORTION.		
As Radius	90°	10.0000000
To Tan. Sun's Decl. N.	$23^{\circ} 28' 30''$	9.6377835
So Cot. of Latitude	$51^{\circ} 28' 30''$	9.7943877
To Cot. Azim. } fr. the N. or S. }	$74^{\circ} 51' 49''$	9.4321712 or Tang. fr. E. or West.

N. B. When the Sun's Declination and Latitude are both North, the Azimuth found by the foregoing Proportions, is from the North; when the Declination is South, and Latitude North, the Azimuth is from the South. — The contrary for the Southern Latitude.

There are many more Questions concerning the Altitude, Latitude, Declination, Azimuth, Amplitude, &c. of the Sun when in the Equinoctial, which are solved by one Proportion of Spherical Trigonometry; which, as they are of a singular Nature, and less useful in Practice than those we have inserted, we have omitted, to furnish Amusement, or Exercise, for the young Astronomer's Invention. Who being supposed already acquainted with the Rules and Demonstration of Spherical Proportion, and with the Idea of Spherical Lines and Angles, as applied to Astronomy and Navigation, we leave such Questions and their Answers to his Investigation: Having inserted the general Proportion of Spherical Lines and Angles, in a Table of Spherical Trigonometry (as is customary in most Books of the Subject here treated on) for his ready Use and Reference to, further on. As likewise the most noted or famous Theorems, Rules, and Proportions in Practice, added, for solving all the Cases in Spherical and plain Trigonometry; without which, not only Astronomy, but Navigation, and all the Arts depending upon these Rules, would be imperfect.

For Demonstration of which infallible Rules, we refer the Reader to Mr. Emerson's most excellent Treatise of Trigonometry, (as clear and demonstrative as the Elements of Euclid) for ample Satisfaction. Sold by Mr. Richardson, in Pater-Noster-Row.

Of the NATURE and CAUSE of TWILIGHT.

WHEN Twilight begins and ends, the Sun is always reckoned about 18° below the Horizon: but exactly begins and ends when the Sunbeams passing (as a Tangent, allowing for Refraction) by the Earth's Surface, through the surrounding Body of the Air, or Atmosphere, to the outer or extreme Parts thereof, from whence (at the Middle of the extreme Atmosphere, or Meeting of the two Tangents of Incidence and Reflection) the Illumination or Illustration there, of the Sun's first or last Rays are reflected, horizontally, to the Eye of an Observer.

As soon as which incident Rays fall short of meeting the horizontal Tangent, at the extreme Parts of the surrounding Atmosphere, in the Evening Twilight ceases, and total Night or Obscurity ensues. And, in the Morning, as soon as the Sun's incident Rays meet the horizontal Tangent at the outer Parts of the Earth's Atmosphere, then Twilight begins, and encreases its Light till open Day appears by the general Illumination of the whole Atmosphere, every Way reflecting Light; besides what Light is reflected by the Earth's Surface, and all terrestrial Objects, which the Sun's Light directly falls upon.

The Atmosphere, varying in its Height and Magnitude, (in the Summer expanding by Heat, bigger, and condensing in Winter by Cold, lower) is one Cause of the Variation of the Beginning and End of Twilights, at the same certain Times of the Year. Another Cause is the variable Quantity of Matter in the Atmosphere fit for reflecting Light, not too rare, or too dense, in this transparent Body. Add to which a third Cause of the Variation of Length, as well as of the Clearness of Twilight, which is the Variation of the Sun's ethereal Atmosphere surrounding his Body, shining before the Sun himself rises, and after he sets, with variable Lustre, augmenting the Clearness of Twilight, more or less, and for a longer or shorter Time. The Duration of Twilight is likewise reckoned somewhat shorter in Mornings than in the Evenings of the same Days, especially in Increase of Twilight: the Morning Twilight is reckoned to begin when Stars of the Sixth Magnitude disappear by its superior Lustre; and the Evening Twilight ends when the same Stars begin to appear, which the Light of the Air before rendered invisible: As the brightest Stars in the whole Heavens vanish out of Sight in the Presence of the superior Lustre of the Sun.

From these Considerations, it will be easy to determine the Beginning and End of Twilight, in all Places, and at all Times of the Year; and also the Day of the shortest Twilight. The Height of the Atmosphere may be measured from the known Depression of the Sun, when Twilight begins and ends. All which are shewn hereafter.

XXIV. QUESTION. To find the Time of Day-Break, in the Morning, and End of Twilight in the Evening, from the Sun's Declination 10° N. and Latitude of the Place $51^{\circ} 32'$ N. being given?

RULE.

Sun's Zen. Dist. at Beg.	0				
and End of Twil.	108	0	Side opp.	∠ reqd.	Logarithms.
Comp. Lat.	38	28	Sides inclg	Lo.S.co.	0.2061683
Sun fr. Pole	80	0	∠ reqd.	Lo.S.co.	0.0066485

	Sum	226	28
From . . .	Half	113	14

Take Com. Lat. δ 1 rem. 74 46	Log. Sine	9.9844660
Take Sum fr. Pole. 2 rem. 33 14	Log. Sine	9.7388201

			Sum	19.9361029
	68° 17' 29"	L. Sine . . Half		9.9680514
Double 136	34 58	∠ from Noon.		
In Time 9 ^h	6 ^m 20 ^s			
12				

Day-Break, 2^h 53^m 40^s Morning, *required.*

Add the Increase of Sun's Declination for double the $\angle$ of Day-break from Noon, viz. for $18^h 12^m 40^s$, to the former Declination 10° N. and repeat the Operation, with the same Zenith Distance and Complement of Latitude, and the Result will give the exact Time of Twilight ending in the Evening; which however may be taken $9^h 6^m 20^s$ that Night, Dist. of Morn. Twilight, from Noon, as nearly.

OPERATION repeated.

Increase of the Sun's Decl. in 18^h 12^m . . . 16' 6"
 (See p. 194. April) Incr. in . . . 24 . . . being 21 14
 Whence, Sun's Decl. at ending of Twil. . . 10° 16' 6" N.
 [for that Day.

Sun's Zen. Dist. at Beg.	0	0	0	Side opp.	∠ reqd.	Logarithms.
and End of Twil.	108	0	0			
Comp. Lat.	38	28	0	Sides incl.	L.S.co.	0.2061683
Sun fr. the Pole	79	43	54	∠ reqd.	L.S.co.	0.0070121

Sum	226	11	54
From . . Half	113	5	57

Take Comp. Lat. 1. rem.	74	37	57	Log. Sine.	9.9841878
Take Sun fr. Pole. 2. rem.	33	22	3	Log. Sine.	9.7403683

68° 33' 50" Log. S. . . Half Sum | 19.9377365
 Double 137 7 40 $\angle$ from Noon. | 9.9688682
 In Time fr. Noon 9^h 8^m 31^s at Night, Twil. ends, req^d.
 In T. fr. Noon, 0 6 20 Twil. begun.

Dif. 2 11 Twilight longer in the Evening
[than in the Morning.

The same Method finds the Beginning and Ending of Twilight for South Declination of the Sun, in any Latitude, the Sun's Zenith Distance, at that Time, being always 1080.

N. B. The Twilight increases from March 2, to its greatest; and then decreases to October 11, at London, again at its shortest. Then it increases to Sun in Vf , then decreases to March 2, then increases as before, &c.

When the Sun descends just 18° below the Horizon, at Midnight, then there is no Darknefs or Night, but all Twilight begins to take Place. This happens when the Sun's Declination is equal to the Difference between the Complement of Latitude of the Place of the same Name with the Sun's Declination, and 18° the Depreffion at Midnight. In the Latitude of $51^{\circ} 32'$ N. the Complement of Latitude is $38^{\circ} 28'$, from which deducting 18° , the Depreffion, there remains $20^{\circ} 28'$ N. Declination of the Sun, when Darknefs difappears, in that Latitude, which is from the 22d of *May* to the 21st of *July* (see P. 195.) N. S. when Night again returns.


But from the Sun's Excursion from, to his Return to, this *Limit*, in the Latitude, *London*, it is all Twilight, from Sun-Setting to Sun-Rising; being the *longest* Duration of Twilight that can happen there.

As this *longest* Twilight happens when the Sun's Point or Place of Declination (*of the same Name with the Latitude*) descends the most obliquely or most parallel with the Horizon, to 18° below it, and consequently descends most slowly; so the *shortest* Twilight always happens, when the Point of the Sun's Declination (*of contrary Name to the Latitude*) descends in the most perpendicular Direction to the Horizon; or descends fastest to 18° Depression in any Latitude: Supposing the Sun's Declination to be invariable, or nearly so, for the Time betwixt Rising and Setting, and 18° Depression, below the Horizon.

Hence, when the Sun has the same Azimuth at 18° Depression below the Horizon, that he had at Setting, his Descent will be perpendicular to the Horizon, and therefore on that Day, and Declination (of contrary Name to that of the elevated Pole) the Twilight will be the *shortest possible*, in any given Latitude. This Consideration furnishes a *New Method* of finding the Time of *shortest Twilight*, as follows.

XXV. QUESTION. To find the Day of the shortest Twilight, from the Latitude $51^{\circ} 32'$ being given?

☞ The Time of the shortest Twilight, is when the Sun's Azimuth, during that Time, increases or decreases the least or slowest, viz. not at all.



In the *Figure*,

Put m and n = Sine and Cof. of $ZS = 108^\circ$,
 p and q = Sine and Cof. ZP = Comp. Lat.
 x = Rad. = $\odot Z$,
 And y = Cofine $\odot P = SP$, Dist. fr. Pole.

Astronomical QUESTIONS (in PRACTICE) answered by Spherical PROPORTION and LOGARITHMS.

Then (by a known Theorem*) $\frac{y}{p} = \text{Cof. } \angle \text{QZP}$. And $\frac{m+1}{mp} = \text{Cof. } \angle \text{SZP}$. Which Angles being equal, their Cofines are also equal, or $\frac{y}{p} = \frac{m+1}{mp}$, whence, $y = \frac{m+1}{m}$, a GENERAL THEOREM, for answering all Questions of this Nature.

Hence, $m+1 : n :: q : y$, the Sine of the Sun's Declination, of contrary Name to the Latitude, required.

Or, in Words at Length,
As the Sine of 72° and Radius added,
Is to the Sine of 18° ,
So is the Sine of the Latitude,
To the Sine of the Sun's Declination,
On the Day of the shortest Twilight.

Or, to the constant Logarithm . . .	9.1997126
Add the Log. Sine of the Latitude $51^\circ 32'$	9.8937452
Sine Sun's Declin. S. reqd. $7^\circ 7' 25''$. .	9.0934578
Answering to March 2d and Oct. 11.	

OTHERWISE, according to the common Method.


As Radius $90^\circ 0'$	10.0000000
To Tang. $\frac{1}{2}$ Depression $9^\circ 0'$	9.1997125
So Sine Latitude $51^\circ 32'$	9.8937452

To Sine Sun's Declin. reqd. . . $7^\circ 7' 25''$ S. 9.0934577
answering to March 2, and Oct. 11, as before; between which Times the Twilights increase; and from the latter to the former of which Times, they decrease. The Days (different to the Increase and Decrease of Twilights) increase from the Sun in $\text{V}\beta$ to his Arrival in $\text{O}\alpha$, and decrease from $\text{O}\alpha$ to $\text{V}\beta$, in all Latitudes.—While the Days increase the Afternoon Half-Days are the longer; but while the Days decrease the Forenoon Half-Days are longer than those of the Afternoon. Which Consideration furnishes a Rule (by Proportion) for determining the Difference of Time, between Forenoon and Afternoon Observations, from Noon, in the Method of corresponding or same Altitudes of the Sun; by having the Distance, in Time, of the two Observations of the same Altitudes of the Sun, (in the Forenoon and Afternoon) given.

XXVI. QUESTION. To find the Height of the Earth's Atmosphere, or of its surrounding Body of Air, from the Sun's Depression 18° , at the Beginning and End of Twilight, being given?

THE Time of the Beginning or End of Twilight being given, the Sun's Depression at that Time, is found nearly as above, viz. 18° , by Spherical Trigonometry.

At which Time, the Sun's last Ray SR, touches the Earth's Surface at E, and is reflected by a Particle of Air from the Top of the Atmosphere, at R, to the Observer's Eye at H, in a horizontal Direction RH. And since HH represents the Horizon, the $\angle \text{HRS}$, = the Sun's Depression, = 18° .



Now, because SR and HR, are both Tangents to the Earth's Surface, at H and E, meeting at the Top of the Atmosphere, R, the Angle HCE, at the Earth's Center, is equal to the Angle SRH = 18° ; whose Half, or the Angle HCR is equal to Half $\angle \text{SRH}$, = 9° ; which would be correctly so, if the Ray SR passed straight through the Atmosphere without Refraction. But because the Sun's Rays are refracted and bent towards CR, (all the Way of their Passage, by the Earth's Surface, from one Part of the external Atmosphere to their Arrival at the Top of the contrary Part thereof) therefore the Angle of Depression will be $\angle \text{RH} = 18^\circ$, and the $\angle \text{CRS}$ will be the Sun's horizontal Refraction = $30'$, at least; whence, the $\angle \text{SRH} = 17^\circ 30'$: the Sun's Rays passing directly in the straight Line $\text{O}\alpha$, till they meet the outside of the Atmosphere in r , and then pass thro' it in the curved Line reR to R, the opposite Part thereof, from whence they are horizontally reflected to the Observer at H.

NOW, $\frac{1}{2} \angle \text{SRH} = \angle \text{RHC} = \angle \text{RCE} = 8^\circ 45'$
The Comp. = $\angle \text{CRH} = 81^\circ 15'$


Say, As $\angle \text{CRH}$ $81^\circ 15'$. . co.	0.0050842
To CH, Earth's Rad. 4000 Miles	3.6020600
So Radius S. 90°	10.0000000

To CR = 4047 Miles.

— 4000 = CD.

Miles 47 = DR, the Height of the Atmosphere, required.

* The Rectangle of the Cofines of two Sides, including an Angle of any Spherical Triangle, plus or minus, the Product of the Sines of those including Sides, and the Cofine of the included Angle (plus when the included Angle is less, and minus when it is more, than a right Angle) is equal to the Cofine of the Side opposite to the included Angle.



Hence in the Spherical Triangle, ABC, when the inside Letters denote the Cofines, and outside Letters denote the Sines, of the given Sides and Angle, included.

Then $bd \pm cae = f$. Hence a (when $\angle A$ less than 90°) = $\frac{f-bd}{ce}$, or (when Angle A is greater than 90°)

$$b = \frac{f-cae}{d}, \text{ or } d = \frac{f-cae}{b} \text{ (when } \angle A \text{ less than } 90^\circ\text{.) But } b = \frac{f+cae}{d}, \text{ and } d = \frac{f+cae}{b}, \text{ when } \angle A \text{ is greater than } 90^\circ\text{.)}$$

Astronomical QUESTIONS (in PRACTICE) answered by Spherical PROPORTION and LOGARITHMS.

But supposing the Sun's Rays, in entering the Atmosphere, at 18° Depression, describe *SIR*, or nearly so.

Then the $\angle SRH = 18^\circ$, and $\angle RCH = 9^\circ$
[and $\angle CRH = 81^\circ$]

Wherefore,

Say, As $\angle CRH$ 81° co. | 0.0053801
To CH , Earth's Rad. 4000 Miles . . | 3.6020600
So Radius $S. 90^\circ$ | 10.0000000

To $CR = 4050$ Miles, ferè. | 3.6074401
— 4000 = CD .

Miles . . 50 = DR , the Height of the Atmosphere,
(at most) required.

XXVII. QUESTION. To find the Declination of the Moon, of a Planet, or Star, from the Longitude (suppose of the Moon) 1762, Paris, April 1, at Noon, $25^\circ 27'$, and Latitude $4^\circ 11' N.$ being given?

PROPORTION.

As Radius $S. 90^\circ$ | 10.0000000
To $S.$ Long. from the nearest equinoct. } $84^\circ 33'$ | 9.9980323
Point Ψ or Δ } $23^\circ 28' 30''$ | 9.6377835
So Tan. Ecl. Obliquity } $23^\circ 28' 30''$ | 9.6377835
To Tan. 4th Arc $23^\circ 22' 49''$ | 9.6358158
From $85^\circ 49'$ o Comp. Ψ Lat.

5th Arc . . Rem. $62^\circ 26' 11''$

If the given Latitude and Sign of the Longitude, be of the same Name, the 4th Arc subtracts from the Complement of the given Latitude for a 5th Arc; if the Latitude given and Sign of the Longitude be of a different Name, then the 4th Arc adds to the Complement of given Latitude for a 5th Arc.

NOW, As Cof. 4th Arc . . . $23^\circ 22' 49''$ co. | 0.0372088
To Cof. 5th Arc . . . $62^\circ 26' 11''$ | 9.6653306
So Cof. Ecl. Obliq. . . $23^\circ 28' 30''$ | 9.9624801
To $S.$ Decl. N. reqd. $27^\circ 32' 32''$ | 9.6650195

EXAMPLE II. 1760, Paris, August 5, Moon's Longitude at Noon, $8^\circ 29' 52''$, Latitude $S. 1^\circ 28'$, required the Declination N.

According to the ABOVE-RULE.

As Radius $S. 90^\circ$ | 10.0000000
To $S.$ Long. fr. next equin. Point Ψ $59^\circ 52'$ | 9.9369456
So Tan. Ecl. Obliquity . . . $23^\circ 28' 30''$ | 9.6377835
To Tan. 4th Arc $20^\circ 35' 11''$ | 9.5747291
Lat. and Decl. Dif. Name . . Add $88^\circ 32'$ o Comp. Ψ Lat.

5th Arc . . . Sum $109^\circ 7' 11''$
180

To Cof. 5th Arc $70^\circ 52' 49''$ | 9.5152685
As Cof. 4th Arc $20^\circ 35' 11''$ co. | 0.0286578
So Cof. Ecl. Obliquity . . . $23^\circ 28' 30''$ | 9.9624801
To $S.$ Declination N. required. . . $18^\circ 43' 8''$ | 9.5064064
Being a Proof of the Truth of M. Dela Caille's Ephemerides.

N.B. The two Examples above contain all the Cases that can happen, in finding the Declination of the Moon, a Planet, or the Stars, from their Longitudes and Latitudes, respectively, given.

XXVIII. QUESTION. To find the Right Ascension of the Moon, and of any Star, or Planet, from the Longitude, Latitude, and Declination thereof, being given?

Required the Right Ascension of the Moon, in the foregoing Instances, Longitude $25^\circ 27' 0''$
Latitude $4^\circ 11' 0'' N.$ } given.
Declination $27^\circ 32' 34'' N.$

PROPORTION.

As Cof. Declination $27^\circ 32' 34''$ co. | 0.0522378
To Cof. Long. fr. next eq. Point Δ $84^\circ 33' 0''$ | 8.9776188
So Cof. Latitude $4^\circ 11' 0''$ | 9.9988414
To Cof. R. Ascension, fr. Δ , reqd. $83^\circ 52' 2''$ | 9.0286980
180 0 0

Or R. A. fr. Ψ . . . $96^\circ 7' 58''$

EXAMPLE II. Longitude Ψ . . $8^\circ 29' 52'' 0''$
Latitude $1^\circ 28' 0'' S.$ } given.
Declination $18^\circ 43' 8'' N.$

As Cof. Declination $18^\circ 43' 8''$ co. | 0.0236021
To Cof. Long. fr. next eq. Point Ψ . . $59^\circ 52' 0''$ | 9.7007158
So Cof. Latitude $1^\circ 28' 0''$ | 9.9998577
To Cof. R. A. fr. next eq. Point } $58^\circ 0' 10''$ | 9.7241756
 Ψ , required }
So for the Refs.

The above Question may be otherwise answered by finding an Angle APE of a Spherical Triangle, whose Sides,

AP the Comp. Declination,
 AE the Comp. Latitude,
 PE constant Dist. of the two Poles of Ecliptic } are
and Equinoctial $23^\circ 28' 30''$ } given.

The Sides including the required $\angle$ being AP, PE .

XXIX. QUESTION. To find the Right Ascension of a Star, or Planet, from the Longitude and Latitude thereof being given?

Here, In $\triangle APE$, are given $\angle E$, Sides AE and PE , to find $\angle P$

Longitude of the Moon . . $25^\circ 27' 0''$ } given.
Latitude $4^\circ 11' 0'' N.$

To find the Right Ascension of the Moon in that Position.

PROPORTION.

As Radius $S. 90^\circ$ | 10.0000000
To $S.$ Long. fr. next eq. Point Δ . . $84^\circ 33' 0''$ | 9.9980302
So Cotan. Latitude N. $4^\circ 11' 0''$ | 11.1358275
To Tan. 4th Arc $85^\circ 47' 54''$ | 11.1339077
Ecliptic Obliquity — $23^\circ 28' 30''$ Sig. Long. and
[Lat. of like
The 5th Arc . . Dif. $62^\circ 19' 24''$ [Name.

When the given Latitude and Sign of the Longitude are of the same Name N. or S. you must subtract the Ecliptic Obliquity from the 4th for a 5th Arc. But if the Latitude and Sign of the Longitude are of a different Denomination, you must add the Ecliptic Obliquity, to the 4th for a 5th Arc.

$\Psi, \delta, \Pi, \varpi, \Omega, \Upsilon$, Northern Signs.
 $\Delta, \cap, \downarrow, \text{♄}, \text{♅}, \text{♆}$, Southern Signs.

Astronomical QUESTIONS (in PRACTICE) answered by Spherical PROPORTION and LOGARITHMS.

Now, As Sine 4th Arc	85° 47' 54" co.	Logarithms.	0.0011696
To Sine 5th Arc	62 19 24		9.9472292
So Tan. Lon. fr. next Eq. Point γ fr. $\triangle$	84 33 0		11.0204135
To Tan. R. A. fr. $\triangle$ required	83 52 2		10.9688125
	180 0 0		

Or Right Ascension from γ 96 7 58
 EXAMPLE II. Longitude γ 8 29° 52' } given.
 Latitude 1 28 S.

To find the Right Ascension γ in that Position?
 By the above Rule.

As Radius Sine	90° 0' 0"	Logarithms.	10.0000000
To Sine Long. from next Eq. Point γ	59 52 0		9.9369456
So Cotan. Latitude South	1 28 0		11.5916963
To Tan. 4th. Arc	88 18 15		11.5286419
Ecliptic Obliquity	+ 23 28 30	Sig. Long. and Lat.	[of dif. Name.]
The 5th Arc	Sum 111 46 45		
	180 0 0		

Now, To Sine 5th Arc	68 13 15	Logarithms.	9.9678384
As Sine 4th Arc	88 18 15 co.		0.0001903
So Tan. Longitude from γ	59 52 0		10.2362298

To Tan. R. A. from γ required 58 0 10
 Star or $\{ \text{☿, ♀, ☿, ♀, ♀, ♀, ♀, ♀} \}$ Lat. S. $\{ \text{R. A.} \}$ Lat. N. $\{ \text{R. A.} \}$
 Planet in $\{ \text{☿, ♀, ☿, ♀, ♀, ♀, ♀, ♀} \}$ Lat. N. $\{ \text{lefts.} \}$ Lat. S. $\{ \text{greater} \}$
 [than Sun R. A. in the Ecliptic, being in the same Signs.]

XXX. QUESTION. To find the Longitude and Latitude of a Star or Planet, from the Right Ascension and Declination thereof, being given?

γ 's Right Ascension from γ 99° 7' 58" } given.
 Declination North 27 32 32 }
 To find her Latitude North and Longitude from Aries.

PROPORTION.

As Radius Sine	90° 0' 0"	Logarithms.	10.0000000
To Tangent Ecliptic Obliquity	23 28 30		9.6377835
So Sine R. A. from next Eq. Point $\triangle$	83 53 2		9.9975074
To Tangent 4th Arc	23 21 18		9.6352909
	+ 62 27 28 comp. given Decl.		[N.]
Fifth Arc	Sum 85 48 46		
Now, As Cofine 4th Arc	23 21 18 co.		0.0371260
To Cofine 5th Arc	85 48 46		2.8634167
So Cofine Ecliptic Obliquity	23 28 30		9.9624801

To Sine Latitude N. required 4 11 N. 8.8630228

N. B. When the Right Ascension and Declination given are of the same Name, N. or S. the Complement of the Declination given adds to the 4th for the 5th Arc: But when the Right Ascension and Declination given are of different Name, the 4th Arc subtracts from the Complement of Declination for the 5th Arc.

Now, to find the Longitude from Libra.

PROPORTION.

As Cofine Latitude N.	4° 11' 0" co.	Logarithms.	0.0011586
To Cofine R. A. from next Eq. Point $\triangle$	83 52 2		9.0287050
So Cofine Declination	27 32 32		9.9477622
To Cofine Lon. from next Eq. Point $\triangle$	84 33 0		8.9776258
	180 0 0		

Longitude from γ 95 27 or $\triangle$ 5° 27' reqd.
 Being the Converse to the Proportions answering to the XXVIII. Question.

EXAMPLE II. γ 's R. Ascension from γ 58° 0' 10" } given.
 Declination 18 43 8 N. }
 To find her Latitude South and Longitude from γ .

PROPORTION as before.

As Radius Sine	90° 0' 0"	Logarithms.	10.0000000
To Tan. Ecliptic Obliquity	23 28 30		9.6377835
So Sine R. A. from γ	58 0 10		9.9284336
To Tangent 4th Arc	20 13 10		9.5662171
	+ 71 16 52 Comp. Decl. γ N.		
Fifth Arc	91 30 2		
As Cofine 4th Arc	20 13 10 co.		0.0276232
To Cofine 5th Arc	91 30 2		8.4180797
So Cofine Ecliptic Obliquity	23 28 30		9.9624801
To Sine Latitude S. required	1 28 0		8.4081830
NOW, To find the Longitude from Aries.			
As Cofine Latitude S.	1° 28' 0" co.		0.0001423
To Cof. R. A. fr. the next Eq. Point γ	58 0 10		9.7241760
So Cofine Declination N.	18 43 8		9.9763979
So Cof. Long. fr. same Eq. Point γ	59 52 0		9.7007162
	or, 8 29 52 0 required.		

In the Triangle APE of the foregoing Question, are given the constant Distance of the Poles of the Equator PE, and Ecliptic 23° 28' 30", the Complement of Declination (or 90° and contrary Declination) AP, with the included $\angle$ of Right Ascension from the Solstitial Colure of $\triangle$ or γ , P, to find the Side opposite to that $\angle$, viz. AE, whose Complement to 90° is the Latitude; or Excess above 90°, the contrary Latitude, required.

And also to find the $\angle$ E. at the Pole of the Ecliptic, included between the said opposite Side AE, last found, (on which the Latitude is measured) and the constant Distance of the Poles (afore said) PE; which $\angle$ E found, will be the Longitude required from the same Solstitial Colure of $\triangle$ or γ .

NOW, The Right Ascension at P, and Longitude at E, from $\triangle$ or γ being known, the Right Ascension and Longitude from γ and $\triangle$ will be also known; the latter being the Complements of the former to 90°

XXX. QUESTION. To find the Longitude of a Star, Planet, or Comet, from the Latitude 2° 59' N. and Declination 20° 1' S. being given?

N. B. This Question is resolved by a Projection on the Solstitial Colure. In a Spherical Triangle APE (AP being $\star$'s Dist. from the N. Pole of the World, AE $\star$'s Dist. from the N. Pole of the Ecliptic, PE Distance of the two Poles, and $\angle$ E the Longitude from $\triangle$, and $\angle$ P, the Right Ascension from γ)

*AP = $\star$'s D. fr. N. }
 Pole World 110° 1' 0" Sides op. $\angle$ reqd. Logarithms.
 AE = $\star$'s Dist. fr. N. }
 Pole Ecliptic 87 1 0 { Sides
 PE = Dist. of the two }
 Poles 23 28 30 { incl. $\angle$ re-
 quird. } L.S.co. 0.0005890
 L.S.co. 0.3997364

Sum 220 30 30
 From . . . Half 110 15 15

Take AE, 1 Rem. 23 14 15 L. S. 9.5960948
 Take PE, 2 Rem. 86 45 45 L. S. 9.9993134

Sum 19.9957336
 Half 9.9978668
 Double is $\angle$ E 168 39 7 Longitude from $\triangle$ following.
 Subtract 5 Signs 150 0 0 $\triangle$, γ , γ , $\triangle$, γ .

Place of the Star, } γ 18 39 7 In 2d. Quadrant of the Ecliptic.
 Planet or Comet. }

Or, 168 39 7 Longitude from $\triangle$ preceding.
 γ 180 0 0

Viz. γ 11 20 53 in 4th Quadrant of the Ecliptic.
 * Are given to find $\angle$ E fr. $\triangle$.

Astronomical QUESTIONS (in PRACTICE) answered by Spherical PROPORTION and LOGARITHMS.

Each Place being $78^{\circ} 39' 7''$ from the nearest Equinoctial Point $\triangle$ and ∇ respectively.

N. B. The Place (of the two here found) must be determined by the near Situation of the Planet or Comet whose Longitude is required, in respect of some known Star, whose Longitude is given.

EXAMPLE II. To find the Longitude of the Pleiades in the 294th Year before Christ, from their Declination at that Time $14^{\circ} 30' N.$ and Latitude $4^{\circ} N.$ being given? See fol. 12, Sherbone; fol. 56, Vincent Wing, Inst. and Page 16, Street.

In a Spherical $\triangle APE$, as before.
 Are given* } *AP = $75^{\circ} 30' 0''$ Side op. $\angle$ required. Logarithms.
 to find $\angle$ } AE = $86^{\circ} 0' 0''$ Sides inclu. } L.S.co. 0.0010592
 P fr. ∇ , or } PE = $23^{\circ} 28' 30''$ $\angle$ required } L.S.co. 0.3997364
 its Supt.
 Sum 184 58 30
 From Half 92 29 15

Take AE. 1 Rem. 6 29 15 L. S. 9.0530263
 Take PE. 2 Rem. 69 0 45 L. S. 9.9701881

Sum 19.4240100
 Double is $\angle E$ 62 1 34 Longitude, from ∇ preceding.
 90 0 0 ∇ from ∇ .

Place of Pleiades } $\nabla 27^{\circ} 58' 26''$ required.
 294 Y. bef. Chr. }
 Pl. 1720 since Xt. 8 26 5 8 Halley.

Sum 2014 Yrs. 28 6 42 Increase of Place.

2014) 101202" (50" 14" 57iv. —
 321
 1007

Annual Motion of the Stars in following Order.

XXXI. QUESTION. To find the Right Ascension of a Star, Planet, or Comet, from the Latitude $2^{\circ} 59' N.$ and Declination $20^{\circ} 1' S.$ being given, in the former or 30th Question?

In a Spherical Triangle APE,
 Are given* } *AE = $87^{\circ} 1' 0''$ Side op. $\angle$ required.
 to find $\angle$ } AP = $110^{\circ} 1' 0''$ Sides inclu. } L.S.co. 0.0270602
 P from ∇ } PE = $23^{\circ} 28' 30''$ $\angle$ required } L.S.co. 0.3997364

Sum 220 30 30
 From Half 110 15 15
 Take AP, 1 Rem. 0 14 15 L. S. 7.6175397
 Take PE, 2 Rem. 86 46 45 L. S. 9.9993134

Sum 18.0436497
 Double is $\angle P$, 12 4 20 R. A. from ∇ preceding.
 180 0 0

167 55 40 R. A. * &c. from ∇ following.
 90 0 0 R. A. ∇ from ∇

Sum 257 55 40 R. A. from ∇ .

Or, 12^h 4' 20" R. A. *, &c. from ∇ following.
 270 0 0 R. A. of ∇ from ∇ .

Sum 282 4 20 R. A. from ∇ following.

N. B. The Right Ascension of the two here found, must be determined by Circumstances, or Phenomena accompanying the above Data; the Vicinity of a Comet or Planet to a known Star of given Right Ascension, &c. But here in the same Case, when the Longitude is from ∇ or $\triangle$, the Right Ascension is from the same Equinoctial Point.

EXAMPLE II. To find the Right Ascension of the Pleiades in the 294th Year before Christ, from their Declination at that Time $14^{\circ} 30' N.$ and Latitude $4^{\circ} N.$ being given, as in the former Question?

In a Spherical Triangle APE,
 Are given* } *AE = $85^{\circ} 0' 0''$ Side op. $\angle$ required. Logarithms.
 to find $\angle$ } AP = $75^{\circ} 30' 0''$ Sides inclu. } L.S.co. 0.0140584
 P fr. ∇ , or } PE = $23^{\circ} 28' 30''$ $\angle$ required } L.S.co. 0.3997364
 its Supt.

Sum 184 58 30
 From Half 92 29 15

Take AP. 1 Rem. 16 59 15 L. S. 9.4656253
 Take PE. 2 Rem. 69 0 45 L. S. 9.9701881

Sum 19.8496082
 Half 9.9248041

Double is the Supplement $\angle P$ } $32^{\circ} 45' 11''$ L. S.
 65 30 22 R. A. Pleiades from ∇ preceding.
 R. A. 90 0 0 ∇ from ∇ .

R. A. Pleiades in the } 24 29 38 from ∇ following required.
 294th Y. bef. Xt. }

The Supplement of the included $\angle$ is, in the Case above, found instead of the included $\angle$ itself; the Sine and Sine of its Supplement to 180° being the same. This Distinction must be carefully regarded, according to the Nature of the three Sides of the Spherical Triangle given, that it may be known from what Part of the Solstitial Colure ∇ or ∇ , the $\angle$ of Right Ascension, or Longitude is determined.

When you would find the Times of Rising and Setting of the Moon, or Planets, or the Time, at any Altitude, their Amplitudes of Rising and Setting, &c. You must substitute the Times of their Passage through the same Meridian, in the preceding Calculations, instead of Noon, the Time of the Sun's Passage.

The Time of a Planet from the Meridian, is reduced to an Arc of the Equator by this PROPORTION.

As the Interval of the Time betwixt two next Passages of the Moon or a Planet, through the same Meridian is to 360° so is the Interval betwixt the given Time, and the Time of nearest Passage through the Meridian, to the Degrees, Minutes and Seconds of the Planets angular Distance, measured on the celestial Equator, from the same Meridian, when the Altitude of the Planet is required.

XXXII. QUESTION. To find the Altitude of Mars, the 12th Feb. 1758, at Six in the Morning; being the same as the 11th Feb. 18^h past Noon?

δ passed the Meridian 1758, Feb. 11 { 11 16 40 } At Paris De la
 12 { 11 11 20 } Caille's Ephem.

Dif. 0 5 20
 24 0 0

Interval of δ Return to the same Merid. 23 54 40

d h m s
 Feb. 11 18 0 0 Time given,
 11 11 16 40 Souths.

δ from Meridian 0 6 43 20 Past.

As $23^h 54^m 40^s : 360^{\circ} :: 6^h 43^m 20^s : 101^{\circ} 12' 29''$
 Correspondent Dist. δ from Meridian, 1758.

Feb. 11^d 18^h

Declination δ $21^{\circ} 52' 0'' N.$ Feb. 11^d 18^h 1758, La Caille,
 48 50 10 N. Height Pole, Paris.

Astronomical QUESTIONS (in PRACTICE) answered by Spherical PROPORTION and LOGARITHMS.

By PROPORTION, QUEST. XI.		Logarithms.
As Radius	90° 0' 0"	10.0000000
To Sine Comp. Dist. δ fr. Meridian	101 12 29	9.2886448
So Tan. Comp. Lat. Paris,	48 50 10	9.9416710
To Tangent Arc X	9 38 43	9.2303158
Dist. δ from Pole +	68 8 0	See Below.
To Sine Complement Arc Y	77 46 43	9.3256996
As Sine Complement Arc X	9 38 43 co.	0.0061844
So Sine Latitude	48 50 10	9.8766969
To Sine Altitude δ required	9 18 10	9.2085809

M. De la Caille made it (See his Ephem.) 9 12 8 only.
Who found the Arc X 9° 47' (See p. 16 La Caille Eph.) instead of 9° 38' 43" as above.

Who finds 101° 12' 25" = 6h 43m 20s instead of 101 12 29 = as above.

* When Distance from the Meridian is above 6h or 90° add X to the Planet's Distance from Pole for Y. if less, subtract X from Pole's Distance from Planet for Y.

N. B. The same Method as that above serves for all other Planets, and also the Moon; if you observe to take out of an Ephemeris the true Interval of Time betwixt the two next Passages of the Moon, or a Planet through the same Meridian, for proportioning their Distances in Time therefrom, either before or after their Passage at a Time given by the Clock.

XXXIII. QUESTION. To find the Right Ascension of the Mid-Heaven Ecliptic at the Meridian, from the Sun's Place, or R. Ascension, and the Time of the Day or Night being given?

RULE. To the Sun's Right Ascension in Degrees, Minutes, &c. found to the Time given, when Right Ascension of Mid-Heaven is required, add the Degrees, &c. of that Time from the former Noon (allowing 15° Degrees to an Hour) and the Sum rejecting 360° if above) will be the R. Ascension of the Mid-Heaven, at the Meridian, required.

EXAMPLE. Required the Right Ascension of Mid-Heaven for 1764, March 30th, 10h 20m in the Forenoon?

Sun's R. A. in Time.

1764 Mar. 29, Noon 34m 6s } De la Caille's Ephemerides.
30, Noon 37 43 }

Increase Sun's R. A. in 24h 3 37

As 24h : 3m 37s :: 22h 20m : 3m 22s Increase.
+ 34 6

Right Ascension in Time 1764, }
Mar. 30th. 10h 20m Morn. } 37 28

By Table p. 28. 37m 0s . . . 9° 15' 0"
0 28 . . . 0 7 0

R. A. of the Sun in Degrees at the Time 9 22 0

(Or See p. 194.)

Now, 22h 20m reduced to Degrees = 335° 0' 0"

Right Ascension of Mid-Heaven required . . Sum 344 22 0
360 0 0

R. A. of Mid-Heaven, short of γ . . . 15 38 0

When the Sum aforesaid is less than 90° it is Right Ascension Mid-Heaven from γ following; if above 90° and less than 180, subtract it from 180, for R. Ascension of Mid-Heaven from α preceding; if more than 180, and less than 270, subtract 180 from it for R. A. of Mid-Heaven from α following: But if it is above 270°, and less than 360, subtract it from 360° for Right Ascension of Mid-Heaven from γ preceding, as in the Example given.

This Question is of singular Use in the Computation of Eclipses, subservient to Navigation.

XXXIV. QUESTION. To find the Point of the Ecliptic, at the Meridian, or Mid-Heaven Longitude, correspondent to the Mid-Heaven Right Ascension 15° 38' from γ preceding; and Ecliptic obliquely 25° 28' 30" being given?

PROPORTION.		Logarithms.
As Radius	90° 0' 0"	10.0000000
To Cosine Ecliptic Obliquity	23 28 30	9.9624801
So Cot. R. A. Mid-Heaven	15 38 0	10.5531022
To Cot. Mid-Heaven fr. γ preceding	16 57 58	10.5155823
	12° 0' 0"	

Longitude Mid-Heaven rem. Sig. . 11 13 2 2 or culminating [Point of the Ecliptic required.]

N. B. The Mid-Heaven Longitude is always found from the same Equinoctial Point γ or α , that the Mid-Heaven Right Ascension is taken from.

XXXV. QUESTION. To find the Meridian Angle, or Angle of the Ecliptic and Meridian, from the Mid-Heaven Right Ascension 15° 38' and Ecliptic Obliquity 23° 28' 30" being given?

PROPORTION.		Logarithms.
As Radius	90° 0' 0"	10.0000000
To Sine Ecliptic Obliquity	23 28 30	9.6002636
So Cosine R. A. Mid-Heaven	15 38 0	9.9836290
To Cosine Meridian $\angle$ required	67 26 33	9.5838926

XXXVI. QUESTION. To find the Mid-Heaven Declination, or Declination of the Point of the Ecliptic at the Meridian, from the Mid-Heaven Right Ascension 15° 38' and Ecliptic Obliquity 23° 28' 30", being given?

PROPORTION.		Logarithms.
As Radius	90° 0' 0"	10.0000000
To Tan. Ecliptic Obliquity	23 28 30	9.6377835
So Sine R. A. Mid-Heaven	15 38 0	9.4305267

To Tan. Declination on the Merid. S. reqd. 6 43 15 9.0683102

N. B. When the Right Ascension of the Mid-Heaven is above 180°, as in the present Case, then the Mid-Heaven Declination of the Ecliptic Point is South. But if the Mid-Heaven Right Ascension is less than 180°, then the Mid-Heaven Longitude being North, the Mid-Heaven Declination will be North.

XXXVII. QUESTION. To find the Altitude of the Mid-Heaven, from the Latitude of the Place 51° 28' 30", and Mid Heaven Declination 6° 43' 15" S. being given?

RULE: If the Latitude and Mid-Heaven Declination are of the same Name, add the Declination to the Complement of Latitude for the Mid-Heaven Altitude.

If the Latitude and Mid-Heaven Declination are of different Names, subtract the Declination from the Complement of Latitude for the Mid-Heaven Altitude.

EXAMPLE. Complement of Latitude . . . 38° 31' 30" N.
Mid-Heaven Declination . . . 6 43 15 S.

Mid-Heaven Altitude required . . 31 48 15

XXXVIII.

Astronomical QUESTIONS (in PRACTICE) answered by Spherical PROPORTION and LOGARITHMS.

XXXVIII. QUESTION. To find the Altitude of the Nonagesimal, or 90th Degree of the Ecliptic from the Horizon (being equal to the Angle of the Ecliptic and Horizon) from the Mid-Heaven Altitude $31^{\circ} 48' 15''$, and the Meridian Angle $67^{\circ} 26' 33''$ being given?

As Radius Sine	$90^{\circ} 0' 0''$	Logarithms.	10.0000000
To Sine Meridian $\angle$	$67^{\circ} 26' 33''$		9.9654345
So Cofine Altitude Mid-Heaven	$31^{\circ} 48' 15''$		9.9293445

To Cofine Altitude 90th Degree required $38^{\circ} 17' 40''$ 9.8947790

Then you may proceed to find the Place of the 90th Degree of the Ecliptic, or its Distance from the Mid-Heaven Longitude, THUS, As Radius to Cofine Altitude Mid-Heaven, So Cofine Merid. Angle to Tan. Dist. Mid-Heaven from the 90th Degree. Which Place is found by adding the Distance to, or subtracting it from the Distance of the Mid-Heaven, according as its Place follows or precedes the Place of the Mid-Heaven.

Viz, If the Mid-Heaven be in $\{ \text{Aries, Taurus, Gemini, Cancer, Leo, Virgo, Libra, Scorpio, Sagittarius, Capricorn, Aquarius, Pisces} \}$, add the Distance of Heaven be in $\{ \text{Aries, Taurus, Gemini, Cancer, Leo, Virgo, Libra, Scorpio, Sagittarius, Capricorn, Aquarius, Pisces} \}$, subtract the Distance of Mid-Heaven from the 90th Degree, to and from the Place of Mid-Heaven, for the Place of the 90th Degree, in North Latitudes, and do the contrary in South Latitudes.

Then the Point of the Ascendant has been usually found, or the Ecliptic Place at the Horizon, at a given Time, for the Purpose of finding the Parallaxic Angle, or that Angle which a Vertical (passing through the Sun, Moon, or Star) makes with the Ecliptic, for computing ECLIPSES and OCCULTATIONS.

But as these Methods are tedious and obsolete, we give the following UNIVERSAL NEW METHOD of finding the PARALLACTIC ANGLE, and of computing the MOON'S PARALLAXES.

XXXIX. QUESTION. To find the apparent Place of the Moon, from her Longitude, or true Place, and Latitude given?

Find the Δ 's Declin. and R. A. (by Quest. XXVII. and XXVIII.) answerable to the Longitude and Latitude given; for in a Spherical Triangle APE, there are given AE the Complement of Latitude; PE, the Distance of the Poles of Equator and Ecliptic; and $\angle E$, the Longitude from Δ or Δ , the Solstitial Colure, to find AP the Complement of Declination, and $\angle P$, the Right Ascension from Δ or Δ ?

Or take out the Declination and Right Ascension to the Time given, from Ephemerides, De la Caille's, Connaissance des Temps, &c. reduced to Greenwich-Meridian.


EXAMPLE. To find the Moon's apparent Place from the true, $11^{\circ} 10' 44''$, for the Year 1737, February 18^d 2^h 25^m 9^s apparent, or 2^h 38^m 36^s mean Time, Afternoon, O. S. about the Beginning of the solar Eclipse, observed at Greenwich.

The Sun's Longitude then	$11^{\circ} 6' 55''$
His Right Ascension correspondent	$342^{\circ} 34' 48''$
Declination correspondent S.	$7^{\circ} 24' 28''$
Moon's Longitude then, by Theory,	$11^{\circ} 10' 44''$
Latitude N.	$0^{\circ} 37' 40''$
Declination correspondent S.	$6^{\circ} 48' 11''$
Right Ascension correspondent	$342^{\circ} 23' 56''$
Time from Noon apparent, in Degrees, $360^{\circ} 17' 15''$	
Hence R. A. of Mid-Heaven	$18^{\circ} 52' 3''$
Mid-Heaven Declination correspondent N.	$7^{\circ} 59' 40''$
Altitude of Mid-Heaven at Greenwich	$46^{\circ} 31' 10''$
Mid-Heaven Zen. Dist. from the Ecliptic Altitude at the Meridian	$43^{\circ} 28' 50''$
Mid-Heaven R. A. — Moon's R. A. is the Moon's Angular Dist. fr. the Meridian, past	$36^{\circ} 28' 7''$
Angle of Ecliptic and Meridian, to R. A. Mid-Heaven, or Mid-Heaven Longitude	$67^{\circ} 51' 20''$

First to find the Moon's Altitude and Azimuth?

In the $\{ \begin{array}{l} \text{Fig. are } \{ \begin{array}{l} \text{ZP} = 38^{\circ} 31' 30'' \text{ Comp. Latitude,} \\ \text{QP} = 96^{\circ} 48' 11'' \text{ Comp. Moon's Declination,} \\ \text{given. } \angle \text{QPZ} = 36^{\circ} 28' 7'' \text{ Moon's angular Dist. fr. Meridian.} \end{array} \end{array} \right.$

To find QZ = Comp. Altitude? And $\angle \text{QZP}$ = the Moon's Azimuth from the North?



N. B. The Line PQ terminates in ZG, above the Intersection of MN and ZG, between r and o.

PROPORTION.

As Radius	S. 90°	Logarithms.	10.0000000
To Cof. angular Dist. from Meridian $\angle \text{QPZ}$	$36^{\circ} 28' 7''$		9.9053547
So Tan. Comp. Lat. ZP	$38^{\circ} 31' 30''$		9.9009941
To Tan. 4th Arc	$32^{\circ} 37' 46''$		9.8063488
Δ Decl. and Lat. Place of diff. Name.	Add $83^{\circ} 11' 49''$	Comp. Decl. [S.	
5th Arc	Sum $115^{\circ} 49' 35''$ or $64^{\circ} 10' 25''$		

N. B. Subtract the 4th Arc from the ~~Cofine Declination~~ ^{polar dist} for the 5th Arc, when the Declination and Latitude of the Place are of a different Name.

As Cof. 4th Arc	$32^{\circ} 37' 46''$ co.		0.0745974
To Cof. 5th Arc	$64^{\circ} 10' 25''$		9.6391334
So Cof. ZP or S. Latitude	$51^{\circ} 28' 30''$		9.8933936
To Cof. Alt. QZ or S. Alt. QG	$23^{\circ} 52' 19''$		9.6071244

NOW, As Cof. Alt. GQ or S. QZ	$66^{\circ} 7' 41''$ co.		0.0388390
To S. Δ Ang. Dist. fr. Merid. $\angle \text{QPZ}$	$36^{\circ} 28' 7''$		9.7740658
So S. Dist. from Pole QP	$96^{\circ} 48' 11''$		9.9969314
To S. Δ 's Azim. } from the S.	$\angle \text{MZQ } 40^{\circ} 11' 48''$		9.8098362

Here, In the Spherical Triangle MZr, the $\angle \text{MZr}$, $\angle \text{rMZ}$, (Supplement of $\angle$ of Mid-Heaven Ecliptic, and Meridian) with the included Side MZ, the Zenith Distance of the Mid-Heaven Ecliptic, to find the Parallaxic Angle, or $\angle \text{MrZ}$, that the Vertical ZG passing through the Moon's Center, makes with the Ecliptic MN, without having Regard to the $\angle \text{HFM}$, of the Horizon and Ecliptic, commonly called the Altitude of the Nonagesimal or 90th Degree of the Ecliptic, to which that Angle of the Horizon and Ecliptic is equal?

PROPORTION.

Astronomical QUESTIONS (in PRACTICE) answered by Spherical PROPORTION and LOGARITHMS.

For the Parallax in Longitude.			Logarithms.
As Radius	90° 0' 0"		10.0000000
To Tan. Par. in Alt.	0 49 39		8.1596689
So Cofine Parallaic $\angle$	43 48 20		9.8583531
To Tangent Parallax in Longitude	35 40		8.0180220
Add in the East and subtract in the West } $\times 11$	10 44	True Long.	
D visible Longitude $\times 10$ 34 54 required.			
For the Parallax in Latitude.			
As Radius	90° 0' 0"		10.0000000
To Sine D's Parallax in Altitude	0 49 39		8.1596689
So Sine Parallaic $\angle$	43 48 20		9.8402398
To Sine Parallax in Latitude	34 23		7.9999087
Sub. from . True Latitude D N. D.	37 40		
Visible Latitude D N.	3 17	required.	

In computing the Moon's Parallax, by Dean Cowper's *short Method*, for the Time of ECLIPSES of the Sun, the Moon's Right Ascension may be taken the same with the Sun's, without sensible Error. And if, instead of the absolute horizontal Parallax of the Moon, you take the Excess of the Moon's horizontal Parallax, above that of the Sun, you will have the Parallax of the Moon from the Sun, in Altitude, Longitude and Latitude.

The Old Method of finding the Parallaic Angle (from whence the Parallaxes in Altitude, Longitude and Latitude are had) is thus,

The Altitude and Longitude of the Nonagesimal or 90 Degree of the Ecliptic from the Horizon, being computed by Rules before delivered, or found from Tables, on Purpose, for a given Latitude.

Say, As Radius to Cofine Dist. D from the 90° of Ecliptic, So is the Tang. Alt. 90° to Tang. 4th Arc.

This Arc taken from the Complement D's Latitude, or Distance from the North Pole of the Ecliptic, leaves a 5th Arc.

Then, As Cofine 4th to Cofine 5th Arc,

So is Cofine Altitude 90° Ecliptic to Sine D's Altitude.

And, As Cofine D's Altitude to Sine Dist. from 90° Ecliptic,

So is Sine Altitude 90° to Cofine Parallaic $\angle$ sought.

But this old Method is far exceeded by Dean Cowper's new and compendious One, before delivered; which however is here given Place to for the practical Astronomer's Satisfaction, that he may compare the different Methods, and choose the best.

Mr. Dunthorne, in his Astronomy, p. 63, recommends the finding the Angle, at the Moon's Center, which the Vertical makes with a Circle of Latitude, and thence the Moon's Parallax, in Longitude, and Latitude, which he says is the more exact. By this we apprehend, that Mr. Dunthorne directs us to find the Angle that the Moon's Orbit makes with the Vertical Circle at the Moon's Center. But since the Longitudes and Latitudes of the Sun, Moon, and Stars, refer to, or are computed to the Ecliptic, the Parallaic Angle and all the Parallaxes of Longitude, Latitude, &c. derived therefrom, must have also Reference to the same Ecliptic, and not to the Lunar Orbit. For it is not the Orbit but Ecliptic Place of the Moon we compute from.

And a Solar Eclipse 18th Feb. 1737, being computed by the Operose Method in *Leadbetter's Uranoscopia*, from p. 147 to p. 153, viz. for finding the Meridian Angle, culminating Point of the Ecliptic, Alt. of Mid-Heaven, Nonagesimal Degree, &c. in the Lunar Orbit, the same Eclipse was also computed with the same Requisites in the Ecliptic; the Eclipse by the last Method was found to come nearest to Observation.

Likewise in the Examples of the Use of Dr. Halley's Tables (d, d 2) the Nonagesimal Degree and Parallaic Angle at the Moon (i. e. corresponding to the Moon's Place) are both taken in the Ecliptic (and not in the Lunar Orbit) tho' the Moon had almost 5 Degrees of Latitude; which our universal Method of finding the Parallaic $\angle$, allows for.

To apply the Parallaxes to the computed Places of the Moon by the Tables.

RULES.

1. When the Moon is to the East of the 90° of the Ecliptic, the Parallax in Longitude must be added to the true for the visible Longitude; but when she is to the Westward thereof it must be subtracted from the true for the visible or apparent Longitude.

2. The above Rule holds good for Right Ascensions, according as the Moon is to the East or West of the Meridian of the Place of Observation.

3. The Parallax in Latitude being added to the true Distance of the Moon from the Pole of the Ecliptic, which is nearest to the Vertex of the Place of Observation, will give the apparent or visible Distance of the Moon from that Pole, whence it will appear, whether this Parallax is to be added to, or subtracted from the true Latitude for the apparent. But between the Tropics, where the Ecliptic is vertical once in 24 Hours, at such Times it is always to be added to the true for the apparent Latitude.

4. In like Manner, the Parallax in Declination being added to the true Distance of the Moon from that Pole of the Equator, nearest to the Vertex of the Place of Observation, will give the apparent or visible Distance from the Pole. Whence it will appear whether this Parallax is to be added or subtracted to or from the true for the apparent Declination. But in Places on the Globe which have no Latitude, the Parallax in Declination is always additive to the true for the apparent Declination.

The NAVIGATOR being Master of applying the MOON'S PARALLAXES, the Longitudes betwixt Places on the Globe, may be determined by observing the Moon's Appulses to a fixed Star, or her Distance from a fixed Star, lying near the Parallel of Latitude where the Moon then is (to avoid the Effect of Error in the Moon's computed Distance from the Star, resulting from computed Error of her Latitude) or else by the Distance of the Moon from the Sun in the first or last Quarter. From any of which Observations, the Latitude of the Place, and Time of an Observation being known, the Distance of the present Meridian of the Place from that of Greenwich, is discovered by finding the correct Time at Greenwich, from our Tables, when such Phenomena could happen: The Difference of which Times being turned into Degrees (allowing 15° to an Hour) will shew the Distance of the two Meridians of those Places from each other, required. Which METHOD, with necessary Examples we have exhibited further on.

PARALLAXES determined from EPHEMERIDES: Relative to the PRECEDING QUESTION of the PARALLACTIC ANGLE, and also to TABLE p. 215, of the EQUATION of the Moon's Horizontal Diameter to the Moon's Diameter in Altitude.

XL. QUESTION. To find the true from the apparent or visible Altitude of the Moon?

EXAMPLE I. Suppose the apparent Altitude of the Moon's Center at Paris, 6th October 1761, at 9h 53m Night, to be 27° 33' 4", what will be the Moon's true Altitude at that Time?

D's Hor. Parallax.

1761 Oct. 6 Noon { 55' 2" }
11 Noon { 54 40 } De la Caille's Ephemerides.

If 5 Ds. decreases 22" : 9h 53m : Decr. 1" 49", or 2"
55' 2"
— 2

Hor. Par. 55 0, at 9h 53m Night.

Astronomical QUESTIONS (in PRACTICE) answered by Spherical PROPORTION and LOGARITHMS.

Again As Radius,	S. 90° 0'	Logarithms. 10.0000000
To Cosine Δ Declination at that Time	12 15	9.9899973
So 29' 53" or Mot. of Equator	S.	7.9391497
To 29' 12" or S.	S.	7.9291470
The Δ 's true Horizontal Diameter required.		

M. De la Caille has the Sem. Diameter of the Moon, 14' 48" $\frac{1}{2}$ or 29' 37" the Diameter. See his Ephem. 1763, Oct. 21. N. S. differing from his own Example by 25" Horizontal Parallax Δ then 54' 46" N. B. The Sines of small Arcs are nearly the same as those Arcs. After the same Manner are correctly computed the true Diameter of the Sun and of each of the Planets.

XLIV. QUESTION. To find the Inclination or Angle of the Moon's Orbit and Equinoctial, from the Longitude of the Moon's Node being given, $26^{\circ} 0'$?

Universal PROPORTION.

As Radius	Sine 90° 0' 0"	10.0000000
To Cosine Declination, correspondent to the Longitude Δ from Υ or Δ ,	23 28 30	9.9624801
So Sine Diff. betwixt the Incl. of the Ecl. with Δ 's Orb and Mer. $\angle$ fr. Δ in Δ to 26° (but S. Sum of said Incl. and Merid. $\angle$ fr. Δ in 26° to Δ) viz. in this Case S. 85 0 25 (Inclin. Δ 's Orb. with the Ecliptic being 4 59 35)	0 25	9.9983488
To Cosine $\angle$ of the Δ 's Orb with the Equinoctial required	23 58 15	9.9608289
Leadbetter in his Astron. 2 vol. makes it (Tab. p. 71)	23 29 0	$\angle$ of Ecliptic and Equi. for Δ 's Orb. & Eq. 29 15 his Error.

EXAMPLE II. To find the Inclination of the Moon's Orb with the Equinoctial, when the Longitude of the Δ 's Node is 2° ?

As Radius	Sine 90° 0' 0"	10.0000000
To Cosine Declination correspondent to Longitude Δ 's Δ	19 44 39	9.9736867
So S. Sum of the Inclination of Ecliptic and Δ 's Orb. and Meridian $\angle$, viz. Sine 82 6 51 (Angle Inclination with Ecliptic, being 5 4 28 and Meridian $\angle$ 77° 2' 23")	5 4 28	9.9958735
To Cosine $\angle$ of the Δ 's Orb. and Equinoctial required	21° 12' 8"	9.9695602
Leadbetter in his Astron. 2 vol. makes it (Tab. p. 72)	21 36 10	

+ 24 2 his Error.

Greatest $\angle$ of Ecliptic with the Moon's Orbit Δ in Υ and Δ } 5° 17' 20"
 $\angle$ of the Ecliptic and Equinoctial in all Positions of the Δ 's Δ } 23 28 30

Greatest and least $\angle$ of the Δ 's Orb with Equin. Δ in Υ 28 45 50
 Δ 18 11 10
 correctly.

The two last Instances of the Inclination of Δ 's Orb with the Equinoctial, are the only two, wherein Leadbetter's Tables, p. 71 and 72, are true, in all the 180 Operations; however he came to be mistaken in the Rules of so essential a Point of Computation.

XLV. QUESTION. To find the Time from Noon of the Conjunction (or any other Aspect) of the Moon with the several Planets, from the Difference of Diurnal Motion of the Moon, and of any Planet, and the Distance of the Moon from the Conjunction or other Aspect of that Planet at Noon, (less than the given Difference of Diurnal Motions) being given?

According to M. De la Caille's Ephem. for Paris.

1764 October 23. Δ 's Longitude Noon 7° 54'
 24 22 44

Δ 's Diurnal Motion 14 50

Sub. Place Δ 23 Oct. from Place Δ 23 Δ Longitude Noon 15 44
 Δ 23 Oct. for their Dist. fr. Conj. Δ 24 15 55

Δ 's Diurnal Motion 0 11

Difference Diurnal Motion Δ à Δ 14 39

Dist. Δ before Δ 23d October Noon 7 50
 viz. fr. both in the same Sig. and Degree

N. B. Always subtract the Place of the swifter from the Place of the slower Body, for their Distance, in proportioning the Time required for the swifter overtaking the slower Body, from their Difference of Motions.

RULE. As Difference of Diurnal Motion of the Moon and any Planet, in Degrees and Minutes, is to 24 Hours, so is the Distance of the Moon from any Planet (or other Aspect) at Noon, in Degrees and Minutes (less than the said Difference of Diurnal Motions) to the Hour and Minute forward from that Noon, when the Conjunction (or other Aspect) of the Moon with that Planet happens.

Thus, As Difference Diurnal Motion Δ à Δ 14° 39' co. 9.3877
 To 24 Hours 0.3979
 So Distance of Δ before Δ at Noon 23d Oct. 7 50 0.8842

To Time of Conjunction Δ Δ past Noon } 12h 50m
 1764 October 23. Oct. 24th 50m Morning, Paris.

DECIMALLY.

As 14° 39' to 24h So 7° 50' to 12h 50m
 = 14.65 = 7.833 . by 6 } = 24
 46,998 . by 4 } = 117.20
 14.65 187,992 (12h.83 = 12h 50m fere, past Noon at Paris, or 12h 40m 40s at Greenwich.

4149

2930

12192

11720

4720

4395

rem. 325

N. B. M. De la Caille (by Inattention) made the Conjunction of the Δ and Δ to happen on 21st Oct. 1764, instead of on 24th, as above. See his Ephemerides for that Year; Col. Des Phenomenes & Observations, p. 255.

Δ 's Latitude at the Time of the above Conjunction 1° S. 50'
 Δ Latitude then 1 N. 16

Sum 3 6

Consequently the Moon passes under Venus in the said Time of Conjunction, at the Distance of 3° 6'. For whatever Planet has most Northern Latitude, or is nearest to our elevated North Pole, is the uppermost or superior Planet, when any two Planets happen in Conjunction, or in the same Sign of the Ecliptic.

Astronomical QUESTIONS applied to PRACTICE.

OPERATION of the Latitudes.

Moon's Lat.	♀ Lat.
1764 Oct. 23 1° S. 9'	1° N. 9'
24 2 25	1 22
As 24 ^h to 1 16	As 24 ^h to 0 13 Difference.
So 12 ^h 50 ^m to 0 41	So 12 ^h 50 ^m to 0 7 To add to 1st Lat.
♂'s Lat. 24 D. 1° S. 50'	1° N. 16' ♀ Latitude.

The Conjunction of the Planets with one another are found in the same Manner. And the other Aspects of the Moon with the several Planets, and Planets with one another, are found in the same Manner also, by proportioning the Distance of the Moon or Planet from its Trine, Quartile, Sextile, or Opposition, at Noon, viz. from 120°, 90°, 60°, or 180° respectively distant from one another at Noon.

EXAMPLE II. To find the Time of the first Planetary Opposition with the Moon in April 1764?

Examining Ephemerides for Paris by De la Caille.

The first Sign of the Moon having Opposition by a Planet, is the ♃ in ♐, and ♄ in the opposite Sign ♈.

1764 April 6, ♃'s Longitude at Noon 12° 11' 16"

7 24 24

♃'s Diurnal Motion 12 8

Sub. ♃'s Place April 6, } 6 ♄ Longitude at Noon 15 11
From ♄'s Place April 6, } 7 ♄ Longitude at Noon 15 25
for their Dist. fr. Opposit.

♄ Diurnal Motion 0 14

Difference Diurnal Motion ♃ à ♄ 11 54

Distance of ♃ from an Opposition with ♄ at Noon, Apr. 6, } 2 55
viz. from an opposite Sign and Degree.

Now, As Difference Diurnal Motion ♃ à ♄ 11° 54' co. Lo. Log. 9.2974
To 24 Hours 0.3979
So Dist. ♃ and ♄ fr. Opposition Apr. 6th 1764 2 55 1.3133

To Time Opposition ♃ ♄ past Noon 1764 Apr. 6. 5^h 53^m 1.0086
at Paris, or 5^h 43^m 40^s [at Greenwich.]

DECIMALLY.

As 11° 54' to 24^h so 20.55^m to 5^h 53^m
= 110.9 = 20.916 by 6 } = 24
17.496 by 4 }
110.9)69.984(5^h 53^m 88 = 5^h 53^m past Noon at Paris, or
595 .. 5^h 43^m 40^s at Greenwich.

1048

952

964

952

Rem. 12

In the same Manner you are to proportion the Distance of the Moon, or swifter moving Planet, from any Aspect of another Planet at Noon: The Aspects of the Moon, with all the Planets, and each with one another being discoverable at Sight, by the annex'd Scheme, or Figure; which is particularly necessary for finding the Oppositions of the Sun and any of the Stars or Planets.

OPERATION of the Latitudes.

Moon's Lat.	♀ Lat.
1764 April 6 4° N. 52'	0° N. 19'
7 5 11	0 21
As 24 ^h to 0 19	As 24 ^h to 0 2 Difference
So 5 ^h 53 ^m to 0 5	So 5 ^h 53 ^m to 0 1 To add to 1st Lat.
♂'s Lat. at the Op. of ♃ and ♄ 4° N. 57'	0° N. 20' ♂ Lat. at the Op. of ♃ and ♄
0 N. 20'	


4° N. 37' Moon superior to ♀.

THE following FIGURE is for discovering the TIMES of the OPPOSITIONS, or other ASPECTS, of the Celestial BODIES, from their PLACES in the ECLIPTIC being given, for any YEAR, and all the Month-Days.

Signs

2 Distance in the Eclip-
3 tic between two Ce-
4 lestial Bodies called
6 } Sextile.
Quartile.
Trine.
Opposition.

Two Bodies being in the same Point of the Ecliptic is called Conjunction. See the Figure below.



EXAMPLE III. To find the Times of the New, Full, and Quarter MOONS for any Month in any Year; the Sun and Moon's Places to each Month and Day in that Year being given?

For the Conjunction, or New Moon.

Place.	♃ Place.	♃ before ♃.
1764 March 2 12° 27' 40"	10° 0'	2° 27' 40"
3 13 27 41	22 13	
Diurnal Motions ♃ and ♃	12 13	ris.
	1 0	

Dif. Diurnal Motion ♃ à ♃ 11 13

Now, As * Dif. Diurnal Motion ♃ à ♃ 11° 13' co. Lo. Log. 9.2717
To 24 Hours 0.3979
So Dist. ♃ before ♃ at Noon from Conjunction 2 28 fere 1.3860

To Time of their Conjunction 5^h 16^m past Noon, March 12th, 1764, at Paris or 5^h 7^m at Greenwich, required.

* When the Planets are retrograde take the Sum instead of Difference of Diurnal Motion.

EMENDATIONS. P. 228, 1 Col. l. 5. read ♈ or ♉ for ♈ and ♉. P. 231, 2 Col. N. B. Sub. 4th from Co. Declination or Dist. from N. Pole, for 5th Arc, when the ♄ from the Meridian is less than 90°; but add the 4th Arc to Co. Declination or Dist. from the N. Pole, for 5th Arc, when ♄ from the Meridian is greater than 90°. P. 231. 1 Col. l. 17. for Distance r. Place of the Mid-Heaven. N. B. Adding the 4th Arc to the Co-Declination, when of contrary Name to the Pole, and taking the Supplement of the Sum for a 5th Arc, is the same as subtracting the said 4th from the Dist. from the Pole, for a 5th Arc. See Examp. p. 231.

Astronomical QUESTIONS applied to PRACTICE.

DECIMALLY.
As $11^{\circ} 13'$ to 24^h for $2^{\circ} 28'$ to $5^h 16^m$
= 11,216 = 2,466 by 6 } = 24
14,796 by 4 } = 24
11,216 59,184 (5^h, 27 = 5^h 16^m as before.
56080
31040
22432
86080
78512

Rem. 7563

For the First Quarter Moon ?

	Place.	Place.	before 3 ^h à D.
1764 March 10	20° 27' 0"	15° 11' 47"	4° 40'
11	21 26 45	28 8	Note, X and II are distant 3 ^h by foregoing Figure.
Diurnal Motions Q & D	59 45	12 21	
		I	
		II 21	

NOW, As Dif. Diurnal Motion D à Sun . $11^{\circ} 21'$ co. 9.2768
To 24 Hours 0.3979
So Dist. Sun before D at Noon, from 3^h, or } 4 40
First Quarter 1.1091

To the Time past Noon in their Distance 3^h or } 9^h 52^m 0.7838
1st Quarter at Paris
Or 9 43 at Green-
[wich req.]

DECIMALLY.
As $11^{\circ} 21'$ to 24^h for $4^{\circ} 40'$ to $9^h 52^m$
= 11,35 = 4,666 by 6 } = 24
27,996 by 4 } = 24
11,35 111,984 (9^h, 86 = 9^h 52 nearly; as before.
10215

9834
90807540
6810

Rem. 730

The slower moving Body of the two, must be left than the Difference of the Diurnal Motions, before the faster moving Body, to find the Time of all the Aspects, required to be known, betwixt the two Bodies.

For the Opposition, or full Moon.

	Sun's Place.	Place.	Sun bef. Opp ⁿ . D
1764 March 17	27° 24' 41"	20° 11' 21"	7° 3' 41"
18	28 24 11	5 21 16	Note, X and II are opposite by the Figure foregoing.
Diur. Motions Sun & D	59 30	14 55	
		I	
		II 55	

Difference Diurnal Motion D à Sun 13 55
NOW, As Dif. Diurnal Motion D à Sun . $13^{\circ} 55'$ co. 9.3654
To 24 Hours 0.3979
So Dist. Sun before D at Noon from Opposition . 7 4 0.9289
To the Time past Noon of her Opposition at Paris, 12^h 11^m 0.6922
Or 12 2 at Green-
[wich req.]

DECIMALLY.
As $13^{\circ} 55'$ to 24^h for $7^{\circ} 4'$ to $12^h 11^m$
= 13,916 = 7,066 by 6 } = 24
42,396 by 4 } = 24
13,916 169,584 (12^h, 18 = 12^h 11^m as before.
13916

30424
2783225920
13916120040
111328

Rem. 8712

N. B. The Difference of Meridians, in Time, between Greenwich and Paris is 9^m 20^s, that the same Phenomenon happens sooner at Greenwich than at Paris.

For the last Quarter Moon.

	Sun's Place.	Place.	Sun before 9 ^h à D.
1764 March 24	4° 20' 44"	2° 12' 28"	1° 52' 44"
25	5 20 0	15 55	Note, Q and W are 9 ^h Distance by Figure foregoing.
Diur Motions Sun & D	59 16	13 27	
		— 59	
Difference Diurnal Motion D à Sun		12 28	

NOW, As Dif. Diurnal Motion D à Sun . $12^{\circ} 28'$ co. 9.3176
To 24 Hours 0.3979
So Dist. Sun before D at Noon from 9^h the last }
or third Quarter 1 53 1.5038

To the Time past Noon in their Dist. of 9^h, or last }
Quarter at Paris 3^h 37^m 1.287
Or 3 28 at Green-
[wich req.]

DECIMALLY.
As $12^{\circ} 28'$ to 24^h for $1^{\circ} 53'$ to $3^h 38^m$
= 12,466 = 1,883 by 6 } = 24
11,298 by 4 } = 24
12,466 45,192 (3^h, 12 = 3^h 37^m.
37398

77940
7479631440
24932

Rem. 6508

N. B. When a Planet is retrograde, the Sum, instead of the Difference of Diurnal Motion, must be taken for all the Aspects.

XLVI. QUESTION. To find the Time of the Moon's, and also a Planet's Southing, Rising and Setting (practically) from Ephemerides, and preceding Tables.

FIRST, For the Moon's Southing, or Passage through the Meridian ?

RULE. Subtract the Sun's Right Ascension in Time, from the Moon's Right Ascension in Time at Noon, on the Day preceding the Time when the D's Southing is required, and the Remainder will be the Time preceding Noon, of the Moon's Distance from the Meridian, when the Sun, at Noon, is then upon it.

NOW, while the Sun (by the Earth's revolving Easterly on it's Axis) is apparently departing from the Meridian to the Westward, as

It must always be noted in finding the Parallaxic $\angle$, P. 232, by D. Cowper's improved compendious Method, 2 Col. whether the Meridian $\angle$ is greater or less than 90°.

Ecliptic Place.

Meridian $\angle$ according to Table { $\odot$, Ω , Υ , ω , $\cap$, A , } Forenoon Observation. { Supplement $\angle$ } Afternoon Observation.
Supplement of Tab. $\angle$ { V , X , X , V , X , II , } Tabular $\angle$
The Meridian $\angle$ above and below the Ecliptic, and one Side and other of the Meridian, being Supplements of each other to 180° respectively.
many.

Astronomical QUESTIONS practically answered.

many Hours as the Moon wanted of coming to the Meridian, she is advancing towards it, and would arrive there, in the Manner of a Star, had the Moon no Easterly Motion, in the Time of the Sun's passing those Hours to the Westward; deducting the contrary Motion or Advance of the Sun's Right Ascension in that Time, to the Eastward (p. 218) all the while meeting the Star, and accelerating it's Approach to the same Meridian.—But by the Moon having a more considerable Easterly Motion of Right Ascension in 24 Hours, than the Sun's Easterly Motion of Right Ascension in that Time, therefore the Moon's Approach or coming to the Meridian is thereby retarded or prolonged; by as much as the Increase of the Moon's Right Ascension in Time Easterly, from Noon, exceeds that of the Sun's in Time the same Way, during her Approach, from Noon, to the Meridian. For while the Sun is going that Time to the Westward, he spends the Difference of Increase of the two Right Ascensions more than the Moon's first Distance in Time, from the Meridian, while the Moon exactly arrives there.—Hence we have this universal and correct RULE, for determining the Time of Passage of the Moon, and of all the PLANETS through the MERIDIAN.

As 24 Hours is to the Difference in Time, of the Moon's Increase of Right Ascension above the Sun's (or Increase of the Sun's Right Ascension above a Planet's) in that Time, so is the Distance of the Moon's from the Sun's Right Ascension (or Planet's Right Ascension from the Sun's Right Ascension) in Time, at the Noon preceding the Time of the Southing required, to the proportional Increase of the Moon's Right Ascension above that of the Sun's, in Time, (or Sun's Right Ascension greater or less than a Planet's in Time) to be added to the Moon's or Planet's Distance from the Meridian at Noon (but subtracted from a Planet's Distance at Noon, if the Sun is the Body of swifter Motion) for the true or accurate Time of Southing of the Moon or Planet sought.

N. B. When the Sun's Right Ascension is greater than the Moon or Planet's Right Ascension, 24 Hours must be added to the latter, that Subtraction may be made of the first. And when the Planet moves Retrograde, the Sum of the Diurnal Right Ascensions of the Sun or Planet, in Time, are to be taken instead of the Difference, in the foregoing Proportions.

M. De la Caille's Eph. Paris	Long.	Lat. S.	Decl. S.
Ex. I. 1762, July 10 Noon	5° 1'	4° 33'	13° 55'
11 Noon	16 55	3 59	8 50
Difference in 24 Hours	11 54	0 34	5 5
	Increase.	Decrease.	Decrease.
☾'s Right Ascension without Lat. 5° 1' by p. 27.	22 ^h 27 ^m 27 ^s	Days 10	☉ R. A. 7 ^h 18 ^m 14 ^s
		11	7 22 19
		July.	
Equation for 4° 33' S. Lat. (Table follg) = 1° 42' 40" +	Time 6 50	Incr. 4 5	
		in 24 Hours.	
☾'s R. A. 10 July, Noon	22 34 17	22 34 17 fr.	

☾'s R. A. Diff. from Sun's R. A. 1762 July 10, Noon 15 16 3
Equation of the ☾'s Southing fr. Noon as below. + 25 15

☾'s R. A. without Latitude to 16° 55' by Table follg	23 ^h 11 ^m 51 ^s	15 41 18
Equation for 3° 59' S. Lat. (Table follg) = 1° 33' +	Time 6 12	☾'s Southing 10th July past Noon, accurately req. at Paris.
		or 15 ^h 32 ^m at Greenwich.
☾'s R. A. 11 July, Noon 1762, 23 18 3		
☾'s Diurnal R. Ascension between 10 and 11 July	43 ^m 46 ^s	Increase.
Sun's Diurnal Right Ascension	4 5	Increase.

Difference of Diurnal Right Ascension . . . 39 41 Increase.

Now say, As 24^h : 39^m 41^s :: 15^h 16^m : 25^m 15^s to be added to Difference ☾'s Right Ascension and ☉'s Right Ascension, as above, for the correct Time of ☾'s Southing required.

N. B. The ☾'s Right Ascension in Time answering to the Longitude and Latitude of the Moon, are more readily found from computed Tables of Right Ascension in Time, answering to Longitude and Latitude at Sight.

Long.	Lat. S.	R. A. ☾
Against { 5° 1' 4° 33' }	Stand { 22 ^h 34 ^m }	
{ 16 55 3 59 }	{ 23 18 }	

Diurnal R. A. ☾	0 44
Diurnal R. A. Sun.	0 4

Diff. Diurnal R. A. . . . 0 40 nearly,

Or the Moon's Right Ascension in Degrees is found by a SINGLE PROPORTION from the DATA; thus,

As Cosine Declination (N. or S.)	☾ 3° 55' co.	Logarithms.
To Cos. Longitude from next Equinox.		0.0129389
tial Point here preceding	☾ 24 59	9.9573346
So Cos. Latitude from the Ecliptic	4 33	9.9986292

To Cos. R. A. from the same Equinox.		
Point, ☾ preceding	21 25 28	9.9689027
In Time = 1 ^h 25 ^m 42 ^s R. A. from	24 0 0	[☾ preceding.]

☾'s Right Ascension from ☾ . . . 22 34 18 following.
Correctly and nearly as above.

Again, As Cos. Declination	8° 50' 0" co.	0.0051819
To Cos. Long. from ☾ preceding	13 5 0	9.9885776
So Cosine Lat. from the Ecliptic	3 59 0	9.9989496

To Cos. R. A. from ☾ preceding	10 28 10	9.9927091
In Time = 1 ^h 41 ^m 53 ^s R. A. from	24 0 0	[☾ preceding.]

☾'s Right Ascension from ☾ . . . 23 18 7 following.
Correctly, and nearly as above.

Thus the Moon's Southing is directly and accurately determined (from the Longitude and Lat. of the Moon given for two succeeding Days) by two easy Operations, besides a small concluding Proportion; without the Trouble of an estimate Southing, as taught by most Authors.

LEADBETTER in his *Astronomy* (p. 264, 265, 1 vol.) by an estimate Method, finds the estimate Southing from the Moon's Age, when from the Places and Latitudes of the Moon for two successive Days, finds the Moon's Place to the estimate Southing by Proportion, and then the Right Ascension of Moon thereto; then he finds the Sun's Place to that estimate Time, whose Difference in Time, he makes the approximate Time of Southing for the true Time, or the Distance the Sun is past the Meridian in Time, when the Moon is upon it. He then adds the daily Difference of the Moon from the Sun, in Time for the Moon's Southing on the successive Day, which is nearly true but not accurate.

BUT if the Moon and Sun's Right Ascension are daily given in Ephemerides, the Difference of those Ascensions, or Moon's Right Ascension—Sun's Right Ascension will always be the Moon's preceding Distance in Time from the Meridian, when the Sun is upon it. And if you add the proportional Distance in Time, thereto, for the Excess of the Increase of the Moon's Right Ascension above the Sun's Right Ascension in that Time, you will accurately and shortly find the Moon's Southing, for any Day; by Help of the Connoissance des Temps, or other Ephemerides, wherein the Moon and Sun's daily Right Ascensions are given. And, if you continually add the daily Increase of the Moon's Right Ascension above the Sun's Right Ascension in Time, to the Time of the former Day's Southing of the Moon, you will correctly (and not nearly) have the Time, later and later, of the Moon's Southing, on successive Days throughout the Year. And the same Rule holds good for finding the Times of the Planets Southing, successively, by adding or subtracting to and from the Times of the former Day's Southing, the Diurnal Difference of Right Ascension of the two Bodies, according as the Sun (apparently) is the slower or faster moving Body than that whose Time of Southing is sought.

Astronomical QUESTIONS practically answered.

Mr. CHARLES BRENT (who has done Service to Computation) in his *Astronomy*, p. 333, finds the near Time of the Moon's Southing, by taking the D 's Right Ascension from Tables, as if she had no Latitude; and then to find a still nearer Time, he makes Equation for her real Latitude according to the Sign the Moon's Place is then in. As his Method is near and easy, we here give it as plain as possible with a small Improvement.

In the foregoing EXAMPLE.

D 's R. A. without Lat. to $5^{\circ} 1'$ 1762, July 10 }
at Noon } = $22^{\text{h}} 27^{\text{m}} 27^{\text{s}}$
Sun's R. A. the same Time, had from Tab. p. 217 . 7 18 24

Near Time of Moon's Southing 1762, July 10 at Paris 15 9 13

RULE. Add to which near Southing $2^{\text{m}} 2^{\text{s}}$ for every Hour thereof being about the Dif. of Motion in Time of D à Sun in an Hour at a Mean . . } 0 30 48

Nearer Time D 's Southing on that Day, past Noon . 15 40 1

EQUATION for Latitude D according to Mr. Brent.

RULE. As greatest Lat. 5° is to 8^{m} Dif. of Time, }
so present Lat. $4^{\circ} 33'$ S. to $7^{\text{m}} 16^{\text{s}}$ + . . . } + 7 16

D 's Southing at Paris 1762, July 10 past, Noon . 15 47 17

RULE for Sign of EQUATION. { 11 Morning $3^{\text{h}} 37^{\text{m}} 7^{\text{s}}$ reqd.

D 's Place. Equation { D 's Place. Equation

Lat. S. . . . } + { Lat. N. . . . } +

Lat. N. . . . } - { Lat. S. . . . } -

N. B. The above easy Method varies in the Instance given but 7^{m} too much, from the true Time of the Moon's Southing; and therefore, if used with the first Equation only (omitting the last) will be worth Regard; coming frequently very near the Truth, at the same Time it is very ready for Practice.

The other more easy, but less near, Method of finding the near Time of the Moon's Southing, is by multiplying the Days of her Age by 8 Tenths, for the Hours and decimal Parts of an Hour of her Southing past Noon, which are turned into Minutes, mentally, by multiplying those Decimal Parts by 6.

The Moon's Age, 1762, July 10 . . . 19 Days.
8

Near Time of D 's Southing on that Day past Noon $15^{\text{h}} 12^{\text{m}}$

Or, July 11^d $3^{\text{h}} 12^{\text{m}}$ Morn.

Thus, having shewn how to find the Southing of the Moon, and all the Planets, accurately, as well as nearly, it follows next,

To find the MOON'S, and also a Planet's Rising and Setting accurately, as well as nearly, for any PLACE, whose Latitude is given.

EXAMPLE II. To find the Time of the Moon's Rising and Setting at Greenwich Observatory, 1762, July 10, the Time of the Moon's Southing at Paris Observatory, having before been determined $15^{\text{h}} 41^{\text{m}} 18^{\text{s}}$, on that Day past Noon?

FIRST, To find the Time of Southing under a different Meridian? Time Greenwich is West of Paris } $9^{\text{m}} 20^{\text{s}}$

As 24^{h} is to the Time of later Southing of the Moon on the following than on the former Day, in the same Place (or to the Difference in Time of R. A. of $\odot$ and D in 24 Hours) so is the Difference in Time between the first Meridian, and any other Meridian lying East or West therefrom, to the Time of sooner or later Southing there (respectively) than at the first Place; reckoning by Time in the second Place.

D Souths Paris, 1762, La Caille's Ephem. } 10 July P.M. $15^{\text{h}} 41^{\text{m}}$
11 } 16 22

Dif. 41

As 24^{h} to 41^{m} so $9^{\text{m}} 20^{\text{s}}$ to 16^{s} Moon souths later at Greenwich in Greenwich Time, than at Paris, in Paris Time.

For, As 24^{h} to $24^{\text{h}} 41^{\text{m}}$ D 's later Southing at Paris, so $9^{\text{m}} 20^{\text{s}}$ Greenwich is West of Paris in Time, to $9^{\text{m}} 36^{\text{s}}$ the Moon is going from Paris to Greenwich Meridian.

1762, July 10 D souths at Paris in Paris Time, } $15^{\text{h}} 41^{\text{m}}$
P. M. }

Time spent by the D going from Paris to Greenwich Meridian Westward of Paris . . . } 9 36

D souths at Greenwich in Paris Time . . . } $15^{\text{h}} 50^{\text{m}} 36^{\text{s}}$
P. M. }

Deduct $9^{\text{m}} 20^{\text{s}}$, Time sooner at Greenwich than at Paris . . . } 9 20

1762 July 10, D souths at Greenwich, in Greenwich Time . . . } $15^{\text{h}} 41^{\text{m}} 16^{\text{s}}$
P. M. as before.

TO REDUCE EPHEMERIDES from one MERIDIAN to ANOTHER. And First from Paris to Greenwich MERIDIAN.

UNIVERSALLY. As 24^{h} is to $9^{\text{m}} 20^{\text{s}}$ so is the following Day's Difference of the Moon's Southing, of the Sun's Place, Sun's Declination, Sun's Right Ascens. Moon's Place, Moon's Latitude and Declination, or daily Difference of Longitude, Latitude, Declination, and also daily Difference of Southing of the Planets, or Stars, to the respective proportional Difference to be added to, or subtracted from, the first Day's Times, Places, &c. at Noon, according as the daily Difference increase or decrease, for the Times, Places, &c. at Greenwich; correspondent to the Times, Places, &c. at Paris, Eastward of Greenwich—The same Manner of proportioning holds for any other Difference of Meridian, instead of $9^{\text{m}} 20^{\text{s}}$ Westward of Paris—OBSERVATORY. And if the Meridian is Eastward, instead of Westward of Paris, or of any other first Meridian, then in reducing Paris Places, Times of Southing, &c. to those for any other fixed Meridian to correspond with a Second Meridian (Eastward of Greenwich, or the first Meridian) you must subtract the Proportional Diff. when the daily Diff. increases, and add it when the daily Diff. decreases, to the first Quantities, respectively, for the Quantities sought, corresponding with the Second Meridian.

And, because $\frac{9^{\text{m}} 20^{\text{s}}}{24 \text{ Hours}} = .0065$ nearly. Therefore, DECIMAL-

LY, Any daily Difference in Paris Ephemerides, being multiplied by 65 and four Figures cut off to the Right Hand in the Product will give the proportional Difference to be added to or subtracted from the first Day's Quantity, for the Quantity corresponding with Greenwich Meridian.

D 's daily Diff. Southing 1762 July 10 and 11, 41^{m}
 $65 \times 41 =$

205

246

2665 = 16^{s} as before.

Add $15^{\text{h}} 41^{\text{m}} 0$

Time of Southing at Greenwich. 15 41 16

Correspond. to Time of Southing at Paris 15 41 0

The Southing of Planets and of the Moon in all Places nearly at the same Time of the Day or Night in each Place, is owing to the Sun Southing in all Places at 12 at Noon, or the same Time exactly in each Place; from which Luminary, (measuring the apparent Time in all Places) the Celestial Bodies vary their Distances, but a small Matter in R. A. in 24 Hours, or diurnal Circuit; the Moon varying her Distance in R. A. in that Time the most.

HENCE, the Moon varies her Time of Southing at Greenwich and at Antigua in the West-Indies, but the Sixth Part of her diurnal Difference later, at the latter than at the former Place. For, 4 Hours Diff. of Meridians is equal the one Sixth of 24 Hours, in which Time the diurnal Difference of Southing is performed. Which Variation of D 's Southing, to an Hour Diurnal Difference, will be but 10 Minutes of Time reckoned later at Antigua than at Greenwich.

Therefore,

Astronomical QUESTIONS practically answered.

Therefore, it follows, that in seeking the Moon's near Time of Rising and Setting, in all Places, you may use the Time of her South- ing in any Place to compute the Rising and Setting by. And, if you would compute the Rising and Setting of the Planets, in any Place, the Time of their South- ing (and especially of the Stars) may be taken, with still greater Nearness to Truth, from Ephemerides to serve for the Time of South- ing in any one Place. The Difference of the Swift-ness of Planets, (Mercury's different South- ing in 24 Hours, or diurnal Circuit, being seldom above 22 Minutes) amount not to above 11 Minutes of Time Difference of South- ing in half the Globe Dis- tance between two Places; and sometimes not to a Quarter of that Quantity.

But, whatever this Difference is, whether of the Moon, or any of the Planet's South- ing, it is readily had from correctly computed Ephe- merides; by which the South- ing for any Place is correctly found, and thence the Rising and Setting of the Moon and Planets are had for that Place, as follows.

	D's Pla.	Lat. S.	D Souths.	D Dec. S.
1762 Ju. 10	5X1	4 33	15 ^h 41 ^m	13° 55'
11	16 55	3 59	16 22	8 50
LaCaille's Eph.				
Differences	Incr. 11 54	De. 34	Incr. 41'	5° 5' Decr.
Reduced to Greenwich, July 10			13° 53'	Noon.
July 10.	15 ^h 41 ^m 16 ^s		10° 34'	Noon.
	D Souths P.M.		D Decl. S.	Greenwich.
			correspondent	

For, as 24^h : 5° 5' Decr. Decl. :: 15^h 41^m P. M. : 3° 19' Decr. Declination.

By Tab. p. 202, 203, Semiduration Arc, corre-
pondent to 10° 34' D S. Decl. and Latitude }
Greenwich 51° 23½' } ± 5^h 9^m 0^s
D's South- ing } 15 41 16

Near Time D's Rising 10 July . . . 10 32 16
Setting . P. M. 20 50 16

D's Declinations to } Rising { 11° 39' } by Proportion.
those mean Times } Setting { 9 28 }

For, as 24^h : 5° 5' Decr. Decl. :: 10^h 32^m : 2° 14'—

13 53

11 39

20 50 : 4° 25'—

13 53

9 28

By Tab. p. 202, 203, Semiduration Arc, corre-
pondent to 11° 39' D S. Decl. Latitude Green-
wich } — 5^h 4^m 0^s
D's South- ing } 15 41 16

Time D's Rising at Greenwich, 10 July 1762, al-
lowing for Refraction; but not Parallax . . . } 10 37 16
past Noon.

By Tab. p. 202, 203, Semiduration Arc, corre-
pondent to 9° 28' D S. Declination Lat. Green-
wich } + 5^h 18^m 0^s
D's South- ing } 15 41 16

Time D's Setting, 10 July 1762, allowing } P. M. 20 59 16
for Refraction; but not Parallax . . . } or 11^d 8 59 16
Morning.

OTHERWISE, To find the true RISING of the PLANETS.

Subtract the Sun's R. A. from the oblique Ascension of the Planet, for the Latitude of the Place, and the Remainder being more than six Hours, deduct 6 Hours therefrom, for the Time of Rising. But, being less than 6 Hours, add 6 Hours thereto, for the Time of Rising.

1762 July 10, oblique Ascension, by Tables for
Latitude, London, to D's Pl. 5X1, Lat. S. 4° }
33' at Noon } 23^h 47^m
R. A. Noon, same Day, De la Caille's Ephem. } — 7 18

16 29
— 6 0

Near Time of D's Rising 1762, July 10 . . . 10 29
past Noon.

Correspondent to which near Time of Rising,
D's Place 10° X 13' Latitude S. 4° 18' oblique Ascen. 23^h 54^m
R. A. } 7 20

16 34
— 6 0

Moon's true Time of Rising 1762 July 10, at }
Greenwich, Refraction here being not allowed for } 10 34
Past Noon, nearly as before.

OF THE MOON'S PARALLAX AND REFRACTION.

THE Horizontal Parallax is not considered in the foregoing Rules for finding the Rising and Setting of the Moon, and therefore those Rules find the Times of the true, and not apparent, Rising and Setting of the Moon. The foregoing Tables of Semidurnal Arcs are computed by allowing for Refraction of the Celestial Bodies; and therefore they very well serve to find the apparent (not the true) Rising and Setting of the Planets.

But, the Moon having great horizontal Parallax (and considerable Parallax in Altitude) that Parallax must always be considered, together with the Refraction, for determining the Times of the ap- parent Rising and Setting of the Sun and Moon; and thence to find whether an Eclipse or Occultation will be visible, or not, in a given Place; which is one considerable Use of what is called visible Time.

The visible Altitudes of the Moon, or other celestial Bodies, must be truly augmented by Parallax, and decreased by Refraction; because those Bodies are apparently depressed by Parallax, and raised by Refrac- tion; which contrary Appearances, or Effects, must therefore be al- ways truly allowed for, (as above) by a contrary Augmentation and Decrease of their apparent, for determining their true Altitudes; and thence the true Time correspondent. So, likewise, on the contrary, the visible or apparent Rising and Setting of the Moon must be de- termined from her given real Place, Parallax and Refraction, on a given Day, by reducing her real Altitude given, to her apparent Ri- sing, by Means of her horizontal Parallax depressing, and Refraction raising, her Body, at the same Time.

OF VISIBLE AND TRUE TIME.

HENCE, when the Moon's Parallax in Altitude, and Refraction are equal or alike, there is no Difference between visible and true Time, viz. between the Time correspondent to the visible Altitude of the Sun, Moon or Star, and the Time correspondent to their real Altitudes, which Times differ, when the Parallax and Refraction of the Sun, Moon, or Star, vary from Equality. This Distinction between vi- sible and true Time therefore respects the visible and true Places or Alti- tudes of the Sun, Moon or Star, in the Heavens; which true Time (only respecting the true Alte, not the visible Time respecting the visible Altitude) is equatable to Mean or equal Time.

THEREFORE the Refractions of the celestial Bodies must, for the foregoing Reasons, be always taken into Consideration, and the Paral- laxes also, where they are considerable, for discovering the true Time; which otherwise, in the Result of Computation, will be the visible Time; considerably different, in some Cases, from that which is true. Which true Time, as it refers to the Sun's Motion, will be what is

otherwise

Astronomical QUESTIONS practically answered.

otherwise called the *apparent Time* by the Sun, to be reduced, (as aforesaid) to *equal or Clock Time*, by the *Equation of Time Table*, p. 19.

The *Time of the visible Conjunction of the Sun and Moon*, seen from the *Earth's Surface*, always differs from the *Time of the true Conjunction*, as would appear from the *Earth's Center*: Except the *Conjunction* happens in the *Nonagesimal Degree of the Ecliptic*, when the *true and visible Time* are the same.

Hence it follows, that if the *true Conjunction* be in the *Oriental Quadrant of the Ecliptic*, or between the *Nonagesimal Degree and Eastern Horizon*, the *Moon's Place* is then apparently advanced *Easterly* by her *Parallax of Longitude*, and the *Time of the visible Conjunction* will then be before the *Time of the true Conjunction*. But, if the *true Conjunction* fall in *Occidental Quadrant of the Ecliptic*, or between the *Nonagesimal Degree of the Ecliptic and Western Horizon*, the *Moon's Place* will then be apparently advanced *Westerly*, by as much as the *Moon's Parallax* is in *Longitude*; and consequently the *Time of the visible Conjunction* will follow the *Time of the true Conjunction*. Which *Distinctions of visible, true and also equal Time*, are of *Importance* to be known in the *Computation of Eclipses*, as well as in other *Parts of Astronomy*.

The foregoing Account of the *Distinctions of Time*, is an Answer to one of our *Correspondents* and *Subscribers* writing thus. "In all the *Methods of finding the Hour of the Night by the Stars*, no mention is made whether the *Equation of Time* is to be taken into Consideration or not, or whether the *Time found* is *apparent or equal*; some People being at a *Loss* in this *Affair*."

N. B. As the *fixed Stars* have no sensible *Parallax*, only their *Refraction* is to be considered in finding the *Hour of the Night*, *apparent solar Time*, by their *apparent Altitude*.

To find the *TRUE TIME* of the *MOON's apparent Rising and Setting* on *July 10th 1762*, at *Greenwich*?

FIRST, for the *MOON's apparent RISING*.

1762 July 10 { Without Regard to Parallax. { Moon's Decl. { Hor. Par. { De la Caille's Ephem.
Rising 10^h 37^m { 11^o 39' { 55' 40" {
Setting 20 59 { 9 28 { 55 36 {
Moon below the visible Horizon at her apparent { 33' 0" - {
Rising by her Horizontal Refraction. {
Moon's Horizontal Parallax then 55 40 + }

Moon's true Altitude above the true Horizon at her { 22 40 + {
apparent Rising {
Now, there are given,

Moon's Zenith Dist. 89^o 37' 0" Side op. $\angle$ required Logarithms.
Comp. Pole's Height 38 31 30 { Sides inclu. } L.S.co. 0.2056123
D's Dist. fr. N. Pole 101 39 0 { $\angle$ required } L.S.co. 0.0090402
Sum 229 47 30
From . . . Half 114 53 45

Take Comp. { 1 rem. 76 22 15 Log. S. 9.9875952
Pole's Height {
Take D's D. { 2 rem. 13 14 45 Log. S. 9.3600812
from N. Pole. {
Sum 19.5623289
Double 37 10 11 L. S. . . . Half 9.7811644
Double 74 20 22 $\angle$ Apparent Rising fr. the Merid.

Now, As 360^o is to the Interval of Time betwixt two next Passages of the Moon, or any Planet, through the same Meridian, so is the angular Distance of that Celestial Body, from the same Meridian, either at apparent Rising, Setting, or any Altitude, to its Distance in Time from the Meridian.—To be subtracted from, or added to the Time of it's Southing, according as it is advancing towards, or has passed the Meridian, for the true Time of Rising, Setting, or at any given Altitude.

1762 July 10 Moon Souths { 15^h 41^m { De la Caille's Eph. at Paris.
11 { 16 22 {

Difference 24^h + 41^m Difference also at Greenwich.
Now, As 360^o to 24^h 41^m so 74^o 20' 22" to 5^h 5^m 49^s—
1762 July 10, D Souths accury. at Greenwich, P.M. 15 41 16

1762 July 10, Moon apparently rises at Green- } 10 35 27
wich P. M. accurately }

N. B. The Difference of the Time of the Moon or a Planet's Southing in each 24 Hours, is only the Difference of Right Ascensions of the Sun, and of that Planet, in Time. The celestial Body Southing sooner or later each Day, according as it's Motion is slower or faster than the Sun's Motion in Right Ascension.

NEXT, for the *MOON's apparent SETTING*.

The Difference of Horizontal Parallax at Moon's Rising and Setting being here but 4" less at Setting; the same Parallax may be used at Setting as for Rising, viz. 55' 40"; whence, D's true Altitude at apparent Setting, is 22' 40"; or 23' very nearly, as before.

GIVEN,
Moon's Zenith Dist. 89^o 37' 0" Side op. $\angle$ required Logarithms.
Comp. Pole's Height 38 31 30 { Sides inc. } L.S.co. 0.2056123
D's Dist. fr. N. Pole 99 28 0 { $\angle$ req'd. } L.S.co. 0.0059551
Sum . 227 36 30
From . . Half . 113 48 15

Take Comp. { 1 rem. 75 16 45 Log. S. 9.9855052
Pole's Height {
Take D's D. { 2 rem. 14 20 15 : Log. S. 9.3938087
fr. No. Pole. {
Sum 19.5908813
Double 77 16 18 $\angle$ at apparent Setting fr. Merid. Half 9.7954406

Now, As 360^o to 24^h 41^m so 77^o 16' 18" to 5^h 17^m 53^s
1762 July 10, Moon Souths at Greenwich ac- } 15 41 16
curately P. M. }

1762 July 10, Moon apparently sets at Greenwich } 20 59 9
accurately P. M. [required.]

Finding the accurate Time of the Moon's apparent Rising and Setting, and also the accurate Time of the Night, (which is performed in the same Manner) to her apparent Altitude, has not been shewn before by any Author that we have seen.

WHENEVER you would determine the Time by the Moon's or a Planet's Altitude, in any Latitude, to a Degree of Accuracy, the Declination of the celestial Body at that Time, must be first determined as near as possible, by an Estimate of the Time; allowing for Refraction, and also Parallax; or the Time will differ considerably from the Truth; and answer no End in determining the Longitude from that Observation. For which Purpose, to facilitate the Operation, correct EPHEMERIDES of the Sun and Moon's Places, Declinations, Right Ascensions, &c. with their Differences, for each Day of the Year, should be carried to Sea, if Promoters of Literature and Science in this Nation can be found, to encourage the correct Computation of such useful Ephemerides, as are encouraged in other Nations, for public Utility. Or, if our ASTRONOMICAL PROFESSORS, who have made a Secret of their Art, can (like those in other Nations) be prevailed upon to undertake the PUBLIC WORK.

If you would know the Time when the Moon's LOWER LIMB ascends the visible Horizon, you must add the D's horizontal Semidiameter to her true Altitude, at apparent Rising, and proceed as before, for her Center.

Astronomical QUESTIONS practically answered.

Or, if you would know the Time when the Moon's upper Limb descends the visible Horizon, you must subtract her horizontal Semidiameter from the true Altitude at apparent Setting, (the same as at Rising) then proceed as in the foregoing Operation.

One near Method to find the Time of true Rising of the Moon, Planet, or Star, for a given Latitude, is by taking the Sun's oblique Ascension in Time, (for that Latitude) at the estimate Time of the Moon's Planet or Star's Rising, from their respective oblique Ascension, in Time, at Rising; adding the Time of Sun-Rise to the Remainder for the near Time of their respective Rising that Day.

This Method is rendered correcter by repeating the Operation to the near Time found; and also by finding the Moon, Planet, or Star's oblique Ascension to the Place and Latitude at that Time.

And the near Time to the Moon, a Planet, or Star's true Setting, is also found by taking the oblique Descension of the Sun from their respective oblique Descension, at the estimate Time of Setting; adding the Time of Sun-setting, on that Day, to the Remainder, for the Time of the respective Setting.

And, if you would be more correct, repeat the Operation to the last found Time: Observing to take the oblique Descension of the Moon, Star, or Planet, correspondent to its Latitude at the mean Time of Setting.

Another near Method of performing the same is as follows.

From the oblique Descension in Time of the Moon, Planet, or Star, subtract the Right Ascension of the Sun, in Time, at the estimate Time of Setting; and, if the Remainder exceeds 6 Hours, subtract 6 Hours therefrom; but if the Remainder be less than 6 Hours, add 6 Hours thereto; the Difference, or Sum, respectively, will be the Time of the Moon, Planet, or Star's Setting, required.

N. B. The oblique Descensions are taken out of Tables of oblique Ascensions, by entering them with the opposite Sign and Degree of the Planet's Places, and contrary Name of Latitude to those Places. Also contrary Sign and Degree of the Sun's Place, without Latitude.

EXAMPLE of the REDUCTION of PLACES from one MERIDIAN to ANOTHER.

	Sun's Pl. Noon.	D's Pl. Noon.	Decl. D.	Moon's Southings.	
1762 July 10	18° 53'	5° 31'	13° 55'	15 ^h 41 ^m	Paris
11	19 0	16 55	8 50	16 22	Observatory.
Diurnal Diff.	+ 57	+ 11 54	- 5 5	+ 41 ^m	

Dis. by, 0065 } Prop. Pt. Prop. Pt. Prop. Pt. Prop. Pt. }
+ 22" + 4' 38" - 1' 59" + 16" }
Reduced to Greenwich by nearest Minutes.

1762 July 10	18° 53'	5° 31'	13° 53'	15 ^h 41 ^m	
11	19 0	17 0	8 48	16 22	Greenwich.
					Diurnal Differences as for Paris above.

N. B. The foregoing is a Specimen of the Reduction of Places from any Meridian to any other Meridian Westward of the first Place, according to Proportion of daily Difference, or Increase or Decrease of those Places from any Day to the next Day forward, for 360° Westward. And therefore, the Proportion of Difference for easterly Longitude from the first Meridian, must be regulated according to the Increase or Decrease from any Day to a former Day, for 360° Longitude Eastward: The Difference for easterly Longitude being contrary to that for westerly Longitude: Increase for Decrease, and Decrease for Increase.

And the Sun, Moon, or a Planet's Declination, or right Ascension, for any Place, to the Westward or Eastward of Greenwich, is determined in the same Manner, by proportioning for the Difference of Meridians Westward or Eastward, according to the Difference of the Declination, or right Ascension, (or any other Requisite) for a Day following or preceding, answering to a diurnal Circuit of the revolving Body.

EXAMPLE II. REDUCTION from PARIS to ANTIGUA-MERIDIAN: 4^h 9^m 20^s Westward of Paris-Observatory.

	☉'s R.A. Noon.	Sun's Dec. N.	D's Decl. S. Noon.	
1762 July 10	7 18 14	22 15 13	13 S 55	Paris.
11	7 22 19	22 7 24	8 50	De la Caille.
Diurnal Diff.	+ 4 5	- 7 49	- 5 5	
As 24 ^h to 4 ^h 9 ^m 20 ^s , so the respective Difs to	Prop. Pt. + 42 ^s	Prop. Pt. - 1' 21"	Prop. Pt. - 52' 47"	Antigua.
	Reduced to Antigua.			

1731 Xr.

	h m s	° ' "	° ' "	
1762 July 10	7 18 56	22 13 52	13 2 13	S.
11	7 23 1	22 6 3	7 57 13	S.
	Diurnal Differences come out, as before, for Paris Observatory.			

EXAMPLE. To find the near Time of the Moon's true Setting, from the foregoing RULES.

	D's Lat. Noon.	
1762 July 10	4° 33'	
11	3 59	Greenwich.
	34	Correspondent.
1762 July 10	Estimate Time. Setting 21 ^h 0 ^m	Sun's Pl. 18° 52' D's Pl. 15° 31' D's Lat. 4° S 4'
		Opposite Places.
		18° 52' 15° 31' 4° N 4'

Opposite Places.	From Tables for 51° 32' Lat. London.	
D 15° 31' Lat 4° N. 4'	Oblique Descension.	h m
☉ 18° 52'	Oblique Descension.	10 25
		- 21 26

N. B. The same Method, with oblique Ascension, serves to find the Moon's true Rising. See p. 240. Rem. h. 12 59 Sun sets 10th July 8 8

Near Time D's Setting 1762, July 10, past Noon, 21 7
Correct Time of the Moon's apparent Setting . . h. 20 59
Difference . . h. 0 8

EXAMPLE II.

	h m
D 15° 31' Lat. 4° N. 4'	Oblique Descension. 10 25
☉ 18° 52'	R. A. 19 22

N. B. When above 12^h remain, 12^h must be added to the Result . . 15 3
- 6 0

Near Time of the Moon's true Setting 1762, on the } 9 3
11th Day of July in the Forenoon, as before, }
+ 12 0

Viz. 10th July, past Noon : . 21 3

The same Methods serve to find the near Time of the true Rising and Setting (not the apparent) of all the Planets; and when you would find the true Time of the true Rising and Setting of the Moon, or any of the Planets, by either Method, you must repeat the Operation to the last found near Time: Determining the Sun's Place, and Planet's Places, and Latitude to that Time, and the oblique and right Ascensions from

thence.

Astronomical QUESTIONS practically answered.

thence. By which Means a *correct* or true Time will result after the *New Operation*.

The *estimate* Time of the Moon's Rising and Setting, or for any of the Planets, may be always had by respectively taking the right Ascension of the Sun from the oblique Ascension and Descension in Time, of the Moon, or Planet, (by Tables for the given Latitude of the Place) and adding or subtracting 6 Hours according as the Remainder be less or more than 6 Hours.

But, *universally*, the *estimate* Time of Rising and Setting of the Moon, or any of the Planets or Stars, is readily determined by subtracting and adding the semi-diurnal Arc, in Time, correspondent to the Latitude of the Place, and the Moon or Planet's Declination at Noon, (but better at Southings) from and to the Time of Southing of the Moon or Planet for that Day, (of 24 Hours) on which the Rising or Setting are required.

See Tables from p. 200 to 203, for this Purpose.

Mr. CHARLES BRENT, (in his *compendious Astronomer*, p. 334.) gives us his Method of finding the near Time of the Moon's true Rising and Setting; after his finding her Southing, as before exhibited, which Method (with our small Improvement) is as follows.

FIRST for the MOON's near DECLINATION to her true Time of SOUTHING.

THIS is found as the Sun's Declination, from the Declination Table, (p. 23.) answerable to the Moon's Place in the Ecliptic, considered without Latitude, first subtracting from and adding to her Place at Southing, $\frac{1}{2}$ a Degree, for every Hour before and after her Southing.

Next, for every Degree of the Moon's Latitude to allow 5' through all the Signs of the $\mathcal{D}$'s Place, in the Ecliptic, according to this RULE.

Multiply the Degrees of *present* Latitude by 5', And say, as 90° are to that Product, so are the Number of Degrees of the Moon's Place, from Cancer, (betwixt Aries and Libra) or Number of Degrees from Capricorn, (betwixt Libra and Aries) to a proportional Quantity, to be always subtracted from the Moon's Latitude, for the Quantity to be added to or subtracted from her Declination, found according to her Place in the Ecliptic, without Latitude, for the Moon's near Declination at Southing, required?

Whether this last found Quantity is to be added to, or subtracted from, the Ecliptic-Declination, you have this RULE.

Lat. } same } Name } with Ec- } + } Quantity
 $\mathcal{D}$. } different } } cliptic De- } - } found.
 } } } clination. } } For Moon's
 } } } } } Declination.

BUT, When the Latitude is greater than the Ecliptic-Declination, that must be deducted from the Latitude, for the Moon's Declination, required; which will then be of the same Name with the Moon's Latitude.

See Tab. following, for the same Purpose.

From New to Full Moon, it is called the Moon's Setting.

From Full to New 'tis called the Moon's Rising.

The Hours of Moon-Light being this Way determined.

You will always know, by the Sun and Moon's Places, approaching to, or having passed, a Conjunction, or Opposition, whether it is the Rising or Setting to be found.

For the Moon moving *swifter* than the Sun, if you subtract his from her Place, in the Ecliptic, the Distance, in Signs, will shew you their *Aspect*, past, or to which they are approaching, according to the following Order of Signs, increasing their Distance from New to Full; but diminishing their Distance, according to preceding Order of Signs, from Full to New Moon. See FIGURE for the Aspects, p. 236.

RULE. With the Moon's Declination, at Southing, just found, enter the Table of Semi-duration Arcs to a given Latitude, (see p. 200

to 203.) and take out the Time of that Arc, which subtract from and add to the true Southing, for the *estimate* Times of Rising and Setting; to which *estimate* Times of Rising and Setting find the Moon's respective Declinations, with which Declinations take out the semi-diurnal Arcs for Rising and Setting. To each of which Arcs add as many Times 2^m 2^s, (about the mean Motion in Time of Moon to Sun per Hour) as there are Hours, and subtract and add the Sum thereof respectively, from and to the Time of Southing, and you will have the near and respective Time of the Moon's Rising and Setting, required.

N. B. The Time of the semi-duration Arc for the Moon's Rising-Declination, with 2^m 2^s for every Hour thereof, added, and Time of the Moon's Semi-duration Arc for her Setting Declination, with 2^m 2^s for every Hour thereof added, will be the Moon's Time of Continuance above the Horizon.

When the Moon's Setting Declination is nearly equal to and of the same Name with the Sun's Setting Declination, the semi-diurnal Arc of the Sun, (the Time of Sun-Setting) and semi-duration Arc of the Moon, being then nearly equal, and to which the Time of Southing and semi-duration Equation to the Moon's Setting Declination, are to be added, for the Moon's Time of Setting, therefore, in that Case (and no other) the Sum of the Time of Southing and of the Semi Duration Equation, will be the Hours of Moonlight after Sun-Setting, from New to Full Moon.

In other Cases, between New and Full Moon, you must subtract the Time of Sun-Setting from the Time of Moon-Setting, and the Remainder will be the Hours of Moon-Light after Sun-Setting. So likewise between Full and New Moon, if you take the Time of Moon-Rising from Sun-Rising, you will have the Hours of Moon-Light, before Sun-Rising.

EXAMPLE to the last near Method.

	Sun's Place.	Moon's Place.	Moon's Lat. S.	Moon Souths.	
1762 July 10	18° 23'	5° 46'	4° 33'	15 ^h 41 ^m	Greenwich Observatory.
Noon . . 11	19 0	17 0	3 59	16 22	
Diurnal Diff's.	+ 57	+ 11 54	- 34	+ 41	

To find the Moon's Declination at Noon, correspondent?

	July	Moon's Place.	Ecl. Declin. S.	$\mathcal{D}$ near Decl. S. correspondent.	
Hence, by Tab. Page 23, and Rule foregoing.	10	5° 46'	9° 39' 19"	13° 55' 43"	Noon.
	11	17 0	5 8 28	8 50 22	
		Diurnal Dif. . .		5 5 21	

FOR,

As 90° to 22^h 7 (= 4° 33' $\times$ 5') so 65° fr. }
 1/2 to } - 16 36 Equation.
 4° 33' $\mathcal{D}$'s Lat. S.

$\mathcal{D}$'s Lat. and Ecliptic Decl. of same Name + 4 16 24 } equad Quant.
 Ecliptic Declination 9 39 19 } S.

Sum . . Moon's near Declination, }
 July 10, Noon } 13° 55' 43" S.

AND,
 As 90° to 20' (= 4° $\times$ 5') so 77° fr. 1/2 to - 17 6 Equation.
 3 59 0 $\mathcal{D}$'s Lat. S.

$\mathcal{D}$'s Lat. and Ecliptic Decl. same Name + 3 41 54 } Equad Quant.
 Ecliptic Declination 5 8 28 } S.

Sum . . Moon's near Declination, }
 July 11, Noon } 8 50 22 S.

As 24^h to 5° 5' diurnal Decr. $\mathcal{D}$'s Decl. so 15 41 to 3 19
 16 22 to 3 28

Astronomical QUESTIONS practically answered.

	Decl. at Southings.	Mean sem. D. Arc.		Declinat. corresp. S.	Sem. D. Arc.
Hence July 10	10° 37'	5 ^h 9 ^m 4 ^s			
Moon souths	15 41				
Estimate Rising July 10	10 ^h 32 ^m			11° 42'	5 ^h 3 ^m
Estimate Setting	20 50			8 35	5 ^h 19
Being near the true.		Past Noon.			
For, As 24 ^h to 5° 5' diur. Decr. D's Decl. so 10 32 to 2 14					
As 1 ^h to 2 ^m 2 ^s at a Mean, so 5 3 to 10 20 50 to 4 25					
Sem. D. Arc Rising 5 3 Sem. D. Arc Setting 5 19					
Correction + 10 Correction + 11 at a Mean.					
Corrected Sem. D. Arc 5 13 Cor'd. S. D. Arc 5 30					
D Souths 15 41 D Souths 15 41					
Near Time D's true } 10 28 Setting 21 11 past Noon.					
Rising }					
But, 24 ^h to 41 ^m true Incr. so 5 3 to 8 36 } true Correc ⁿ to S.					
5 19 to 9 4 } Duration Arcs.					
Hence, 5 3 5 19					
+ 9 9 true Correc ⁿ .					
True S. D. Arc Rising 5 12 Tr. S. D. Arc Set. 5 28					
D Souths 15 41 D Souths 15 41					
Correct Time Moon's } 10 29 Correct T. D's } 21 9					
true Rising }					
true Setting }					

From the foregoing repeated Operations, the Moon's semi-duration Arc is seen to vary considerably by a Change of her Declination; making the semi-duration Arc longer or shorter, and consequently her Rising or Setting sooner or later, according as her South Declination, at Rising and Setting, happens to be less and more, and more and less, than at Southings; or her North Declination at Rising and Setting, more and less, or less and more, than her Declination at Southings.

And as there is no direct Way of coming at the Moon's Declination at Rising and Setting, but by repeated Trials and Operations, (however accurate the semi-duration Arc may be found from a given, or supposed Declination of the Moon, at Rising or Setting, by Trigonometry, (or Tables for that Purpose) this last near Method, by a mean Declination at Southings, will discover the near Rising and Setting of the Moon for all Latitudes, the best and soonest of any Method whatsoever: neglecting the Correction of the Rising and Setting semi-duration Arcs, which is often compensated for in the mean Declination, taken at the Moon's Southings.

Hence, from what is done above,

	D's Declin.	Mean S.
	at Southings.	Dur. Arc.
1762 July 11	5° 22'	5 ^h 37 ^m 4 ^s
Moon souths	16 22	La Caille's Eph.

Rises 10 45 } nearly.
Sets 21 59 }

Since the Moon's true semi-duration Arcs (from Rising to Southings and from Southings to Setting) depend on the accurate Determination of the Moon's Declination, at Rising, Southings, and Setting, (doubtfully obtained within few Minutes of Truth, because of the Declination's very variable Motion) there is not a more difficult Question, in practical Astronomy, to rightly resolve, than the foregoing, Of finding the correct Time of the Moon's true or apparent Rising and Setting?

And therefore we have been the more particular and full in explaining this useful Question on that Account. For the true Declination at apparent Rising and Setting must be previously had, as well as at the true Rising and Setting, before the angular Distance from the Meridian, respecting the apparent and real semi-duration Arcs, can be rightly found. And hence to determine in any Year, On what Day of the Month in that Year, in a given Latitude, the greatest Difference between the Moon's Rising and Southings, and Southings and Setting, will happen, or when the Rising and Setting semi-duration Arcs will be most unequal, (or most equal if you please) would be a still more difficult Question, unknown in practical Astronomy: And was never yet resolved by the famous Author of the Miscellaneous Tracts and Physical Astronomy.

XLVII. QUESTION. To find the Point of the Ascendent, or Degree of the Ecliptic, ascending the Eastern Horizon, from the Latitude of the Place and Time of the Day, or Night, (past the Meridian) being given?

EXAMPLE. In the Year 1764, October 14, at 10 at Night, what Point of the Ecliptic, at London, will be then rising, or ascending the Eastern Horizon?

Sun's Place at that Time, London	22° 1' 19"
Sun's R. Ascension then, correspondent	200 21 26
Time from Noon 10 Hours in Degrees, add	150 0 0
Sum, R. Ascension of Mid-Heaven	350 21 26
Add	90 0 0
Sum (rejecting 360°) the oblique Ascension of the Ascendent	80 21 26
	past Aries.

NOW, by PROPORTION.

As Radius	S. 90° 0' 0"	Logarithms.
To Cos. Obl. Ascension ascendent fr. ☿ or ♀ (here ☿)	80 21 26	9.2240274
So Cotan. Lat. Place	51 32 0	9.9000865
To Cotan. of 4th Arc	82 25 10	9.1241139
Here oblique Ascension ascendent is nearest following	+23 28 30	

5th Arc . . . 105 53 40, or 74° 6' 20".
N. B. Here the 5th Arc exceeds 90°.

Oblique Ascension of Ascend. } nearer to ☿ than ♀ following or preceding. } Sum of the } For a }
Ecliptic } 5th }
Obliquity } Arc.

Oblique Ascension of Ascend. } nearer to ♀ than ☿ following or preceding. } Difference of } For a }
Ecliptic } 5th }
Obliquity } Arc.

The Projection shows the Truth. This Answer is the Solution to the 7th Case of oblique spherical Triangles, where 2 Angles and an included Side are given to find the Side opposite to one of them.

Viz. { $\angle =$ Co. Latitude } included oblique } to find Dist. Eclip-
Given { $\angle$ Obliquity Ecliptic } Ascension from } tic from same Equ.
☿ or ♀. } Point.

Or { $\angle =$ Lat. + 90° } included oblique } to find Dist. Eclip-
Given { $\angle$ Obliquity Ecliptic } Ascension from } tic from same Equ.
☿ or ♀. } Point.

NOW,

Astronomical QUESTIONS practically answered.

NOW, As Cos. 5th Arc	74° 6' 20" co.	0.5624619
To Cos. 4th Arc	82 25 10	9.1203106
So Tan. Obl. Ascen. Ascend.	80 21 26	10.7697163

Logarithms.

Here 5th Arc { To Tan. of the Distance of the Ascendant from the same or contrary equ. Point from which the oblique Ascension is reckoned . . . } 70 34 4 10.4524888

WHEN the 5th Arc is less than 90°, then you reckon the Distance of the Ascendant on the Ecliptic, from the same equinoctial Point that the oblique Ascension of the Ascendant is reckoned from; but if the 5th Arc is greater than 90°, you reckon the Distance of the Ascendant from the contrary equinoctial Point, to that from whence the oblique Ascension of the Ascendant is reckoned. For when the 5th Arc is greater than 90°, the Distance of the Ascendant from the same equinoctial Point from whence the oblique Ascension of the Ascendant is reckoned, will be also greater than 90°; and the Sines, Tangents, &c. of Arcs, and of their Supplements, being the same, therefore the Answer will be from the contrary equinoctial Point, to give the Distance from the same equinoctial Point greater than 90°.

Hence when the oblique Ascension is from an equinoctial Point, preceding, the Answer will be from the contrary equinoctial Point, following.

By the foregoing necessary Distinctions the Place of the Ascendant of the Ecliptic, or Sun's Place at Rising, correspondent to the oblique Ascension or Descension, of a Rising or Setting Star, is truly determined.

If the Point of the Ascendant at the Horizon (or Arc of the Ecliptic between the Meridian and Horizon) were required when the Beginning of Aries, or Libra, culminates, or is on the Meridian above the Horizon, the foregoing Proportions, for determining generally the Point of the Ascendant, from ♈ or ♎, fail, and the following Proportions, in these two Instances, take Place.

EXAMPLE II. To find the Place of the Ascendant, at London, when the Beginning of Aries is on the Meridian above the Horizon?

Libra is then on the inferior Part of the Meridian below the Horizon, at London, and consequently are given the Part of the Meridian below the Horizon, 38° 28', = Complement of Latitude, the ∠ of Ecliptic and Meridian 66° 31' 30", and ∠ of Meridian and Horizon 90°, to find the Arc of Ecliptic betwixt the Meridian and Horizon. — Hence, by solving a right angled spherical Triangle, of one Leg, and an adjacent Angle to find the Hypotenuse, being the inferior Arc of the Ecliptic betwixt the Horizon and Meridian, we have this

PROPORTION.		
As Cosine Meridian ∠	66° 31' 30"	9.6002636
To Radius S.	90 0 0	10.0000000
So Tang. of Comp. Latitude	38 28 0	9.9000865
To Tan. Ecliptic between the Horizon } and ∠ preceding.	63 22 16	10.2998229
From ♈ to ∠	180 0 0	

Place of Ascendant . . 116 37 44 fr. ♈ following viz, ∠ 26 37 44, required.

EXAMPLE III. To find the Place of the Ascendant at London, when the Beginning of Libra is on the Meridian, above the Horizon?

Aries is then on the inferior Part of the Meridian, below the Horizon, at London, and consequently are given, the Part of the Meridian above the Horizon 38° 28' = the Complement of Latitude, 66° 31' 30" the ∠ of the Ecliptic and Meridian; and 90° the ∠

of the Meridian and Horizon; to find the superior Arc of the Ecliptic between the Meridian and Horizon?

By solving a right-angled spherical Triangle of one Leg and an adjacent ∠ given, to find the Hypotenuse, being the superior Arc of the Ecliptic betwixt the Meridian and Horizon, we have, as before, this

PROPORTION.		
As Cosine Meridian Angle	66° 31' 30"	9.6002636
To Radius : S.	90 0 0	10.0000000
So Tangent of Comp. Lat.	38 28 0	9.9000865
To Tang. Ecliptic betw. the Meridian } (from ∠) and Horizon; following	63 22 16	10.2998229

Place of Ascendant following, [or ♈ 3° 22' 16", required.

N. B. When ∠ or ♎ culminates, the Arc of the Ecliptic betwixt the Meridian and Eastern or Western Horizon is evidently a Quadrant; the Meridian being then the solstitial Colure. Therefore, when ∠ culminates, the Place of the Ascendant will be the Beginning of ♈; and that of the Descendant the Beginning of ♏. And when ♎ culminates, the Place of the Ascendant will be the Beginning of ♏, and that of the Descendant the Beginning of ♈: the Places of Ascendant and Descendant being always opposite; and the Arcs of the Ecliptic between the Meridian and Eastern and Western Horizon are the Supplements of each other to 180°.

The Places of the Ascendant, at the Beginning of each Sign culminating, is nearly as follows, by the foregoing Rules and Proportions, or this RULE.

As S. ∠ of Ecliptic and Meridian to S. Altitude of Mid-Heaven, so Radius to S. Arc between Meridian and Horizon.

Mid-Heaven.	Place of Ascendant.	Arc of Ecliptic betw. Merid. and East. Horiz.
Begin. ♈	∠ 26° 38'	116° 38'
♈	♈ 16 30	106 30
♉	♉ 7 21	97 21
♊	♊ 0 0	90 0
♋	♋ 22 38	82 38
♌	♌ 13 30	73 30
♍	♍ 3 22	63 22
♎	♎ 25 22	55 15
♏	♏ 27 10	57 10
♐	♐ 0 0	90 0
♑	♑ 2 50	122 50
♒	♒ 4 45	124 45

HENCE, it would be a proper Question to determine the Place of the Ascendant, and what Sign and Degree of the Ecliptic culminates, when the greatest and least Distances of the Ecliptic are included betwixt the Meridian and Horizon.

N. B. The above eastern Arcs will be a Guide in determining others that are ambiguous.

The Place of the Ascendant may be otherwise found, from the ∠ of the Ecliptic and Meridian, Mid-Heaven, Zenith Distance, correspondent to right Ascension of Mid-Heaven, and 90° from Zenith to the Ascendant, given, in an oblique spherical Triangle, to find the Arc of Ecliptic betwixt the Meridian and Horizon?

XLVIII. QUESTION. To find the Day of the Year in a given Latitude, when a Star, or Planet, will rise with the Sun, (or rises cosmically) from the oblique Ascension of that Star or Planet being given?

AS every Star, or Planet, rises with the Point of the Ecliptic, having the same oblique Ascension with that Star, or Planet, it follows, that when the Sun is in that Point of the Ecliptic, the same Star, or Planet, will rise with the Sun, or cosmically.

EXAMPLE.

Astronomical QUESTIONS practically answered.

EXAMPLE. Required the Day when the bright Star in the Eagle rises with the Sun, at London, 1760?

By P. 210, N^o. 126.

Corrected				
to 1760	Declination N. bright Star in Eagle	8	15	8
Begin.	Right Ascension of that Star	294	46	3
Ascensional Dif. in Degs (by Quest. VI.) subtract		10	31	6
Oblique Ascension * from γ , following		284	14	57
From γ , preceding		75	45	3

NOW, by the preceding Question XLVII.

As Radius	S. 90° 0' 0"	Logarithms.	10.0000000
To Cos. obl. Ascen. * fr. next equin. }			
Point (here γ preceding)	75 45 3		9.3911808
So Cotang. Lat. Place, London	51 32 0		9.9000865
To Cotang. of 4th Arc	78 56 7		9.2912673
Obliq. Ascen. $\odot$ or * nearest }			
γ preceding.	add + 23 28 30	Eclip. Obliq.	

NOW, As Cos.	5th Arc	102 24 37	
Or	77 35 23 co.	0.6677423	
To Cos.	4th Arc	78 56 7	9.2831151
So Tang. obl. Ascen. * or $\odot$	75 45 3	10.5952481	

To Tang. of Pl. of Ascend. fr. Δ follows	74 7 32	10.5461055	
---	---------	------------	--

Or contrary eq. Pt. fr. whence obl. Ascen. is reckoned; because 5th Arc is more than 90°.

Which Place answers to the 6th of December, 1760, N. S. when the bright Star in the Eagle and the Sun rise together; and will continue to rise together nearly on that Day for a Number of Years, at London, or most Places near the Middle of England.

XLIX. QUESTION. To determine the Day of the Year, 1760, at London, when the aforesaid bright Star in the Eagle sets at the Time the Sun rises, (or sets cosmically) from the oblique Descension of that Star, given?

Right Ascension of the Star as before	294 46 3	
Ascensional Dif. in Degrees as before, add	10 31 6	
Sum, Oblique Descension * fr. γ following, given	305 17 9	
Add	180 0 0	
Oblique Ascension of the Ascendant, then (rejecting 360°) from γ following	125 17 9	
Supplement — obl. Ascen. from Δ preceding	54 42 51	

Now, proceed to find the Place of the Ascendant at Sun-Rising, when the Star sets, from the oblique Ascension of the Ascendant given, as before.

As Radius	S. 90° 0' 0"	10.0000000	
To Cos. obl. Asc. fr. next eq. Pt. Δ pre.	54 42 51	9.7616691	
So Cotan. Lat. London	51 32 0	9.9000865	
To Cotan. 4th Arc	65 20 52	9.6617556	
Ob. Ascen. of $\odot$ nearest Δ preced. sub.	23 28 30		
As Cos.	5th Arc	41 52 22 co.	0.1280602
To Cos.	4th Arc	65 20 52	9.6202500
So Tan. obl. Ascen ^a Ascend ^t fr. Δ preced.	54 42 51	10.1501703	

To Tan. of Pl. of the Ascend. fr. Δ preced.	38 21 49	9.8984805	
Or same Equ. Point from whence obl. Ascen. Ascend ^t is reckoned; be-			
cause the 5th Arc is less than 90°.	141° 38' 11" from γ followg.		
	viz. Ω 21° 38' 11"		
	Sun's Place, required.		

Which Place answers to the 14th Aug. 1760, N. S. when the Sun rises, as the bright Star in the Eagle sets, at London, and will continue to rise and set together nearly on that Day for a Number of Years, at London, or most Places near the Middle of England.

L. QUESTION. To find the Day, in the Year 1760, at London, when the aforesaid bright Star in the Eagle, will rise, as the Sun sets, (or rise achronically) from the oblique Ascension of that Star being given?

THE Answer to this Question is contained in the Answer to the former Question, being only to find the Point of the Descendant, or Setting-Point of the Ecliptic, opposite to the Rising-Point thereof, or Pl. of the Ascendant, found to the cosmical Rising of a Star with the Sun. Which Place of the Descendant will be the Sun's Place, at Setting, answerable to the achronical Rising of that Star.

Thus, the Place of Ascendant at cosmical Rising of the bright Star in the Eagle is found $\uparrow$ 14° 7' 32"

Whence the opposite Place of Sun-Setting, at achronical Rising of the same Star $\downarrow$ II 14° 7' 32"

Which Place is answerable to June the 4th, 1760, N. S. when the bright Star in the Eagle rises achronically, required.

This Star will continue to rise achronically, on that Day, at most Places near the Middle of England, for a Number of Years.

LI. QUESTION. To find the Day, in the Year 1760, at London, when the aforesaid bright Star in the Eagle will set as the Sun sets, (or set achronically) from the oblique Descension of that Star being given?

THE Answer to this Question is contained in the Answer to XLIX. Question.

The same Point of the Ecliptic, or Place of Descendant, that sets with the aforesaid Star, is the Sun's Place, at Setting, when that Star sets (achronically) as the Sun sets. Therefore, the opposite Point of the Ecliptic, to that of the Ascendant, when the same Star sets cosmically, or at Sun-Rising, will evidently be the Place of the Descendant, or Sun's Place in the Ecliptic, at his Setting when the Star sets, achronically.

Thus, the Place of the Ascendant, or of Sun-Rising, at cosmical Setting of the bright Star in the Eagle, being found Ω 21° 38' 11"

The opposite Place to Sun-Rising, at cosmical Setting, is the Place of the Ecliptic at achronical Setting, of the same Star, viz. ω 21° 38' 11"

Which Place answers to February 9, 1760, N. S. when the bright Star in the Eagle sets achronically, required.

This Star will continue to set achronically on that Day, in most Places near the Middle of England, for a Number of Years.

A TABLE of the DAYS in the YEAR, *New Style*, when the *cosmical* and *acronical* Rising and Setting of 42 eminent FIXED STARS happen at London.

N. B. Cosmical Rising and Setting of a Star is at Sun-Rise; Acronical Rising and Setting of a Star is at Sun-set.		Cosmical.		Acronical.	
No.	STARS NAMES.	Rising at Sun Rise.	Setting at Sun Rise.	Rising at Sun Set.	Setting at Sun Set.
T H E					
1	First in the Wing of Pegasus, <i>Marchab</i>	January 12	September 24	July 17	March 22
2	Right Shoulder of <i>Æquarius</i>	20	August 29	24	February 24
3	Extreme in the Wing of Pegasus	February 8	October 5	16	April 2
4	Last in the Tail of the Goat	18	August 11	24	February 7
5	Bright * in the Head of <i>Aries</i>	March 15	November 4	September 18	May 2
6	In the preceding Horn of the <i>Ram</i> , or γ	21	October 30	23	April 27
7	In <i>Whale's</i> Tail	April 23	September 15	October 27	March 13
8	Brightest in the <i>Pleiades</i>	May 4	November 23	November 6	May 21
9	Brightest in the <i>Whale's</i> Mouth, <i>Mencar</i>	31	October 26	30	April 23
10	Northern Horn of the <i>Bull</i> , or Foot of <i>Auriga</i>	26	November 21	26	May 20
11	<i>Fomabaunt</i>	21	August 4	21	January 31
12	Northern Eye of the <i>Bull</i>	31	November 24	December 2	May 23
13	Bright * in the Belly of the <i>Whale</i>	June 3	September 23	5	March 21
14	Southern Eye of the <i>Bull</i> , <i>Aldebaran</i>	8	November 24	10	May 23
15	Southern Horn of the <i>Bull</i>	16	December 11	17	June 10
16	<i>Castor</i>	20	February 14	21	August 17
17	<i>Pollux</i>	July 3	January 30	January 1	3
18	Middle Star in <i>Orion's</i> Belt	14	November 17	10	May 16
19	Little Dog, <i>Procyon</i>	30	December 18	25	June 17
20	Bright * in the <i>Hare's</i> Thigh	August 2	October 20	29	April 26
21	Bright * in the great Dog's Mouth, <i>Syrius</i>	11	November 16	February 6	May 14
22	<i>Lion's</i> Heart	20	February 24	15	August 29
23	<i>Lion's</i> Back	21	April 27	16	October 30
24	<i>Hydra's</i> Heart	31	December 28	26	June 28
25	Bright * in the <i>Lion's</i> Tail, <i>Deneb</i>	September 2	April 26	29	October 29
26	<i>Vindematrix</i>	21	May 18	March 18	November 20
27	<i>Arcturus</i>	26	June 23	23	December 22
28	Bright * in the <i>Virgin's</i> Girdle	30	April 30	27	November 2
29	Bright * in the <i>Crown</i>	October 9	July 20	28	January 17
30	<i>Virgin's</i> Spike	15	April 7	April 12	October 11
31	Right Shoulder of <i>Hercules</i>	17	July 23	14	January 20
32	Left Shoulder of <i>Hercules</i>	21	August 5	20	February 1
33	Head of <i>Hercules</i>	November 1	July 22	30	17
34	<i>Swan's</i> Bill	10	September 1	May 8	27
35	Right Shoulder of <i>Ophiuchus</i> , <i>Serpentarius</i>	16	July 18	14	January 16
36	Inferior Wing of the <i>Swan</i>	16	September 26	15	March 16
37	<i>Vulture's</i> Tail	22	August 12	20	February 8
38	Right Knee of <i>Ophiuchus</i> , or of <i>Serpentarius</i>	27	June 18	21	December 18
39	<i>Scorpion's</i> Heart	December 3	May 16	June 2	November 17
40	Bright * in the <i>Eagle</i>	6	August 14	4	February 9
41	Bright * in the Thigh of Pegasus, <i>Scheat</i>	22	October 7	22	April 3
42	Bright * in the Head of <i>Andromeda</i>	January 5	22	July 7	January 5

EXPLANATION.

The SUN, in his annual Motion through the Ecliptic, gets to the Eastward of the Fixed Stars; therefore, In North Latitudes of Places, Stars having South Declination,

- 1st { From Cosmical Setting (at Sun Rise) Stars set be- } rise after } are visible, above the Horizon, from their Rising to Setting during
 { To Acronical Rising (at Sun Set) fore Sun Rise } Sun-Set } that Interval.

EXAMPLE I. Scorpion's Heart visible from its Rising to Setting from May 16, to June 2, following.

- 2d { From Acronical Setting (at Sun Set) Stars set be- } rise after } are invisible, above the Horizon, from their Rising to Setting,
 contrarily { To Cosmical Rising (at Sun Rise) fore Sun Set } Sun Rise } during that Interval.

EXAMPLE II. Scorpion's Heart invisible from its Rising to Setting, in the Day from November 17, to December 3, following.

- 3d { From Cosmical Rising (at Sun Rise,) Stars rise be- } set after } are constantly visible, above the Horizon, at some Time of the
 { To Acronical Setting (at Sun Set) fore Sun Rise } Sun Set } Night.

EXAMPLE III. Scorpion's Heart visible sometime between Rising and Setting, from December 3, to November 17, following.
 In North Latitudes of Places, Stars having North Declination.

- 4th { From Acronical Rising (at Sun Set) Stars rise be- } set after } are visible till they approach so near the Sun, as to be hid in his
 { To Cosmical Rising (at Sun Rise) fore Sun Set } Sun Rise } Rays.

EXAMPLE IV. The Pleiades visible from November 6, to May 4, following, or till they approach so near the Sun as to be hid in his Rays; which when a Star recovers from, and is seen in the Morning, before Sun Rise, is called its *Heliacal* Rising; as the near Approach of a Star to the Sun, in the Evening, so as to become inconspicuous, is called, the *Heliacal* Setting of that Star—The Time to be determined as follows.

Astronomical QUESTIONS practically answered.

LII. QUESTION. To find the Time of a Star's heliacal Rising, from the Latitude of the Place and Star's oblique Ascension, with the Sun's Depression below the Horizon, at the Time of that Star's heliacal Rising (after its cosmical Rising) being given?

OBSERVATION shews that the *smallest Stars* are invisible while the upper Hemisphere receive any of the Sun's Rays; and therefore are first seen in the Evening, at the End of Evening Twilight, and last seen in the Morning, at the Beginning of Morning Twilight, when the Sun, at either of those Times, is 18° below the Horizon. And Stars of different Magnitudes are observed to appear first and last in the Evening and Morning, respectively, when the Sun's Depression, below the Horizon, is as follows.

Stars of the	Magnitude appear first and last in the Evening and Morning, when the Sun is	Deg.	Below the Horizon.
1	2	12	
2	3	13	
3	4	14	
4	5	15	
5	6	16	
6		17	

EXAMPLE. The Month is required in the Year 1760, when the bright Star in the Eagle (of the 2d Magnitude) will rise heliacally, at London?

N. B. It will rise heliacally, when the Sun has described an Arch of the Ecliptic, further to the Eastward, from his Place at cosmical Rising, correspondent to 13° of his Depression below the Horizon, at the Time of the same Star's Rising. Consequently heliacal is after cosmical Rising, and is when the same Star first begins to appear, after being hid under the Sun's Rays.

Oblique Ascension of the bright Star in the Eagle,	0	1	4
before found	284	14	57
Subtract 90°	90	0	0
Remain R. A. of Mid-Heaven at the Star's Rising	194	14	57
R. A. from following	14	14	57
Longitude of Mid-Heaven, correspondent fr. following (by Quest. 34.)	15	28	32
Meridian $\angle$ correspondent (by Tab. p. 23, or 35 Quest.)	67	17	18
Declination of Mid-Heaven S. correspondent (by Tab. p. 23, or 36 Quest.)	6	6	6
Mid-Heaven Zenith Distance, London	57	38	6

Now there are given, in an oblique spherical Triangle, the Meridian $\angle$, Mid-Heaven Zenith Distance, as above, and the Sun's Zenith-Distance 103° , at the Time of the Star's heliacal Rising, to find the Arch of the Ecliptic between the Meridian and the Sun, at that Time?

Viz. Two Sides, (the Mid-Heaven Zenith Distance and Sun's Zenith Distance) with an Angle (the Meridian Angle) opposite to one of them given (the Sun's Zenith-Distance) to find the third Side, intercepted between the Meridian and Sun.

PROPORTION.

As Radius	S. 90°	Logarithms.
To Tan. Mid-Heaven Zen. Distance	57 38 6	10.0000000
So Cos. Meridian $\angle$	67 17 18	10.1980733
To Tang. 4th Arc	31 21 1	9.5866930
		9.7847663

As Cos. Mid-Heaven Zenith Distance	57 38 6 co.	Logarithms.
To Cos. Sun's Zenith Distance	77 0 0	0.2713939
Therefore, Supplement of the Answer must be taken	for 103 0 0	9.3520880
So Cos. 4th Arc	31 21 1	9.9314592
To Cosine of	68 58 8	9.5549411
Its Supplement, 5th Arc	111 1 52	
Dif. between 4th and 5th Arcs	79 40 51	Ecliptic Arc, [required.

Because the Meridian Angle, next to which the Perpendicular falls, is above 90° fr. Place Mid-Heaven, or less than 90° , When Mid-Heaven Longitude is in $\{ \begin{matrix} \text{♈, ♉, ♊, ♋, ♌, ♍, ♎, ♏, ♐, ♑, ♒, ♓} \end{matrix} \}$ * Dif. Sum } 4th and 5th Arcs for the Ecliptic Arc between the Merid. and Sun.

* For East of the Meridian, or Rising; but Sum and Dif. for West of it, or Setting.

When Supplement of an Arc above 90° is used (as 77° for 103°) the Supplement of the Answer (but not otherwise) must be taken, as $111^\circ 1' 52''$ for $68^\circ 58' 8''$, above.

OTHERWISE, to find the Ecliptic Arc below the Horizon, the superior Arc being found, in the Longitude of the Ascendant at the Star's cosmical Rising.

As Sine Sun's Zenith Distance	77°	co.	0.0112761
To S. Meridian Angle	67 17 18		9.9649473
So S. Mid-Heaven Zenith Distance	57 38 6		9.9266793
To Sine $\angle$ of the Vertical and Ecliptic, below the Horizon	53 5 50		9.9029027
Or Parallax $\angle$, at the Ecliptic.			

NOW, As Radius	90°	10.0000000
To Cotan. Depression	13° 0' 0"	10.6366359
So Cos. Paral. $\angle$ below Horiz.	53 5 50	9.7784834
To Cotan. of an Arc of Eclip. below the Horizon, correspondent	21 1 52	10.4151193
Add $\angle$ 14 7 32		Sun's Pl. at *'s [cosmical Rif.
5 9 24		Sun's Pl. at *'s [heliacal Rising.

Or, As S. $\angle$ of the Horizon and Ecliptic, To Sine of the Sun's Depression, at heliacal Rising or Setting of a Star, or Planet, So is Radius, To Sine of Ecliptic Arc below the Horizon, correspondent.

The foregoing Place of the Sun answers to the 26th of December, N. S. 1760, when the bright Star in the Eagle rises heliacally, required; the Sun being then 13° below the Horizon. Which Star will continue to rise heliacally on that Day, for a Number of Years, after having rose cosmically, at the same Time with the Sun, (and consequently invisible) on the 6th Day of the same Month. So that on the 26th of December the bright Star in the Eagle begins to appear, after being hid in the Sun's Rays, from and before December 6, at rising with the Sun. After the heliacal Rising December 26, it continues to appear in the Morning before Sun-Rise.

Astronomical QUESTIONS practically answered.

Longitude of the Ascendant foregoing, } 0 ' "
 at the Star's Rising, p. 246. . . } 254 7 32 fr. ♀ following.
 Longitude of Mid-Heaven, then . . . } 195 28 32 fr. ♀ following.

Ecliptic Arc from Meridian to Horizon, at Star's Rising . . . } 58 39 0
 Add Ecliptic Arc fr. Horizon to the Sun depressed last found . . . } 21 1 52

Ecliptic Arc from the Meridian to the Sun, as first found . . . } 79 40 52
 Add Mid-Heaven Longitude . . . } 195 28 32

To ♄, 9 Signs, Subtract . . . } 275 9 24 fr. ♀ following.
 270

Remains Sun's Place . . . } ♄ 5 9 24
 at the Star's heliacal Rising, as before.

LIII. QUESTION. To find the Time of a Star's heliacal Setting, from the Latitude of the Place, and Star's oblique Descension with the Sun's Depression below the Horizon, at the Time of that Star's heliacal Setting (before its achronical Setting) being given?

EXAMPLE. The Month Day is required, in 1760, when the bright Star in the Eagle will set heliacally, at London?

N. B. It will set heliacally, when the Sun has yet to describe an Arc of the Ecliptic further to the Eastward, till he comes to his Place, at the Star's achronical Setting from his present Place of 13° below the Horizon, or so much more to the Westward, than he will be at the Star's achronical Setting: And therefore the heliacal is before the Time of the achronical Setting.

The first Method, in the former Question, determines the heliacal Rising and Setting at once, independent of the cosmical Rising, and achronical Setting, which Method is repeated in determining the heliacal Setting, at once, as follows.

Oblique Descension of the bright Star in the Eagle, before found, p. 246. at Setting, achronically . . . } 305 17 9
 Add 90° . . . } 90 0 0
 Sum (rejecting 360° R. A. of Mid-Heaven from ♀ following) . . . } 35 17 9
 Longitude of Mid-Heaven correspondent from ♀ following (by Quest. 34) . . . } 37 39 4
 Meridian ∠ correspondent (by Tab. p. 23, or 35 Quest.) . . . } 71 1 28
 Declination Mid-Heaven N. correspondent (by Tab. p. 23, or 36 Quest.) . . . } 14 5 0
 Mid-Heaven Zenith Dist. London . . . } 37 27 0
 Angle of Ecliptic and Horizon . . . } 54 53 55

By PROPORTION.

As Radius 90°

To Tang. Mid-Heaven Zenith Dist. . . 37 27 0
 So Cof. Merid. Angle correspondent . . 71 1 28

To Tang. 4th Arc 13 59 7

Logarithms.

10.0000000

9.8841956

9.5121035

9.3962991

And, As Cof. Mid-H. Zen. Dist. . . 37 27 0 co. } 0.1002428
 To Cof. Sun's Zen. Dist. at 13° Depref. 77 0 0 } 9.3520880
 Supplement of Arcs must be taken . . 103 0 0
 So Cof. 4th Arc 13 59 7 } 9.9869320

To Cofine 74 2 26 } 9.4392628
 Supplement 5th Arc 105 57 34
 Dif. between 4th and 5th Arc . . . 91 58 27 } Ecliptic Arc [required.]

Place } ♄, ♏, ♍, ♌, ♍, ♎, ♏, ♐, ♑ } Sum. } 4th and 5th Arcs
 Mid-Heaven. } ♄, ♏, ♍, ♌, ♍, ♎, ♏, ♐, ♑ } Dif. } for West Side Meridian or Setting.

Or to find the Arch of the Ecliptic below the Horizon, intercepted between it and the Sun's Place.

As S. ∠ of Horizon and Ecliptic . . 54° 53' 55" co. } 0.0871746
 To S. Sun's Depression 13 0 0 } 9.3520880
 So Radius S. 90 0 0 } 10.0000000
 To S. Ecliptic Arc below the Horizon 15 57 34 } 9.4392626

To find the ARC OF THE ECLIPTIC between the Meridian and Horizon?

As S. ∠ of Ecliptic and Horizon . . 54° 53' 55" co. } 0.0871746
 To Cof. Zen. Dist. of Mid-Heaven, or } 37 27 0 } 9.8997572
 S. Alt. of Mid-Heaven Sine 90° } 10.0000000
 So Radius Sine 90°

To S. Arc of Ecliptic between Meridian and Horizon . . . } 76 0 53 } 9.9869318
 Arc below the Horizon before found 15 57 34

Sum, Ecliptic Arc betwixt the Merid. and Sun, at the ♄'s heliacal Setting } 91 58 27, as before.
 Deduct fr. Long. Mid-Heaven, [adding 360°.

Long. of Mid-Heaven 37 39 4 from ♀.
 Remain, Long. Sun, at the Star's Setting heliacally . . . } 305 40 37
 To ♄ 10° Deduct 300 0 0 } Sun's Place ♄ 5 40 37
 Or Sun's Place at achronical Setting, before found, p. 246. . . } ♄ 21 38 11
 Arch of the Ecliptic below the Horizon, at 13° Sun's Depression, before achronical Setting . . sub. } 15 57 34

Sun's Place Remains in ♄ 5 40 37 as before.

Which Place answers to January 25, 1760, when the bright Star in the Eagle sets heliacally, in most Places about the Middle of England, and will continue to set heliacally, on that Day, for a Number of Years. So that December 26 (the Time of this Star's heliacally Rising) is the Time of its first Appearance, in the Morning, after being hid under the Sun's Rays in the Day Time; which Star, from that Time, continues to appear in the Morning, before Sun-Rise; also appears in the Evening, after Sun-set, till the 25th of January following, when it last appears in the Evening, setting heliacally. It then comes again into the Sun's Rays, as he further advances to the Eastward, setting achronically with him, on February 9th, and then sets before him.

Astronomical QUESTIONS practically answered.

The Times of heliacal Rising and Setting of any other fixed Star, may be determined by the foregoing general Method, which equally serves for finding the heliacal Rising and Setting of the Planets, by observing the Sun's Depressions at those Times, as follows.

h	can be seen	Deg.	
21	at Rising	11	
3	or Setting	10	below
3	when the	11	the
8	Sun is	5	Horizon.
D		10	
		5	

Likewise ♀ may be seen in the Day. ♀ was seen in the Day, April 22, 1715. D may be also seen in the Day.

The chief Use of the foregoing cosmical, achronical, and heliacal Rising and Setting of the Stars and Planets (otherwise called the Poetical Rising and Setting thereof by the Antients) is to direct us when and where to look for them, to make our Observations. — But a PAIR of Mr. Senex's GLOBES, sold by Mr. MARTIN, in Fleetstreet, London, according to his late Improvements, will satisfy us in these, and many other useful Enquiries with most Ease and Expedition; where great Accuracy is not required. Which GLOBES no Astronomer ought to be without, as they will serve to confirm the Truth, or discover an Error, in his Computations.

OF THE MOON.

TO find the Moon's right Ascension at her given Time of Southing, on any given Day, the right Ascension of the Sun, in Time, at the preceding Noon of that Day being also given?

RULE. The Time of the Moon's Southing added to right Ascension of the Sun, in Time, at the Time of her Southing, will be then equal to the right Ascension of the Mid-heaven, or right Ascension of the Moon then upon the Meridian.

Consequently, by Transposition. If from right Ascension of the Moon, in Time, on a given Day you deduct the right Ascension of the Sun, in Time, (both at the Moon's Southing) the Remainder will be the exact Time of the Moon's Southing, past Noon, on that Day.

Hence the Reason of first seeking the Moon's estimate Southing, by the common Methods of finding her true Southing.

EXAMPLE. To find the Moon's right Ascension at the Time of her Southing, October 1, 1760?

1760 Oct. 1	R. A. ☉.	12 ^h 31 ^m 59 ^s	Moon souths 1760 Oct. 1.	18 ^h 38 ^m past Noon.
Noon . . . 2		12 35 37	Sun's R. A. then	12 34 48 from below.
La Caille's Eph.	Dif. 3 38		Moon's R. A. then } Sum	7 12 48 = 108° 12' (by Tab. p. 28.) from Aries following.
			or R. A. Mid-heaven } Rejecting 24 Hours.	
Now, As 24 ^h : 3 ^m 38 ^s Incr. R. A. :: 18 ^h 38 ^m : 2 ^m 49 ^s + }				
			Sun's R. A. Oct. 1. Noon 12 ^h 31 59	
Sun's R. A. at Moon's next Southing 18 ^h 38 ^m }			12 34 48, in Time.	
past Noon.				

TO find the Moon's right Ascension at Noon, preceding the given Time of Moon's Southing?

Sept. 30	☾ souths.	17 ^h 39 ^m	
1760 Oct. 1		18 38	
Diurnal Dif. R. A. ☾ à ☉.	Dif. 59 ^m		
Diurnal Dif. R. A. Sun	3 38 fr. above.		
Diurnal Dif. R. A. Moon	62 38		
By La Caille's Eph. 1760, Oct. 1.	☾'s Pla. Lat. Decl.		
	5° 26' 26" N. 4' 25° N. 26'		
As 24 ^h : 62 ^m 38 ^s :: 18 ^h 38 ^m ☾'s Southing since Noon to 48 ^m 53 ^s .			
To be deducted from R. A. Moon, at Southing 7 ^h 12 48			
Rem. Moon's R. A. 1760. Oct. 1. Noon . . . 6 ^h 23 ^m 55 ^s			
By Tab. p. 28, in Degrees	95° 58' 45"		
Whence, by Rule, Quest. 28. p. 227. R. A. ☾ 96° 0 52 fr. Aries.			
Dif. — 1 7 Error.			

N. B. The diurnal Difference of the right Ascension of the ☾ à ☉ from 12 to 12 is not exactly the same as the Difference of ☾ à ☉ from Southing to next Southing, thereby causing a small Error, computing by the Difference of Southing, as appears above.

EQUATION of the Ecliptic Right ASCENSION, to the Right ASCENSION with LATITUDE.

Longi- tude.	Signs					Signs								
	Lat.	{ North $\gamma, \delta, \Pi, \}$ } South $\alpha, \mu, \text{f}, \}$				0 1 2 6 7 8	Lat.	{ South $\gamma, \delta, \Pi, \}$ } North $\alpha, \mu, \text{f}, \}$				0 1 2 6 7 8		
		1°	3°	5°	7°			9°	1°	3°	5°		7°	9°
	—	—	—	—	—	+	+	+	+	+				
γ	0 23	1 11	1 59	2 47	3 37	0 24	1 11	1 59	2 47	3 35				
3	0 23	1 11	1 59	2 47	3 37	0 23	1 11	1 59	2 47	3 35				
6	0 23	1 11	1 59	2 47	3 36	0 23	1 11	1 59	2 47	3 34				
9	0 23	1 11	1 58	2 48	3 36	0 23	1 11	1 58	2 46	3 32				
12	0 23	1 10	1 58	2 48	3 35	0 23	1 11	1 58	2 45	3 30				
15	0 23	1 10	1 57	2 46	3 34	0 23	1 10	1 57	2 43	3 28				
18	0 23	1 9	1 55	2 44	3 32	0 23	1 9	1 55	2 41	3 26				
21	0 23	1 9	1 53	2 42	3 30	0 22	1 9	1 53	2 39	3 23				
24	0 22	1 8	1 51	2 40	3 28	0 22	1 8	1 52	2 36	3 20				
27	0 22	1 6	1 50	2 37	3 25	0 22	1 6	1 50	2 33	3 15				
δ	0 21	1 5	1 49	2 35	3 22	0 22	1 4	1 47	2 28	3 10				
3	0 20	1 3	1 47	2 31	3 18	0 21	1 3	1 44	2 25	3 5				
6	0 20	1 2	1 44	2 28	3 13	0 20	1 2	1 42	2 21	2 59				
9	0 20	1 1	1 42	2 24	3 9	0 19	0 59	1 38	2 15	2 54				
12	0 19	0 58	1 38	2 19	3 3	0 19	0 57	1 34	2 10	2 45				
15	0 18	0 55	1 33	2 13	2 55	0 19	0 55	1 31	2 6	2 39				
18	0 17	0 55	1 30	2 8	2 48	0 18	0 52	1 27	2 0	2 31				
21	0 17	0 51	1 26	2 2	2 39	0 17	0 49	1 21	1 54	2 22				
24	0 16	0 48	1 21	1 55	2 29	0 16	0 47	1 17	1 47	2 13				
27	0 15	0 45	1 16	1 48	2 20	0 15	0 44	1 12	1 39	2 4				
Π	0 14	0 41	1 10	1 40	2 10	0 14	0 41	1 7	1 32	1 55				
3	0 13	0 39	1 5	1 31	1 59	0 12	0 37	0 58	1 25	1 45				
6	0 11	0 34	0 58	1 22	1 48	0 11	0 33	0 49	1 16	1 34				
9	0 10	0 30	0 52	1 14	1 36	0 10	0 30	0 42	1 8	1 23				
12	0 8	0 26	0 44	1 3	1 24	0 9	0 26	0 35	0 58	1 12				
15	0 7	0 22	0 38	0 54	1 11	0 7	0 21	0 31	0 49	1 0				
18	0 6	0 18	0 30	0 43	0 58	0 6	0 17	0 22	0 40	0 49				
21	0 4	0 13	0 23	0 33	0 44	0 5	0 13	0 15	0 30	0 37				
24	0 3	0 9	0 15	0 23	0 31	0 3	0 9	0 9	0 20	0 24				
27	0 1	0 4	0 7	0 11	0 16	0 1	0 4	0 4	0 10	0 13				
30	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0				
0	+	+	+	+	+	—	—	—	—	—				
Longi- tude.	Lat.	{ South $\gamma, \delta, \Pi, \}$ } North $\alpha, \mu, \text{f}, \}$	1°	3°	5°	7°	9°	Lat.	{ North $\gamma, \delta, \Pi, \}$ } South $\alpha, \mu, \text{f}, \}$	1°	3°	5°	7°	9°

EQUATION to be subtracted fr. the Lat. of a Planet, for the EQUATION of the Ecliptic Declin. to the Declin. with that Planet's LATITUDE.

Lat. N. } $\gamma, \delta, \Pi, \alpha, \mu, \text{f}$ } + } 1 st S. } $\gamma, \delta, \Pi, \alpha, \mu, \text{f}$ } - } Eq.						Longi- tude.
1 ^o	2 ^o	3 ^o	4 ^o	5 ^o	6 ^o	
—	—	—	—	—	—	—
/	/	/	/	/	/	o
5	10	15	20	25	30	30
5	10	15	20	25	30	27
5	10	15	20	25	30	24
5	10	15	20	25	29	21
5	10	15	19	24	29	18
5	10	14	19	24	28	15
5	9	14	18	23	27	12
4	9	14	18	22	26	9
4	8	13	17	21	25	6
4	8	13	17	20	24	3
4	8	12	16	19	23	X μ
4	8	11	15	18	22	27
4	7	10	14	17	21	24
3	6	9	13	16	20	21
3	6	9	12	15	18	18
3	6	8	11	14	17	15
3	5	7	10	13	16	12
3	5	6	9	12	14	9
2	4	6	8	10	13	6
2	4	5	7	9	11	3
2	3	4	6	7	9	m N
2	2	3	5	6	8	27
2	2	3	4	5	6	24
1	2	2	3	4	5	21
1	1	2	2	3	4	18
1	1	1	1	2	3	15
1	1	1	1	1	2	12
0	0	0	0	0	1	9
0	0	0	0	0	0	6
0	0	0	0	0	0	3
0	0	0	0	0	0	f N
—	—	—	—	—	—	o
1 ^o	2 ^o	3 ^o	4 ^o	5 ^o	6 ^o	Long.

Lat. S. { $\gamma, \delta, \Pi, \alpha, \mu, \text{f}, \}$ }
N. { $\gamma, \delta, \Pi, \alpha, \mu, \text{f}, \}$ }

The following **Practical Rules**, from whence the above Equations are deduced, are in IMITATION of the Shortness and Usefulness of the **Dean of Durham's METHOD** in finding the **Parallactic Angle**.

CONSTRUCTION. For Lat. { North } Long. { $\gamma, \delta, \Pi, \}$ }
South { $\alpha, \mu, \text{f}, \}$ } 1. 2. }
Eq. 6. 7. 8. }

Or Lat. { South } Long. { $\gamma, \delta, \Pi, \}$ }
North { $\alpha, \mu, \text{f}, \}$ } 9. 10. 11. }
Eq. 3. 4. 5. }

PROPORTION.

As Radius S. 90°
To Cof. $\angle$ of Eclip. and Meridian (γ and α) . . . 66° 31' 30"
So Tan. Lat. 9 0 0
To Tan. Dif. R. A. required 3° 36' 36"

Logarithms.
10.0000000
9.6002636
9.1997125
8.7999761

For Lat. { S } Lon. { $\gamma, \delta, \Pi, \}$ }
N { $\alpha, \mu, \text{f}, \}$ } 0. 1. 2. }
Eq. 6. 7. 8. } Or Lat. { N } Lon. { $\gamma, \delta, \Pi, \}$ }
S { $\alpha, \mu, \text{f}, \}$ } 9. 10. 11. }
Eq. 3. 4. 5. }

As Radius S. 90°
To Cof. $\angle$ of Eclip. and Meridian (γ and α) . . . 66° 31' 30"
So S. Latitude 9 0 0
To S. Dif. R. A. required 3 34 22

Logarithms.
10.0000000
9.6002636
9.1943324
8.7945960

Example. To find the Right Ascension and Declination answerable to Longitude Π 5° 38', and Latitude 4° 38' 33" S.
Ecliptic R. A. (by Tab. p. 24) 63° 43' 15" } Eclip. Decl. (by Tab. p. 23) 21° 16' 32" N.
Eq. of R. A. to Lon. & Lat. + 44 0 } Equated Lat. by next Tab. — 4 34 33
R. A. required 64 27 15 } Declination required 16 42 0 N.

Conf. For Lat. N. { $\gamma, \delta, \Pi, \alpha, \mu, \text{f}, \}$ }
S. { $\gamma, \delta, \Pi, \alpha, \mu, \text{f}, \}$ } + } 1st
Eq.

PROPORTION. Logarithms.
As Rad. S. 90° 10.0000000
To S. $\angle$ of Eclip. and Meridian (γ and α) . . . 66° 31' 30" 9.9624801
So S. Lat. 6 0 0 9.0192346
To S. Difference Declin. req. 5 30 7 8.9817147

For Lat. S. { $\gamma, \delta, \Pi, \alpha, \mu, \text{f}, \}$ }
N. { $\gamma, \delta, \Pi, \alpha, \mu, \text{f}, \}$ } — } 1st
Eq.

As Rad. S. 90° 10.0000000
To S. $\angle$ of Eclip. and Meridian (γ and α) . . . 66° 31' 30" 9.9624801
So Tan. of Lat. 6 0 0 9.0216202
To Tang. Dif. Decl. req. 5 30 24 8.9841003

Ex. To find the Declination answerable to Long. 28° 11' m , and Lat. 2° 34' N.
Ecliptic Declination 19° 47' 5" S. by Tab. p. 23.
— 4' from the Lat. } = 2 30 0 } Lat. equated,
2° 34' N. } by Table above.

Remain 17 17 S. Declination reqd.

A TABLE of near LONGITUDE and RIGHT ASCENSION of the MID-HEAVEN, with the near Longitude of the NONAGESIMAL or 90 DEGREE of the ECLIPTIC from the Horizon, correspondent. For Latitude LONDON, $51^{\circ} 32'$.

Long.	Υ R. A. Mid-Hea.	Longitude Non. Deg. Υ .	δ R. A. Mid-Hea.	Longitude Non. Deg. δ .	Π R. A. Mid-Hea.	Longitude Non. Deg. Π .	ϖ R. A. Mid-Hea.	Longitude Non. Deg. ϖ .	Ω R. A. Mid-Hea.	Longitude Non. Deg. Ω .	Υ R. A. Mid-Hea.	Longitude Non. Deg. Υ .
0	0 0	26 42	27 54	16 31	57 48	7 12	90	0 0	122	22 38	152	13 30
1	0 55	27 23	28 51	17 11	58 51	8 6	91	0 47	123	23 32	153	14 9
2	1 50	28 2	29 49	17 51	59 54	8 50	92	1 33	124	24 5	154	14 49
3	2 45	28 42	30 46	18 31	60 57	9 34	93	2 19	125	24 48	155	15 29
4	3 40	29 21	31 44	19 12	62 0	10 18	94	3 5	126	25 32	156	16 9
5	4 35	29 51	32 42	19 52	63 3	11 3	95	3 51	127	26 16	157	16 48
6	5 30	30 40	33 40	20 32	64 6	11 47	96	4 37	128	26 58	158	17 28
7	6 25	31 20	34 39	21 13	65 10	12 31	97	5 23	129	27 41	159	18 8
8	7 21	32 0	35 37	21 24	66 13	13 16	98	6 9	130	28 22	160	18 47
9	8 16	32 39	36 36	22 25	67 17	14 1	99	6 55	131	29 6	161	19 27
10	9 11	33 19	37 35	23 16	68 18	14 46	100	7 40	132	29 47	162	20 6
11	10 6	33 59	38 34	23 57	69 45	15 31	101	8 26	133	30 30	162	20 46
12	11 2	34 38	39 33	24 38	70 29	16 16	103	9 12	134	1 13	163	21 26
13	11 57	35 17	40 32	25 19	71 34	17 1	104	9 57	135	1 55	164	22 5
14	12 53	35 57	41 32	26 0	72 38	17 46	105	10 43	136	2 36	165	22 45
15	13 48	36 36	42 31	26 42	73 43	18 32	106	11 28	137	3 18	166	23 24
16	14 44	37 15	43 31	27 24	74 47	19 17	107	12 14	138	4 0	167	24 4
17	15 40	37 55	44 31	28 6	75 52	20 4	108	12 59	139	4 41	168	24 43
18	16 36	38 35	45 32	28 47	76 57	20 49	109	13 45	140	5 22	169	25 23
19	17 31	39 14	46 32	29 30	78 2	21 35	110	14 30	141	6 4	170	26 2
20	18 27	39 54	47 33	29 51	79 7	22 20	111	15 14	142	6 45	171	26 41
21	19 24	40 33	48 33	30 55	80 12	23 6	112	15 59	143	7 26	172	27 22
22	20 20	41 12	49 34	31 37	81 17	23 51	113	16 44	144	8 6	173	28 1
23	21 16	41 52	50 35	32 20	82 22	24 37	114	17 29	145	8 47	174	28 40
24	22 13	42 32	51 37	3 2	83 28	25 23	115	18 14	146	9 28	174	29 20
25	23 9	43 12	52 38	3 46	84 33	26 9	116	18 58	147	10 8	175	29 50
26	24 6	43 52	53 40	4 29	85 38	26 35	118	19 42	148	10 48	176	30 39
27	25 2	44 32	54 42	5 12	86 44	27 41	119	20 26	149	11 29	177	1 19
28	26 0	45 12	55 44	5 55	87 49	28 27	120	21 10	150	12 9	178	2 0
29	26 57	45 51	56 46	6 39	88 55	29 13	121	21 54	151	12 50	179	2 30
30	27 54	46 31	57 48	7 22	90 0	30 0	122	22 38	152	13 30	180	3 19

Long.	ω R. A. Mid-Hea.	Longitude Non. Deg. ω .	Υ R. A. Mid-Hea.	Longitude Non. Deg. Υ .	δ R. A. Mid-Hea.	Longitude Non. Deg. δ .	Π R. A. Mid-Hea.	Longitude Non. Deg. Π .	ϖ R. A. Mid-Hea.	Longitude Non. Deg. ϖ .	Ω R. A. Mid-Hea.	Longitude Non. Deg. Ω .
0	180	3 19	208	25 15	238	27 10	270	0 0	302	2 50	332	4 45
1	181	3 59	209	26 5	239	28 37	271	2 37	303	4 14	333	5 35
2	182	4 40	210	26 56	240	29 6	272	5 19	304	5 37	334	6 23
3	183	5 20	211	27 47	241	30 37	273	7 55	305	6 58	335	7 12
4	184	6 2	212	28 39	242	31 11	274	10 29	306	8 17	336	8 0
5	185	6 48	213	29 31	243	32 48	275	13 2	307	9 33	337	8 48
6	186	7 24	214	30 24	244	34 27	277	15 37	308	10 49	338	9 35
7	186	8 5	215	31 19	245	36 8	278	18 7	309	12 3	339	10 22
8	187	8 46	216	32 13	246	38 52	279	20 35	310	13 14	340	11 7
9	188	9 20	217	33 10	247	41 40	280	23 0	311	14 24	341	11 52
10	189	10 10	218	34 6	248	44 30	281	25 22	312	15 32	342	13 37
11	190	10 52	219	35 5	249	47 20	282	27 43	313	16 40	342	13 23
12	191	11 35	220	36 1	250	50 10	283	30 0	314	17 46	343	14 7
13	192	12 18	221	37 0	252	53 18	284	32 16	315	18 53	344	14 51
14	193	12 50	222	38 0	253	56 20	285	34 27	316	19 56	345	15 35
15	194	13 43	223	39 2	254	59 30	286	36 33	317	20 58	346	16 17
16	195	14 26	224	40 6	255	62 32	287	38 39	318	22 0	347	17 1
17	196	15 10	225	41 10	256	65 44	288	40 43	319	23 0	348	17 44
18	197	15 54	226	42 14	257	68 58	289	42 42	320	23 59	349	18 26
19	198	16 39	227	43 21	258	72 17	291	44 40	321	24 58	350	19 9
20	198	17 23	228	44 29	259	75 38	292	46 31	322	25 55	351	19 52
21	199	18 8	229	45 37	260	79 0	293	48 20	323	26 51	352	20 33
22	200	18 54	230	46 46	261	82 24	294	50 5	324	27 47	353	21 15
23	201	19 39	231	47 58	262	85 53	295	52 52	325	28 41	354	21 56
24	202	20 26	232	49 11	263	89 23	296	54 33	326	29 36	354	22 37
25	203	21 14	233	50 27	265	92 59	297	56 13	327	30 29	355	23 18
26	204	22 1	234	51 43	266	96 20	298	58 49	328	31 32	356	23 59
27	205	22 49	235	53 3	267	99 5	299	61 22	329	32 14	357	24 39
28	206	23 37	236	54 24	268	102 39	300	63 54	330	33 5	358	25 11
29	207	24 26	237	55 47	269	105 20	301	66 23	331	34 56	359	25 6
30	208	25 15	238	57 10	270	108 0	302	68 30	332	35 45	360	26 42

N. B. The Nonagesimal or 90 Degree of the Ecliptic from the Horizon, is the highest Point of the Ecliptic at that Time, whose Altitude is therefore always equal to the $\angle$ of the Ecliptic and Horizon, which is also equal to the Zenith-Distance of the elevated Pole of the Ecliptic, at that Time.

A TABLE for determining the near Time, in any Day, when the SUN, MOON, a PLANET, or STAR, is in the highest Point of the Ecliptic. For Latitude London.

Argument. SUN's Place at Noon, or Planet's Longitude at Southing.											
Sun's Place.	1 st		2 ^d		3 ^d		4 ^d		5 ^d		
	Point of Ascendant	R. A. in Time.	Point of Ascendant	R. A. in Time.	Point of Ascendant	R. A. in Time.	Point of Ascendant	R. A. in Time.	Point of Ascendant	R. A. in Time.	
0	0 /	h m	0 /	h m	0 /	h m	0 /	h m	0 /	h m	
0	26 42	0 0	16 31	1 52	7 22	3 51	0 0	6 0	22 38	8 9	
1	27 23	0 4	17 11	1 55	8 6	3 55	0 47	6 4	23 22	8 13	
2	28 2	0 7	17 51	1 59	8 50	4 0	1 33	6 9	24 5	8 17	
3	28 42	0 11	18 31	2 3	9 34	4 4	2 19	6 13	24 48	8 21	
4	29 21	0 15	19 12	2 7	10 18	4 8	3 5	6 17	25 32	8 25	
5	0 1	0 18	19 52	2 11	11 3	4 12	3 51	6 22	26 16	8 29	
6	0 40	0 22	20 32	2 15	11 47	4 16	4 37	6 26	26 58	8 34	
7	1 20	0 26	21 13	2 19	12 31	4 21	5 23	6 31	27 41	8 38	
8	2 0	0 29	21 54	2 22	13 16	4 25	6 9	6 35	28 22	8 42	
9	2 39	0 33	22 25	2 26	14 1	4 29	6 55	6 39	29 6	8 46	
10	3 19	0 37	23 16	2 30	14 46	4 33	7 40	6 44	29 47	8 50	
11	3 59	0 40	23 57	2 34	15 31	4 38	8 26	6 48	0 30	8 54	
12	4 38	0 44	24 38	2 38	16 16	4 42	9 12	6 52	1 13	8 58	
13	5 17	0 48	25 19	2 42	17 1	4 46	9 57	6 57	1 55	9 2	
14	5 57	0 51	26 0	2 46	17 46	4 51	10 43	7 1	2 36	9 6	
15	6 36	0 55	26 42	2 50	18 32	4 55	11 28	7 5	3 18	9 10	
16	7 15	0 59	27 24	2 54	19 17	5 0	12 14	7 9	4 0	9 14	
17	7 55	1 2	28 6	2 58	20 4	5 4	12 59	7 14	4 41	9 18	
18	8 35	1 6	28 47	3 2	20 49	5 8	13 45	7 18	5 22	9 21	
19	9 14	1 10	29 30	3 6	21 35	5 12	14 30	7 22	6 4	9 26	
20	9 54	1 14	0 13	3 10	22 20	5 16	15 14	7 27	6 45	9 30	
21	10 33	1 18	0 55	3 14	23 6	5 21	15 59	7 31	7 26	9 34	
22	11 12	1 21	1 37	3 18	23 51	5 25	16 44	7 35	8 6	9 38	
23	11 52	1 25	2 20	3 22	24 37	5 29	17 29	7 39	8 47	9 41	
24	12 32	1 29	3 2	3 26	25 23	5 34	18 14	7 44	9 28	9 43	
25	13 12	1 33	3 46	3 31	26 9	5 38	18 58	7 48	10 8	9 49	
26	13 52	1 36	4 29	3 35	26 55	5 43	19 42	7 52	10 48	9 53	
27	14 32	1 40	5 12	3 39	27 41	5 47	20 26	7 56	11 29	9 57	
28	15 12	1 44	5 55	3 43	28 27	5 51	21 10	8 0	12 9	10 1	
29	15 51	1 48	6 39	3 47	29 13	5 56	21 54	8 5	12 50	10 5	
30	16 31	1 52	7 22	3 51	30 0	6 0	22 38	8 9	13 30	10 8	

Sun's Place.	6 ^d		7 ^d		8 ^d		9 ^d		10 ^d		11 ^d	
	Point of Ascendant	R. A. in Time.	Point of Ascendant	R. A. in Time.	Point of Ascendant	R. A. in Time.	Point of Ascendant	R. A. in Time.	Point of Ascendant	R. A. in Time.	Point of Ascendant	R. A. in Time.
0	0 /	h m	0 /	h m	0 /	h m	0 /	h m	0 /	h m	0 /	h m
0	3 19	12 0	25 15	13 52	27 10	15 51	0 0	18 0	2 11	20 9	4 25	22 8
1	3 59	12 4	26 5	13 55	28 37	15 55	2 37	18 4	4 14	20 13	5 35	22 12
2	4 40	12 7	26 56	13 59	0 6	16 0	5 19	18 4	5 37	20 17	6 23	22 16
3	5 20	12 11	27 47	14 3	1 37	16 4	7 55	18 13	6 58	20 21	7 12	22 20
4	6 2	12 15	28 39	14 7	3 11	16 8	10 29	18 17	8 17	20 25	8 0	22 24
5	6 48	12 18	29 31	14 11	4 48	16 12	13 2	18 22	9 33	20 29	8 48	22 27
6	7 24	12 22	0 24	14 15	6 27	16 16	15 37	18 26	10 49	20 33	9 35	22 31
7	8 5	12 26	1 19	14 19	8 8	16 21	18 7	18 31	12 3	20 38	10 22	22 35
8	8 46	12 29	2 13	14 22	9 52	16 25	20 35	18 35	13 14	20 42	11 7	22 39
9	9 28	12 33	3 10	14 26	11 40	16 29	23 0	18 39	14 24	20 46	11 52	22 43
10	10 10	12 37	4 6	14 30	13 30	16 33	25 22	18 44	15 32	20 50	12 37	22 46
11	10 52	12 40	5 5	14 34	15 20	16 38	27 43	18 48	16 40	20 54	13 23	22 50
12	11 35	12 44	6 1	14 38	17 19	16 42	30 0	18 52	17 46	20 58	14 7	22 54
13	12 18	12 48	7 0	14 42	19 18	16 46	2 16	18 57	18 53	21 2	14 51	22 57
14	12 50	12 51	8 0	14 46	21 20	16 51	4 27	19 1	19 56	21 6	15 35	23 1
15	13 43	12 55	9 2	14 50	23 30	16 55	6 33	19 5	20 58	21 10	16 17	23 5
16	14 26	12 59	10 6	14 54	25 32	16 59	8 39	19 9	22 0	21 14	17 1	23 9
17	15 10	13 3	11 9	14 58	27 44	17 3	10 43	19 14	23 0	21 18	17 44	23 12
18	15 54	13 6	12 14	15 2	29 58	17 8	12 45	19 18	23 59	21 22	18 26	23 16
19	16 39	13 10	13 21	15 6	2 17	17 12	14 40	19 22	24 58	21 26	19 9	23 20
20	17 23	13 14	14 29	15 10	4 38	17 16	16 31	19 27	25 55	21 30	19 52	23 23
21	18 8	13 18	15 37	15 14	7 0	17 21	18 20	19 31	26 51	21 34	20 33	23 27
22	18 54	13 21	16 46	15 18	9 24	17 25	20 5	19 35	27 47	21 38	21 15	23 31
23	19 39	13 25	17 58	15 22	11 53	17 29	21 52	19 39	28 41	21 41	21 56	23 34
24	20 26	13 19	19 11	15 26	14 23	17 34	23 33	19 44	29 36	21 45	22 37	23 38
25	21 14	13 33	20 27	15 31	16 59	17 38	25 13	19 48	0 29	21 49	23 18	23 42
26	22 1	13 36	21 23	15 35	19 30	17 43	26 49	19 52	1 22	21 53	23 59	23 45
27	22 47	13 40	23 3	15 39	22 5	17 47	28 22	19 56	2 14	21 57	24 39	23 49
28	23 36	13 44	24 24	15 43	24 29	17 51	29 54	20 0	3 5	22 1	25 21	23 53
29	24 26	13 48	25 47	15 47	27 20	17 56	1 23	20 5	3 56	22 5	26 2	23 56
30	25 15	13 52	27 10	15 51	30 0	18 0	2 50	20 9	4 45	22 8	26 42	24 0

Construction. The above Table contains the right Ascensions, in Time, of the Mid-Heaven, correspondent to the several Ascendant Point of the Ecliptic at the Horizon, and the right Ascension of the Mid-Heaven, in Degrees, at Noon, or Southing.

RULES and EXAMPLES for finding when the SUN, MOON, PLANETS, and STARS, will be in the Nonagesimal DEGREE.

And, therefore, the Difference of right Ascension, in Time, correspondent to the right Ascension of the *Mid-Heaven* to the Ascendant at Noon, or Southing, of the Sun, Moon, Planet or Star, and the right Ascension of the Mid-Heaven, correspondent to the Ascendant of the Nonagesimal Degree, will be the Time from Noon, or Southing, when the Sun, Moon, Planet, or Star, will be in the Nonagesimal Degree, or highest Point of the Ecliptic, required.

EXAMPLE I. Required the Time when the SUN will be in the Nonagesimal Degree, July 15, 1764?

RULE. Take out the right Ascension in Time to the Sun's Place at Noon, or Place of the Moon, Planet, or Star, at Southing. Then add 3 Signs to the Sun's Place at Noon, or to the Moon's Planet, or Star's Place at Southing, (taken also for the Place of the Nonagesimal Degree) and the Sum will be the Point of the Ecliptic, or Ascendant at the Horizon of London, (supposing no Change of Place from Noon, or Southing, to the Nonagesimal Degree) correspondent to which Ascendant take out the right Ascension of the Mid-Heaven, in Time, and place it under the former right Ascension of the Mid-Heaven, at Noon, or Southing, and the Difference will be the Time of the Nonagesimal Degree, past or before Noon for the Sun, or past or before Southing for the Planets, required. If which Time be added to or subtracted from the Time of Southing of the Planets, according as it is before or after Southing, you will have the Time of their being in the Nonagesimal Degree, respectively, required.

Sun's Place at Noon $23^{\circ} 18' 41''$ Right Ascension Mid-Heaven correspondent $7^h 40^m 40^s$
Add $3^{\circ} 0' 0''$

The Ascendant to the Nonagesimal Deg. $23 18 41$ R. A. Mid-Heaven correspondt. by the Table $8 12 44$
having nearly the Sun's Place at Noon, at that Time. }
Dif. R. Ascension or Time, past Noon, when the Sun is in the 90 Deg. or highest Point of Ecliptic, for that Day. } $32^m 4^s$

N. B. When the right Ascension of Mid-Heaven at Noon, or Southing in Time, is less than the right Ascension of Mid-Heaven to the Ascendant of the Nonagesimal Degree, then the Difference of right Ascensions, or Time of Nonagesimal Degree is past Noon, or Southing; but if the right Ascension of the Mid-Heaven, at Noon, or Southing, in Time, is more than the right Ascension of Mid-Heaven to the Ascendant of the Nonagesimal Degree, then the Difference of right Ascensions or Time of Nonagesimal Degree is before Noon, or Southing.

EXAMPLE II. To find the Time of the Sun being in the Nonagesimal Degree, 17th of April, 1764?

Sun's Place at Noon $27^{\circ} 54' 19''$ Right Ascension Mid-Heaven correspondent $1^h 43^m 38^s$
Add . . . $3^{\circ} 0' 0''$

The Ascendant to the Place of the Nonag. Degree having nearly the same Place with the Sun at Noon . . . } $27 54 19$ Right Ascension Mid-Heaven correspondent, by the Table $0 6 23$
Dif. R. Ascensions, or Time before Noon, when the Sun is in the 90 Deg. — $1 37 15$
 $12 0 0$

N. B. You may always deduct the R. A. of Mid-Heaven at Noon, from the R. A. of Mid-Heaven to the Ascendant, to the Nonag. Deg. for the Time past the present or former Day's Noon.

Viz. Hour before Noon . . . $10 22 45$
Or Time past former Noon . . . $22 22 15$
by deducting the Right Ascension Mid-Heaven at Noon, fr. R. A. to Mid-H. of Ascend. of the Nonag. Deg. first adding 24 Hours.

EXAMPLE III. To find the Time when the Moon will be in the Nonagesimal Degree, January 23, 1763?

Moon's Place at Noon $15^{\circ} 24'$ R. A. Mid-Heaven correspondent $19^h 6^m 36^s$
Add $3^{\circ} 0' 0''$

The Ascendant to the Place of Nonag. Deg. being supposed the same then with the Moon's Place. } $15 24$ R. A. Mid-Heaven correspondent $18 25 43$
Dif. R. A. censions or Time before the Moon souths when she is in the Nonages. Deg. — $40 53$
Moon souths Jan. 23, 1763, Noon . . . $12 0 0$

N. B. The Change of the Moon's Place before or after her Southing to the Nonages. Degree should be allowed for, when considerable.

Moon in the Nonagesimal Degree, required. $11 19 7$
Forenoon.

EXAMPLE

RULES and EXAMPLES for finding when the SUN, MOON, PLANETS, or STARS, will be in the Nonagesimal DEGREE.

EXAMPLE IV. To find the Time when the Moon will be in the Nonagesimal Degree, November 19, 1763?

Moon's Place at Noon $8^{\circ} 13' 23''$ R. A. Mid-Heaven, correspondent $2^h 44^m$
Add $3^{\circ} 0' 0''$

The Ascendant to the Place of the Nonag. Deg. supposed the same with the Moon's Place at Noon. } $4^{\circ} 13' 23''$ R. A. Mid-Heaven correspondent $0^h 41^m$

Dif. R. Ascensions, or Time before Moon souths, when she is in Nonag. } $-2^m 3^s$

But the Moon not being in the same Place, Say, As $24^h : 12^{\circ} 7'$ } Moon souths Nov. 19, 1763, London . . 11 22 P. M.
Moon's diur. Mot. :: $2^h 4^m$ estimate Time at Nonag. Deg. before Southing : $1^{\circ} 2'$ Change of Moon's Place less than } Moon in the Nonag. Degree, required . . 9 19 Night.
at Southing.

Hence, the Ascendant then will be $4^{\circ} 13' 23''$
 $- 1^{\circ} 2' 0''$

$3^{\circ} 11' 23''$ R. A. Mid-Heaven correspondent $- 37^m$

Time before Moon souths when she is in Nonages. $2^m 7^s$
Moon in the Nonagesimal Deg. $9^m 15^s$ Night,
more nearly.

EXAMPLE V. To find the Time when Mars will be in the Nonagesimal Degree, September 28, 1762?

Place $\S$ at Noon $1^{\circ} 13' 32''$ R. A. Mid-Heaven correspondent $16^h 48^m \frac{1}{2}$
Add $3^{\circ} 0' 0''$

The Ascendant to the Place of Nonag. Deg. } $1^{\circ} 13' 32''$ R. A. Mid-Heaven correspondent $17^h 32^m \frac{1}{4}$
being the same then supposed with Pl. $\S$ }

Dif. R. Ascensions, or Time past Southing, when $\S$ is in the Nonages. Degree . . $0^h 43^m \frac{3}{4}$
Add, $\S$ souths Sept. 28, 1762, London $4^h 28^m$ P. M.

$\S$ in Nonagesimal Deg. P. M. $5^h 11^m \frac{3}{4}$
[required.]

The same Method serves for finding the Time when any Star is in the Nonagesimal Degree; the foregoing Examples being sufficient for all Cases.

Here follow

Sexagesimal Tables:

Proportioning the

Difference of EQUATIONS

At S I G H T.

SEXAGESIMAL TABLE (continued) for proportioning the DIFFERENCE of EQUATIONS at Sight; being the PRODUCT of Sexagesimals divided by 60.

"X"	iv	iv	iv	iv	iv	iv	iv	iv	iv	iv	iv	iv	iv	iv	iv	iv	iv
"X"	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "
"X"	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "
" "	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30		
1	0 16	0 17	0 18	0 19	0 20	0 21	0 22	0 23	0 24	0 25	0 26	0 27	0 28	0 29	0 30		
2	0 32	0 34	0 36	0 38	0 40	0 42	0 44	0 46	0 48	0 50	0 52	0 54	0 56	0 58	1 0		
3	0 48	0 51	0 54	0 57	1 0	1 3	1 6	1 9	1 12	1 15	1 18	1 21	1 24	1 27	1 30		
4	1 4	1 8	1 12	1 16	1 20	1 24	1 28	1 32	1 36	1 40	1 44	1 48	1 52	1 56	2 0		
5	1 20	1 25	1 30	1 35	1 40	1 45	1 50	1 55	2 0	2 5	2 10	2 15	2 20	2 25	2 30		
6	1 36	1 42	1 48	1 54	2 0	2 6	2 12	2 18	2 24	2 30	2 36	2 42	2 48	2 54	3 0		
7	1 52	1 59	2 6	2 13	2 20	2 27	2 34	2 41	2 48	2 55	3 2	3 9	3 16	3 23	3 30		
8	2 8	2 16	2 24	2 32	2 40	2 48	2 56	3 4	3 12	3 20	3 28	3 36	3 44	3 52	4 0		
9	2 24	2 33	2 42	2 51	3 0	3 9	3 18	3 27	3 36	3 45	3 54	4 3	4 12	4 21	4 30		
10	2 40	2 50	3 0	3 10	3 20	3 30	3 40	3 50	4 0	4 10	4 20	4 30	4 40	4 50	5 0		
11	2 56	3 7	3 18	3 29	3 40	3 51	4 2	4 13	4 24	4 35	4 46	4 57	5 8	5 19	5 30		
12	3 12	3 24	3 36	3 48	4 0	4 12	4 24	4 36	4 48	5 0	5 12	5 24	5 36	5 48	6 0		
13	3 28	3 41	3 54	4 7	4 20	4 33	4 46	4 59	5 12	5 25	5 38	5 51	6 4	6 17	6 30		
14	3 44	3 58	4 12	4 26	4 40	4 54	5 8	5 22	5 36	5 50	6 4	6 18	6 32	6 46	7 0		
15	4 0	4 15	4 30	4 45	5 0	5 15	5 30	5 45	6 0	6 15	6 30	6 45	7 0	7 15	7 30		
16	4 16	4 32	4 48	5 4	5 20	5 36	5 52	6 8	6 24	6 40	6 56	7 12	7 28	7 44	8 0		
17	4 32	4 49	5 6	5 23	5 40	5 57	6 14	6 31	6 48	7 5	7 22	7 39	7 56	8 13	8 30		
18	4 48	5 6	5 24	5 42	6 0	6 18	6 36	6 54	7 12	7 30	7 48	8 6	8 24	8 42	9 0		
19	5 4	5 23	5 42	6 1	6 20	6 39	6 58	7 17	7 36	7 55	8 14	8 33	8 52	9 11	9 30		
20	5 20	5 40	6 0	6 20	6 40	7 0	7 20	7 40	8 0	8 20	8 40	9 0	9 20	9 40	10 0		
21	5 36	5 57	6 18	6 39	7 0	7 21	7 42	8 3	8 24	8 45	9 6	9 27	9 48	10 9	10 30		
22	5 52	6 14	6 36	6 58	7 20	7 42	8 4	8 26	8 48	9 10	9 32	9 54	10 16	10 38	11 0		
23	6 8	6 31	6 54	7 17	7 40	8 3	8 26	8 49	9 12	9 35	9 58	10 21	10 44	11 7	11 30		
24	6 24	6 48	7 12	7 36	8 0	8 24	8 48	9 12	9 36	10 0	10 24	10 48	11 12	11 36	12 0		
25	6 40	7 5	7 30	7 55	8 20	8 45	9 10	9 35	10 0	10 25	10 50	11 15	11 40	12 5	12 30		
26	6 56	7 22	7 48	8 14	8 40	9 6	9 32	9 58	10 24	10 50	11 16	11 42	12 8	12 34	13 0		
27	7 12	7 39	8 6	8 33	9 0	9 27	9 54	10 21	10 48	11 15	11 42	12 9	12 36	13 3	13 30		
28	7 28	7 56	8 24	8 52	9 20	9 48	10 16	10 44	11 12	11 40	12 8	12 36	13 4	13 32	14 0		
29	7 44	8 13	8 42	9 11	9 40	10 9	10 38	11 7	11 36	12 5	12 34	13 3	13 32	14 1	14 30		
30	8 0	8 30	9 0	9 30	10 0	10 30	11 0	11 30	12 0	12 30	13 0	13 30	14 0	14 30	15 0		
31	8 16	8 47	9 18	9 49	10 20	10 51	11 22	11 53	12 24	12 55	13 26	13 57	14 28	14 59	15 30		
32	8 32	9 4	9 36	10 8	10 40	11 12	11 44	12 16	12 48	13 20	13 52	14 24	14 56	15 28	16 0		
33	8 48	9 21	9 54	10 27	11 0	11 33	12 6	12 39	13 12	13 45	14 18	14 51	15 24	15 57	16 30		
34	9 4	9 38	10 12	10 46	11 20	11 54	12 28	13 2	13 36	14 10	14 44	15 18	15 52	16 26	17 0		
35	9 20	9 55	10 30	11 5	11 40	12 15	12 50	13 25	14 0	14 35	15 10	15 45	16 20	16 55	17 30		
36	9 36	10 12	10 48	11 24	12 0	12 36	13 12	13 48	14 24	15 0	15 36	16 12	16 48	17 24	18 0		
37	9 52	10 29	11 6	11 43	12 20	12 57	13 34	14 11	14 48	15 25	16 2	16 39	17 16	17 53	18 30		
38	10 8	10 46	11 24	12 2	12 40	13 18	13 56	14 34	15 12	15 50	16 28	17 6	17 44	18 22	19 0		
39	10 24	11 3	11 42	12 21	13 0	13 39	14 18	14 57	15 36	16 15	16 54	17 33	18 12	18 51	19 30		
40	10 40	11 20	12 0	12 40	13 20	14 0	14 40	15 20	16 0	16 40	17 20	18 0	18 40	19 20	20 0		
41	10 56	11 37	12 18	12 59	13 40	14 21	15 2	15 43	16 24	17 5	17 46	18 27	19 8	19 49	20 30		
42	11 12	11 54	12 36	13 18	14 0	14 42	15 24	16 6	16 48	17 30	18 12	18 54	19 36	20 18	21 0		
43	11 28	12 11	12 54	13 37	14 20	15 3	15 46	16 29	17 12	17 55	18 38	19 21	20 4	20 47	21 30		
44	11 44	12 28	13 12	13 56	14 40	15 24	16 8	16 52	17 36	18 20	19 4	19 48	20 32	21 16	22 0		
45	12 0	12 45	13 30	14 15	15 0	15 45	16 30	17 15	18 0	18 45	19 30	20 15	21 0	21 45	22 30		
46	12 16	13 2	13 48	14 34	15 20	16 6	16 52	17 38	18 24	19 10	19 56	20 42	21 28	22 14	23 0		
47	12 32	13 19	14 6	14 53	15 40	16 27	17 14	18 1	18 48	19 35	20 22	21 9	21 56	22 43	23 30		
48	12 48	13 36	14 24	15 12	16 0	16 48	17 36	18 24	19 12	20 0	20 48	21 36	22 24	23 12	24 0		
49	13 4	13 53	14 42	15 31	16 20	17 9	17 58	18 47	19 36	20 25	21 14	22 3	22 52	23 41	24 30		
50	13 20	14 10	15 0	15 50	16 40	17 30	18 20	19 10	20 0	20 50	21 40	22 30	23 20	24 10	25 0		
51	13 36	14 27	15 18	16 9	17 0	17 51	18 42	19 33	20 24	21 15	22 6	22 57	23 48	24 39	25 30		
52	13 52	14 44	15 36	16 28	17 20	18 12	19 4	19 56	20 48	21 40	22 32	23 24	24 16	25 8	26 0		
53	14 8	15 1	15 54	16 47	17 40	18 33	19 26	20 19	21 12	22 5	22 58	23 51	24 44	25 37	26 30		
54	14 24	15 18	16 12	17 6	18 0	18 54	19 48	20 42	21 36	22 30	23 24	24 18	25 12	26 6	27 0		
55	14 40	15 35	16 30	17 25	18 20	19 15	20 10	21 5	22 0	22 55	23 50	24 45	25 40	26 35	27 30		
56	14 56	15 52	16 48	17 44	18 40	19 36	20 32	21 28	22 24	23 20	24 16	25 12	26 8	27 4	28 0		
57	15 12	16 9	17 6	18 3	19 0	19 57	20 54	21 51	22 48	23 45	24 42	25 39	26 36	27 33	28 30		
58	15 28	16 26	17 24	18 22	19 20	20 18	21 16	22 14	23 12	24 10	25 8	26 6	27 4	28 2	29 0		
59	15 44	16 43	17 42	18 41	19 40	20 39	21 38	22 37	23 36	24 35	25 34	26 33	27 32	28 31	29 30		
60	16 0	17 0	18 0	19 0	20 0	21 0	22 0	23 0	24 0	25 0	26 0	27 0	28 0	29 0	30 0		

EXAMPLE II. Dif. Equat. $29' 23''$ }
By Dif. Arg. $51' 41''$ }

Product divided by 60 = 25' 19" Propor^l. Dif.
[required.]

For, $51' \times 29' = 24' \quad 39''$ *iii*
 $\quad \quad \quad 23'' = \quad \quad 19 \quad 33$
 $41'' \times 29' = \quad \quad 19 \quad 49$ *iv*
 $\quad \quad \quad 23'' = \quad \quad 15 \quad 43$ }

According to the Table above.

☞ Hence, to find the proportional Sexagesimal true to Seconds, the Fourths may be rejected, or omitted, in the Computation, and as far as Thirds only retained. — See further on.

[illegible]

According to the Table above.

29' 16" 47"

L1a

A TABLE of the VISIBLE ECLIPSES of the SUN, for 21 Principal CITIES of EUROPE.

CITIES NAMES.	1760. June 13, Morning.				1762. October 17, Morning.				1764. April 1, Forenoon.			
	Begin- ning.	Middle.	End.	Digits eclipsed.	Begin- ning.	Middle.	End.	Digits eclipsed.	Begin- ning.	Middle.	End.	Digits eclipsed.
	h m	h m	h m	d '	h m	h m	h m	d '	h m	h m	h m	d '
Amsterdam	7 1	7 42	8 42	4 48	7 8	8 14	9 14	6 0	9 43	11 8	0 43	11 1
Avignon	6 43	7 28	8 33	6 35	7 30	8 13	8 59	3 0	9 25	10 53	0 29	9 46
Berlin	7 35	8 31	9 28	5 24	7 44	8 52	9 59	7 4	10 14	11 52	1 32	10 3
Bologna in It.	7 10	8 4	9 10	7 41	7 52	8 45	9 41	4 33	9 59	11 31	1 5	8 52
Copenhagen	7 41	8 28	9 25	5 0	7 52	8 52	9 54	7 48	10 27	11 57	1 25	10 37
Cracow	7 55	8 46	9 59	6 56	8 24	9 28	10 31	6 40	10 56	0 26	1 56	8 34
Edinburgh	7 20	8 7	9 0	4 30	Set Ecl.	7 42	8 55	6 32	9 10	10 33	0 2	10 22
Geneva	6 58	7 48	8 54	7 0	7 39	8 25	9 23	4 5	9 44	11 15	0 50	9 19
Leipzig	7 31	8 19	9 22	5 50	7 51	8 50	9 54	6 45	10 17	11 48	1 19	9 57
Lisbon	5 41	6 24	7 18	6 27	7 0	Limb of Moon	pourra mordre.		9 57	11 26	1 3	7 30
London	6 41	7 14	8 17	4 10	6 59	7 50	8 48	5 20	9 16	10 43	0 15	Annular.
Lyons	6 44	7 30	8 33	6 25	7 25	8 10	9 5	4 0	9 27	10 56	0 33	10 10
Madrid	6 3	6 48	7 47	6 24	7 4	7 28	8 3	1 0	8 32	10 1	11 36	10 53
Montpellier	6 40	7 26	8 30	6 12	7 26	8 10	8 53	3 0	9 17	10 47	0 24	10 3
Naples	7 15	8 14	9 22	9 0	8 11	9 9	0 56	3 40	10 19	11 44	1 16	7 44
Paris	6 43	7 28	8 26	5 21	7 12	8 3	8 56	4 42	9 22 1/2	10 50	0 24	11 3
Petersburg	9 4	9 57	11 0	5 55	9 9	10 17	11 28	10 45	0 4	1 28	2 51	9 33
Rome	7 9	8 6	9 13	8 21	8 0	8 44	9 44	3 40	10 2	11 31	1 6	8 9
Turin	6 55	7 46	8 49	6 48	7 35	8 23	9 20	4 8	9 39	11 12	0 45	9 30
Vienna	7 37	8 30	9 42	7 6	8 10	9 6	10 12	6 0	10 36	0 1	1 38	8 39
Upsal	8 10	9 2	9 54	4 58	8 15	9 18	10 26	9 38	11 9	0 36	1 59	10 30

A TABLE of the VISIBLE ECLIPSES of the MOON, for Greenwich OBSERVATORY.
And likewise the INVISIBLE ECLIPSES of SUN and MOON.

1760.	1761.	1762.	1763.	1764.
ECLIPSES 4.	ECLIPSES 4.	ECLIPSES 4.	ECLIPSES 2.	ECLIPSES 4.
Two of the Sun, and 2 of the Moon.	Two of the Sun, and 2 of the Moon.	Two of the Sun, and 2 of the Moon.	Two of the Sun only.	Two of the Sun, and 2 of the Moon.
I. Of the Moon, May 29. Beg. 9 ^h 13 ^m 37 ^s } Night Mid. 9 36 37 } vis. at End 9 59 37 } Greenwich. Duration 46 0 } Quantity eclipsed 24' of a Digit on the Southern Limb.	I. Of the Moon, May 18. Beg. 8 ^h 31 ^m 40 ^s } Night. Imm. 9 38 22 } vis. at Mi. 10 25 9 } Greenwich. Em. 11 11 36 } End 12 18 38 } Dur. 3 46 58 } Quantity eclipsed 17 ^d 26' on the North Part.	I. Of the Sun, April 24, visible in the Southern Parts of the Earth. II. Of the Moon, May 8, visible in the West Indies. Beg. 2 ^h 21 ^m 34 ^s } Morn. Mid. 3 53 54 } Green- End 5 26 14 } wick. Quantity eclipsed 9 ^d 42' on the Southern Part of the Moon.	I. Of the Sun, April 13, visible in Africa, and towards the extreme Southern Parts of Europe. II. Of the Sun, Oct. 7, visible in South America, and not visible in Europe.	I. Of the Moon, March 17. Beg. 10 ^h 45 ^m 19 ^s } Night. Mid. 12 8 42 } vis. at End 1 32 5 } Green- Durat. 2 46 46 } wick. Quantity eclipsed 8 ^d 31' on the Southern Part of the Moon.
II. Of the Sun, (as above) June 13, in 1st Table.	II. Of the Sun, June 3, visible in Lapland and Norway.	III. Of the Sun, Oct. 17, visible, as above, in the 1st Table.	SEE	II. Of the Sun, April 1, visible at London, as in Table above.
III. Of the Moon, Nov. 22. Beg. 7 ^h 55 ^m 16 ^s } Night Mid. 9 8 46 } vis. at End 10 22 16 } Green- Dur. 2 27 0 } wick. Quantity eclipsed 6 Dig. 21' on the Northern Part of the Moon.	III. Of the Moon, Nov. 12. Beg. 10 ^h 0 ^m 3 ^s } Fore- Imm. 10 58 50 } noon. Mid. 11 47 57 } Green- Em. 11 51 26 } wick. End 1 35 51 } Quantity eclipsed 19 ^d 58', visible at Japan and China.	IV. Of the Moon, Nov. 1, visible, at Greenwich. Beg. 7 ^h 22 ^m 26 ^s } Night. Mid. 8 45 0 } End 10 7 34 } Quantity eclipsed 6 ^d 53' on the Southern Part of the Moon.	THE RULES OF COMPUTATION FURTHER ON.	III. Of the Moon, Sept. 10, visible in America. Beg. 6 ^h 1 ^m 8 ^s } Morning. Mid. 7 12 33 } at End 8 23 58 } Green- Dur. 2 22 50 } wick. Quantity eclipsed 4 ^d 48' on the Northern Part of the Moon.
IV. Of the Sun, Dec. 7. Middle 2 ^h 8 ^m Afternoon, invisible in Europe, but visible in South America.	IV. Of the Sun, Nov. 26. visible in the Southern Parts of the Globe, but not visible in Europe.			IV. Of the Sun, Sept. 25, visible only in the Southern Parts of the Earth.

A TABLE of the New, Full and QUARTER MOONS for Greenwich OBSERVATORY, from Noon.

MONTHS.	Days.	1760.	Days.	1761.	Days.	1762.	Days.	1763.	Days.	1764.
		h m		h m		h m		h m		h m
JANUARY.	2	4 51 Full.		h m	3	0 35 First Q.	6	5 29 Last Q.	2	22 2 New.
	9	17 56 Last Q.	5	21 44 New.	9	21 57 Full.	13	14 49 New.	10	19 24 First Q.
	17	18 32 New.	13	14 12 First Q.	16	15 9 Last Q.	21	20 39 First Q.	18	12 34 Full.
	24	19 11 First Q.	20	6 14 Full.	24	16 11 New.	28	23 59 Full.	25	9 51 Last Q.
	31	18 24 Full.	27	11 26 Last Q.						
FEBRUARY.			4	15 39 New.	1	16 6 First Q.	4	15 47 Last Q.	1	12 43 New.
	8	15 20 Last Q.	11	23 58 First Q.	8	8 3 Full.	12	10 15 New.	9	14 34 First Q.
	16	6 42 New.	18	17 47 Full.	15	6 50 Last Q.	20	13 49 First Q.	17	1 16 Full.
	23	3 17 First Q.	26	7 50 Last Q.	23	10 39 New.	27	10 20 Full.	23	18 11 Last Q.
MARCH.										
	1	9 17 Full.	6	6 59 New.	3	2 58 First Q.	6	4 9 Last Q.	2	5 7 New.
	9	11 55 Last Q.	13	7 44 First Q.	9	18 11 Full.	14	4 48 New.	10	9 43 First Q.
	16	20 3 New.	20	6 13 Full.	17	0 30 Last Q.	22	3 43 First Q.	17	12 3 Full.
	23	11 48 First Q.	28	4 14 Last Q.	25	3 34 New.	28	19 30 Full.	24	3 28 Last Q.
	31	1 7 Full.							31	22 30 New.
APRIL.										
	8	5 35 Last Q.	4	19 31 New.	1	11 22 First Q.	4	18 26 Last Q.	9	1 10 First Q.
	15	5 11 New.	11	14 29 First Q.	8	4 42 Full.	12	22 28 New.	15	22 38 Full.
	21	19 10 First Q.	18	19 47 Full.	15	19 9 Last Q.	20	13 13 First Q.	22	14 14 Last Q.
	29	17 27 Full.	26	22 37 Last Q.	30	19 9 First Q.	27	5 24 Full.	30	15 29 New.
MAY.										
	7	19 21 Last Q.	4	5 37 New.	7	16 0 Full.	4	12 24 Last Q.	8	12 47 First Q.
	14	13 5 New.	10	21 1 First Q.	15	13 17 Last Q.	12	13 23 New.	15	4 54 Full.
	21	7 56 First Q.	18	10 23 Full.	23	4 42 New.	19	18 1 First Q.	22	3 2 Last Q.
	29	9 26 Full.	26	14 0 Last Q.	29	22 23 First Q.	26	13 40 Full.	30	6 53 New.
JUNE.										
	6	5 17 Last Q.	2	19 44 New.	6	4 19 Full.	3	3 12 Last Q.	6	21 13 First Q.
	12	20 18 New.	9	6 6 First Q.	14	5 56 Last Q.	11	1 25 New.	13	12 14 Full.
	19	20 33 First Q.	17	1 14 Full.	21	13 18 New.	18	0 43 First Q.	20	18 11 Last Q.
	28	0 19 Full.	25	2 26 Last Q.	28	4 3 First Q.	24	23 23 Full.	28	20 18 New.
JULY.										
	5	12 16 Last Q.	1	20 31 New.	5	18 0 Full.	2	20 23 Last Q.	6	3 11 First Q.
	12	3 53 New.	8	14 32 First Q.	13	20 55 Last Q.	10	11 31 New.	12	20 15 Full.
	19	11 37 First Q.	16	16 38 Full.	20	20 33 New.	17	5 37 First Q.	20	10 27 Last Q.
	27	13 27 Full.	24	12 1 Last Q.	27	12 40 First Q.	24	10 52 Full.	28	8 4 New.
AUGUST.										
	3	17 50 Last Q.	7	3 26 First Q.	4	8 11 Full.	1	13 17 Last Q.	4	8 7 First Q.
	10	12 49 New.	15	7 18 Full.	12	9 43 Last Q.	8	20 3 New.	11	6 15 Full.
	18	4 37 First Q.	22	19 57 Last Q.	19	2 33 New.	15	11 39 First Q.	19	3 59 Last Q.
	26	1 32 Full.	29	11 17 New.	25	22 24 First Q.	31	4 53 Last Q.	26	18 58 New.
SEPTEMBER.										
	1	23 18 Last Q.	5	19 43 First Q.	3	0 5 Full.	7	4 19 New.	2	13 31 First Q.
	8	23 50 New.	13	21 28 Full.	10	20 33 Last Q.	13	20 18 First Q.	9	19 4 Full.
	16	23 24 First Q.	21	2 48 Last Q.	17	11 33 New.	21	17 1 Full.	17	21 44 Last Q.
	24	12 21 Full.	27	21 24 New.	24	12 45 First Q.	29	18 49 Last Q.	25	5 9 New.
OCTOBER.										
	1	5 56 Last Q.	5	14 43 First Q.	2	16 30 Full.	6	13 2 New.	1	20 30 First Q.
	8	14 0 New.	13	11 6 Full.	10	5 31 Last Q.	13	8 9 First Q.	9	10 52 Full.
	16	18 12 First Q.	20	9 32 Last Q.	16	21 14 New.	21	10 58 Full.	17	14 49 Last Q.
	23	22 46 Full.	27	10 31 New.	24	6 34 First Q.	29	6 44 Last Q.	24	14 43 New.
	30	14 34 Last Q.							31	6 32 First Q.
NOVEMBER.										
	7	7 2 New.	4	10 51 First Q.	1	8 53 Full.	4	22 43 New.	8	4 39 Full.
	15	11 19 First Q.	11	23 48 Full.	8	13 18 Last Q.	11	23 55 First Q.	16	6 51 Last Q.
	22	9 24 Full.	18	17 3 Last Q.	15	9 18 New.	20	5 6 Full.	23	0 45 New.
	29	1 59 Last Q.	26	2 35 New.	23	2 44 First Q.	27	12 10 Last Q.	29	20 35 First Q.
DECEMBER.										
	7	2 8 New.	4	6 32 First Q.	7	20 53 Last Q.	4	9 32 New.	7	23 47 Full.
	15	2 13 First Q.	11	11 20 Full.	14	23 35 New.	11	4 12 First Q.	15	19 25 Last Q.
	21	19 27 Full.	18	2 33 Last Q.	23	9 12 First Q.	19	21 15 Full.	22	10 57 New.
	28	17 0 Last Q.	25	20 54 New.	30	12 43 Full.	27	1 42 Last Q.	29	11 48 First Q.

The USE of Table, p. 264. To reduce the Moon's Place from Greenwich to Antigua Meridian 60° W. Diff. Longitude?

D's Pl. Greenwich.

By the Table.

Example. 1763 Nov. 1, Noon 13° 17' 51" Against 14° Diur. Dif. under 60° Dif. Long. W. . . . 2° 20'

2, Noon 28 36 } 45' 7 30"

Diur. Dif. 14° 45'

Equation of D's Place . . . + 2 27 30

1763 Nov. 1, } D's Place at Greenwich . . 13 51 0

1 Noon.

} D's Place at Antigua, reqd. 11 16 18 30

N.B. The same Method serves for
☉: Declin. R. A. &c.

A TABLE of the Semi-diurnal Arcs of the SUN to every Degree of his Longitude in the Ecliptic, for London.

Argument. SUN's Longitude.

Lon.	3°	4°	5°	6°	7°	8°	Lon.
0	+	+	+	+	+	+	0
0	8 13	7 50	6 59	6 0	5 1	4 10	30
1	8 13	7 49	6 58	5 58	4 59	4 8	29
2	8 12	7 47	6 56	5 56	4 57	4 7	28
3	8 12	7 46	6 54	5 54	4 55	4 5	27
4	8 11	7 45	6 52	5 52	4 53	4 4	26
5	8 11	7 43	6 50	5 50	4 52	4 3	25
6	8 10	7 42	6 48	5 48	4 50	4 2	24
7	8 10	7 41	6 46	5 46	4 48	4 1	23
8	8 9	7 40	6 44	5 44	4 46	4 0	22
9	8 9	7 39	6 42	5 42	4 45	3 59	21
10	8 8	7 38	6 40	5 40	4 43	3 58	20
11	8 8	7 37	6 38	5 38	4 41	3 57	19
12	8 7	7 36	6 36	5 36	4 39	3 56	18
13	8 7	7 35	6 34	5 34	4 37	3 55	17
14	8 6	7 33	6 32	5 32	4 36	3 54	16
15	8 6	7 31	6 30	5 30	4 34	3 54	15
16	8 5	7 30	6 28	5 28	4 32	3 53	14
17	8 4	7 28	6 26	5 26	4 30	3 53	13
18	8 4	7 26	6 24	5 24	4 29	3 52	12
19	8 3	7 23	6 22	5 22	4 27	3 51	11
20	8 2	7 20	6 20	5 20	4 25	3 51	10
21	8 1	7 17	6 18	5 18	4 23	3 50	9
22	8 0	7 15	6 16	5 16	4 21	3 50	8
23	7 59	7 13	6 14	5 14	4 20	3 49	7
24	7 57	7 11	6 12	5 12	4 18	3 49	6
25	7 56	7 9	6 10	5 10	4 16	3 48	5
26	7 55	7 7	6 8	5 8	4 15	3 48	4
27	7 53	7 5	6 6	5 6	4 13	3 48	3
28	7 52	7 3	6 4	5 4	4 12	3 47	2
29	7 51	7 1	6 2	5 2	4 11	3 47	1
30	7 50	6 59	6 0	5 1	4 10	3 47	0
Lon.	+	+	+	+	+	+	Lon.
0	II 25	8 15	V 05	X 115	105	95	0

A TABLE of the Semidiurnal Arcs of the MOON to every Degree of her Longitude in the Ecliptic, at Southing, without Lat. for London.

Argument. MOON's Longitude.

Lon.	3°	4°	5°	6°	7°	8°	Lon.
0	+	+	+	+	+	+	0
0	8 29	8 6	7 13	6 12	5 11	4 18	30
1	8 29	8 5	7 12	6 10	5 9	4 16	29
2	8 28	8 3	7 10	6 8	5 7	4 15	28
3	8 28	8 2	7 8	6 6	5 5	4 13	27
4	8 27	8 1	7 6	6 4	5 3	4 12	26
5	8 27	7 59	7 4	6 2	5 2	4 11	25
6	8 26	7 58	7 2	6 0	5 0	4 10	24
7	8 26	7 57	7 0	5 58	4 58	4 9	23
8	8 25	7 56	6 58	5 56	4 56	4 8	22
9	8 25	7 55	6 56	5 54	4 55	4 7	21
10	8 24	7 54	6 54	5 52	4 53	4 6	20
11	8 24	7 53	6 52	5 50	4 51	4 5	19
12	8 23	7 52	6 50	5 48	4 49	4 4	18
13	8 23	7 51	6 48	5 46	4 47	4 3	17
14	8 22	7 49	6 46	5 44	4 46	4 2	16
15	8 22	7 47	6 43	5 42	4 44	4 2	15
16	8 21	7 45	6 41	5 39	4 42	4 1	14
17	8 20	7 43	6 39	5 37	4 40	4 1	13
18	8 20	7 41	6 37	5 35	4 38	4 0	12
19	8 19	7 38	6 35	5 33	4 36	3 59	11
20	8 18	7 35	6 33	5 31	4 34	3 59	10
21	8 17	7 31	6 31	5 29	4 32	3 58	9
22	8 16	7 29	6 29	5 27	4 30	3 58	8
23	8 15	7 27	6 26	5 24	4 29	3 57	7
24	8 13	7 25	6 24	5 22	4 27	3 57	6
25	8 12	7 23	6 22	5 20	4 25	3 56	5
26	8 11	7 21	6 20	5 18	4 23	3 56	4
27	8 9	7 19	6 18	5 16	4 21	3 56	3
28	8 8	7 17	6 16	5 14	4 20	3 55	2
29	8 7	7 15	6 14	5 12	4 19	3 55	1
30	8 6	7 13	6 12	5 11	4 18	3 55	0
Lon.	+	+	+	+	+	+	Lon.
0	II 25	8 15	V 05	X 115	105	95	0

This Table is a Correction of the Sun's Semi-diur. Arcs by mean Mot. D to 0.

A TABLE shewing, by the Moon's Age, the near Time of her Rising and Setting, in England and Ireland, when she is in the Beginning, Middle, or End of each Sign in the Ecliptic (supposing her to have no Lat.) her mean Place and Age being known, at Sight, by Tab. p. 151, 152.

(Age.)	3 ^s	4 ^s	5 ^s	6 ^s	7 ^s	8 ^s
	Mid.	End Be.	Mid.	End Be.	Mid.	End.
	8h 21m	8h 6m	7h 47m	7h 13m	6h 43m	6h 12m
	Sets.	Sets.	Sets.	Sets.	Sets.	Sets.
1	9 A 10	8 54	8 A 35	8 A 1	7 A 31	7 A 0
2	9 58	9 42	9 23	8 49	8 19	7 48
3	10 46	10 30	10 11	9 37	9 7	8 36
4	11 34	11 18	10 59	10 25	9 55	9 24
5	0 M 22	0 M 6	11 47	11 13	10 43	10 12
6	1 10	0 54	0 M 35	0 M 1	11 31	11 0
7	1 58	1 42	1 23	0 49	0 M 19	11 48
8	2 46	2 30	2 11	1 37	1 7	0 M 36
9	3 34	3 18	2 59	2 25	1 55	1 24
10	4 22	4 6	3 47	3 13	2 43	2 12
11	5 10	4 54	4 35	4 1	3 31	3 0
12	5 58	5 42	5 23	4 49	4 19	3 48
13	6 46	6 30	6 11	5 37	5 7	4 36
14	7 34	7 18	6 59	6 25	5 55	5 24
	Rifes.	Rifes.	Rifes.	Rifes.	Rifes.	Rifes.
15	3 A 38	3 A 54	4 A 13	4 A 47	5 A 17	5 A 48
16	4 26	4 42	5 1	5 35	6 5	6 36
17	5 14	5 30	5 49	6 23	6 53	7 24
18	6 2	6 18	6 37	7 11	7 41	8 12
19	6 50	7 6	7 25	7 59	8 29	9 0
20	7 38	7 54	8 13	8 47	9 17	9 48
21	8 26	8 42	9 1	9 35	10 5	10 36
22	9 14	9 30	9 49	10 23	10 53	11 24
23	10 2	10 18	10 37	11 11	11 41	0 M 10
24	10 50	11 6	11 25	11 59	0 M 29	1 0
25	11 38	11 54	0 M 13	0 M 47	1 17	1 48
26	0 M 26	0 M 42	1 1	1 35	2 5	2 36
27	1 14	1 30	1 49	2 23	2 53	3 24
28	2 2	2 18	2 37	3 11	3 41	4 12
29	2 50	3 6	3 25	3 59	4 29	5 0
50	3 38	3 54	4 13	4 47	5 17	5 48
D	Mid.	Be. End	Mid.	Be. End	Mid.	Be. End
A	11 2 ^s		8 1 ^s		7 0 ^s	6 11 ^s

EQUATION of the true Horizontal DIAMETER of the MOON to the apparent Horizontal DIAMETER, in Parts of the EQUATOR.

Argument. MOON'S DECLINATION.

True Horizontal Diameter of the Moon.

Dec. of $\odot$	29' 30"	30' 0"	30' 30"	31' 0"	31' 30"	32' 0"	32' 30"	33' 0"	33' 30"	34' 0"
0	+	+	+	+	+	+	+	+	+	+
1	0	0	0	0	0	0	0	0	0	0
2	0	1	0	1	0	1	0	1	0	1
3	0	2	0	2	0	2	0	3	0	3
4	0	4	0	4	0	4	0	5	0	5
5	0	7	0	7	0	7	0	7	0	7
6	0	10	0	10	0	10	0	11	0	11
7	0	13	0	13	0	14	0	15	0	15
8	0	17	0	17	0	18	0	19	0	20
9	0	21	0	22	0	23	0	24	0	25
10	0	27	0	27	0	28	0	29	0	30
11	0	33	0	34	0	35	0	36	0	37
12	0	40	0	41	0	42	0	43	0	44
13	0	47	0	48	0	49	0	50	0	51
14	0	54	0	55	0	56	0	57	0	58
15	1	2	1	3	1	4	1	5	1	6
16	1	11	1	12	1	13	1	14	1	15
17	1	21	1	22	1	23	1	24	1	25
18	1	31	1	32	1	33	1	34	1	35
19	1	42	1	43	1	44	1	45	1	46
20	1	54	1	55	1	56	1	57	1	58
21	2	6	2	8	2	10	2	12	2	14
22	2	19	2	21	2	23	2	26	2	28
23	2	33	2	35	2	38	2	41	2	44
24	2	47	2	50	2	53	2	56	2	59
25	3	2	3	5	3	9	3	12	3	15
26	3	18	3	22	3	25	3	29	3	32
27	3	36	3	40	3	43	3	47	3	50
28	3	55	3	59	4	3	4	7	4	11
29	4	14	4	19	4	23	4	27	4	31
30	4	34	4	39	4	43	4	48	4	52

CONSTRUCTION.

To find the Equation to 30' true Semi-Diam. of the Moon in Parts of the Equator?

As Col. Declination	30°	co.	0.0624694
To Radius	S. 90		10.0000000
So for the Ref. So true Horiz. Diam. $\odot$	30'		7.9408419
To Apparent Diameter	34 39"		8.0033113
Diff. The Equation	+ 4' 39"		required.

NEXT, To find the Time of the Moon's Passage thro' the Meridian?

As 360° is to $\odot$'s apparent horizontal Diam. in Parts of the Equator, so is the Interval of Time betwixt the Moon's two next Passages thro' the same Meridian, (or Dif. of the Sun and Moon's R. A. in Time, for 24 Hours) to the Time of the Moon's Passage thro' that Meridian.

EXAMPLE. To find the Time of the Moon's Passage thro' the Meridian, December 30, 1764, at Greenwich?

$\odot$'s Declination 30 Dec. at Noon, 7° 31' N. True Horiz. Parallax then, 54' 21" For Horiz. Parall. 21 Dec. 56' 52" 1' 7" True Hor. Parallax } 54 16
26... 55 28 } at passing the Merid. }

Dif. in 5 Days 1 24 : 4 : 1' 7" }
 $\odot$ Passes Merid. Greenwich 30 Dec. 1674 6h 18m } Equation . . + 0 15
31 . . . 6 59 } $\odot$ app. Horiz. Diam. 29 30

Now, As 360° to 29' 30", So 24h 41m to 2m 1s, Time of $\odot$ passing the Meridian required.

N. B. De La Caille's Ephem. has true Horiz. Diam. 29' 23" }
Time of $\odot$'s Passage thro' the Meridian 2m 2s }

The Moon's Horizontal Parallax to $\odot$'s Horiz. Diameter is 55' 40" to 30'

Equation of the true Diameter of $\odot$ to the ap. Diam. in Alt.

Argument. $\odot$'s Altitude.

$\odot$'s Alt.	Moon in Apog.	Perig.
+	+	+
0	0	0
3	2	2
6	3	4
9	5	6
12	6	7
15	8	9
18	9	11
21	10	12
24	11	15
27	12	16
30	14	18
33	15	19
36	16	21
39	18	23
42	19	24
45	20	25
48	21	27
51	22	28
54	23	29
57	23	30
60	24	31
63	25	32
66	26	33
69	26	34
72	27	34
75	27	35
78	27	35
81	28	36
84	28	36
87	28	36
90	29	36

Equat. or Reduct. of Paris to Greenwich Places.

Argument. Differences.

Dif.	Equat.	Dif.	Equat.
+	+	+	+
0	0	0	0
1	0 23	31	12 3
2	0 47	32	12 27
3	1 10	33	12 50
4	1 33	34	13 13
5	1 56	35	13 37
6	2 20	36	14 0
7	2 43	37	14 23
8	3 7	38	14 47
9	3 30	39	15 10
10	3 53	40	15 33
11	4 17	41	15 57
12	4 40	42	16 20
13	5 3	43	16 43
14	5 27	44	17 7
15	5 50	45	17 30
16	6 13	46	17 53
17	6 37	47	18 17
18	7 0	48	18 40
19	7 23	49	19 3
20	7 47	50	19 27
21	8 10	51	19 50
22	8 33	52	20 13
23	8 57	53	20 37
24	9 20	54	21 0
25	9 53	55	21 13
26	10 7	56	21 47
27	10 30	57	22 10
28	10 53	58	22 33
29	11 16	59	22 57
30	11 40	60	23 20

N. B. The above small Table is similar to that at p. 215, here recited for Use.

$\odot$ tr. Horiz. Diam. 29' 15" Equation + 0 15

$\odot$'s apparent Hor. Diam. } 29 30
 $\odot$'s Altitude } 46° 2' Greenwich
Equation at a Means . } + 0 22

$\odot$'s ap. Diam. in Altitude. } 29 52
Correspondent to which, in first Tab. p. 29, answers 1m 59s
Answerable to which $\odot$ mean Mot. } 1 5

Sum 30 57
In Time, by Tab. p. 29. } 2m 3s
Or 360° to 29' 52" :: 24h 41m : 2m 3s as before.

EXAMPLE I.

$\odot$'s Place, 1764, Aug. 9, Noon 18° 16' 29'
At Paris 10 . . . 2 17

Dif. 13 48
To reduce the $\odot$'s Places to Greenwich for those Days.

Argument. 13° + 5' 3" 48' + 0 19

Sum, Equation + 5 22
Noon { Aug. 9 18° 16' 34'
10 2 22
 $\odot$'s Places at Greenwich, reqd.

EXAMPLE II.

$\odot$'s Declination. Noon { 1763, Oct. 4 12° 44' N
At Paris 5 5 58 N

Dif. 6 46
To reduce these Declinations to Greenwich?

Argument. 6° - 2' 20" 46' - 0 18

Sum, Equation - 2 38
Noon { Oct. 4 12 41 N
5 5 55 N
At Greenwich, required.

N. B. The Moon's semi-diurnal Arc, as in 2d Table of the left Hand Page, is added to or subtracted from the mean Time of her Setting, according to her Age, for her Setting and Rising, as at Bottom of the left Hand Page, in the 3d Table.

A TABLE shewing the PROPORTIONAL DIFFERENCE to be added or subtracted to or from the SUN, MOON, or PLANET's Place, Declination, R. A. &c. at a first Meridian for the respective Place, Declination, R. A. &c. to a given Difference of West or East Longitude.

Arguments. Diurnal Difference more or less; and Difference of Longitude.

Diurnal Diff.	10°			20°			30°			40°			50°			60°			70°			80°			90°			100°			110°			120°			
	+	-		+	-		+	-		+	-		+	-		+	-		+	-		+	-		+	-		+	-		+	-					
	°	'	iv	°	'	iv	°	'	iv	°	'	iv	°	'	iv	°	'	iv	°	'	iv	°	'	iv	°	'	iv	°	'	iv	°	'	iv				
1	°	'	iv	°	'	iv	°	'	iv	°	'	iv	°	'	iv	°	'	iv	°	'	iv	°	'	iv	°	'	iv	°	'	iv	°	'	iv				
0	0	1	iv	0	1	iv	0	1	iv	0	1	iv	0	1	iv	0	1	iv	0	1	iv	0	1	iv	0	1	iv	0	1	iv	0	1	iv				
1	0	1	40	0	3	20	0	5	0	0	6	40	0	8	20	0	10	0	0	11	40	0	13	20	0	15	0	0	16	40	0	18	20	0	20	0	
2	0	3	20	0	6	40	0	10	0	0	13	20	0	16	40	0	20	0	0	23	20	0	26	40	0	30	0	0	33	20	0	36	40	0	40	0	
3	0	5	0	0	10	0	0	15	0	0	20	0	0	25	0	0	30	0	0	35	0	0	40	0	0	45	0	0	50	0	0	55	0	1	0	0	
4	0	6	40	0	13	20	0	20	0	0	26	40	0	33	20	0	40	0	0	46	40	0	53	20	0	1	0	0	1	6	40	1	13	20	1	20	0
5	0	8	20	0	16	40	0	25	0	0	33	20	0	41	40	0	50	0	0	58	20	1	6	40	1	15	0	1	23	20	1	31	40	1	40	0	
6	0	10	0	0	20	0	0	30	0	0	40	0	0	50	0	1	0	0	1	10	0	1	20	0	1	30	0	1	40	0	1	50	0	2	0	0	
7	0	11	40	0	23	20	0	35	0	0	46	40	0	58	20	1	10	0	1	21	40	1	33	20	1	45	0	1	56	40	2	8	20	2	20	0	
8	0	13	20	0	26	40	0	40	0	0	53	20	1	6	40	1	20	0	1	33	20	1	46	40	2	0	0	2	12	20	2	26	40	2	40	0	
9	0	15	0	0	30	0	0	45	0	1	0	0	1	15	0	1	30	0	1	45	0	2	0	0	2	15	0	2	30	0	2	45	0	3	0	0	
10	0	16	40	0	33	20	0	50	0	1	6	40	1	23	20	1	40	0	1	56	40	2	13	20	2	30	0	2	46	40	3	3	20	3	20	0	
11	0	18	20	0	36	40	0	55	0	1	13	20	1	31	40	1	50	0	2	8	20	2	26	40	2	45	0	3	3	20	3	21	40	3	40	0	
12	0	20	0	0	40	0	1	0	0	1	20	0	1	40	0	2	0	0	2	20	0	2	40	0	3	0	0	3	20	0	3	40	0	4	0	0	
13	0	21	40	0	43	20	1	5	0	1	26	40	1	48	20	2	10	0	2	31	40	2	53	20	3	15	0	3	36	40	3	58	20	4	20	0	
14	0	23	20	0	46	40	1	10	0	1	33	20	1	56	40	2	20	0	2	43	20	3	6	40	3	30	0	3	53	20	4	16	40	4	40	0	
15	0	25	0	0	51	0	1	15	0	1	40	0	2	5	0	2	30	0	2	55	0	3	20	0	3	45	0	4	10	0	4	35	0	5	0	0	
16	0	26	40	0	54	20	1	20	0	1	46	40	2	13	20	2	40	0	3	6	40	3	33	20	4	0	0	4	26	40	4	53	20	5	20	0	
17	0	28	20	0	57	40	1	25	0	1	53	20	2	21	40	2	50	0	3	18	20	3	46	40	4	15	0	4	43	20	5	11	40	5	40	0	
18	0	30	0	1	0	0	1	30	0	1	0	0	2	30	0	3	0	0	3	30	0	4	0	0	4	30	0	5	0	0	5	30	0	6	0	0	
19	0	31	40	1	3	20	1	35	0	2	6	40	2	38	20	3	10	0	3	41	40	4	13	20	4	45	0	5	16	40	5	48	20	6	20	0	
20	0	33	20	1	6	40	1	40	0	2	13	20	2	46	40	3	20	0	3	53	20	4	26	40	5	0	0	5	33	20	6	6	40	6	40	0	
21	0	35	0	1	10	0	1	45	0	2	20	0	2	55	0	3	30	0	4	5	0	4	40	0	5	15	0	5	50	0	6	25	0	7	0	0	
22	0	36	40	1	13	20	1	50	0	2	26	40	3	3	20	3	40	0	4	16	40	4	53	20	5	30	0	6	6	40	6	43	20	7	20	0	
23	0	38	20	1	16	40	1	55	0	2	33	20	3	11	40	3	50	0	4	28	20	5	6	40	5	45	0	6	23	20	7	1	40	7	40	0	
24	0	40	0	1	20	0	2	0	0	2	40	0	3	20	0	4	0	0	4	40	0	5	20	0	6	0	0	6	40	0	7	20	0	8	0	0	
25	0	41	40	1	23	20	2	5	0	2	46	40	3	28	20	4	10	0	4	51	40	5	33	20	6	15	0	6	56	40	7	38	20	8	20	0	
26	0	43	20	1	26	40	2	10	0	2	53	20	3	36	40	4	20	0	5	3	20	5	46	40	6	30	0	7	13	20	7	56	40	8	40	0	
27	0	45	0	1	30	0	2	15	0	3	0	0	3	45	0	4	30	0	5	15	0	6	0	0	6	45	0	7	30	0	8	15	0	9	0	0	
28	0	46	40	1	33	20	2	20	0	3	6	40	3	53	20	4	40	0	5	26	40	6	13	20	7	0	0	7	46	40	8	33	20	9	20	0	
29	0	48	20	1	36	40	2	25	0	3	13	20	4	1	40	4	50	0	5	38	20	6	26	40	7	15	0	8	3	20	8	51	40	9	40	0	
30	0	50	0	1	40	0	2	30	0	3	20	0	4	10	0	5	0	0	5	50	0	6	40	0	7	30	0	8	20	0	9	10	0	10	0	0	
31	0	51	40	1	43	20	2	35	0	3	26	40	4	18	20	5	10	0	6	1	40	6	53	20	7	45	0	8	36	40	9	28	20	10	20	0	
32	0	53	20	1	46	40	2	40	0	3	33	20	4	26	40	5	20	0	6	13	20	7	6	40	8	0	0	8	53	20	9	46	40	10	40	0	
33	0	55	0	1	50	0	2	45	0	3	40	0	4	35	0	5	30	0	6	25	0	7	20	0	8	15	0	9	10	0	10	5	0	11	0	0	
34	0	56	40	1	53	20	2	50	0	3	46	40	4	43	20	5	40	0	6	36	40	7	33	20	8	30	0	9	26	40	10	23	20	11	20	0	
35	0	58	20	1	56	40	2	55	0	3	53	20	4	51	40	5	50	0	6	48	20	7	46	40	8	45	0	9	43	20	10	41	40	11	40	0	
36	1	0	0	2	0	0	3	0	0	4	0	0	5	0	0	6	0	0	7	0	0	8	0	0	9	0	0	10	0	0	11	0	0	12	0	0	
37	1	1	40	2	3	20	3	5	0	4	6	40	5	8	20	6	10	0	7	11	40	8	13	20	9	15	0	10	16	40	11	18	20	12	20	0	
38	1	3	20	2	6	40	3	10	0	4	13	20	5	16	40	6	20	0	7	23	20	8	26	40	9	30	0	10	33	20	11	36	40	12	40	0	
39	1	5	0	2	10	0	3	15	0	4	20	0	5	25	0	6	30	0	7	35	0	8	40	0	9	45	0	10	50	0	11	55	0	13	0	0	
40	1	6	40	2	13	20	3	20	0	4	26	40	5	33	20	6	40	0	7	46	40	8	53	20	10	0	0	11	6	40	12	13	20	13	20	0	
41	1	8	20	2	16	40	3	25	0	4	33	20	5	41	40	6	50	0	7	58	20	9	6	40	10	15	0	11	23	20	12	31	40	13	40	0	
42	1	10	0	2	20	0	3	30	0	4	40	0	5	50	0	7	0	0	8	10	0	9	20	0	10	30	0	11	40	0	12	50	0	14	0	0	
43	1	11	40	2	23	20	3	35	0	4	46	40	5	58	20	7	10	0	8	21	40	9	33	20													

CONSTRUCTION. As 360° to any diurnal Difference, so 10°, 20°, 30°, &c. Dif. Long. to the proportional Dif. to be added to or subtracted from the Quantity for the 1st Meridian, according as the Quantity for 2d Day is more or less than the Quantity for the 1st Meridian, the Longitude being West of 1st Meridian. Which proportional Difference is to be subtracted instead of added, or added instead of subtracted, when the Difference of Longitude is easterly from the 1st Meridian. See Bottom of p. 261 for Exam. here not being Room.

EQUATION of the DIFFERENCE of Right Ascensions, in Time, at Noon, between the SUN, MOON, PLANETS, and STARS, to the nearer Time of their respective Southing at Greenwich.

Dif.	Argument.	Difference of the Sun's R. A. in Time, at Noon, from the R. A. in Time of the Moon, a Planet or Star, at Noon.				CONSTRUCTION of the adjacent TABLE.			
R.A. at Noon.		δ	η	μ	ϕ	ψ	θ	κ	
		+	-	-	-	+	+	-	
h	m s	m s	m s	m s	m s	m s	m s	m s	
m	s th	s th	s th	s th	s th	s th	s th	s th	
s	th fo	th fo	th fo	th fo	th fo	th fo	th fo	th fo	
1	0 2 2	0 10	0 9	0 5	0 6	0 31	0 10		
2	0 4 4	0 19	0 18	0 9	0 12	1 2	0 20		
3	0 6 6	0 29	0 27	0 14	0 18	1 33	0 30		
4	0 8 8	0 38	0 36	0 18	0 25	2 4	0 39		
5	0 10 10	0 49	0 45	0 23	0 31	2 35	0 49		
6	0 12 11	0 57	0 54	0 28	0 37	3 6	0 59		
7	0 14 13	1 7	1 3	0 32	0 43	3 38	1 9		
8	0 16 15	1 16	1 12	0 37	0 49	4 9	1 19		
9	0 18 17	1 26	1 21	0 42	0 55	4 40	1 29		
10	0 20 19	1 35	1 30	0 46	1 2	5 11	1 39		
11	0 22 21	1 45	1 39	0 51	1 8	5 42	1 49		
12	0 24 23	1 54	1 48	0 55	1 14	6 13	1 58		
13	0 26 25	2 4	1 58	1 0	1 20	6 44	2 8		
14	0 28 27	2 14	2 7	1 5	1 26	7 15	2 18		
15	0 30 29	2 23	2 16	1 9	1 32	7 46	2 28		
16	0 32 31	2 33	2 25	1 14	1 39	8 17	2 38		
17	0 34 32	2 42	2 34	1 19	1 45	8 48	2 48		
18	0 36 34	2 52	2 43	1 23	1 51	9 19	2 58		
19	0 38 36	3 1	2 52	1 28	1 57	9 51	3 8		
20	0 40 38	3 10	3 1	1 32	2 3	10 22	3 17		
21	0 42 40	3 20	3 10	1 37	2 9	10 53	3 27		
22	0 44 42	3 29	3 19	1 42	2 16	11 24	3 37		
23	0 46 44	3 39	3 28	1 46	2 22	11 55	3 47		
24	0 48 46	3 49	3 37	1 51	2 28	12 26	3 57		
25	0 50 48	3 59	3 46	1 56	2 34	12 57	4 7		
26	0 52 50	4 8	3 55	2 0	2 40	13 28	4 17		
27	0 54 51	4 18	4 4	2 5	2 46	13 59	4 27		
28	0 56 53	4 27	4 13	2 9	2 53	14 30	4 36		
29	0 58 55	4 38	4 22	2 14	2 59	15 1	4 46		
30	1 0 57	4 46	4 31	2 19	3 5	15 32	4 56		
31	1 2 59	4 56	4 40	2 23	3 11	16 4	5 6		
32	1 5 2	5 5	4 49	2 28	3 17	16 35	5 16		
33	1 7 3	5 15	4 58	2 33	3 23	17 6	5 26		
34	1 9 5	5 24	5 7	2 37	3 30	17 37	5 35		
35	1 11 7	5 34	5 16	2 42	3 36	18 8	5 45		
36	1 13 8	5 43	5 25	2 46	3 42	18 39	5 55		
37	1 15 10	5 53	5 35	2 51	3 48	19 10	6 5		
38	1 17 12	6 3	5 44	2 56	3 54	19 41	6 15		
39	1 19 14	6 12	5 53	3 0	4 0	20 12	6 25		
40	1 21 16	6 22	6 2	3 5	4 7	20 43	6 35		
41	1 23 18	6 31	6 11	3 10	4 13	21 14	6 45		
42	1 25 20	6 41	6 20	3 14	4 19	21 45	6 54		
43	1 27 22	6 50	6 29	3 19	4 25	22 17	7 4		
44	1 29 24	6 59	6 38	3 23	4 31	22 48	7 14		
45	1 31 26	7 9	6 47	3 28	4 37	23 19	7 24		
46	1 33 28	7 18	6 56	3 33	4 44	23 50	7 34		
47	1 35 29	7 28	7 5	3 37	4 50	24 21	7 44		
48	1 37 32	7 38	7 14	3 42	4 56	24 52	7 54		
49	1 39 4	7 48	7 23	3 47	5 2	25 23	8 4		
50	1 41 6	7 57	7 32	3 51	5 8	25 54	8 13		
51	1 43 8	8 7	7 41	3 56	5 14	26 25	8 23		
52	1 45 10	8 16	7 50	4 0	5 21	26 56	8 33		
53	1 47 12	8 27	7 59	4 5	5 27	27 27	8 43		
54	1 49 14	8 35	8 8	4 10	5 33	27 58	8 53		
55	1 51 16	8 45	8 17	4 14	5 39	28 30	9 3		
56	1 53 17	8 54	8 26	4 19	5 45	29 1	9 13		
57	1 55 19	9 4	8 35	4 24	5 51	29 32	9 23		
58	1 57 21	9 13	8 44	4 28	5 58	29 43	9 32		
59	1 59 23	9 23	8 53	4 33	6 4	30 34	9 42		
60	2 1 25	9 32	9 2	4 37	6 10	31 5	9 52		

N. B. The above Table also finds the Dif. of Time to be added to or subtracted from the Time of Southing at the first Meridian, for the Time of Southing at any Place of given East or West Longitude, respectively, the Sun's being the swifter Motion; but Dif. of Time to be subtracted and added for East and West Long. in Time, the Sun's being the slower Motion; add for Southing of δ , η , and ψ ; and sub. for μ , ϕ , and θ ; and the contrary Long. E. The Equation being taken from the Hours, Minutes, and Seconds, Difference of Longitude, East or West from the first Meridian.

CONSTRUCTION of the adjacent TABLE.

M. Diurnal Motion.	D. Mot. Time.	
δ 12° 11' 26" 41"	48m 46s	Taken as the mean
η 0 57 7 46	3 49	Differences
μ 0 54 9 2	3 37	of R. A. of
ϕ 0 27 41 40	1 51	those Bodies
ψ 0 36 59 29	2 28	for 24 Hours
θ 3 6 24 14	12 26	respectively.
κ 0 59 8 10	3 57	

From which diurnal Differences of mean Motion, in Time, taken as mean Differences of right Ascension, the Quantities of Equation to the first Time of near Southing, is respectively proportioned, in the several adjacent Columns.

RULE. These Equations add to or subtract from the Time of near Southing, at Greenwich, for the nearer Time there, according as the Motion of the celestial Body (marked at the Head of each Column) is swifter or slower than the Sun's Motion, respectively.

EXAMPLE I. To find the nearer Time of the Moon's Southing, from the Difference of right Ascension of the Sun, from right Ascension of the Moon, at Noon, being given?

By p. 238. δ R. A. — κ R. A. 1762, } h m s
 July 10, Noon, Greenwich . . . } 15 16 3
 By Table on the Left 15h . . . Equation + 30 29
 16m . . . Equation + 0 33

δ 's nearer Time of Southing at Greenwich, 1762, July 10, P. M. . . } 15 47 5
 Being an Improvement on Mr. BRENT'S METHOD.
 Correct Time of δ 's Southing, P. M. at } 15 41 18
 Greenwich, by the same Page 238 . . }

Error . . — 5m 47s

This METHOD of finding the near Time of the Moon's Southing never differs from the Truth but in a few Minutes. As the same Method finds the Southing of the Stars within a few Seconds of Time; and also the Southing of all the Planets scarcely differing a Minute from Truth.

EXAMPLE II. To find the nearer Time of Aldebaran's Southing from the Difference of right Ascension of the Sun from right Ascension of Aldebaran, at Noon, being given?

By p. 218. R. A. of Aldebaran — R. A. } h m s
 of κ or + Com. R. A. κ . 1760, }
 Dec. 30, Noon . . . } 9 41 25

By Tab. on the Left 9h . . . Equation — 1 29
 41m . . . Equation — 0 7

Total Equation — 1 36
 Rem. Near Time Southing of Aldebaran } 9 39 49
 1760, Dec. 30, at Greenwich, P. M. }
 Correct Time of Aldebaran's Southing, by } 9 39 38
 same Page 218 . . . }

Error — 11s

Antigua 4h West Long. . . : Equation — 39s
 correspondent to Aldeb. Southing at Greenwich.

Aldebaran nearly souths at Antigua . . . 9 39 10
 Correctly souths at Antigua, by p. 218 . . . 9 38 54

The same Method holds } Error + 16s
 good for the Planets. }

HERE FOLLOW

LOGISTICAL LOGARITHMS.

The NATURE and USE of LOGISTICAL LOGARITHMS.

THE Difference between the common Logarithm of 3600" (being the Seconds in a Degree) and the common Logarithm of any other Number of Seconds, under or above 3600, is called the Logistical Logarithm of that Number of Seconds. These logistical Logarithms begin at 60', and increase negatively under and affirmatively above 60'.

The principal Use of these logistical Logarithms is for the ready proportioning Sexagesimal Quantities of all Kinds; and especially those in the Ratio of 1 Degree Increase of Argument to a given Increase or Decrease of Equation, correspondent: though they have all the Use of common Logarithms as far as the Number of Seconds in the following Table (p. 267) extend.

The DETERMINATION and USE of the logistical Logarithms from the common LOGARITHMS.

EXAMPLE I. To find the Equation of the SUN's Center for 1st 6^o 27' 38" mean Anomaly of the Sun?

Argument.	Equat. ☉'s Cent.	As 60' 0" = 3600"	Com. Log.	Same	Com. Log.	Lo. Log.	extra	As 60' 0"
1 st 6 ^o	1 ^o 6' 57"		3.5563025	As 1	L. 1	0.0000	000	As 60' 0"
1 7	1 8 33							
		To +1 36 = 96	1.9822712	To 96	L. 96—L. 3600	—1.5740	313	To 1 36
Incr. 1 ^o	Incr. 1' 36"	So 27 38 = 1658	3.2195845	To 3600				
		To +0 44 = 44	1.6455532	So 1658	L. 1658—L. 3600	—0.3367	180	So 27 38
		First Equation 1 ^o 6' 57"		To 44	L. 44—L. 3600	—1.9107	493	To 0 44
		Equation of Sun's Center 1 ^o 7' 41" reqd.		To 3600				

EXAMPLE II. To find the SUN's Right Ascension answerable to the Sun's Place 27^o 37' 41"?

Argument.	R. A. Sun.	As 60' 0" = 3600"	Com. Log.	Same	Com. Log.	Lo. Log.	extra	As 60' 0"
27 ^o	266 ^o 43' 47"		3.5563025	As 1	L. 1	0.0000	000	As 60' 0"
28	267 49 11							
		To 65 24 = 3924	3.5937290	To 3924	L. 3924—L. 3600, co.	—9.9625	735	65 24
Incr. 1 ^o	Incr. 1 5 24	So 37 41 = 2261	3.3543006	To 3600				
		To 41 4 = 2464	3.3917271	So 2261	L. 2261—L. 3600	—0.2020	019	37 41
		First Equation 266 ^o 43' 47"		To 2464	L. 2464—L. 3600	—0.1645	754	41 4
		R. A. Sun 267 24 51 reqd.		To 3600				

By the above two Operations it appears, that the logistical Logarithms, for all Sexagesimals, under 60', come out *negative; and all logistical Logarithms for Sexagesimals above 60' come out affirmative, or with contrary Signs to one another. So that if the logistical Logarithms for Quantities under 60' be considered affirmative (as they are considered in the logistical Tables in general) the logistical Logarithms for Quantities above 60' must be considered as negative; and therefore we have given their arithmetical Complement, to make them correspond with the logistical Logarithms for Quantities under 60'; only in taking them out, you must remember to prefix the Index +9 or 1 to all logistical Logarithms for Numbers above 60'; the Indices for all Numbers under 60' being given in the following Table; viz. below 60', down to 6', the Index is 0; and below 6' down to 1", the Indices are either 1, 2, or 3, as expressed. So that you have Nothing to do but use the logistical Tables like the common Tables of Logarithms, in taking out the Logarithms to the Numbers, and finding the Number correspondent to any Logarithm.

TO find the logistical Logarithm of any Number of Degrees and Minutes, not exceeding 90^o, or of Minutes and Seconds, not above 90', or of any Number not exceeding 5400?

AT the Top, or Bottom of the Table find the Degrees or Minutes, and in the same Column against the Minutes or Seconds, respectively, on the Side of the Page, you will find the logistical Logarithm, required. — In the second Row, across the Page, from the Top, find the Number next less than the Number given, and in the same Column, against the Difference greater than that Number, to make up the Number sought, on the Side of the Page, you will find the logistical Logarithm thereof, required.

When two of the given Terms are greater than 5400, take their Halves, Thirds, &c. being in the same Ratio. — But if only one of the given Terms exceed 5400, take its Half, Third, &c. and multiply the 4th Term resulting, by 2, 3, &c. to correspond with the given Ratio.

If Hours are in the Terms of the Proportion, their logistical Logarithms may be taken out of the Table as Minutes, and the logistical Logarithm to the Minutes of an Hour may be taken out as for Seconds.

The log. Logarithms of two small Arcs (being nearly as their Sines or Tangents) may be used in trigonometrical Proportions, with the common Log. Sines, or Tangents, of two of the other Terms, being large Arcs; but then instead of the Log. Sines or Tangents of those large Arcs you must take their arithmetical Complements, and the contrary; because (properly) the logistical Logarithms in our Table are all negative, (but being all of one Kind are considered affirmative) and so in using them with affirmative common Logarithms the latter must be reduced to Negatives. Or else our negative logistical Logarithms must be reduced to Affirmatives, (like common Logarithms) by taking their arithmetical Complements, when they are used with common Logarithms, without taking their arithmetical Complements. But this Proportion may be as speedily resolved by common Logarithm Sines to every Second of small Arcs (serving indifferently for Log. Tangents of those Arcs) with the Log. Sines or Tangents of large Arcs.

EXAMPLES.

As 60' 0"	Lo. Log.	As 70 ^o 19' co.	L. Log.	As 24 ^h co.	Lo. Log.	As 1 ^o 0' co.	Lo. Log.	As 4721 co.	Lo. Log.
To 75 18	1.9014	To 34 50	0.2362	To 76' 34"	1.8941	To 13' 7"	0.6603	To 60' . . .	0.0000
So 76 11	1.8963	So 29 31	0.3081	So 13 ^h 53 ^m	0.6357	So 6 ^o 19'	0.9777	So 2948	0.0868
To 95 36	1.7977	To 14 37	0.6132	To 44' 17"	0.1319	To 82' 52"	9.8598	To 37' 28"	0.2045
Or, As 60' 0"	0.0000	As 15' 37" co.	1.4154	As S. 20' 35" co.	2.2227608	As 20' 35" co.	1.5354	As 20' 35"	0.4646
To 75 18	1.9014	To 8 ^h 7'	0.8688	To S. 14' 27"	7.6235927	To 14' 27"	0.6183	To 14 27 co	1.3817
So 1 ^o or 38 51	0.1973	So 46' 11"	0.1137	So S. 37 ^o 19'	9.7826301	So S. 37 ^o 19' co.	0.2173	So S. 37 ^o 19'	9.7826
To 1 ^o or 47 48	0.0987	To 24 ^h 0	0.3979	To S. 25 ^o 11'	9.6289836	To S. 25 ^o 11' co.	0.3710	To S. 25 11	9.6289
Doub. 95 36 as above.						S. 9.6290			

LOGISTICAL

LOGISTICAL LOGARITHMS.

0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	0	
1	0	60	120	180	240	300	360	420	480	540	600	660	720	780	840	1
0		1.7782	1.4771	1.3010	1.1761	1.0792	0000	9331	8751	8239	7782	7368	6990	6642	6320	0
1	3.5563	1.7710	1.4735	1.2986	1.1743	1.0777	9988	9320	8742	8231	7774	7361	6984	6637	6315	1
2	3.2553	1.7639	1.4699	1.2962	1.1725	1.0763	9976	9310	8733	8223	7767	7354	6978	6631	6310	2
3	3.0792	1.7570	1.4664	1.2939	1.1707	1.0749	9964	9300	8724	8215	7760	7348	6972	6625	6305	3
4	2.9542	1.7501	1.4629	1.2915	1.1689	1.0734	9952	9289	8715	8207	7753	7341	6966	6620	6300	4
5	2.8573	1.7434	1.4594	1.2891	1.1671	1.0720	9940	9279	8706	8199	7745	7335	6960	6614	6294	5
6	2.7782	1.7368	1.4559	1.2868	1.1654	1.0706	9928	9269	8697	8191	7738	7328	6954	6609	6289	6
7	2.7112	1.7302	1.4525	1.2845	1.1636	1.0692	9916	9259	8688	8183	7731	7322	6948	6603	6284	7
8	2.6532	1.7238	1.4491	1.2821	1.1619	1.0678	9905	9249	8679	8175	7724	7315	6942	6598	6279	8
9	2.6021	1.7175	1.4457	1.2798	1.1601	1.0663	9893	9238	8670	8167	7717	7309	6936	6592	6274	9
10	2.5563	1.7112	1.4424	1.2775	1.1584	1.0649	9881	9228	8661	8159	7710	7302	6930	6587	6269	10
11	2.5149	1.7050	1.4390	1.2753	1.1566	1.0635	9869	9218	8652	8152	7703	7296	6924	6581	6264	11
12	2.4771	1.6990	1.4357	1.2730	1.1549	1.0621	9858	9208	8643	8144	7696	7289	6918	6576	6259	12
13	2.4424	1.6930	1.4325	1.2707	1.1532	1.0608	9846	9198	8635	8136	7688	7283	6912	6570	6254	13
14	2.4102	1.6871	1.4292	1.2685	1.1515	1.0594	9834	9188	8626	8128	7681	7276	6906	6565	6248	14
15	2.3802	1.6812	1.4260	1.2663	1.1498	1.0580	9823	9178	8617	8120	7674	7270	6900	6559	6243	15
16	2.3522	1.6755	1.4228	1.2640	1.1481	1.0566	9811	9168	8608	8112	7669	7264	6894	6554	6238	16
17	2.3259	1.6698	1.4196	1.2618	1.1464	1.0552	9800	9158	8599	8104	7660	7257	6888	6548	6233	17
18	2.3010	1.6642	1.4165	1.2596	1.1447	1.0539	9788	9148	8591	8097	7653	7251	6882	6543	6228	18
19	2.2775	1.6587	1.4133	1.2574	1.1430	1.0525	9777	9138	8582	8089	7646	7244	6877	6538	6223	19
20	2.2553	1.6532	1.4102	1.2553	1.1413	1.0512	9765	9128	8573	8081	7639	7238	6871	6532	6218	20
21	2.2341	1.6478	1.4071	1.2531	1.1397	1.0498	9754	9119	8565	8073	7632	7232	6865	6527	6213	21
22	2.2139	1.6425	1.4040	1.2510	1.1380	1.0484	9742	9109	8556	8066	7625	7225	6859	6521	6208	22
23	2.1946	1.6372	1.4010	1.2488	1.1363	1.0471	9731	9099	8547	8058	7618	7219	6853	6516	6203	23
24	2.1761	1.6320	1.3979	1.2467	1.1347	1.0458	9720	9089	8539	8050	7611	7212	6847	6510	6198	24
25	2.1584	1.6269	1.3949	1.2445	1.1331	1.0444	9708	9079	8530	8043	7604	7206	6841	6505	6193	25
26	2.1413	1.6218	1.3919	1.2424	1.1314	1.0431	9697	9070	8522	8035	7597	7200	6836	6500	6188	26
27	2.1249	1.6168	1.3890	1.2403	1.1298	1.0418	9686	9060	8513	8027	7590	7193	6830	6494	6183	27
28	2.1091	1.6118	1.3860	1.2382	1.1282	1.0404	9675	9050	8504	8020	7583	7187	6824	6489	6178	28
29	2.0939	1.6069	1.3831	1.2362	1.1266	1.0391	9664	9041	8496	8012	7577	7181	6818	6484	6173	29
30	2.0792	1.6021	1.3802	1.2341	1.1249	1.0378	9652	9031	8487	8004	7570	7175	6812	6478	6168	30
31	2.0649	1.5973	1.3773	1.2320	1.1233	1.0365	9641	9021	8479	7997	7563	7168	6807	6473	6163	31
32	2.0512	1.5925	1.3745	1.2300	1.1217	1.0352	9630	9012	8470	7989	7556	7162	6801	6467	6158	32
33	2.0378	1.5878	1.3716	1.2279	1.1201	1.0339	9619	9002	8462	7981	7549	7156	6795	6462	6153	33
34	2.0248	1.5832	1.3688	1.2259	1.1186	1.0326	9608	8992	8453	7974	7542	7149	6789	6457	6148	34
35	2.0122	1.5786	1.3660	1.2239	1.1170	1.0313	9597	8983	8445	7966	7535	7143	6784	6451	6143	35
36	2.0000	1.5740	1.3632	1.2218	1.1154	1.0300	9586	8973	8437	7959	7528	7137	6778	6446	6138	36
37	1.9881	1.5695	1.3604	1.2198	1.1138	1.0287	9575	8964	8428	7951	7522	7131	6772	6441	6133	37
38	1.9765	1.5651	1.3576	1.2178	1.1123	1.0274	9564	8954	8420	7944	7515	7124	6766	6435	6128	38
39	1.9652	1.5607	1.3549	1.2159	1.1107	1.0261	9553	8945	8411	7936	7508	7118	6761	6430	6123	39
40	1.9542	1.5563	1.3522	1.2139	1.1091	1.0248	9542	8935	8403	7929	7501	7112	6755	6425	6118	40
41	1.9435	1.5520	1.3495	1.2119	1.1076	1.0235	9532	8926	8395	7921	7494	7106	6749	6420	6113	41
42	1.9331	1.5477	1.3468	1.2099	1.1061	1.0223	9521	8917	8386	7914	7488	7100	6743	6414	6108	42
43	1.9228	1.5435	1.3441	1.2080	1.1045	1.0210	9510	8907	8378	7906	7481	7093	6738	6409	6103	43
44	1.9128	1.5393	1.3415	1.2061	1.1030	1.0197	9499	8898	8370	7899	7474	7087	6732	6404	6099	44
45	1.9031	1.5351	1.3388	1.2041	1.1015	1.0185	9488	8888	8361	7891	7467	7081	6726	6398	6094	45
46	1.8935	1.5310	1.3362	1.2022	1.0999	1.0172	9478	8879	8353	7884	7461	7075	6721	6393	6089	46
47	1.8842	1.5269	1.3336	1.2003	1.0984	1.0160	9467	8870	8345	7877	7454	7069	6715	6388	6084	47
48	1.8751	1.5229	1.3310	1.1984	1.0969	1.0147	9456	8861	8337	7869	7447	7063	6709	6383	6079	48
49	1.8661	1.5189	1.3284	1.1965	1.0954	1.0135	9446	8851	8328	7862	7441	7057	6704	6377	6074	49
50	1.8573	1.5149	1.3259	1.1946	1.0939	1.0122	9435	8842	8320	7855	7434	7050	6698	6372	6069	50
51	1.8487	1.5110	1.3233	1.1927	1.0924	1.0110	9425	8833	8312	7847	7427	7044	6692	6367	6064	51
52	1.8403	1.5071	1.3208	1.1908	1.0909	1.0098	9414	8824	8304	7840	7421	7038	6687	6362	6059	52
53	1.8320	1.5032	1.3183	1.1889	1.0894	1.0085	9404	8814	8296	7832	7414	7032	6681	6357	6055	53
54	1.8239	1.4994	1.3158	1.1871	1.0880	1.0073	9393	8805	8288	7825	7407	7026	6676	6351	6050	54
55	1.8159	1.4956	1.3133	1.1852	1.0865	1.0061	9383	8796	8279	7818	7401	7020	6670	6346	6045	55
56	1.8081	1.4918	1.3108	1.1834	1.0850	1.0049	9372	8787	8271	7811	7394	7014	6664	6341	6040	56
57	1.8004	1.4881	1.3083	1.1816	1.0835	1.0036	9362	8778	8263	7803	7387	7008	6659	6336	6035	57
58	1.7929	1.4844	1.3059	1.1797	1.0821	1.0024	9351	8769	8255	7796	7381	7002	6653	6331	6030	58
59	1.7855	1.4808	1.3034	1.1779	1.0806	1.0012	9341	8760	8247	7789	7374	6996	6648	6325	6025	59
60	1.7782	1.4771	1.3010	1.1761	1.0792	1.0000	9331	8751	8239	7782	7368	6990	6642	6320	6021	60
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	

LOGISTICAL LOGARITHMS.

°	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	°
15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	15	
0	6021	5740	5477	5229	4994	4771	4559	4357	4164	3979	3802	3632	3468	3310	3158	0
1	6016	5736	5473	5225	4990	4768	4556	4354	4161	3976	3799	3629	3465	3307	3155	1
2	6011	5731	5469	5221	4986	4764	4552	4351	4158	3973	3796	3626	3463	3305	3153	2
3	6006	5727	5464	5217	4983	4760	4549	4347	4155	3970	3793	3623	3460	3302	3150	3
4	6001	5722	5460	5213	4979	4757	4546	4344	4152	3967	3791	3621	3457	3300	3148	4
5	5997	5718	5456	5209	4975	4753	4542	4341	4149	3964	3788	3618	3454	3297	3145	5
6	5992	5713	5452	5205	4971	4750	4539	4338	4145	3961	3785	3615	3452	3294	3143	6
7	5987	5709	5447	5201	4967	4746	4535	4334	4142	3958	3782	3612	3449	3292	3140	7
8	5982	5704	5443	5197	4964	4742	4532	4331	4139	3955	3779	3610	3446	3289	3138	8
9	5977	5700	5439	5193	4960	4739	4528	4328	4136	3952	3776	3607	3444	3287	3135	9
10	5973	5695	5435	5189	4956	4735	4525	4325	4133	3949	3773	3604	3441	3284	3133	10
11	5968	5691	5430	5185	4952	4732	4522	4321	4130	3946	3770	3601	3438	3282	3130	11
12	5963	5686	5426	5181	4949	4728	4518	4318	4127	3943	3768	3598	3436	3279	3128	12
13	5958	5682	5422	5177	4945	4724	4515	4315	4124	3940	3765	3596	3433	3276	3125	13
14	5954	5677	5418	5173	4941	4721	4511	4311	4120	3937	3762	3593	3431	3274	3123	14
15	5949	5673	5414	5169	4937	4717	4508	4308	4117	3934	3759	3590	3428	3271	3120	15
16	5944	5669	5409	5165	4933	4714	4505	4305	4114	3931	3756	3587	3425	3269	3118	16
17	5939	5664	5405	5161	4930	4710	4501	4302	4111	3928	3753	3585	3423	3266	3115	17
18	5935	5660	5401	5157	4926	4707	4498	4298	4108	3925	3750	3582	3420	3264	3113	18
19	5930	5655	5397	5153	4922	4703	4494	4295	4105	3922	3747	3579	3417	3261	3110	19
20	5925	5651	5393	5149	4918	4699	4491	4292	4102	3919	3745	3576	3415	3259	3108	20
21	5920	5646	5389	5145	4915	4696	4488	4289	4099	3917	3742	3574	3412	3256	3105	21
22	5916	5642	5384	5141	4911	4692	4484	4285	4096	3914	3739	3571	3409	3253	3103	22
23	5911	5637	5380	5137	4907	4689	4481	4282	4092	3911	3736	3568	3407	3251	3101	23
24	5906	5633	5376	5133	4903	4685	4477	4279	4089	3908	3733	3565	3404	3248	3098	24
25	5902	5629	5372	5129	4900	4682	4474	4276	4086	3905	3730	3563	3401	3246	3096	25
26	5897	5624	5368	5125	4896	4678	4471	4273	4083	3902	3727	3560	3399	3243	3093	26
27	5892	5620	5364	5122	4892	4675	4467	4269	4080	3899	3725	3557	3396	3241	3091	27
28	5888	5615	5359	5118	4889	4671	4464	4266	4077	3896	3722	3555	3393	3238	3088	28
29	5883	5611	5355	5114	4885	4668	4460	4263	4074	3893	3719	3552	3391	3236	3086	29
30	5878	5607	5351	5110	4881	4664	4457	4260	4071	3890	3716	3549	3388	3233	3083	30
31	5874	5602	5347	5106	4877	4660	4454	4256	4068	3887	3713	3546	3386	3231	3081	31
32	5869	5598	5343	5102	4874	4657	4450	4253	4065	3884	3710	3544	3383	3228	3078	32
33	5864	5594	5339	5098	4870	4653	4447	4250	4062	3881	3708	3541	3380	3225	3076	33
34	5860	5589	5335	5094	4866	4650	4444	4247	4059	3878	3705	3538	3378	3223	3073	34
35	5855	5585	5331	5090	4863	4646	4440	4244	4055	3875	3702	3535	3375	3220	3071	35
36	5850	5580	5326	5086	4859	4643	4437	4240	4052	3872	3699	3533	3372	3218	3069	36
37	5846	5576	5322	5082	4855	4639	4434	4237	4049	3869	3696	3530	3370	3215	3066	37
38	5841	5572	5318	5079	4852	4636	4430	4234	4046	3866	3693	3527	3367	3213	3064	38
39	5836	5567	5314	5075	4848	4632	4427	4231	4043	3863	3691	3525	3365	3210	3061	39
40	5832	5563	5310	5071	4844	4629	4424	4228	4040	3860	3688	3522	3362	3208	3059	40
41	5827	5559	5306	5067	4841	4625	4420	4224	4037	3857	3685	3519	3359	3205	3056	41
42	5823	5554	5302	5063	4837	4622	4417	4221	4034	3855	3682	3516	3357	3203	3054	42
43	5818	5550	5298	5059	4833	4618	4414	4218	4031	3852	3679	3514	3354	3200	3052	43
44	5813	5546	5294	5055	4830	4615	4410	4215	4028	3849	3677	3511	3351	3198	3049	44
45	5809	5541	5290	5051	4826	4611	4407	4212	4025	3846	3674	3508	3349	3195	3047	45
46	5804	5537	5285	5048	4822	4608	4404	4209	4022	3843	3671	3506	3346	3193	3044	46
47	5800	5533	5281	5044	4819	4604	4400	4205	4019	3840	3668	3503	3344	3190	3042	47
48	5795	5528	5277	5040	4815	4601	4397	4202	4016	3837	3665	3500	3341	3188	3039	48
49	5790	5524	5273	5036	4811	4597	4394	4199	4013	3834	3663	3497	3338	3185	3037	49
50	5786	5520	5269	5032	4808	4594	4390	4196	4010	3831	3660	3495	3336	3183	3034	50
51	5781	5516	5265	5028	4804	4590	4387	4193	4007	3828	3657	3492	3333	3180	3032	51
52	5777	5511	5261	5025	4800	4587	4384	4189	4004	3825	3654	3489	3331	3178	3030	52
53	5772	5507	5257	5021	4797	4584	4380	4186	4001	3822	3651	3487	3328	3175	3027	53
54	5768	5503	5253	5017	4793	4580	4377	4183	3998	3820	3649	3484	3325	3173	3025	54
55	5763	5498	5249	5013	4789	4577	4374	4180	3995	3817	3646	3481	3323	3170	3022	55
56	5758	5494	5245	5009	4786	4573	4370	4177	3991	3814	3643	3479	3320	3168	3020	56
57	5754	5490	5241	5005	4782	4570	4367	4174	3988	3811	3640	3476	3318	3165	3018	57
58	5749	5486	5237	5002	4778	4566	4364	4171	3985	3808	3637	3473	3315	3163	3015	58
59	5745	5481	5233	4998	4775	4563	4361	4167	3982	3805	3635	3471	3313	3160	3013	59
60	5740	5477	5229	4994	4771	4559	4357	4164	3979	3802	3632	3468	3310	3158	3010	60
15	16	17	18	19	20	21	22	23	24	25	26	27	28	29		

LOGISTICAL LOGARITHMS.

°	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	'
1	1800	1860	1920	1980	2040	2100	2160	2220	2280	2340	2400	2460	2520	2580	2640	1
0	3010	2868	2730	2596	2467	2341	2218	2099	1984	1871	1761	1654	1549	1447	1347	0
1	3008	2866	2728	2594	2465	2339	2216	2098	1982	1869	1759	1652	1547	1445	1345	1
2	3005	2863	2725	2592	2462	2337	2214	2096	1980	1867	1757	1650	1546	1443	1344	2
3	3003	2861	2723	2590	2460	2335	2212	2094	1978	1865	1755	1648	1544	1442	1342	3
4	3001	2859	2721	2588	2458	2333	2210	2092	1976	1863	1754	1647	1542	1440	1340	4
5	2998	2856	2719	2585	2456	2331	2208	2090	1974	1862	1752	1645	1540	1438	1339	5
6	2996	2854	2716	2583	2454	2328	2206	2088	1972	1860	1750	1643	1539	1437	1337	6
7	2993	2852	2714	2581	2452	2326	2204	2086	1970	1858	1748	1641	1537	1435	1335	7
8	2991	2849	2712	2579	2450	2324	2202	2084	1968	1856	1746	1640	1535	1433	1334	8
9	2989	2847	2710	2577	2448	2322	2200	2082	1967	1854	1745	1638	1534	1432	1332	9
10	2986	2845	2707	2574	2445	2320	2198	2080	1965	1852	1743	1636	1532	1430	1331	10
11	2984	2842	2705	2572	2443	2318	2196	2078	1963	1850	1741	1634	1530	1428	1329	11
12	2981	2840	2703	2570	2441	2316	2194	2076	1961	1849	1739	1633	1528	1427	1327	12
13	2979	2838	2701	2568	2439	2314	2192	2074	1959	1847	1737	1631	1527	1425	1326	13
14	2977	2835	2698	2566	2437	2312	2190	2072	1957	1845	1736	1629	1525	1423	1324	14
15	2974	2833	2696	2564	2435	2310	2188	2070	1955	1843	1734	1627	1523	1422	1322	15
16	2972	2831	2694	2561	2433	2308	2186	2068	1953	1841	1732	1626	1522	1420	1321	16
17	2969	2828	2692	2559	2431	2306	2184	2066	1951	1839	1730	1624	1520	1418	1319	17
18	2967	2826	2689	2557	2429	2304	2182	2064	1950	1838	1728	1622	1518	1417	1317	18
19	2965	2824	2687	2555	2426	2302	2180	2062	1948	1836	1727	1620	1516	1415	1316	19
20	2962	2821	2685	2553	2424	2300	2178	2061	1946	1834	1725	1619	1515	1413	1314	20
21	2960	2819	2683	2551	2422	2298	2176	2059	1944	1832	1723	1617	1513	1412	1313	21
22	2958	2817	2681	2548	2420	2296	2174	2057	1942	1830	1721	1615	1511	1410	1311	22
23	2955	2815	2678	2546	2418	2294	2172	2055	1940	1828	1719	1613	1510	1408	1309	23
24	2953	2812	2676	2544	2416	2291	2170	2053	1938	1827	1718	1612	1508	1407	1308	24
25	2950	2810	2674	2542	2414	2289	2169	2051	1936	1825	1716	1610	1506	1405	1306	25
26	2948	2808	2672	2540	2412	2287	2167	2049	1934	1823	1714	1608	1504	1403	1304	26
27	2946	2805	2669	2538	2410	2285	2165	2047	1933	1821	1712	1606	1503	1402	1303	27
28	2943	2803	2667	2535	2408	2283	2163	2045	1931	1819	1711	1605	1501	1400	1301	28
29	2941	2801	2665	2533	2405	2281	2161	2043	1929	1817	1709	1603	1499	1398	1300	29
30	2939	2798	2663	2531	2403	2279	2159	2041	1927	1816	1707	1601	1498	1397	1298	30
31	2936	2796	2660	2529	2401	2277	2157	2039	1925	1814	1705	1599	1496	1395	1296	31
32	2934	2794	2658	2527	2399	2275	2155	2037	1923	1812	1703	1598	1494	1393	1295	32
33	2931	2792	2656	2525	2397	2273	2153	2035	1921	1810	1702	1596	1493	1392	1293	33
34	2929	2789	2654	2522	2395	2271	2151	2033	1919	1808	1700	1594	1491	1390	1291	34
35	2927	2787	2652	2520	2393	2269	2149	2032	1918	1806	1698	1592	1489	1388	1290	35
36	2924	2785	2649	2518	2391	2267	2147	2030	1916	1805	1696	1591	1487	1387	1288	36
37	2922	2782	2647	2516	2389	2265	2145	2028	1914	1803	1694	1589	1486	1385	1287	37
38	2920	2780	2645	2514	2387	2263	2143	2026	1912	1801	1693	1587	1484	1383	1285	38
39	2917	2778	2643	2512	2384	2261	2141	2024	1910	1799	1690	1585	1482	1382	1283	39
40	2915	2775	2640	2510	2382	2259	2139	2022	1908	1797	1689	1584	1481	1380	1282	40
41	2912	2773	2638	2507	2380	2257	2137	2020	1906	1795	1687	1582	1479	1378	1280	41
42	2910	2771	2636	2505	2378	2255	2135	2018	1904	1794	1686	1580	1477	1377	1278	42
43	2908	2769	2634	2503	2376	2253	2133	2016	1903	1792	1684	1578	1476	1375	1277	43
44	2905	2766	2632	2501	2374	2251	2131	2014	1901	1790	1682	1577	1474	1373	1275	44
45	2903	2764	2629	2499	2372	2249	2129	2012	1899	1788	1680	1575	1472	1372	1274	45
46	2901	2762	2627	2497	2370	2247	2127	2010	1897	1786	1678	1573	1470	1370	1272	46
47	2898	2760	2625	2494	2368	2245	2125	2009	1895	1785	1677	1571	1469	1368	1270	47
48	2896	2757	2623	2492	2366	2243	2123	2007	1893	1783	1675	1570	1467	1367	1269	48
49	2894	2755	2621	2490	2364	2241	2121	2005	1891	1781	1673	1568	1465	1365	1267	49
50	2891	2753	2618	2488	2362	2239	2119	2003	1889	1779	1671	1566	1464	1363	1266	50
51	2889	2750	2616	2486	2359	2237	2117	2001	1888	1777	1670	1565	1462	1362	1264	51
52	2887	2748	2614	2484	2357	2235	2115	1999	1886	1775	1668	1563	1460	1360	1262	52
53	2884	2746	2612	2482	2355	2233	2113	1997	1884	1774	1666	1561	1459	1359	1261	53
54	2882	2744	2610	2480	2353	2231	2111	1995	1882	1772	1664	1559	1457	1357	1259	54
55	2880	2741	2607	2477	2351	2229	2109	1993	1880	1770	1663	1558	1455	1355	1257	55
56	2877	2739	2605	2475	2349	2227	2107	1991	1878	1768	1661	1556	1454	1354	1256	56
57	2875	2737	2603	2473	2347	2225	2105	1989	1876	1766	1659	1554	1452	1352	1254	57
58	2873	2735	2601	2471	2345	2223	2103	1987	1875	1765	1657	1552	1450	1350	1253	58
59	2870	2732	2599	2469	2343	2220	2101	1986	1873	1763	1655	1551	1449	1349	1251	59
60	2868	2730	2596	2467	2341	2218	2099	1984	1871	1761	1654	1549	1447	1347	1249	60
	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	

LOGISTICAL LOGARITHMS.

°	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	'
"	2700	2760	2820	2880	2940	3000	3060	3120	3180	3240	3300	3360	3420	3480	3540	"
0	1249	1154	1061	0969	0880	0792	0706	0621	0539	0458	0378	0300	0223	0147	0073	0
1	1248	1152	1059	0968	0878	0790	0704	0620	0537	0456	0377	0298	0221	0146	0072	1
2	1246	1151	1057	0966	0877	0789	0703	0619	0536	0455	0375	0297	0220	0145	0071	2
3	1245	1149	1056	0965	0875	0787	0702	0617	0535	0454	0374	0296	0219	0143	0069	3
4	1243	1148	1054	0963	0874	0786	0700	0616	0533	0452	0373	0294	0218	0142	0068	4
5	1241	1146	1053	0962	0872	0785	0699	0615	0532	0451	0371	0293	0216	0141	0067	5
6	1240	1145	1051	0960	0871	0783	0697	0613	0531	0450	0370	0292	0215	0140	0066	6
7	1238	1143	1050	0959	0869	0782	0696	0612	0529	0448	0369	0291	0214	0139	0064	7
8	1237	1141	1048	0957	0868	0780	0694	0610	0528	0447	0367	0289	0213	0137	0063	8
9	1235	1140	1047	0956	0865	0779	0693	0609	0526	0446	0366	0288	0211	0136	0062	9
10	1233	1138	1045	0954	0865	0777	0692	0608	0525	0444	0365	0287	0210	0135	0061	10
11	1232	1137	1044	0953	0863	0776	0690	0606	0524	0443	0363	0285	0209	0134	0060	11
12	1230	1135	1042	0951	0862	0774	0689	0605	0522	0442	0362	0284	0208	0132	0058	12
13	1229	1134	1041	0950	0860	0773	0687	0603	0521	0440	0361	0283	0206	0131	0057	13
14	1227	1132	1039	0948	0859	0772	0686	0602	0520	0439	0359	0282	0205	0130	0056	14
15	1225	1130	1037	0947	0857	0770	0685	0601	0518	0438	0358	0280	0204	0129	0055	15
16	1224	1129	1036	0945	0856	0769	0683	0599	0517	0436	0357	0279	0202	0127	0053	16
17	1222	1127	1034	0944	0855	0767	0682	0598	0516	0435	0356	0278	0201	0126	0052	17
18	1221	1126	1033	0942	0853	0766	0680	0596	0514	0434	0354	0276	0200	0125	0051	18
19	1219	1124	1031	0941	0852	0764	0679	0595	0513	0432	0353	0275	0199	0124	0050	19
20	1217	1123	1030	0939	0850	0763	0678	0594	0512	0431	0352	0274	0197	0122	0049	20
21	1216	1121	1028	0938	0849	0762	0676	0592	0510	0430	0350	0273	0196	0121	0047	21
22	1214	1119	1027	0936	0847	0760	0675	0591	0509	0428	0349	0271	0195	0120	0046	22
23	1213	1118	1025	0935	0846	0759	0673	0590	0507	0427	0348	0270	0194	0119	0045	23
24	1211	1116	1024	0933	0844	0757	0672	0588	0506	0426	0346	0269	0192	0117	0044	24
25	1209	1115	1022	0932	0843	0756	0670	0587	0505	0424	0345	0267	0191	0116	0042	25
26	1208	1113	1021	0930	0841	0754	0669	0585	0503	0423	0344	0266	0190	0115	0041	26
27	1206	1112	1019	0929	0840	0753	0668	0584	0502	0422	0342	0265	0189	0114	0040	27
28	1205	1110	1018	0927	0838	0751	0666	0583	0501	0420	0341	0264	0187	0112	0039	28
29	1203	1109	1016	0926	0837	0750	0665	0581	0499	0419	0340	0262	0186	0111	0038	29
30	1201	1107	1015	0924	0835	0749	0663	0580	0498	0418	0339	0261	0185	0110	0036	30
31	1200	1105	1013	0923	0834	0747	0662	0579	0497	0416	0337	0260	0184	0109	0035	31
32	1198	1104	1012	0921	0833	0746	0661	0577	0495	0415	0336	0258	0182	0107	0034	32
33	1197	1102	1010	0920	0831	0744	0659	0576	0494	0414	0335	0257	0181	0106	0033	33
34	1195	1101	1008	0918	0830	0743	0658	0574	0493	0412	0333	0256	0180	0105	0031	34
35	1193	1099	1007	0917	0828	0741	0656	0573	0491	0411	0332	0255	0179	0104	0030	35
36	1192	1098	1005	0915	0827	0740	0655	0572	0490	0410	0331	0253	0177	0103	0029	36
37	1190	1096	1004	0914	0825	0739	0654	0570	0489	0408	0329	0252	0176	0101	0028	37
38	1189	1095	1002	0912	0824	0737	0652	0569	0487	0407	0328	0251	0175	0100	0027	38
39	1187	1093	1001	0911	0822	0736	0651	0568	0486	0406	0327	0250	0174	0099	0025	39
40	1186	1091	0999	0909	0821	0734	0649	0566	0484	0404	0326	0248	0172	0098	0024	40
41	1184	1090	0998	0908	0819	0733	0648	0565	0483	0403	0324	0247	0171	0096	0023	41
42	1182	1088	0996	0906	0818	0731	0647	0563	0482	0402	0323	0246	0170	0095	0022	42
43	1181	1087	0995	0905	0816	0730	0645	0562	0480	0400	0322	0244	0169	0094	0021	43
44	1179	1085	0993	0903	0815	0729	0644	0561	0479	0399	0320	0243	0167	0093	0019	44
45	1178	1084	0992	0902	0814	0727	0642	0559	0478	0398	0319	0242	0166	0091	0018	45
46	1176	1082	0990	0900	0812	0726	0641	0558	0476	0396	0318	0241	0165	0090	0017	46
47	1174	1081	0989	0899	0811	0724	0640	0557	0475	0395	0316	0239	0163	0089	0016	47
48	1173	1079	0987	0897	0809	0723	0638	0555	0474	0394	0315	0238	0162	0088	0015	48
49	1171	1078	0986	0896	0808	0721	0637	0554	0472	0392	0314	0237	0161	0087	0013	49
50	1170	1076	0984	0894	0806	0720	0635	0552	0471	0391	0313	0235	0160	0085	0012	50
51	1168	1074	0983	0893	0805	0719	0634	0551	0470	0390	0311	0234	0158	0084	0011	51
52	1167	1073	0981	0891	0803	0717	0633	0550	0468	0388	0310	0233	0157	0083	0010	52
53	1165	1071	0980	0890	0802	0716	0631	0548	0467	0387	0309	0232	0156	0082	0008	53
54	1163	1070	0978	0888	0801	0714	0630	0547	0466	0386	0307	0230	0155	0080	0007	54
55	1162	1068	0977	0887	0799	0713	0628	0546	0464	0384	0306	0229	0153	0079	0006	55
56	1160	1067	0975	0885	0798	0711	0627	0544	0463	0383	0305	0228	0152	0078	0005	56
57	1159	1065	0974	0884	0796	0710	0626	0543	0462	0382	0304	0227	0151	0077	0004	57
58	1157	1064	0972	0883	0795	0709	0624	0541	0460	0381	0302	0225	0150	0075	0002	58
59	1156	1062	0971	0881	0793	0707	0623	0540	0459	0379	0301	0224	0148	0074	0001	59
60	1154	1061	0969	0880	0792	0706	0621	0539	0458	0378	0300	0223	0147	0073	0000	60
	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	

LOGISTICAL LOGARITHMS.

0	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	0
1	3600	3660	3720	3780	3840	3900	3960	4020	4080	4140	4200	4260	4320	4380	4440	1
0	0000	9928	9858	9788	9720	9652	9586	9521	9456	9393	9331	9269	9208	9148	9089	0
1	9999	9927	9856	9787	9719	9651	9585	9520	9455	9392	9329	9268	9207	9147	9088	1
2	9998	9926	9855	9786	9717	9650	9584	9519	9454	9391	9328	9267	9206	9146	9087	2
3	9996	9925	9854	9785	9716	9649	9583	9518	9453	9390	9327	9266	9205	9145	9086	3
4	9995	9923	9853	9784	9715	9648	9582	9516	9452	9389	9326	9265	9204	9144	9085	4
5	9994	9922	9852	9782	9714	9647	9581	9515	9451	9388	9325	9264	9203	9143	9084	5
6	9993	9921	9851	9781	9713	9646	9579	9514	9450	9387	9324	9263	9202	9142	9083	6
7	9992	9920	9849	9780	9712	9645	9578	9513	9449	9386	9323	9262	9201	9141	9082	7
8	9990	9919	9848	9779	9711	9643	9577	9512	9448	9385	9322	9261	9200	9140	9081	8
9	9989	9918	9847	9778	9710	9642	9576	9511	9447	9384	9321	9260	9199	9139	9080	9
10	9988	9916	9846	9777	9708	9641	9575	9510	9446	9383	9320	9259	9198	9138	9079	10
11	9987	9915	9845	9775	9707	9640	9574	9509	9445	9381	9319	9258	9197	9137	9078	11
12	9986	9914	9844	9774	9706	9639	9573	9508	9444	9380	9318	9257	9196	9136	9077	12
13	9984	9913	9842	9773	9705	9638	9572	9507	9443	9379	9317	9256	9195	9135	9076	13
14	9983	9912	9841	9772	9704	9637	9571	9506	9442	9378	9316	9255	9194	9134	9077	14
15	9982	9910	9840	9771	9703	9636	9570	9505	9440	9377	9315	9254	9193	9133	9075	15
16	9981	9909	9839	9770	9702	9635	9569	9504	9439	9376	9314	9253	9192	9132	9074	16
17	9980	9908	9838	9769	9701	9633	9567	9502	9438	9375	9313	9252	9191	9131	9073	17
18	9978	9907	9837	9767	9699	9632	9566	9501	9437	9374	9312	9251	9190	9130	9072	18
19	9977	9906	9835	9766	9698	9631	9565	9500	9436	9373	9311	9250	9189	9129	9071	19
20	9976	9905	9834	9765	9697	9630	9564	9499	9435	9372	9310	9249	9188	9128	9070	20
21	9975	9903	9833	9764	9696	9629	9563	9498	9434	9371	9309	9248	9187	9128	9069	21
22	9974	9902	9832	9763	9695	9628	9562	9497	9433	9370	9308	9247	9186	9127	9068	22
23	9972	9901	9831	9762	9694	9627	9561	9496	9432	9369	9307	9246	9185	9126	9067	23
24	9971	9900	9830	9761	9693	9626	9560	9495	9431	9368	9306	9245	9184	9125	9066	24
25	9970	9899	9829	9759	9692	9625	9559	9494	9430	9367	9305	9244	9183	9124	9065	25
26	9969	9897	9827	9758	9690	9624	9558	9493	9429	9366	9304	9243	9182	9123	9064	26
27	9968	9896	9826	9757	9689	9622	9557	9492	9428	9365	9303	9241	9181	9122	9063	27
28	9966	9895	9825	9756	9688	9621	9555	9491	9427	9364	9302	9240	9180	9121	9062	28
29	9965	9894	9824	9755	9687	9620	9554	9490	9426	9363	9301	9239	9179	9120	9061	29
30	9964	9893	9823	9754	9686	9619	9553	9488	9425	9362	9300	9238	9178	9119	9060	30
31	9963	9892	9822	9753	9685	9618	9552	9487	9424	9361	9299	9237	9177	9118	9059	31
32	9962	9890	9820	9751	9684	9617	9551	9486	9422	9360	9298	9236	9176	9117	9058	32
33	9960	9889	9819	9750	9683	9616	9550	9485	9421	9359	9297	9235	9175	9116	9057	33
34	9959	9888	9818	9749	9681	9615	9549	9484	9420	9358	9296	9234	9174	9115	9056	34
35	9958	9887	9817	9748	9680	9614	9548	9483	9419	9356	9294	9233	9173	9114	9055	35
36	9957	9886	9816	9747	9679	9612	9547	9482	9418	9355	9293	9232	9172	9113	9054	36
37	9956	9885	9815	9746	9678	9611	9546	9481	9417	9354	9292	9231	9171	9112	9053	37
38	9954	9883	9813	9745	9677	9610	9545	9480	9416	9353	9291	9230	9170	9111	9052	38
39	9953	9882	9812	9744	9676	9609	9544	9479	9415	9352	9290	9229	9169	9110	9051	39
40	9952	9881	9811	9742	9675	9608	9542	9478	9414	9351	9289	9228	9168	9109	9050	40
41	9951	9880	9810	9741	9674	9607	9541	9477	9413	9350	9288	9227	9167	9108	9049	41
42	9950	9879	9809	9740	9672	9606	9540	9476	9412	9349	9287	9226	9166	9107	9048	42
43	9948	9877	9808	9739	9671	9605	9539	9475	9411	9348	9286	9225	9165	9106	9047	43
44	9947	9876	9807	9738	9670	9604	9538	9473	9410	9347	9285	9224	9164	9105	9046	44
45	9946	9875	9805	9737	9669	9603	9537	9472	9409	9346	9284	9223	9163	9104	9045	45
46	9945	9874	9804	9736	9668	9601	9536	9471	9408	9345	9283	9222	9162	9103	9044	46
47	9944	9873	9803	9734	9667	9600	9535	9470	9407	9344	9282	9221	9161	9102	9043	47
48	9942	9872	9802	9733	9666	9599	9534	9469	9406	9343	9281	9220	9160	9101	9042	48
49	9941	9870	9801	9732	9665	9598	9533	9468	9405	9342	9280	9219	9159	9100	9042	49
50	9940	9869	9800	9731	9664	9597	9532	9467	9404	9341	9279	9218	9158	9099	9041	50
51	9939	9868	9798	9730	9662	9596	9530	9466	9402	9340	9278	9217	9157	9098	9040	51
52	9938	9867	9797	9729	9661	9595	9529	9465	9401	9339	9277	9216	9156	9097	9039	52
53	9937	9866	9796	9728	9660	9594	9528	9464	9400	9338	9276	9215	9155	9096	9038	53
54	9935	9865	9795	9727	9659	9593	9527	9463	9399	9337	9275	9214	9154	9095	9037	54
55	9934	9863	9794	9725	9658	9592	9526	9462	9398	9336	9274	9213	9153	9094	9036	55
56	9933	9862	9793	9724	9657	9590	9525	9461	9397	9335	9273	9212	9152	9093	9035	56
57	9932	9861	9792	9723	9656	9589	9524	9460	9396	9334	9272	9211	9151	9092	9034	57
58	9931	9860	9790	9722	9655	9588	9523	9459	9395	9333	9271	9210	9150	9091	9033	58
59	9929	9859	9789	9721	9653	9587	9522	9457	9394	9332	9270	9209	9149	9090	9032	59
60	9928	9858	9788	9720	9652	9586	9521	9456	9393	9331	9269	9208	9148	9089	9031	60
	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	

LOGISTICAL LOGARITHMS.

°	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	°
'	4500	4560	4620	4680	4740	4800	4860	4920	4980	5040	5100	5160	5220	5280	5340	'
0	9031	8973	8917	8861	8805	8751	8697	8644	8591	8539	8488	8437	8387	8337	8288	0
1	9030	8972	8916	8860	8804	8750	8696	8643	8590	8538	8487	8436	8386	8336	8287	1
2	9029	8971	8915	8859	8803	8749	8695	8642	8589	8537	8486	8435	8385	8335	8286	2
3	9028	8971	8914	8858	8802	8748	8694	8641	8589	8537	8485	8434	8384	8335	8286	3
4	9027	8970	8913	8857	8802	8747	8693	8640	8588	8536	8484	8433	8383	8334	8285	4
5	9026	8969	8912	8856	8801	8746	8692	8639	8587	8535	8484	8433	8383	8333	8284	5
6	9025	8968	8911	8855	8800	8745	8692	8638	8586	8534	8483	8432	8382	8332	8283	6
7	9024	8967	8910	8854	8799	8745	8691	8638	8585	8533	8482	8431	8381	8331	8282	7
8	9023	8966	8909	8853	8798	8744	8690	8637	8584	8532	8481	8430	8380	8330	8281	8
9	9022	8965	8908	8852	8797	8743	8689	8636	8583	8531	8480	8429	8379	8330	8281	9
10	9021	8964	8907	8851	8796	8742	8688	8635	8582	8530	8479	8428	8379	8329	8280	10
11	9020	8963	8906	8850	8795	8741	8687	8634	8582	8530	8478	8428	8378	8328	8279	11
12	9019	8962	8905	8849	8794	8740	8686	8633	8581	8529	8477	8427	8377	8327	8278	12
13	9018	8961	8904	8848	8793	8739	8685	8632	8580	8528	8477	8426	8376	8326	8277	13
14	9017	8960	8903	8847	8792	8738	8684	8631	8579	8527	8476	8425	8375	8325	8276	14
15	9016	8959	8903	8847	8792	8737	8684	8631	8578	8526	8475	8424	8374	8325	8276	15
16	9015	8958	8902	8846	8791	8736	8683	8630	8577	8525	8474	8423	8373	8324	8275	16
17	9015	8957	8901	8845	8790	8736	8682	8629	8576	8524	8473	8423	8373	8323	8274	17
18	9014	8956	8900	8844	8789	8735	8681	8628	8575	8524	8472	8422	8372	8322	8273	18
19	9013	8955	8899	8843	8788	8734	8680	8627	8575	8523	8472	8421	8371	8322	8273	19
20	9012	8954	8898	8842	8787	8733	8679	8626	8574	8522	8471	8420	8370	8321	8272	20
21	9011	8953	8897	8841	8786	8732	8678	8625	8573	8521	8470	8419	8369	8320	8271	21
22	9010	8952	8896	8840	8785	8731	8677	8624	8572	8520	8469	8418	8368	8319	8270	22
23	9009	8952	8895	8839	8784	8730	8677	8624	8571	8519	8468	8418	8368	8318	8269	23
24	9008	8951	8894	8838	8783	8729	8676	8623	8570	8518	8467	8417	8367	8317	8268	24
25	9007	8950	8893	8837	8782	8728	8675	8622	8569	8518	8467	8416	8366	8317	8268	25
26	9006	8949	8892	8837	8781	8727	8674	8621	8568	8517	8466	8415	8365	8316	8267	26
27	9005	8948	8891	8836	8781	8727	8673	8620	8568	8516	8465	8414	8364	8315	8266	27
28	9004	8947	8890	8835	8780	8726	8672	8619	8567	8515	8464	8413	8363	8314	8265	28
29	9003	8946	8889	8834	8779	8725	8671	8618	8566	8514	8463	8413	8363	8313	8265	29
30	9002	8945	8888	8833	8778	8724	8670	8617	8565	8513	8462	8412	8362	8312	8264	30
31	9001	8944	8888	8832	8777	8723	8669	8617	8564	8513	8461	8411	8361	8312	8263	31
32	9000	8943	8887	8831	8776	8722	8668	8616	8563	8512	8460	8410	8360	8311	8262	32
33	8999	8942	8886	8830	8775	8721	8667	8615	8562	8511	8460	8409	8359	8310	8261	33
34	8998	8941	8885	8829	8774	8720	8666	8614	8561	8510	8459	8408	8358	8309	8260	34
35	8997	8940	8884	8828	8773	8719	8666	8613	8561	8509	8458	8408	8358	8308	8260	35
36	8996	8939	8883	8827	8772	8718	8665	8612	8560	8508	8457	8407	8357	8307	8259	36
37	8995	8938	8882	8826	8771	8718	8664	8611	8559	8507	8456	8406	8356	8307	8258	37
38	8994	8937	8881	8825	8771	8717	8663	8610	8558	8506	8455	8405	8355	8306	8257	38
39	8993	8936	8880	8825	8770	8716	8662	8610	8557	8506	8455	8404	8354	8305	8257	39
40	8992	8935	8879	8824	8769	8715	8661	8609	8556	8505	8454	8403	8353	8304	8256	40
41	8992	8935	8878	8823	8768	8714	8661	8608	8556	8504	8453	8403	8353	8304	8255	41
42	8991	8934	8877	8822	8767	8713	8660	8607	8555	8503	8452	8402	8352	8303	8254	42
43	8990	8933	8876	8821	8766	8712	8659	8606	8554	8502	8451	8401	8351	8302	8253	43
44	8989	8932	8875	8820	8765	8711	8658	8605	8553	8501	8450	8400	8350	8301	8252	44
45	8988	8931	8875	8819	8764	8710	8657	8604	8552	8501	8450	8400	8399	8300	8252	45
46	8987	8930	8874	8818	8763	8709	8656	8603	8551	8500	8449	8398	8349	8299	8251	46
47	8986	8929	8873	8817	8762	8709	8655	8603	8550	8499	8448	8398	8348	8299	8250	47
48	8985	8928	8872	8816	8761	8708	8654	8602	8549	8498	8447	8397	8347	8298	8249	48
49	8984	8927	8871	8815	8761	8707	8654	8601	8549	8497	8446	8396	8346	8297	8248	49
50	8983	8926	8870	8814	8760	8706	8653	8600	8548	8496	8445	8395	8345	8296	8247	50
51	8982	8925	8869	8813	8759	8705	8652	8599	8547	8495	8445	8394	8345	8295	8247	51
52	8981	8924	8868	8813	8758	8704	8651	8598	8546	8494	8444	8393	8344	8294	8246	52
53	8980	8923	8867	8812	8757	8703	8650	8597	8545	8494	8443	8392	8343	8294	8245	53
54	8979	8922	8866	8811	8756	8702	8649	8596	8544	8493	8442	8392	8342	8293	8244	54
55	8978	8921	8865	8810	8755	8702	8648	8596	8543	8492	8441	8391	8341	8292	8244	55
56	8977	8920	8864	8809	8754	8701	8647	8595	8542	8491	8440	8390	8340	8291	8243	56
57	8976	8919	8863	8808	8753	8700	8646	8594	8542	8490	8440	8389	8340	8291	8242	57
58	8975	8918	8862	8807	8752	8699	8645	8593	8541	8489	8439	8388	8339	8290	8241	58
59	8974	8917	8861	8806	8752	8698	8645	8593	8540	8489	8438	8388	8338	8289	8240	59
60	8973	8917	8861	8805	8751	8697	8644	8591	8539	8488	8437	8387	8337	8288	8239	60
	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	

HERE

HERE FOLLOWS THE Seaman's Ready Computer:

BEING AN IMPROVEMENT in *Tabular* NAVIGATION.

Whereby *any Person* understanding *Addition*, is qualified to keep a
SHIP's Reckoning at SEA; in *Latitude*, *Departure*,
and *LONGITUDE*:

As by which *easy* TABLE, the *Ship's Place*, at any Time, may be speedily taken out,
(like the SUN's mean *Place* from *Astronomical Tables* :)

And all *practical Questions* in NAVIGATION and *Plane TRIGONOMETRY* are readily
answered, without *Logarithm-Sines*, *Tangents*, or other *artificial Numbers* :

Of which see Examples further on.

By this sufficiently *near Method* of travelling at Sea, assisted by *Helps* contained in this Work, for determining
the *Longitudes* from *Greenwich*, *Spheroidical* Navigation is exploded, as useless.

FOR, as Mr. EMERSON judiciously observes, at P. vi. *Preface* to his NAVIGATION, (containing the exact
Rules and their *Demonstrations* on the Principles of *spherical* Navigation) to which Our NAVIGATION is *supplemental*.
“ It would be an idle Refinement to deduce a Table of meridional Parts from the Spheroid. That it would require all the Parallel
“ of Latitude and Arches of the Meridian to be altered accordingly; creating an endless Deal of Trouble for no real Advan-
“ tage. That, if we consider that no Ship can steer true to a Degree, or hardly to a Quarter Point of the Compass, nor
“ can measure her Distance sailed by the Log, to any such Degree of Exactness, but that there will always be a considerable
“ Variation from the precise Truth, in Regard to the Ship's Place, for which Reason she is obliged, as often as possible, to
“ correct her Latitude and Longitude by Observation. Therefore (he concludes) it would be quite trifling to descend to such
“ minute Differences as can have no sensible Effect in Practice. That, since such is the Difference betwixt spherical and
“ spheroidical Navigation, if we must needs wander from the Truth, we had better do it in a plain and easy Way than in a
“ rigid and difficult one.”

Mr. EMERSON then proceeds to lay down his Rules of spherical Navigation (agreeing with ours) as short and
plain as possible; “ Disencumbering the Seaman of every Thing long and tedious, and giving him as much Satisfaction as he
“ will ever find, or can wish for.”

To which may be added. That every Person's Experience (having travelled by Sea) will inform him of the Truth
of the foregoing Assertions. And a Person we know is perfectly convinced thereof, in three Voyages he made to
the West Indies, and one to Greenland. That the Errors by *Lee Way*, and mistaken Course, and Distance run by the
Log, from their various Causes (of Wind, Sails set, high Seas, Currents, &c.) are with Doubt and Difficulty
corrected. But a *spheroidical Navigator*, having never been a Voyage, if he goes on to keep a Ship's Reckoning,
will find himself involved in greater Errors than those he pretends to correct; since there is always much more
Difference in Mens Judgments about settling the *Lee-Way*, *Currents*, *Course*, and *Distance*, than there is betwixt
spherical and *spheroidical* Navigation.

OUR short and easy Method of computing a Ship's Way through the Ocean, is an Improvement from a Table we have
lately seen at the End of Sherwin's *Mathematical Tables*, revised by Gardiner; but did not see till we contrived our
own Table, Computer No. I. for that Purpose (constructed from natural Sines) in a more general Way, as follows.
Computers No. I. II. are quite a NEW INVENTION.

The SEAMAN'S READY COMPUTER. N^o. 1.

Course		10000		20000		30000		40000		50000		60000		70000		80000		90000		Course	
		D. Lat. Dep.		D. Lat. Dep.		D. Lat. Dep.		D. Lat. Dep.		D. Lat. Dep.		D. Lat. Dep.		D. Lat. Dep.		D. Lat. Dep.		D. Lat. Dep.			
0	1	09998	00175	19997	00349	29995	00524	39994	00698	49992	00873	59991	01047	69989	01222	79988	01396	89986	01571	89	
	2	09994	00349	19988	00698	29982	01047	39976	01396	49970	01745	59963	02094	69957	02443	79951	02792	89945	03141	88	
	3	09988	00491	19976	00981	29964	01472	39952	01963	49940	02453	59928	02944	69916	03435	79904	03925	89892	04410	...	
	4	09986	00523	19973	01047	29959	01570	39945	02093	49931	02617	59918	03140	69904	03664	79890	04187	89877	04710	87	
1	5	09976	00698	19951	01395	29927	02093	39903	02790	49878	03488	59854	04185	69829	04883	79805	05580	89781	06278	86	
	6	09962	00872	19924	01743	29886	02615	39848	03486	49810	04358	59772	05229	69734	06101	79696	06972	89658	07844	85	
	7	09952	00980	19904	01960	29856	02940	39807	03921	49759	04901	59711	05881	69663	06861	79615	07841	89567	08822	...	
	8	09945	01045	19890	02091	29836	03136	39781	04181	49726	05226	59671	06272	69617	07317	79562	08362	89507	09408	84	
2	9	09925	01219	19851	02437	29776	03656	39702	04875	49627	06093	59553	07312	69478	08531	79404	09750	89329	10968	83	
	10	09903	01392	19805	02783	29708	04175	39611	05567	49513	06959	59416	08350	69319	09742	79221	11134	89124	12526	82	
	11	09892	01467	19784	02935	29675	04402	39567	05869	49459	07337	59351	08804	69242	10271	79134	11738	89026	13206	...	
	12	09877	01564	19754	03129	29631	04693	39508	06257	49384	07822	59261	09386	69138	10950	79015	12515	88892	14079	81	
3	13	09848	01736	19696	03473	29544	05209	39392	06946	49240	08682	59088	10419	68937	12155	78785	13892	88633	15628	80	
	14	09816	01908	19633	03816	29449	05724	39265	07632	49081	09540	58898	11449	68714	13357	78530	15265	88346	17173	79	
	15	09808	01951	19616	03902	29424	05853	39231	07804	49039	09754	58847	11705	68655	13656	78463	15607	88271	17558	...	
	16	09781	02079	19563	04158	29344	06237	39126	08116	48907	10396	58689	12475	68470	14554	78252	16633	88033	18712	78	
4	17	09744	02250	19487	04499	29231	06749	38975	08998	48718	11248	58462	13497	68206	15746	77950	17996	87693	20246	77	
	18	09703	02419	19406	04838	29108	07258	38812	09677	48515	12096	58218	14515	67921	16935	77624	19354	87327	21773	76	
	19	09700	02438	19401	04860	29101	07280	38801	09719	48502	12149	58202	14579	67902	17009	77602	19438	87303	21868	...	
	20	09659	02588	19319	05176	28978	07765	38637	10353	48296	12941	57956	15529	67615	18117	77274	20706	86933	23294	75	
5	21	09613	02756	19225	05513	28838	08269	38450	11025	48063	13782	57676	16538	67288	19295	76901	22051	86513	24807	74	
	22	09569	02903	19139	05806	28708	08709	38278	11611	47847	14514	57416	17417	66986	20320	76555	23223	86125	26126	...	
	23	09563	02924	19126	05847	28689	08771	38252	11695	47815	14619	57378	17542	66941	20466	76504	23390	86067	26313	73	
	24	09511	03090	19021	06180	28532	09271	38042	12361	47553	15451	57063	18541	66574	21631	76084	24721	85595	27812	72	
6	25	09455	03256	18910	06511	28366	09767	37821	13023	47276	16278	56731	19534	66186	22790	75642	26045	85097	29301	71	
	26	09415	03369	18831	06738	28246	10107	37662	13476	47077	16844	56493	20213	65908	23582	75324	26951	84739	30320	...	
	27	09397	03420	18794	06840	28191	10261	37588	13681	46985	17101	56382	20521	65779	23941	75175	27362	84572	30782	70	
	28	09336	03584	18672	07167	28007	10751	37343	14335	46679	17918	56015	21502	65351	25086	74686	28669	84222	32253	69	
7	29	09272	03746	18544	07492	27816	11238	37087	14984	46359	18730	55631	22476	64903	26222	74175	29969	83447	33715	68	
	30	09239	03827	18478	07654	27716	11480	36955	15307	46194	19134	55433	22961	64672	26788	73910	30615	83149	34441	...	
	31	09205	03907	18410	07815	27615	11722	36820	15629	46025	19337	55230	23444	64435	27351	73640	31258	82845	35166	67	
	32	09135	04067	18270	08135	27406	12202	36542	16269	45677	20337	54813	24404	63948	28472	73084	32539	82215	36606	66	
8	33	09063	04226	18126	08452	27189	12679	36252	16905	45315	21131	54378	25357	63442	29583	72505	33809	81568	38036	65	
	34	09020	04276	18080	08551	27120	12827	36160	17102	45199	21378	54239	25653	63279	29929	72319	34204	81359	38430	...	
	35	08988	04384	17976	08767	26964	13151	35952	17535	44940	21919	53928	26302	62916	30686	71904	35070	80891	39453	64	
	36	08910	04540	17820	09080	26730	13620	35640	18160	44550	22699	53460	27239	62370	31779	71280	36319	80191	40859	63	
9	37	08829	04695	17659	09389	26488	14084	35318	18779	44147	23474	52977	28168	61806	32863	70636	37558	79465	42252	62	
	38	08819	04714	17638	09428	26458	14142	35277	18856	44096	23570	52915	28284	61734	32998	70554	37712	79373	42426	...	
	39	08746	04848	17492	09696	26239	14544	34985	19392	43731	24240	52477	29085	61223	33937	69970	38785	78716	43633	61	
	40	08660	05000	17320	10000	25981	15000	34641	20000	43301	25000	51961	30000	60622	35000	69622	40000	77942	45000	60	
10	41	08577	05141	17155	10282	25732	15423	34309	20564	42886	25705	51464	30846	60041	35987	68618	41128	77196	46269	...	
	42	08572	05150	17143	10301	25715	15451	34287	20602	42858	25752	51430	30902	60002	36052	68573	41203	77145	46353	59	
	43	08480	05299	16961	10598	25441	15896	33922	21197	42402	26496	50883	31795	59363	37094	67843	42394	76324	47693	58	
	44	08387	05446	16773	10893	25160	16339	33547	21786	41934	27232	50320	32678	58707	38125	67094	43571	75480	49018	57	
11	45	08315	05556	16629	11111	24944	16667	33259	22223	41573	27778	49888	33334	58203	38890	66518	44446	74832	50001	...	
	46	08290	05592	16581	11184	24871	16776	33162	22368	41452	27960	49742	33552	58033	39144	66323	44735	74613	50327	56	
	47	08192	05736	16383	11472	24575	17207	32766	22943	40958	28679	49149	34415	57347	40150	65532	45886	73724	51622	55	
	48	08090	05878	16180	11756	24271	17634	32361	23511	40451	29389	48541	35267	56631	41145	64721	47023	72812	52901	54	
12	49	08032	05957	16064	11914	24096	17871	32128	23828	40160	29785	48192	35742	56224	41699	64257	47656	72289	53613	...	
	50	07986	06018	15973	12036	23959	18054	31945	24073	39932	30091	47918	36109	55904	42127	63891	48145	71877	54163	53	
	51	07880	06157	15760	12313	23640	18470	31520	24626	39401	30783	47281	36940	55161	43096	63041	49253	70921	55409	52	
	52	07771	06293	15543	12586	23314	18880	31086	25173	38857	31466	46629	37759	54400	44052	62172	50346	69943	56639	51	
13	53	07730	06344	15460	12688	23190	19032	30920	25376	38650	31720	46381	38064	54111	44408	61841	50751	69571	57095	...	
	54	07660	06428	15321	12856	22981	19284	30642	25712	38302	32139	45963	38567	53623	44995	61284	51423	68944	57851	50	
	55	07547	06561	15094	13121	22641	19682	30188	26242	37736	32803	45283	39363	52830	45924	60377	52485	67924	59045	49	
	56	07431	06691	14863	13383	22294	20074	29726	26765	37157	33457	44589	40148	52020	46839	59452	53530	66883	60222	48	
14	57	07410	06716	14819	13431	22229	20147	29638	26862	37048	33578	44457	40294	51867	47009	59276	53725	66680	60440	...	
	58	07314	06820	14628	13640	21941	20460	29254	27280	36568	34100	43881	40920	51195	47740	58508	54560	65822	61380	47	
	59	07193	06947	14387	13894	21580	20840	28774	27786	35967	34733	43160	41679	50354	48626	57547	55573	64741	62519	46	
	60	07071	07071	14142	14142	21243	21213	28284	28284	35355	35355	42426	42426	49497	49497	56569	56569	63640	63640	45	
Course		10000		20000		30000		40000		50000		60000		70000		80000		90000			

USE of the SEAMAN'S READY COMPUTER.

The Distances sailed, or Hypothenuses, at the Top and Bottom of the Table, may signify any of the Numbers below.

10000	20000	30000	40000	50000	60000	70000	80000	90000
1000	2000	3000	4000	5000	6000	7000	8000	9000
100	200	300	400	500	600	700	800	900
10	20	30	40	50	60	70	80	90
1	2	3	4	5	6	7	8	9

Under or above which stand the correspondent Dif. of Lat. and Departure, or Base and Perpend. against the Course or $\angle$, on one Side of the Table, to be taken in equal whole Numbers. Those to the Right of these being Decimals.

THEREFORE, the Difference of Latitude and Departure, (or Base and Perpendicular) answerable to the given Distance sailed (or Hypothenuse of a right angled Triangle) on a given Course, or Angle, must be taken out by Parts, and added together; taking out first for the highest Number of Tens, then for next Tens less, and so down to Units; according to the Numbers of Figures in the Distance sailed, (or Hypothenuse of a Triangle) to be taken out for.

PROPOSITION I. To find the Difference of Latitude and Departure from the Course and Distance given? Or to find the Base and Perpendicular from an acute Angle and the Hypothenuse of a Triangle being given?

EXAMPLE I. Required the Difference of Latitude and Departure, when a Ship has sailed 85 Miles, on a S. W. and by S. Course $\frac{1}{2}$ Point Westerly; being $3\frac{1}{2}$ Points to the Westward?

	Dist.	Dif. Lat. S.	Depart. W.
Against $3\frac{1}{2}$ Points under 80		61.841	50.751
5		3.868	3.172
Miles 85		65.709	53.923 Answer.

But two Decimals to the Right of any whole Number will be sufficient in taking out; except where Accuracy is required, in Trigonometry.

EXAMPLE II. A Ship having sailed 116 Miles, on a Course of 41 Degrees from the North, easterly, required her Difference of Latitude and Departure?

	Dist.	Dif. Lat. N.	Dep. E.
Against 41° under 100		75.47	65.61
10		7.54	6.56
6		4.53	3.94
Miles 116		87.54	76.11 Answer.

A TABLE shewing the Angles that every Point of the Compass makes with the Meridian: For turning Rhumbs, or the Course, steered by the Ship, into Degrees, and the contrary.

North		Points.	D. M. S.	South	
		$0\frac{1}{4}$	2 48 45		
		$0\frac{1}{2}$	5 37 30		
		$0\frac{3}{4}$	8 26 15		
N. by E.	N. by W.	1	11 15 0	S. by E.	S. by W.
		$1\frac{1}{4}$	14 3 45		
		$1\frac{1}{2}$	16 52 30		
		$1\frac{3}{4}$	19 41 15		
N. N. E.	N. N. W.	2	22 30 0	S. S. E.	S. S. W.
		$2\frac{1}{4}$	25 18 45		
		$2\frac{1}{2}$	28 7 30		
		$2\frac{3}{4}$	30 56 15		
N. E. by N.	N. W. by N.	3	33 45 0	S. E. by S.	S. W. by S.
		$3\frac{1}{4}$	36 33 45		
		$3\frac{1}{2}$	39 22 30		
		$3\frac{3}{4}$	42 11 15		
Nor. East	Nor. West	4	45 0 0	Sou. East	Sou. West
		$4\frac{1}{4}$	47 48 45		
		$4\frac{1}{2}$	50 37 30		
		$4\frac{3}{4}$	53 26 15		
N. E. by E.	N. W. by W.	5	56 15 0	S. E. by E.	S. W. by W.
		$5\frac{1}{4}$	59 3 45		
		$5\frac{1}{2}$	61 52 30		
		$5\frac{3}{4}$	64 41 15		
E. N. E.	W. N. W.	6	67 30 0	E. S. E.	W. S. W.
		$6\frac{1}{4}$	70 18 45		
		$6\frac{1}{2}$	73 7 30		
		$6\frac{3}{4}$	75 56 15		
E. by N.	W. by N.	7	78 45 0	E. by S.	W. by S.
		$7\frac{1}{4}$	81 33 45		
		$7\frac{1}{2}$	84 22 30		
		$7\frac{3}{4}$	87 11 15		
East	West	8	90 0 0	East	West

EXAMPLE III. Required the Base and Perpendicular of a right angled Triangle, having an acute Angle equal to 23°, and its Hypothenuse equal to 1543?

	Hypoth.	Base.	Perp.
Against 23° under 1000		920.5	390.70
500		460.25	195.37
40		36.820	15.629
3		2.761	1.172
Answer 1543		1420.332	602.871

EXAMPLE IV. Required the Base and Perpendicular of a right angled plain Triangle, having an acute Angle equal to 66°, and its Hypothenuse equal to 77?

	Dist.	Base.	Perp.
Against 66° under 70		63.948	28.472
7		6.395	2.847
Answer 77		70.343	31.319

USE of the SEAMAN'S READY COMPUTER.

PROPOSITION II. *To find the Difference of Latitude and Departure from several Courses and Distances sailed; or to work a Ship's Traverse; to find the Miles in a Degree of Latitude, or take a Map of an Estate, or County?*

MAKE a *Traverse-Table*, as on the right Hand, and place therein each Course and Distance. Then find the Difference of Latitude and Departure to each Course, by the foregoing Proposition, placing them in their proper Columns.

NOW, the Difference of the Sums of the *Northings* and *Southings*, will be the whole Difference of Latitude; and the Difference of the Sums of *Eastings* and *Westings*, will be the whole Departure.

EXAMPLE. *The Difference of Latitude and Departure of a Ship sailing several Courses and Distances, will be found as in the adjoining Table, by the former Proposition.*

Courses.	Dist. Miles.	Diff. Latitude.		Departure.	
		North.	South.	East.	West.
N. 85° W.	70	06.101			69.734
S. S. W. $\frac{1}{4}$ W.	37		31.736		19.022
S. 40° W.	84		64.348		53.994
S. S. E. $\frac{1}{2}$ E.	79		69.671	37.241	
S. 49° E.	148		97.096	111.697	
N. E. $\frac{1}{2}$ E.	86	54.557		66.479	
		60.658	262.851 60.658	215.417 142.750	142.750
		Dif. Lat.	202.193 Miles.	72.667 Miles.	Depart.

The above PROPOSITION finds the Miles in a Degree of Latitude, and is extensively useful in surveying Estates, Counties, or large Tracts of Land. Mr. Norwood, in the Year 1635, made Use of it in determining the Difference of Latitude (according to the Method in the foregoing Traverse-Table) between the Parallels of London and York of given Latitudes, by celestial Observation; and so determines, very near the Truth, the Miles in a Degree of Latitude. (See his *Seaman's Practice*, since reprinted for W. Mount and T. Page, on Tower-Hill.) For, he measuring the several Distances between all the Turnings of the Road, from London to York, as the Road led him, at the same Time he observed the several Bearings betwixt those Turnings, by his *Circumferentor*, he then determined, by Trigonometry, the several Differences of Latitude and Departure; (which is done at Sight by our *Easy Computer*) and thence he determined the Distance, or Miles, Difference of Latitude, between the Parallels of London and York, and also Distance of those Meridians. And consequently the Miles in a Degree of Latitude, by Proportion of the Miles in the Degrees, Difference of Latitude, by Observation. If which Bearings and Distances were again taken and measured, more correctly, according to the last improved Methods of Surveying, there is no Doubt, but that the Distance between the Parallels of London and York (on so large an Arc of the Meridian) might be determined with as much Accuracy, as that Distance (measured by the French) was determined between the two Parallels near the Polar Circle.

In surveying Counties (or even Estates) by the Bearings and Distances of their Circuits, the Use of the *Needle* (not always keeping Parallel with itself) is not to be trusted to; but the several Bearings and Distances must be taken and measured by the *correcter Methods* known. And, in plotting the Survey of a County, by *Circuit Station Lines*, (though consisting of never so many) they may be reduced to a small Number, for the first Plan or Closing. And the like Method should be pursued in plotting the intermediate Circuit Stations betwixt each two plotted Stations in the first Plan, and so on; whereby every Circuit Station will be more correctly placed than by any other Method; and the resulting Out line or Plan, will be as near the Truth as possible.

At every Bearing taken, each Distance between the several Circuit-Stations, may be measured in Chains, of 100 Feet, or by any other Measure, (to be inserted in a Station or Traverse-Table, as above) correspondent to which, the several Differences of Latitude and Departure, or Northings and Southings, and Eastings and Westings, will be found in the same Measure, taken out of the *ready Computer*, for plotting each Station, and the whole Tract of Land.

REMARK. The Meridional Distance, and Departure (of Easting and Westing) are manifestly different. In sailing towards the Pole, the Departure is the greater, because of the nearer and nearer Approach of the Meridians cross'd over by the same Course, for one Time. But in sailing towards the Equator the Departure is less than the meridional Distance; because the Meridians cross'd over by the same Course, for one Time, grow wider, or recede more and more from one another: The Departure being of a middle Quantity (especially on one Course) between the Meridional Distances of two Places, for the Parallel departed from and for that arrived in. The Longitude is invariable, between any two Places, or Parallels, and different from the Meridional Distance and Departure. The Meridional Distance is different in different Parallels; and the Departure is only the same, for a single Course, between two Parallels; because many different Courses (in long Runs) will have many manifest Differences in the Meridional Distance, on which the Sum of the several Departures depend. This makes no Error between short Parallels of Latitude, (from Day to Day's Reckoning) on the plain Chart.

USE of the SEAMAN'S COMPUTER.

PROPOSITION III. To find the Course and Distance, from the Difference of Latitude and Departure given?

RULE I. Find the given Difference of Latitude and Departure together, in some of the Columns of the Seaman's READY COMPUTER Number I, or find the nearest Numbers to them, and even therewith, on the Side of the Page, is found the Course; and the Distance is found at the Top or Bottom of the Page, in the Column of given Latitude and Departure.

EXAMPLE. To find the Course and Distance answering to 700 Miles Difference of Latitude Southerly, and 388 Miles Departure Westerly?

In the Column 80000 you will find 699.70 Miles Difference of Latitude, and 387.85 Miles Departure, the nearest Numbers to those given, even with which tabular Numbers you find the Course S. 29° westerly; and, at the Top of the Column of the given Latitude and Departure, you find 800 (taking an equal Number of Figures) the Distance correspondent, required.

RULE II. If the given Difference of Latitude and Departure cannot be found together or in near Numbers, you must divide the less by the greater Number (Difference of Latitude by Departure, or Departure by Difference of Latitude, according as which happens to be the greater) and ascertaining a few Figures in the Quotient, find the nearest Number answering thereto, in the Column under 10000, of COMPUTER Number II. following, (which, in all Cases, you will readily find, without Stop, or Difficulty, by casting the Eye along the Column) and even with the said Quotient-Figures stands the Course, on the Side; and the Distance is found by taking it out of the said Ready Computer Number II. answerable to the given Difference of Latitude.

EXAMPLE. In the former Instance, 700)388(.554 which Quote is immediately found against 29°, the Course South-westerly, as before. And against which Course, under 700, Difference of Latitude, (in the Column 70000) will be found 800.45 or 800 Miles for the Distance, required.

But, UNIVERSALLY. Required the Course and Distance answerable to 971 Miles, Difference of Latitude, South; and 1117 Miles Departure, easterly?

1117)971.000(.869 which Quote is immediately found against 49°, Column 10000, in the Seaman's Computer, Number II. following; which is the required Course, South-easterly.

8936	
7740	
6702	
10380	
10053	
327	

WHEN (of the Difference of Latitude and Departure) the Departure is the greater, the Course is above 45°; but when the Difference of Latitude is the greater, the Course is less than 45°.

NOW, by the Seaman's Ready Computer, Number II. following, always take out for the greater,

Course S.	Depart. E.	Distance.	Hence, also the two Legs of a plane right-angled Triangle being given, the acute Angles and Hypothenuse may be found in the same Manner.
Against 49°	Under 1000	1325.0	
or 41	100	132.50	
	10	13.250	
	7	9.275	
	Miles 1117	1480.025	Answer.

N. B. When the greater Side (either Latitude or Departure) and an Angle is given (or the Course) the less Side (Departure or Latitude) and also the Hypothenuse (or Distance run) may be found by Computer, Number II. But the less Side and an Angle being given, the REST cannot be found but by Computer, Number III. following.

PROPOSITION IV. Having the greater of either Difference of Latitude or Departure, and also the Course given, to find the less of either, of Difference of Latitude or Departure, and also Distance sailed.

This Proposition is universally answered as in the foregoing Example, by the Seaman's Computer, Number II.

EXAMPLE II. Course under 45° (viz. N. 36° westerly) and Difference of Latitude 349 Miles, to find the Departure and Distance sailed?

Course.	Dif. Lat. N.	Depart. W.	Distance.
Against 36°	Under 300	217.96	370.82
	40	29.062	49.443
	9	6.539	11.125
	Miles 349	253.561	431.388
		Answer.	

Hence, the less Side and Hypothenuse, of any right-angled plane Triangle, are found in the same Manner, from having an acute Angle, and a greater Side, or Leg given.

PROPOSITION V. Having the less of either the Difference of Latitude or Departure, and also the Course given to find the greater of either the Difference of Latitude or Departure, and also the Distance sailed?

This Proposition is solved in all Respects exactly like the former, but by a different Computer, Number III. following, by taking out the greater LEG and Distance to the less LEG and the given Course, instead of less LEG and Distance to the greater LEG, and given Course, by Computer, Number II.

The *Seaman's* Ready COMPUTER. N^o. II. A New Improvement.

Course		Given Difference of LATITUDE. Course under 45°. — Or greater LEG given, next the given $\angle$, under 45°.																Course			
		10000		20000		30000		40000		50000		60000		70000		80000		90000			
Pts.	D.	Dep.	Dist.	Dep.	Dist.	Dep.	Dist.	Dep.	Dist.	Dep.	Dist.	Dep.	Dist.	Dep.	Dist.	Dep.	Dist.	Dep.	Dist.	D.	Pts.
1	00175	10002	00349	20003	00524	30005	00698	40006	00873	50008	01047	60009	01222	70011	01396	80012	01571	90014	89		
2	00349	10006	00698	20012	01048	30018	01397	40024	01746	50030	02095	60037	02444	70043	02794	80049	03143	90055	88		
3	00491	10012	00983	20024	01474	30036	01965	40048	02456	50060	02947	60072	03438	70084	03940	80096	04421	90108	87	7	3
4	00524	10014	01048	20027	01572	30041	02096	40055	02626	50069	03144	60102	03668	70096	04192	80110	04717	90123	87		
5	00699	10024	01399	20039	02098	30073	02797	40097	03496	50122	04196	60146	04895	70171	05594	80195	06293	90220	86		
6	00873	10038	01750	20076	02625	30114	03500	40153	04374	50191	05249	60229	06124	70267	06999	80306	07874	90344	85		
7	00983	10048	01970	20097	02955	30145	03940	40193	04925	50242	05909	60290	06894	70339	07879	80387	08864	90436	84		
8	01051	10055	02102	20110	03153	30165	04204	40220	05255	50275	06306	60330	07357	70386	08408	80441	09459	90496	84		
9	01228	10075	02456	20150	03684	30225	04911	40300	06139	50375	07367	60450	08595	70525	09823	80600	11051	90675	83		
10	01405	10098	02811	20197	04216	30295	05622	40393	07027	50491	08372	60589	09838	70688	11243	80786	12645	90884	82		
11	01483	10109	02967	20219	04450	30320	05933	40438	07417	50547	08900	60657	10384	70766	11867	80875	13350	90985	81	7	3
12	01584	10125	03168	20249	04752	30374	06335	40499	07919	50623	09503	60740	11087	70873	12671	80997	14255	91122	81		
13	01763	10154	03527	20308	05290	30463	07053	40617	08816	50771	10580	60926	12343	71089	14106	81234	15869	91388	80		
14	01944	10187	03888	20374	05831	30561	07775	40749	09719	50936	11663	61123	13607	71310	15550	81497	17494	91684	79		
15	01989	10196	03978	20392	05967	30588	07956	40784	09946	50980	11935	61175	13924	71371	15913	81567	17902	91763	78		
16	02126	10223	04251	20447	06377	30670	08502	40894	10628	51117	12753	61340	14879	71564	17004	81787	19130	92011	78		
17	02309	10263	04617	20526	06926	30789	09235	41052	11543	51315	13852	61578	16161	71841	18469	82104	20778	92367	77		
18	02493	10306	04986	20612	07480	30918	09973	41224	12466	51531	14960	61837	17453	72143	19946	82444	22440	92755	76		
19	02505	10309	05010	20618	07515	30927	10019	41236	12524	51545	15029	61854	17534	72163	20039	82472	22544	92780	76	6	3
20	02679	10353	05359	20706	08038	31058	10718	41411	13397	51764	16077	62117	18756	72466	21436	82822	24115	93175	75		
21	02867	10403	05735	20806	08602	31209	11470	41612	14337	52015	17205	62418	20072	72821	22940	83224	25807	93627	74		
22	03033	10450	06067	20900	09100	31350	12334	41800	15167	52250	18201	62700	21234	73150	24265	83600	27301	94050	73		
23	03057	10457	06115	20914	09172	31271	12229	41828	15286	52285	18344	62741	21401	73198	24458	83655	27516	94112	73		
24	03249	10515	06298	21029	09748	31544	12997	42058	16246	52573	19495	63088	22744	73602	25994	84117	29245	94631	72		
25	03443	10576	06887	21152	10330	31729	13773	42305	17216	52881	20660	63457	24103	74033	27546	84610	30989	95186	71		
26	03578	10621	07156	20242	10734	31863	14312	42483	17890	53104	21468	63725	25040	74346	28624	84967	32202	95588	70	6	3
27	03640	10642	07279	21284	10919	31925	14539	42567	18198	53209	21838	63851	25478	74492	29118	85134	33757	95776	70		
28	03839	10711	07677	21423	11516	32134	15355	42846	19193	53557	23032	64269	26870	74980	30709	85692	34548	96403	69		
29	04040	10785	08081	21572	12121	32356	16161	43141	20201	53927	24242	64712	28281	74497	32322	86283	36362	97068	68		
30	04142	10824	08284	21648	12426	32472	16569	43296	20911	54120	24853	64944	28995	75767	33137	86591	37279	97415	68		
31	04245	10864	08489	21727	12734	32591	16979	43454	21224	54318	25468	65182	29713	76045	33957	86909	38203	97772	67		
32	04452	10946	08905	21893	13357	32839	17809	43785	22261	54732	26714	65678	31166	76625	35618	87571	40071	98517	66		
33	04663	11034	09326	22068	13986	33101	18652	44135	23315	55169	27978	66203	32641	77236	37304	88270	41967	99304	65		
34	04770	11062	09459	22124	14189	33186	18919	44248	23648	55310	28378	66372	33108	77434	37837	88496	42567	99558	65	5	3
35	04877	11126	09755	22252	14632	33378	19509	44504	24387	55630	29254	66756	34121	77882	39019	89008	43850	10.0134	64		
36	05095	11223	10190	22447	15286	33670	20381	44893	25476	56116	30572	67340	35667	78563	40762	89786	45927	10.1009	63		
37	05317	11326	10634	22651	15951	33977	21268	45303	26585	56628	31902	67954	37219	79280	42536	90606	47853	10.1931	62		
38	05345	11339	10660	22676	16035	34017	21380	45356	26726	56694	32071	68033	37416	79372	42761	90711	48106	10.2050	62		
39	05543	11433	11080	22867	16639	34301	22172	45734	27715	57168	33258	68601	38801	80045	44344	91468	49887	10.2502	61		
40	05773	11547	11547	23094	17320	34641	23094	46188	28867	57733	34641	69382	40414	80829	46788	92376	51961	10.3923	60		
41	05994	11655	11988	23317	17981	34970	23975	46635	29969	58293	35962	69952	41956	81613	47950	93270	53944	10.4928	59		
42	06009	11666	12017	23333	18026	34999	24034	46665	30043	58332	36052	69998	42060	81664	48069	93331	54077	10.4997	59		
43	06249	11792	12497	23584	18746	35375	24995	47167	31243	58959	37492	70751	43743	82544	49990	94334	56238	10.6126	58		
44	06494	11924	12988	23847	19483	35771	25976	47695	32470	59618	38964	71542	45458	83465	51952	95389	58446	10.7313	57		
45	06682	12027	13364	24054	20045	36081	26727	48108	33409	60134	40091	72161	46772	84188	53454	96215	60136	10.8242	56		
46	06745	12062	13490	24124	20235	36186	26980	48248	33725	60311	40470	72373	47215	84435	53960	96497	60705	10.8560	56		
47	07002	12208	14004	24415	21006	36623	28008	48831	35010	61039	42012	73246	49014	85454	56016	97262	63018	10.9870	55		
48	07265	12361	14531	24721	21796	37082	29062	49443	36327	61803	43593	74164	50858	86525	58123	98885	65389	11.1246	54		
49	07416	12450	14833	24900	22250	37350	29666	49800	37083	62250	44495	74700	51916	87150	59332	99600	66749	11.2050	53		
50	07536	12521	15071	25043	22607	37564	30142	50085	37678	62607	45213	75128	52749	87649	60284	10.0171	67820	11.2692	53	4	
51	07813	12690	15626	25380	23439	38071	31251	50761	39064	63451	46877	76141	54690	88831	62503	10.1521	70306	11.4212	52		
52	08098	12868	16196	25739	24293	38603	32391	51470	40489	64318	48387	77206	56685	90072	64783	10.2941	72881	11.5808	51		
53	08207	12936	16414	25872	24620	38809	32827	51746	41034	64682	49241	77619	57447	90355	65654	10.3491	73861	11.6428	51		
54	08391	13054	16782	26008	25173	39162	33564	52216	41955	65270	50346	78024	58737	91378	67128	10.4432	75519	11.7486	50		
55	08693	13250	17388	26506	26079	39750	34771	53000	43464	66250	52157	79500	60850	92750	69543	10.6000	78236	11.9256	49		
56	09004	13456	18008	26913	27012	40369	36016	53825	45020	67282	54024	80738	63028	94194	72032	10.7651	81036	12.1107	48		
57	09063	13496	18127	26992	27190	40488	36254	53985	45317	67481	54381	80977	63444	94473	72508	10.7969	81371	12.1465	48		
58	09325	13673	18650	27347	27975	41020	37301	54693	46626	68366	55951	82040	65276	95713	74601	10.9386	83926	12.3055	47		
59	09657	1																			

CONSTRUCTION. 1, 2, 3, 4, 5, 6, 7, 8, 9 Times the natural *Tangents* and *Secants* to Radius 10000.

The

The Seaman's Ready COMPUTER. N^o. III. A New Improvement.

Z		Base.	Hyp.	Base.	Hyp.	Base.	Hyp.	Base.	Hyp.	Base.	Hyp.	Base.	Hyp.	Z
Given DEPARTURE. Course under 45°. — Or Left LEG, opposite an Z given, under 45°.														
Course		10000		20000		30000		40000		50000		60000		Course
Pts.	D.	D. Lat.	Diff.	D. Lat.	Diff.	D. Lat.	Diff.	D. Lat.	Diff.	D. Lat.	Diff.	D. Lat.	Diff.	Pts.
1	1	57.2900	57.2987	114.5799	114.5974	171.8699	171.8961	229.1598	229.1948	286.4498	286.4934	343.7398	343.7921	89
2	2	28.6363	28.6537	57.2725	57.3074	85.9088	85.9611	114.5450	114.6148	143.1813	143.2685	171.8175	171.9222	88
3	3	20.3556	20.3802	40.7112	40.7603	61.0668	61.1404	81.4224	81.5206	101.7780	101.9007	122.1336	122.2809	87
4	4	19.0811	19.1073	38.1623	38.2146	57.2434	57.3220	76.3245	76.4293	95.4057	95.5366	114.4868	114.6439	86
5	5	14.3007	14.3356	28.6013	28.6712	42.9020	43.0068	57.2027	57.3423	71.5033	71.6779	85.8040	86.0135	85
6	6	11.4300	11.4737	22.8601	22.9474	34.2902	34.4211	45.7202	45.8948	57.1502	57.3686	68.5803	68.8423	84
7	7	10.1532	10.2024	20.3064	20.4048	30.4596	30.6073	40.6128	40.8097	50.7660	51.0121	60.9192	61.2145	83
8	8	9.5144	9.5668	19.0287	19.1335	28.5431	28.7003	38.0575	38.2671	47.5718	47.8338	57.0862	57.4006	82
9	9	8.1443	8.2055	16.2887	16.4110	24.4330	24.6165	32.5774	32.8220	40.7217	40.2775	48.8661	49.2330	81
10	10	7.1154	7.1853	14.2307	14.3706	21.3461	21.5559	28.4615	28.7412	35.5768	35.9265	42.6922	43.1118	80
11	11	6.7415	6.8152	13.4829	13.6305	20.2244	20.4457	26.9658	27.2600	33.7073	34.0761	40.4487	40.8914	79
12	12	6.3138	6.3925	12.6275	12.7849	18.9413	19.1774	25.2550	25.5698	31.5688	31.9622	37.8825	38.3547	78
13	13	5.6713	5.7588	11.3426	11.5175	17.0138	17.2763	22.6851	23.0351	28.3564	28.7939	34.0277	34.5526	77
14	14	5.1446	5.2408	10.2891	10.4817	15.4337	15.7225	20.5782	20.9634	25.7228	26.2042	30.8673	31.4451	76
15	15	5.0273	5.1258	10.0547	10.2517	15.0820	15.3775	20.1094	20.5033	25.1367	25.6292	30.1640	30.7550	75
16	16	4.7046	4.8097	9.4093	9.6195	14.1139	14.4292	18.8135	19.2389	23.5232	24.0486	28.2278	28.8584	74
17	17	4.3315	4.4454	8.6630	8.8908	12.9944	13.3362	17.3259	17.7816	21.6574	22.2271	25.9889	26.6725	73
18	18	4.0108	4.1336	8.0215	8.2671	12.0323	12.4007	16.0431	16.5343	20.0539	20.6678	24.0647	24.8014	72
19	19	3.9922	4.1156	7.9844	8.2311	11.9767	12.3467	15.9689	16.4623	19.9611	20.5778	23.9533	24.6934	71
20	20	3.7321	3.8637	7.4641	7.7274	11.1962	11.5911	14.9280	15.4548	18.6603	19.3185	22.392	23.1822	70
21	21	3.4874	3.6280	6.9741	7.2559	10.4622	10.8839	13.9496	14.5118	17.4371	18.1398	20.9245	21.7677	69
22	22	3.2966	3.4449	6.5931	6.8898	9.8897	10.3347	13.1862	13.7796	16.4828	17.2245	19.7794	20.6694	68
23	23	3.2709	3.4203	6.5417	6.8406	9.8126	10.2609	13.0834	13.6812	16.3543	17.1015	19.6251	20.5218	67
24	24	3.0777	3.2361	6.1554	6.4721	9.2321	9.7082	12.3107	12.9443	15.3884	16.1803	18.4661	19.4164	66
25	25	2.9042	3.0716	5.8084	6.1431	8.7126	9.2147	11.6168	12.2862	14.5211	15.3578	17.4253	18.4293	65
26	26	2.7948	2.9683	5.5896	5.9367	8.3844	8.9050	11.1793	11.8733	13.9741	14.8416	16.7689	17.8100	64
27	27	2.7475	2.9238	5.4950	5.8276	8.1424	8.7714	10.9899	11.6952	13.7374	14.6190	16.4849	17.5428	63
28	28	2.6051	2.7904	5.2102	5.5809	7.8153	8.3713	10.4203	11.1617	13.0254	13.9521	15.6305	16.7426	62
29	29	2.4751	2.6695	4.9502	5.3389	7.4253	8.0084	9.9003	10.6779	12.3754	13.3473	14.8505	16.0168	61
30	30	2.4142	2.6131	4.8234	5.2261	7.2426	7.8394	9.6569	10.4525	12.0711	13.0656	14.4853	15.6788	60
31	31	2.3559	2.5593	4.7117	5.1186	7.0676	7.6779	9.4234	10.2372	11.7793	12.7965	14.1351	15.3558	59
32	32	2.4460	2.4586	4.4921	4.9172	6.7381	7.3758	8.9841	9.8354	11.2302	12.2930	13.4762	14.7516	58
33	33	2.1445	2.3662	4.2890	4.7324	6.4335	7.0986	8.5780	9.4648	10.7225	11.8310	12.8670	14.1972	57
34	34	2.1143	2.3389	4.2286	4.6778	6.3435	7.1666	8.4573	9.3555	10.5716	11.6944	12.6856	14.0333	56
35	35	2.0503	2.2812	4.1006	4.5623	6.1509	6.8435	8.2012	9.1247	10.2515	11.4059	12.3018	13.6770	55
36	36	1.9626	2.2027	3.9252	4.4054	5.8878	6.6081	7.8504	8.8108	9.8131	11.0134	11.7757	13.2161	54
37	37	1.8807	2.1301	3.7615	4.2601	5.6422	6.3902	7.5229	8.5202	9.4036	10.6503	11.2844	12.7803	53
38	38	1.8709	2.1234	3.7417	4.2427	5.6126	6.3641	7.4835	8.4854	9.3543	10.6068	11.2252	12.7281	52
39	39	1.8040	2.0627	3.6081	4.1253	5.4121	6.1880	7.2162	8.2507	9.0202	10.3133	10.8243	12.3760	51
40	40	1.7321	2.0000	3.4641	4.0000	5.1062	6.0000	6.9282	8.0000	8.6603	10.0000	10.3923	12.0000	50
41	41	1.6684	1.9451	3.3368	3.8903	5.0052	5.8354	6.6736	7.7805	8.3420	9.7257	10.0110	11.6708	49
42	42	1.6643	1.9416	3.3286	3.8832	4.9928	5.8245	6.6571	7.7664	8.3214	9.7080	9.9857	11.6496	48
43	43	1.6003	1.8871	3.2007	3.7742	4.8010	5.6612	6.4013	7.5483	8.0017	9.4354	9.6020	11.3225	47
44	44	1.5399	1.8361	3.0797	3.6722	4.6196	5.5082	6.1595	7.3443	7.6993	9.1804	9.2362	11.0165	46
45	45	1.4996	1.8000	2.9993	3.5999	4.4988	5.3998	5.9984	7.1998	7.9980	8.9998	8.9976	10.7997	45
46	46	1.4826	1.7883	2.9651	3.5666	4.4477	5.3445	5.9302	7.1332	7.4128	8.9415	8.8954	10.7297	44
47	47	1.4281	1.7434	2.8563	3.4860	4.2844	5.2303	5.7126	6.9738	7.1407	8.7172	8.5689	10.4607	43
48	48	1.3764	1.7013	2.7528	3.4026	4.1291	5.1033	5.5055	6.8052	6.8819	8.5065	8.2583	10.2078	42
49	49	1.3483	1.6787	2.6967	3.3574	4.0450	5.0361	5.3934	6.7160	6.7417	8.3915	8.0901	10.0722	41
50	50	1.3270	1.6616	2.6541	3.3233	3.9811	4.9846	5.3082	6.6466	6.6352	8.3082	7.9623	9.9698	40
51	51	1.2799	1.6243	2.5599	3.2485	3.8398	4.8728	5.1198	6.4972	6.3997	8.1213	7.7696	9.7458	39
52	52	1.2345	1.5890	2.4698	3.1780	3.7047	4.7670	4.9396	6.3561	6.1745	7.9451	7.4094	9.5341	38
53	53	1.2185	1.5763	2.4370	3.1527	3.6555	4.7290	4.8740	6.3053	6.0925	7.8816	7.3110	9.4580	37
54	54	1.1918	1.5557	2.3835	3.1114	3.5753	4.6672	4.7670	6.2229	5.9588	7.7786	7.1505	9.3343	36
55	55	1.1504	1.5243	2.3007	3.0485	3.4511	4.5728	4.6015	6.0970	5.7518	7.6213	6.9021	9.1455	35
56	56	1.1106	1.4945	2.2212	2.9890	3.3318	4.4834	4.4425	5.9779	5.5531	7.4724	6.6637	8.9669	34
57	57	1.1033	1.4891	2.2067	2.9781	3.3100	4.4672	4.4133	5.9562	5.5167	7.4454	6.6000	8.9344	33
58	58	1.0724	1.4663	2.0447	2.9326	3.2171	4.3988	4.2895	5.8651	5.3618	7.3314	6.4342	8.7977	32
59	59	1.0355	1.4396	2.0711	2.8791	3.1066	4.3187	4.1421	5.7582	5.1777	7.1978	6.2132	8.6373	31
60	60	1.0000	1.4142	2.0000	2.8284	3.0000	4.2426	4.0000	5.6569	5.0000	7.0711	6.0000	8.4853	30
Pts.	D.	Dep.	Diff.	Dep.	Diff.	Dep.	Diff.	Dep.	Diff.	Dep.	Diff.	Dep.	Diff.	Pts.
Course		10000		20000		30000		40000		50000		60000		Course
Given Diff. of LATITUDE. Course above 45°. — Or Left LEG next an Z given above 45°.														

CONSTRUCTION. 1, 2, 3, 4, 5, 6, 7, 8, 9 Times the natural Cotangents and Cosecants, to Radius 10000.

COMPUTER.

COMPUTER N^o. III. Continued.

Z		Bafe.		Hyp.		Bafe.		Hyp.		Bafe.		Hyp.		Z	
Given Departure. Course under 45°.															
Course.		70000				80000				90000				Course.	
Pts.	D.	D. Lat.		Diff.		D. Lat.		Diff.		D. Lat.		Diff.		D.	Pts.
1	...	401.0297	401.0908	458.3197	458.3894	515.6097	515.6882	89	...	...	...	...	...	...	...
2	...	200.4538	200.5759	229.0900	229.2296	257.7263	257.8833	88	...	...	...	...	...	...	...
3	...	142.4892	142.6611	162.8448	163.0412	183.2004	183.4214	...	7 3/4	...	...	...	...	...	...
4	...	133.5679	133.7513	152.6491	152.8586	171.7302	171.9659	87	...	...	...	...	...	...	...
5	...	100.1047	100.3491	114.4053	114.6847	128.7060	129.0202	86	...	...	...	...	...	...	...
6	...	80.0103	80.3160	91.4404	91.7897	102.8704	103.2634	85	...	...	...	...	...	...	...
7	...	71.0724	71.4169	81.2255	81.6194	91.3787	91.8218	...	7 1/2	...	...	...	...	...	...
8	...	66.6006	66.9674	76.1149	76.5341	85.6292	86.1009	84	...	...	...	...	...	...	...
9	...	57.0104	57.4385	65.1548	65.6440	73.2991	73.8495	83	...	...	...	...	...	...	...
10	...	49.8076	50.2971	56.9230	57.4824	64.0383	64.6677	82	...	...	...	...	...	...	...
11	...	47.1502	47.7066	53.9316	54.5218	60.6701	61.3370	...	7 1/4	...	...	...	...	...	...
12	...	44.1963	44.7472	50.5100	51.1396	56.8238	57.5321	81	...	...	...	...	...	...	...
13	...	39.6990	40.3114	45.3702	46.0702	51.0415	51.8289	80	...	...	...	...	...	...	...
14	...	36.0119	36.6859	41.1564	41.9267	46.3010	47.1676	79	...	...	...	...	...	...	...
15	...	35.0913	35.8308	40.2187	41.0066	45.2460	46.1330	...	7	...	...	...	...	...	...
16	...	32.9324	33.6681	37.6370	38.4779	42.3417	43.2876	78	...	...	...	...	...	...	...
17	...	30.3203	31.1179	34.6518	35.5633	38.9833	40.0087	77	...	...	...	...	...	...	...
18	...	28.0755	28.9350	32.0862	33.0685	36.0971	37.2020	76	...	...	...	...	...	...	...
19	...	27.5456	28.8089	31.9378	32.9245	35.9300	37.0401	...	6 3/4	...	...	...	...	...	...
20	...	26.1244	27.0459	29.8564	30.9096	33.5884	34.7733	75	...	...	...	...	...	...	...
21	...	24.4119	25.3957	27.8993	29.0236	31.3867	32.6516	74	...	...	...	...	...	...	...
22	...	23.0759	24.1143	26.3724	27.5584	29.6690	31.0041	...	6 1/2	...	...	...	...	...	...
23	...	22.8959	23.9421	26.1668	27.3624	29.4377	30.7827	73	...	...	...	...	...	...	...
24	...	21.5438	22.6525	24.6225	25.8885	27.6992	29.1246	72	...	...	...	...	...	...	...
25	...	20.3295	21.5009	23.2337	24.5724	26.1379	27.6440	71	...	...	...	...	...	...	...
26	...	19.5637	20.7783	22.3585	23.7466	25.1533	26.7150	...	6 1/4	...	...	...	...	...	...
27	...	19.2322	20.4666	21.9798	23.3904	24.7273	26.3142	70	...	...	...	...	...	...	...
28	...	18.2356	19.5330	20.8407	22.3234	23.4458	25.1139	69	...	...	...	...	...	...	...
29	...	17.3256	18.6863	19.8007	21.3557	22.2758	24.0253	68	...	...	...	...	...	...	...
30	...	16.8995	18.2919	19.3137	20.9050	21.7279	23.5181	...	6	...	...	...	...	...	...
31	...	16.4910	17.9151	18.8468	20.4744	21.2027	23.0337	67	...	...	...	...	...	...	...
32	...	15.7223	17.2102	17.9683	19.6687	20.2143	22.1273	66	...	...	...	...	...	...	...
33	...	15.0115	16.5634	17.1560	18.9296	19.3005	21.2958	65	...	...	...	...	...	...	...
34	...	14.8003	16.3722	16.9146	18.7110	19.0289	21.0499	...	5 3/4	...	...	...	...	...	...
35	...	14.3521	15.9682	16.4024	18.2494	18.4527	20.5305	64	...	...	...	...	...	...	...
36	...	13.7383	15.4188	15.7009	17.6215	17.6635	19.8242	63	...	...	...	...	...	...	...
37	...	13.1651	14.9104	15.0458	17.0404	16.9265	19.1700	62	...	...	...	...	...	...	...
38	...	13.0961	14.8495	14.9669	16.9708	16.8378	19.0922	...	5 1/2	...	...	...	...	...	...
39	...	12.6283	14.4387	14.4324	16.5013	16.2364	18.5640	61	...	...	...	...	...	...	...
40	...	12.1244	14.0000	13.8564	16.0000	15.5885	18.0000	60	...	...	...	...	...	...	...
41	...	11.6788	13.6159	13.3472	15.5611	15.0156	17.5062	...	5 1/4	...	...	...	...	...	...
42	...	11.6500	13.5912	13.3142	15.5328	14.9785	17.4744	59	...	...	...	...	...	...	...
43	...	11.2023	13.2096	12.8027	15.0966	14.4030	16.9837	58	...	...	...	...	...	...	...
44	...	10.7791	12.8525	12.3189	14.6886	13.8588	16.5247	57	...	...	...	...	...	...	...
45	...	10.4972	12.6000	11.9968	14.3996	13.4964	16.1996	...	5	...	...	...	...	...	...
46	...	10.3779	12.5180	11.8605	14.3063	13.3439	16.0946	56	...	...	...	...	...	...	...
47	...	99970	12.2041	11.4252	13.9476	12.8533	15.6910	55	...	...	...	...	...	...	...
48	...	96347	11.9091	11.0110	13.6104	12.3874	15.3117	54	...	...	...	...	...	...	...
49	...	94384	11.7509	10.7868	13.4296	12.1351	15.1083	...	4 3/4	...	...	...	...	...	...
50	...	92893	11.6315	10.6164	13.2931	11.9434	14.9548	53	...	...	...	...	...	...	...
51	...	89596	11.3699	10.2395	12.9942	11.5195	14.6184	52	...	...	...	...	...	...	...
52	...	86443	11.1231	98792	12.7121	11.1141	14.3013	51	...	...	...	...	...	...	...
53	...	85295	11.0343	97480	12.6106	10.9665	14.1870	...	4 1/2	...	...	...	...	...	...
54	...	83423	10.8901	95340	12.4458	10.7258	14.0015	50	...	...	...	...	...	...	...
55	...	80526	10.6698	92029	12.1940	10.3533	13.7183	49	...	...	...	...	...	...	...
56	...	77748	10.4613	88849	11.9550	99955	13.4503	48	...	...	...	...	...	...	...
57	...	77235	10.4235	88266	11.9126	99300	13.4016	...	4 1/4	...	...	...	...	...	...
58	...	75166	10.2640	85789	11.7302	96513	13.1965	47	...	...	...	...	...	...	...
59	...	72487	10.0769	82842	11.5165	93198	12.9560	46	...	...	...	...	...	...	...
60	...	70000	98995	80000	11.3137	90000	12.7279	45	...	...	...	...	...	...	...
Pts.	D.	Dep.		Diff.		Dep.		Diff.		Dep.		Diff.		D.	Pts.
Course.		70000				80000				90000				Course.	
Given Dif. Latitude. Course above 45°.															

USE of the Seaman's COMPUTER.

EXAMPLE I. A Ship's Departure is 54 Miles on a Course of 40° South westerly, required her Difference of Latitude and Distance sailed?

By the Computer, Number III.
 Course. | Dep. W. | D. Lat. S. | Dist.
 Agt. 40° | Und. 50 | 59.588 | 77.786
 4 | 4.767 | 6.222
 Miles 54 | 64.355 | 84.008
 Answer

EXAMPLE II. A Ship's Difference of Latitude is 61 Miles North, on a Course North westerly 85°, required her Departure and Distance sailed?

Course. | Dif. Lat. N. | Dep. W.
 Against 85° | 60 | 685.803
 or 5 | 1 | 11.430
 Miles 61 | 697.233
 Dist.
 688.423
 11.473
 699.896 Answer.

PROPOSITION VI. To find the Course and Departure from the Distance and Difference of Latitude being given?

Example. The Distance being 84 Miles, and Difference of Latitude 64.3 Miles, to find the Course and Departure?

First for the Course. RULE. Divide the less by the greater Number, and seek the Decimal Quotient, (about 3 Figures being sufficient) in the Column of Dif. Lat. under 10000, of the Computer Number I. even with which, on the Side, stands the Course.

Thus, 84)64.3(.765 which Quota is found in the said Column against 40° the Course required.

Now, with the given Course and Distance as above find the Departure, by the same Computer Number I.

Course. | Dist. | Depart.
 Against 40° | Under 80 | 51.423
 4 | 2.571
 Miles 84 | 53.994 or 54
 Miles Departure, required

USE of the SEAMAN'S COMPUTER.

PROPOSITION VII. To find the Course and Difference of Latitude, from the Distance and Departure given?

Example. The Distance being 84 Miles, and the Departure 54 Miles, to find the Course and Difference of Latitude?

First for the Course. **RULE.** Divide the less by the greater Number, and seek the Decimal Quotient (of about 3 Figures) in the Column of Departure under 10000 of the Computer Number I. even with which on the Side stands the Course.

Thus, 84)54(.643 nearly; which *Quote* is found in the said Column against 40°, the Course required.

Now, with the given Course, and Distance, as above, find the Difference of Latitude by the same Computer Number I.

Course.	Dist.	Dif. Lat.
Against 40°	Under 80	61.284
	4	3.064
Miles 84		64.348 Dif. Lat. required.

PROPOSITION VIII. To find the Difference of Longitude from the middle Latitude and Departure being given?

The Complement of middle Latitude must be taken as the Course, and the Distance answerable to the given Departure on that Course will be the Difference of Longitude.

RULE. With the given Departure, at the Top or Bottom of the Computer, (Number II. if the Complement of middle Latitude be less than 45°, but Number III. if greater than 45°) take out the Distance, against the Course (equal to the Complement of middle Latitude) on the Side thereof; and you will have the Difference of Longitude, required.

EXAMPLE I. Our Ship being Yesterday at Noon in Latitude 37° 5' N. and this Day at Noon in Latitude 38° 55' N. (the middle Latitude being 38°, and Complement thereof 52°) our Departure 71 Miles Westerly, required our Difference of Longitude?

By Computer Number II. the Complement of middle Latitude being above 45°, viz. 52°.

Course.	Depart.	Dist.
Against 52°	Under 70 W.	88.831
or 38	1	1.269
Miles 71		90.100 Diff. Long. req.

EXAMPLE II. Our Ship being Yesterday at Noon in Latitude 46° 9' N. and this Day at Noon in 47° 51' N. (the middle Latitude being 47°, and Complement thereof 43°) our Departure easterly, 112 Miles, required the Difference of Longitude?

By Computer Number III. the Complement of middle Latitude being under 45°, viz. 43°.

Course.	Dep. E.	Dist.
Against 43°	Under 100	146.63
or 47	10	16.263
	2	2.932
Miles 112		165.825 Dif. Long. req.

N. B. If the Complement of middle Latitude is not in whole Degrees, but in Degrees and odd Minutes, you must find the Longitude for the Complement of middle Latitude a Degree less, and greater than the Complement given; adding a proportional Part of the Difference of those Longitudes for the odd Minutes to the Longitude for the less Degree or Complement of middle Latitude, and you will have the Difference of Longitude, required. But the Long. to the nearest whole Deg. of Mid. Lat. will be sufficient.

PROPOSITION VIII. To find the continual Course between two Places, and their Distance on that Course, from the Latitudes and Longitudes of those Places being given?

RULE. Find (by Computer Number I.) the Departure answering a Distance, equal to the Minutes Difference of Longitude given; then by the Difference of Latitude given, and the Departure just now found, find the Course and Distance by PROPOSITION III.

EXAMPLE. The Lizard lies in Latitude 49° 50' N. and 4° 45' W. Longitude from Greenwich; Cape Finister in the Latitude of 43° 10' and 9° 45' W. Longitude, from Greenwich; Required the constant Course, and Distance thereon, between the Lizard and Cape Finister?

Hence, the Dif. of Latitude is 6° 40' or 400 Minutes.
Dif. of Longitude 5 0 W. or 300 Minutes.

Middle Latitude 46° 30'
Complement 43 30

By Computer Number I.

Course.	Dist.	Depart.
Against 43°	Under 300	204.60
Against 44	300	208.40

Dif. 3.80
Half 1.9 +

Course 43° 30' Depart. 206.5 Miles.
Dep. 206.5

--- = 5016; which *Quote*, in Computer II. stands, in Column 10000, against 27° Course S. westerly.

By Computer, Number III.

Course.	Dep. W.	Dist.	Dif. Lat.
Against 27°	Under 200	440.54	392.52
	6	13.21	11.77
Miles 206		453.75	*404.29

Answer required.

* Here it is observable that the Difference of Latitude comes out agreeing (so far as the Nearness of the Course is taken) with the given Difference of Latitude; a Proof of the Truth of the foregoing Operation.

And thus all Cases relating to conducting a Ship over the Ocean are readily answered by the Seaman's Computer only: being an entire New, easy, and practical METHOD of NAVIGATION.

WITH Regard to the foregoing (VIIIth.) *Proposition*, the constant or continued Course steered by the Ship betwixt two Places, must be carefully distinguished from the direct Bearing of one of those Places from the other:

For any Course, on the Surface of a Globe, making the same constant Angle with the several Meridians, crossed over by the Ship's Way, is a *spiral Track* (of the Logarithmic Kind) which being continued on will terminate in that Pole of the Earth towards which the Ship's Course is directed: Except when the two Places lie under the same Meridian, directly North and South, or on the Equinoctial directly East and West of each other, on the Arch of a great Circle.

The direct Bearing of any Place from another (generally misunderstood for the continued or constant Course all the Way betwixt those Places) is the Angle of Position on the Arch of a great Circle, or nearly such, that one Place, on the convex Surface of the Globe, directly bears from the other.

The constant Course that a Ship steers betwixt any two Places on the convex Surface, being on a *spiral Line*, or nearly so, (as before observed) is in a contrary Direction sailing back, to what it was in sailing forward betwixt those Places.

The direct Bearing on the Arch of a great Circle marks the nearest Distance betwixt any two Places, on the globular Convexity; but this Bearing is continually variable all the Way.

The continually variable Angles in sailing back are not contrary to those in sailing forward, betwixt those Places: as is easily shewn by the Rules of *Spherical Trigonometry*. This has been proposed as a *Paradox*, by some, though an obvious Truth.

THEREFORE this great Circle Sailing, in keeping so many different Courses, as must be kept in sailing over the nearest Distance betwixt any two Places on the Convexity, is hardly practicable, (especially when those Places lie very remote from each other) because there is not only the Difficulty to overcome, in computing all the Variety of Courses necessary to be steered, but the Difficulty of putting them in Practice against all Impediments or Accidents whatsoever, of Winds and Weather, that may conspire to prevent it.

Nor can *elliptical Sailing*, by the Arch of an Ellipsis on the Ellipsoid, or flatted Globe next the Poles, be better put in Practice, for sailing the nearest Distance betwixt any two Places, on Account of the like Obstacles, that may prevent that Kind of Sailing taking so complicated an Effect.

Therefore practical or *spiral Sailing*, or that which has long been in Use, and lately improved, according to some constant Course and Distance run on that Course, or other Things given for finding the Requisites, as near the Truth as possible, is here laid down and exemplified, in all the Variety of Cases that can happen, for the Purpose of the *Practical Navigator*.

After we have considered, from our own Experience, (in several Voyages at Sea) the many Impediments to putting the true Theory of Navigation in Practice, with all the Advantages to Geography, from the late Discoveries made of the true Figure and Dimensions of the Earth as a Spheroid, (by the Measure of a Degree of the Earth's Meridian in Peru and Lapland) we would no Ways depreciate those Experiments, as lately published by Mr. Martin, in his Folio Work, (intituled NEW PRINCIPLES of GEOGRAPHY and NAVIGATION) so far as they have real Use and Advantage.

And as that A U T H O R says with Regard to the Theory itself, "The best Way of setting Falshood and Error in a proper Light, is to compare and confront them with Truth," (the Truth of the *spheroidal Theory* being admitted, however false the *Spheroid* given) so in Regard to putting such Theory into Practice with its superior Advantages to Mercator's and other Sailing, heretofore in Use, will best appear by comparing Examples, computed according to *spheroidal* and other Methods of Navigation.

The Error arising from the Distance (measured by the Log) and Course (steered by the Compass) being so doubtfully ascertained, will be always of far greater Concern to the Truth of a Ship's Reckoning than the Difference between *spherical* and *spheroidal* Navigation. — Till which Course and Distance can be more correctly ascertained, the smaller Correction in Theory will have (as we apprehend) but a very small Use in the Nearness of Practice. And all the Correction that can be expected of Advantage must chiefly arise from a correct Observation of the Sun, or other celestial Bodies, for ascertaining the true Latitude and Longitude of the Ship.

See all Mr. Martin's Examples of *Spheroidal Sailing* following, answered by our Seaman's Computer, (without Scale, Compasses, Chart, or Canon) and compared with the Answers by Mercator's Sailing.

OUR NEW METHOD of Computation (without Scale, Compasses, Chart, or Canon) exemplified in *Answers* to the Five CASES of SAILING, as proposed by Mr. *MARTIN* in his NEW PRINCIPLES of NAVIGATION, according to sailing by the SPHEROID; confronted and compared with the ANSWERS to those CASES, according to MERCATOR-SAILING, by the SPHERE. With REMARKS and INFERENCES drawn on these different METHODS of SAILING.

I. CASE. To find the Course and Distance run, from the Difference of Latitude and Longitude, given?

Example. A Ship sails from a Port in Latitude 38° , and arrives at another Port in Latitude 5° , and finds her Difference of Longitude to be 43° ; required from thence her Course and Distance sailed?

By Tables, P. 290, 288.

Latitude.	Spheroidal. Mer. Parts.	Spherical. Mer. Parts.
Under 38°	2452	2468
5	298	300
Dif. 33°	2154	2168
= $1980'$	Mer. Dif. Lat.	Mer. Dif. Lat.

By Computer N^o. III.

Course.	Dif. Lat.	Dist. run.
NOW, against 50°	1000	1555,72
	900	1400,15
	80	124,46

By Mercator : . 1980 3080,33
By Spheroidal Sailing . . 3071, from below.

9' Diff. in sailing 33° Dif. Lat.

[whereas a small Error in the Course will cause a vastly greater Diff. in the Distance.

Dif. Longitude $43^{\circ} = 2580'$

First, $\frac{2154}{2580} = ,8349$, against which, under Col. 10000 Computer N^o. II. stands $50^{\circ}, 8'$ the Course; (the Dif. Longitude being more than the Dif. Latitude.) } according to spheroidal Sailing.

Or, $\frac{2168}{2580} = ,8402$. . Against which under Col. 10000 same Computer, stands $49^{\circ} 58'$, or nearly 50° Course. } according to Mercator-Sailing.

This CASE is now reduced to one of plain Sailing, the Course and Difference of Latitude given to find the Distance run?

By Table P. 294.

Lat.	Length of Arcs.
38°	2266,35
5	297,76

33° 1968,59 true Dif. Lat. by the [spheroidal Figure of the Earth.]

By Computer N^o. III.

Course.	Dif. Lat.	Distance run.
Against 50°	1000	1555,72
	900	1400,15
	60	93,34
	9	14,00
	1969	3063,21
	1968,6	3071.

According to Mr. Martin's Navigation, p. 52. The Course being $50^{\circ} 8'$. . . 1968,6 [8' only. Dif. 7',79, or about

REMARK. Since no Ship can steer to $8'$, or nearer than to 1° , in her Course; therefore OUR COMPUTERS N^o. I. II. III. for each Degree of Course, are sufficiently exact for the practical Purpose of Navigation, according to Plain, Middle-Latitude, Mercator, or Spheroidal Sailing. For the Error in the Ship's Reckoning, occasioned by the Course, and Distance run, that cannot be kept or determined to Exactness, or hardly to a Nearness, will always be considerably greater than the Error or Difference, arising from the DIFFERENT METHODS of SAILING, whether by old or new Principles.

INFERENCE. That unless Improvement can be made in keeping the Course, and measuring the Distance run, to a greater Exactness than has been hitherto practicable, or known by Navigators, the minuter Improvement of spheroidal Navigation will be so far an idle Refinement, (as observed by Mr. Emerson in the Preface to his Naviga-

SPHERICAL and SPHEROIDAL NAVIGATION compared.

tion) that it will be a *fruitless* and vain Attempt. For, since the *Course* steered, and *Distance* run, by the Ship, are generally *precarious*, and largely different (*more or less*) from Truth, the Correction of the *left Error*, from the Difference between the spherical and true Figure of the Earth, will be of no Advantage to the Ship's *Reckoning*, for long Runs, since the larger Error, from the Course and Distance sailed, will be as often *increased* thereby as diminished. For which Reason this small Correction may sometimes prove an *Ignis fatuus*, in long Runs, instead of a *Taper*, to guide the Mariner. Besides a Ship's *Reckoning*, kept from Day to Day, by *Mercator*, or *Spheroidal* Sailing, will have an insignificant or insensible Difference in Practice; since the Difference of meridional Parts, of both Kinds, for a *short Run*, are nearly the same.

Therefore, all Improvement of *real* Advantage in Navigation chiefly consists in Correction of the Ship's Place, by *astronomical Observation*, or in some Method, to determine thereby the *Longitude*, as well as *Latitude*, of the Ship; to be applied with the *common Methods* of Navigation in Practice.

But, we have given the meridional Parts of the Earth as a *Spheroid*, in a Table following, *confronted* with another Table of the meridional Parts of the Earth as a *Sphere*, for the Navigator's Curiosity and Satisfaction. Who may thereby make *Trial* of either, and their Difference, in computing the Ship's Place, as we have here exemplified, or put in Practice, for his Use, in long Runs; which seldom or never happen in PRACTICAL NAVIGATION.

Therefore Mr. *Murdoch*, and Mr. *Martin* after him, having exemplified *spheroidal* (or *speculative*) Navigation, in long Runs, by Cases never happening in Practice, have thereby concealed the INUTILITY OF THAT DOCTRINE (except to Geography) as much as possible.

II. CASE. To find the Difference of LONGITUDE and DISTANCE sailed from the Difference of LATITUDE and COURSE given?

Example. A Ship sails from a Place in 25° North Latitude to another in 30° of South Latitude, upon a Course of 43° , to find the Difference of Longitude and Distance sailed?

By Tables, P. 289, 290.

Lat.	Spheroidal. Mer. Parts.	Spherical. Mer. Parts.
Under 25°	1539	1550
30	1875	1888
Dif. Lat. Sum 55° = 3300'	3414	3438

} According to the Spheroid.

By Computer No. II.

Course.	Mer. Dif.	Dif. Long. for Depart.
Against 43°	3000	2797,50
	400	373,01
	10	9,32
	4	3,73
	3414	3183,56
		= $53^{\circ} 4'$ Dif. Longitude.

By Mr. Martin's Navigation, P. 53. . . . 3184,1 = $53^{\circ} 4'$.

By Tab. P. 294.	Length of Arcs.
25°	1489,79
30	1788,26
55°	3278,05 true Dif. Lat. by the spheroidal Figure of the Earth.

By our Computer, No. II.	
Against 43°	3000
	400
	30
	8
	3438
	3205,94
	= $53^{\circ} 26'$ Dif. Long.

Now, To find the Distance run from the true Difference of Latitude 3278', or from the common Difference 3300' given; a plain-sailing Case.

By Computer, Number II.

Course.	Dif. Lat.	Dist.
Against 43°	3000	4102,00
	200	273,46
	70	95,71
	8	10,94
	3278	4482,11
By Mr. Martin's Navigation, P. 53. . . .		4482,11 by the Spheroid.

Or, Against 43°

Dif. Lat.	Dist.
3000	4102,00
300	410,20
3300	4512,20
	4482,11
	Distance run by com. Method. by Spheroid.
	But 30',09 Dif. Dist. in sailing 55° Dif. Latitude.

SPHERICAL and SPHEROIDAL NAVIGATION compared.

III. CASE. To find the Difference of LONGITUDE and the COURSE, from the Difference of LATITUDE and DISTANCE sailed, being given?

Example. A Ship sailed from the Latitude 25° North, to Latitude 30° South, and her Distance sailed measured 4482,1 Miles, as in the foregoing Example, to find the Course and Difference of Longitude?

Given, $\frac{3278 \text{ Miles Dif. Lat.}}{4482,1 \text{ Dist. run}} = ,7313$, and stands in Computer, No. I. in Col. 10000, against 43° Course, req^d.

Meridional Dif. Long. 3414 , or 3438 , found before.
By Computer, No. III.

Courfe.	Spheroidal Mer. Parts.	Dif. Long. for Dep.
Against 43°	3000	2797.5
	400	373.01
	10	9.325
	4	3.730
	3414	3183.565

According to
Spheroidal
Sailing.

Courfe.	Spherical Mer. Parts.	Dif. Long. for Dep.
Against 43°	3000	2797.50
	400	373.01
	30	27.97
	8	7.46
	3438	3205.94

By Mr. Martin's Nav. P. 54. $3184.1 = 53^{\circ} 4'$ Dif. Long. as before.

$= 53^{\circ} 26'$ Dif. Long. as before.

.5 Diff.
owing to the spheroidal meridional Parts,
being taken 3414 , instead of 3414.6 .

IV. CASE. To find the Difference of LATITUDE and DISTANCE run, from the COURSE and Difference of LONGITUDE, given?

Example. A Ship sails from a Port in Latitude $51^{\circ} 15'$, steering a Course of $35^{\circ} 30'$ with the Meridian, till she finds her Difference of Longitude $28^{\circ} 30' = 1710$ Miles, to find her Difference of Latitude and Distance run?

By Computer, Number III.

Courfe.	Dif. Long. for Dep.	Mer. Parts.
$35^{\circ} 30'$	1000	1402,2
	700	981,58
	10	14,02
	1710	2397,80

Mer. Parts of
Dif. of Lat.

Lat.	Spheroidal Mer. Parts.	Spheric Mer. Parts.
Now, $51^{\circ} 15'$	3572,5	3592,5
Subtract	2397,8	2397,8

Mer. Parts of Lat. arrived in $= 1174,7$
Martin's Navigation, P. 54. $\dots 19 \cdot 21$ $= 19^{\circ} 21'$ Latitude required.
Same.

How small the Difference by spheroidal and Mercator-Sailing, in $31^{\circ} 54'$ or $31^{\circ} 44'$ Dif. Latitude.

N. B. If Merid. Parts of the Dif. of Lat. are greater than the Mer. Parts of the Lat. given, you must deduct the latter from the former, and the Remainder will answer to the Lat. on the contrary Side of the Equator.

If the Latitude had been $22^{\circ} 40'$ (instead of $51^{\circ} 15'$)

Spheroid. Meridional Parts = 1387
2398 Spheric. Meridional Parts = 1397
2398

Answering to Latitude $16^{\circ} 44' S.$ $\dots 1011$ Dif.
22 40

Dif. Lat. $\dots 39 \ 24$ Spheroidal Sailing.

Lat. $16^{\circ} 27' S.$ $\dots 1001$ Dif.
22 40

Dif. Lat. $39 \ 7$ Spherical Sailing.

How small the Difference, only $17'$, Refinement, in sailing betwixt so distant a Parallel: and therefore what will be the Odds betwixt spheroidal and Mercator Sailing for daily Reckonings?

SPHERICAL and SPHEROIDAL NAVIGATION compared.

NOW to find the Distance sailed from the Course and Difference of Latitude given; being a Case of plain-Sailing?

...ing :

By Tab. P. 294.	Lat.	Length Arcs	} In Mr. Martin's Navigation the Arcs are erroneously given, as follow	{ Egregious Error! 2999.85 1271.2	} Confronted with Truth on the Left.
	51° 15' 19 21	3059.81 1152.91			

31 54 = 19 14'	1906.90 about 1907'	} Which Author has taken out for 50° 15' instead of 51° 15' 21 20 of for 19 21

By Computer, Number II.

Course.	Dif. Lat.	Dist.
Against 35° 30'	1000	1228.42
	900	1105.58
	7	8.59
	1907	2342.59

By Mr. Martin's Nav. P. 55.

Dist. run.
2123.

By same Computer, Number II.

Course.	Dif. Lat.	Dist.
Or, Against 35° 30'	1000	1228.42
	900	1105.58
	10	12.28
	4	4.91
	1914	2351.19

219.57 + Error of Mr. Martin's Navigation.

219.57 + Error of Mr. Martin's Navigation.

V. CASE. To find the Difference of LATITUDE and Difference of LONGITUDE from the Course and Distance run, being given?

THIS is the most frequent and useful Case occurring in Navigation, though considered and introduced last of all by the Author of *New Principles of Geography and Navigation*, whose Questions are proposed all along (like the Reverend Mr. Murdoch's, the original Proprietor of this new Improvement) quite out of Practice, or such as never occur in practical Sailing. — So that if Questions in this new Method, had been proposed to work Traverses, or a Ship's daily Reckoning, as is the chief Business of Navigators, (instead of working thousand Mile Runs) the IDLE REFINEMENT of this new Navigation (observed by so able a Judge as Mr. EMERSON) would appear obvious to the meanest Capacity in Science. And no other Advantage can arise from considering the Earth as a Spheroid but to Geography, Map-making, and small Correction of celestial Observation: The Advantage thereof to Navigation plainly appearing to be none. Therefore, in Imitation of Mr. Emerson, whose Judgment and Method in all he writes is exemplary, we have given superior Place to the meridional Parts of the SPHERE, as being very sufficient for all the Purposes of practical Navigation: at the same Time they correspond with the charming Elegance and correct Solution of all Questions of Spherical NAVIGATION, by the Logarithm-Tangents of the half Complements of Latitudes, as demonstrated and exemplified by the late famous Dr. HALLEY, in *Philos. Transact.* Number 219.

Example to the foregoing Case. A Ship runs 4482.1 Miles upon a Course of 43° from the Meridian, Southward, from a Port lying in Latitude 25° N. to determine from thence her Difference of Latitude and Longitude?

FIRST for the Difference of Latitude, by Computer Number I.

Course.	Dist.	Dif. Lat.	} Being a Case of PLAIN-SAILING.
Against 43°	4000	2925.4	
	400	292.54	
	80	58.51	
	2	1.46	
	.1	.07	
	4482.1	3277.98	Dif. Lat.

By Mr. Martin's Navigation, P. 55. . . . 3278. Dif. Lat. according to his new Principles of Navigation.

Deduct the Length of the Arc 25°.	} 1489.79	} Or 3278'
by Tab. P. 294.		
Remaining Arc on other Side of the	} 1788.21	} 1778 = 29 38'
Equator = 30°		
+ 25		

Dif. Lat. . . 55° required. Or $\frac{3278}{60} = 54° 38'$ Dif. Lat. without Refinement.

SPHERICAL and SPHEROIDAL NAVIGATION compared.

NOW to find the Difference of Longitude, from the Course 43° , and Difference of Latitude 55° (or $54^\circ 38'$) being given?

				By Computer, Number II.				
		Lat.	Spheroid. Mer. Parts.	Spheric. Mer. Parts.	Course.	Mer. D.	Dif. Lon. for Dep.	According to spheroidal Sailing = 53° 4' Lon. gitude re- quired.
By Tab. P. 289, 290.	25° N.		1539	1550	Against 43°	3000	2797.50	
	30 S.		1875	1888		400	373.01	
Sum		55°	3414	3438		10	9.32	
		D. Lat.	Mer. D. Lat.	Mer. D. Lat.		4	3.73	
						3414	3183.56	

According to spheroidal Sailing = $53^\circ 4'$ Longitude required.

Mr. Martin's Navigation, P. 55. . . . 3184.

		Lat.	Spheric. Mer. Parts	Course.	Merid. D. Lat.	Dif. Long. for Dep.			Course.	Merid. Dif. Lat.	D. Long. for Dep.	According to Mercator Sailing.	
By Tab.													
P. 289	25° N.		1550	43°	3000	2797.50			43°	3000	2797.50		
	29 38 S.		1863		400	373.01				400	373.01		
					10	9.32				30	27.97		
Sum	54 38		3413		3	2.79				8	7.46		
						3182.62							
 How near this Answer = 53° 3' to the													
											3438	3205.94	[= 53° 25' Dif. Long. reqd.]

How near this Answer = $53^\circ 3'$ to the refined Answer in Mr. Martin's Navigation, and yet this Nearness is found by meridional Parts of the Sphere.

[= $53^\circ 26'$ Dif. Long. reqd.]

FROM what is exemplified in the foregoing Pages, it may be observed, that our COMPUTERS, No. I. II. III. equally serve for all the METHODS OF NAVIGATION. And that when Longitude is concerned, the meridional Difference of Latitude must be used instead of common Difference of Latitude, used in Cases of Plain Sailing. The only Difference between plain and Mercator, or even spheroidal, Sailing, is in two similar plain Triangles: the plain-sailing Triangle being contained in the Mercator-sailing Triangle. The plain-sailing triangular Leg, opposite to the Course, measuring what is called Departure; and the Mercator triangular Leg, opposite to the same Course, measuring the Longitude, in the Equator, or on the contrary Side thereof, to the first Place of given Latitude, sailed from.

FOR Satisfaction of the Proportion of the Lines on the convex-Surface of the Earth, as a Sphere, used in the different Methods of Navigation, and also the Properties of plain and spherical Triangles, see further on.

We have also shewn, further on, the Error of Mr. B. Martin's METHOD, in his *New Principles of Geography and Navigation*, of finding the Dimensions of the Earth as an oblate Spheroid, who makes Use of an incorrect Radius of Curvature (for solving those several Problems) determined by supposing the Earth a correct Sphere, and then draws his Conclusions relating to the Properties of the spheroidal Form of the Earth. We have shewn the Insufficiency of his Method by several Examples confronted with the Result of M. Maupertuis' (coinciding with Don Juan's) Method of determining the Ratio of the Earth's Diameters, from the Measure of a Degree of Latitude on the elliptic Meridian, at different and remote Places; in Lapland, France, and the at Equator.

Upon the whole of which it appears, that the Spheroid followed by B. Martin's *New Chart*, according to Maupertuis and Murdoch, is twice as much from the true Spheroid as that is from the Sphere. So that if Don Juan's Spheroid (of Diameters 266 to 265) is true, Murdoch's or Martin's Chart-Navigation are doubly erroneous compared with the Navigation of the Sphere.

Hence it follows that Mr. B. Martin has depreciated Mr. EMERSON's Judgment and Spherical Navigation, without Reason, or answering Mr. E's Reasons given. Who, with as much Reason, might depreciate the Globes, Spectacles, Vision-Glasses, and Mathematical Instruments not sold by Himself, while he boasts the Invention or Improvement of his own. Yet we are informed that Senex's Globes were first sold to Ferguson and then to Him. And that Globes, Spectacles, Vision-Glasses, and Mathematical Instruments sold by Mr. Cole, Cuff, and others in Fleetstreet, are really as perfect as those sold by B. Martin.

MERIDIONAL PARTS on Mercator's CHART, corresponding with the Earth's Convex Surface of a Sphere.

Degrees of LATITUDE.																
	0°	1°	2°	3°	4°	5°	6°	7°	8°	9°	10°	11°	12°	13°	14°	
0	0	60	120	180	240	300	361	421	481	542	603	664	725	787	848	0
2	2	62	122	182	242	302	363	423	483	544	605	666	727	789	850	2
4	4	64	124	184	244	304	365	425	485	546	607	668	729	791	852	4
6	6	66	126	186	246	306	367	427	488	548	609	670	731	793	855	6
8	8	68	128	188	248	308	369	429	490	550	611	672	733	795	857	8
10	10	70	130	190	250	310	371	431	492	552	613	674	735	797	859	10
12	12	72	132	192	252	312	373	433	494	554	615	676	737	799	861	12
14	14	74	134	194	254	314	375	435	496	556	617	678	740	801	863	14
16	16	76	136	196	256	316	377	437	498	558	619	680	742	803	865	16
18	18	78	138	198	258	318	379	439	500	560	621	682	744	805	867	18
20	20	80	140	200	260	320	381	441	502	562	623	684	746	807	869	20
22	22	82	142	202	262	322	383	443	504	564	625	686	748	809	871	22
24	24	84	144	204	264	324	385	445	506	566	627	688	750	811	873	24
26	26	86	146	206	266	326	387	447	508	568	629	690	752	813	875	26
28	28	88	148	208	268	328	389	449	510	570	631	692	754	815	877	28
30	30	90	150	210	270	330	391	451	512	573	633	695	756	817	879	30
32	32	92	152	212	272	332	393	453	514	575	635	697	758	820	881	32
34	34	94	154	214	274	334	395	455	516	577	638	699	760	822	883	34
36	36	96	156	216	276	336	397	457	518	579	640	701	762	824	886	36
38	38	98	158	218	278	338	399	459	520	581	642	703	764	826	888	38
40	40	100	160	220	280	340	401	461	522	583	644	705	766	828	890	40
42	42	102	162	222	282	342	403	463	524	585	646	707	768	830	892	42
44	44	104	164	224	284	344	405	465	526	587	648	709	770	832	894	44
46	46	106	166	226	286	346	407	467	528	589	650	711	772	834	896	46
48	48	108	168	228	288	348	409	469	530	591	652	713	774	836	898	48
50	50	110	170	230	290	350	411	471	532	593	654	715	776	838	900	50
52	52	112	172	232	292	352	413	473	534	595	656	717	778	840	902	52
54	54	114	174	234	294	354	415	475	536	597	658	719	781	842	904	54
56	56	116	176	236	296	356	417	477	538	599	660	721	783	844	906	56
58	58	118	178	238	298	358	419	479	540	601	662	723	785	846	908	58
60	60	120	180	240	300	361	421	481	542	603	664	725	787	848	910	60

MERIDIONAL PARTS on a CHART, corresponding with the Earth's Convex Surface of a Spheroid. Diameters 266 to 265.

Degrees of LATITUDE.																
	0°	1°	2°	3°	4°	5°	6°	7°	8°	9°	10°	11°	12°	13°	14°	
0	0	60	119	179	238	298	358	418	478	538	599	659	720	781	842	0
2	2	62	121	181	240	300	360	420	480	540	601	661	722	783	844	2
4	4	64	123	183	242	302	362	422	482	542	603	663	724	785	846	4
6	6	66	125	185	244	304	364	424	484	544	605	665	726	787	848	6
8	8	68	127	187	246	306	366	426	486	546	607	667	728	789	850	8
10	10	70	129	189	248	308	368	428	488	548	609	669	730	791	852	10
12	12	72	131	191	250	310	370	430	490	550	611	671	732	793	854	12
14	14	74	133	193	252	312	372	432	492	552	613	673	734	795	857	14
16	16	76	135	195	254	314	374	434	494	554	615	675	736	797	859	16
18	18	78	137	197	256	316	376	436	496	556	617	677	738	799	861	18
20	20	80	139	199	258	318	378	438	498	558	619	679	740	801	863	20
22	22	82	141	201	260	320	380	440	500	560	621	681	742	803	865	22
24	24	84	143	203	262	322	382	442	502	562	623	683	744	805	867	24
26	26	86	145	205	264	324	384	444	504	564	625	685	746	807	869	26
28	28	88	147	207	266	326	386	446	506	566	627	687	748	809	871	28
30	30	90	149	209	268	328	388	448	508	568	629	689	750	812	873	30
32	32	92	151	211	270	330	390	450	510	570	631	692	752	814	875	32
34	34	94	153	213	272	332	392	452	512	572	633	694	754	816	877	34
36	36	96	155	215	274	334	394	454	514	574	635	696	756	818	879	36
38	38	98	157	217	276	336	396	456	516	576	637	698	759	820	881	38
40	40	100	159	219	278	338	398	458	518	578	639	700	761	822	883	40
42	42	102	161	221	280	340	400	460	520	580	641	702	763	824	885	42
44	44	104	163	223	282	342	402	462	522	582	643	704	765	826	887	44
46	46	106	165	225	284	344	404	464	524	584	645	706	767	828	889	46
48	48	108	167	227	286	346	406	466	526	586	647	708	769	830	891	48
50	50	110	169	229	288	348	408	468	528	588	649	710	771	832	893	50
52	52	112	171	231	290	350	410	470	530	590	651	712	773	834	896	52
54	54	114	173	233	292	352	412	472	532	592	653	714	775	836	898	54
56	56	116	175	235	294	354	414	474	534	594	655	716	777	838	900	56
58	58	118	177	237	296	356	416	476	536	596	657	718	779	840	902	58
60	60	120	179	239	298	358	418	478	538	598	659	720	781	842	904	60

CONSTRUCTION. For the Meridional Parts of the Sphere. RULE. Multiply the Tabular Logarithm Tangent (according to Briggs's Form) of the Sum of Half the Degrees of Latitude and 45°, by the constant Number 7915.7, or 7916, and the Product will be the Meridional Parts for the given Latitude, when the Form of the Earth is considered as a perfect Sphere.

EXAMPLE. To find the Meridional Parts of the Sphere, for the Latitude of 60° ?

$30^\circ + 45^\circ = 75^\circ$, whose Logarithm Tangent = .5719475 = Log. Rad. — Log. 15°.
 Or Log. Radius Minus Log. Tangent of half the Complement of Latitude multiplied by 7916 }
 will give the Meridian Parts for the Sphere

Product = 4527.364 or 4527 Meridional Parts required.

MERIDIONAL

MERIDIONAL PARTS on Mercator's CHART, corresponding with the Earth's Convex Surface of a Sphere.

Degrees of LATITUDE.

	15°	16°	17°	18°	19°	20°	21°	22°	23°	24°	25°	26°	27°	28°	29°	
0	910	973	1035	1098	1161	1225	1289	1354	1419	1484	1550	1616	1684	1751	1820	0
2	912	975	1037	1100	1163	1227	1291	1356	1421	1486	1552	1619	1686	1753	1822	2
4	914	977	1039	1102	1166	1229	1293	1358	1423	1488	1554	1621	1688	1756	1824	4
6	917	979	1041	1104	1168	1231	1296	1360	1425	1491	1557	1623	1690	1758	1826	6
8	919	981	1044	1107	1170	1234	1298	1362	1427	1493	1559	1625	1692	1760	1828	8
10	921	983	1046	1109	1172	1236	1300	1364	1429	1495	1561	1628	1695	1762	1831	10
12	923	985	1048	1111	1174	1238	1302	1367	1431	1497	1563	1630	1697	1765	1833	12
14	925	987	1050	1113	1176	1240	1304	1369	1434	1499	1565	1632	1699	1767	1835	14
16	927	989	1052	1115	1178	1242	1306	1371	1436	1501	1568	1634	1701	1769	1838	16
18	929	991	1054	1117	1180	1244	1308	1373	1438	1504	1570	1636	1704	1771	1840	18
20	931	993	1056	1119	1183	1246	1311	1375	1440	1506	1572	1639	1706	1774	1842	20
22	933	996	1058	1121	1185	1248	1313	1377	1442	1508	1574	1641	1708	1776	1845	22
24	935	998	1060	1123	1187	1251	1315	1379	1445	1510	1576	1643	1710	1778	1847	24
26	937	1000	1062	1125	1189	1253	1317	1382	1447	1512	1579	1645	1713	1781	1849	26
28	939	1002	1064	1128	1191	1255	1319	1384	1449	1515	1581	1648	1715	1783	1851	28
30	941	1004	1067	1130	1193	1257	1321	1386	1451	1517	1583	1650	1717	1785	1854	30
32	944	1006	1069	1132	1195	1259	1323	1388	1453	1519	1585	1652	1719	1787	1856	32
34	946	1008	1071	1134	1197	1261	1326	1390	1456	1521	1587	1654	1722	1790	1858	34
36	948	1010	1073	1136	1200	1263	1328	1392	1458	1523	1590	1657	1724	1792	1861	36
38	950	1012	1075	1138	1202	1266	1330	1395	1460	1526	1592	1659	1726	1794	1863	38
40	952	1014	1077	1140	1204	1268	1332	1397	1462	1528	1594	1661	1728	1797	1865	40
42	954	1016	1079	1142	1206	1270	1334	1399	1464	1530	1596	1663	1731	1799	1868	42
44	956	1018	1081	1144	1208	1272	1336	1401	1466	1532	1599	1665	1733	1801	1870	44
46	958	1021	1083	1147	1210	1274	1338	1403	1469	1534	1601	1668	1735	1803	1872	46
48	960	1023	1085	1149	1212	1276	1341	1405	1471	1537	1603	1670	1737	1806	1874	48
50	962	1025	1088	1151	1214	1278	1343	1408	1473	1539	1605	1672	1740	1808	1877	50
52	964	1027	1090	1153	1216	1281	1345	1410	1475	1541	1607	1674	1742	1810	1879	52
54	966	1029	1092	1155	1219	1283	1347	1412	1477	1543	1610	1677	1744	1812	1881	54
56	968	1031	1094	1157	1221	1285	1349	1414	1480	1545	1612	1679	1747	1815	1884	56
58	971	1033	1096	1159	1223	1287	1351	1416	1482	1548	1614	1681	1749	1817	1886	58
60	973	1035	1098	1161	1225	1289	1354	1419	1484	1550	1616	1684	1751	1820	1888	60

MERIDIONAL PARTS on a CHART, corresponding with the Earth's Convex Surface of a Spheroid. Diameters 266 to 265.

Degrees of LATITUDE.

	15°	16°	17°	18°	19°	20°	21°	22°	23°	24°	25°	26°	27°	28°	29°	
0	904	966	1028	1090	1153	1216	1280	1344	1408	1473	1539	1605	1672	1739	1807	0
2	906	968	1030	1092	1155	1218	1282	1346	1411	1476	1541	1607	1674	1741	1809	2
4	908	970	1032	1094	1157	1220	1284	1348	1413	1478	1543	1610	1676	1743	1811	4
6	910	972	1034	1096	1159	1223	1286	1350	1415	1480	1546	1612	1678	1746	1814	6
8	912	974	1036	1099	1161	1225	1288	1353	1417	1482	1548	1614	1681	1748	1816	8
10	914	976	1038	1101	1164	1227	1291	1355	1419	1484	1550	1616	1683	1750	1818	10
12	916	978	1040	1103	1166	1229	1293	1357	1421	1487	1552	1618	1685	1753	1821	12
14	918	980	1042	1105	1168	1231	1295	1359	1424	1489	1554	1621	1687	1755	1823	14
16	920	982	1044	1107	1170	1233	1297	1361	1426	1491	1557	1623	1690	1757	1825	16
18	922	984	1046	1109	1172	1235	1299	1363	1428	1493	1559	1625	1692	1759	1827	18
20	924	986	1048	1111	1174	1237	1301	1365	1430	1495	1561	1627	1694	1762	1830	20
22	926	988	1051	1113	1176	1240	1303	1368	1432	1497	1563	1629	1696	1764	1832	22
24	928	990	1053	1115	1178	1242	1305	1370	1434	1500	1565	1632	1699	1766	1834	24
26	930	992	1055	1117	1180	1244	1308	1372	1437	1502	1568	1634	1701	1768	1836	26
28	933	995	1057	1119	1182	1246	1310	1374	1439	1504	1570	1636	1703	1771	1839	28
30	935	997	1059	1122	1185	1248	1312	1376	1441	1506	1572	1638	1705	1773	1841	30
32	937	999	1061	1124	1187	1250	1314	1378	1443	1508	1574	1641	1708	1775	1843	32
34	939	1001	1063	1126	1189	1252	1316	1380	1445	1511	1576	1643	1710	1777	1846	34
36	941	1003	1065	1128	1191	1254	1318	1383	1447	1513	1579	1645	1712	1780	1848	36
38	943	1005	1067	1130	1193	1257	1320	1385	1450	1515	1581	1647	1714	1782	1850	38
40	945	1007	1069	1132	1195	1259	1323	1387	1452	1517	1583	1649	1717	1784	1852	40
42	947	1009	1071	1134	1197	1261	1325	1389	1454	1519	1585	1652	1719	1786	1855	42
44	949	1011	1073	1136	1199	1263	1327	1391	1456	1522	1587	1654	1721	1789	1857	44
46	951	1013	1076	1138	1201	1265	1329	1393	1458	1524	1590	1656	1723	1791	1859	46
48	953	1015	1078	1140	1204	1267	1331	1396	1460	1526	1592	1658	1725	1793	1862	48
50	955	1017	1080	1143	1206	1269	1333	1398	1463	1528	1594	1661	1728	1796	1864	50
52	957	1019	1082	1145	1208	1271	1335	1400	1465	1530	1596	1663	1730	1798	1866	52
54	959	1021	1084	1147	1210	1274	1338	1402	1467	1532	1598	1665	1732	1800	1869	54
56	961	1024	1086	1149	1212	1276	1340	1404	1469	1535	1601	1667	1734	1802	1871	56
58	963	1026	1088	1151	1214	1278	1342	1406	1471	1537	1603	1670	1737	1805	1873	58
60	966	1028	1090	1153	1216	1280	1344	1408	1473	1539	1605	1672	1739	1807	1875	60

CONSTRUCTION. For the Meridional Parts of the Spheroid. RULE. Multiply the natural Sine of the Latitude (to Radius 1) by the Focal Distance from the Center of the Ellipsis, in Parts of 1, the Semi-Transverse, and the Product will be the natural Sine of an Arc. The Logarithm Tangent of the Half Complement of which Arc to 90°, being multiplied by the Product of the said Focal Distance and constant Number 7915,7, the LAST PRODUCT will be the Difference left between the Meridional Parts of the Sphere and Spheroid for the same Latitude.

EXAMPLE. To find the Meridional Parts of the oblate Spheroid, or flattened Sphere, for Latitude 60°; the Semi-Diameter of the Equator to be Semi-Diameter of the Axis being as 266 to 265.

Sine 60° = .8660254 } Product } 4° 18' } Comp. } Half } Log. Cotang. Conf. N°. Foc. Dist. } Sph. Merid. Parts } 4527
 X by Foc. Dist. .08663 } Nat. Sine. } 85° 42' } 42° 51' } .0326241 by 7915,7 X .08663 } Merid. Parts required } 4505
 = 22,37 Dif. Merid. Parts.

MERIDIONAL PARTS on Mercator's CHART, corresponding with the Earth's Convex Surface of a Sphere.

Degrees of LATITUDE.																
	30°	31°	32°	33°	34°	35°	36°	37°	38°	39°	40°	41°	42°	43°	44°	
0	1888	1958	2028	2100	2171	2244	2318	2393	2468	2545	2622	2702	2782	2863	2946	0
2	1891	1960	2031	2102	2174	2247	2320	2395	2471	2547	2625	2704	2784	2866	2949	2
4	1893	1963	2033	2104	2176	2249	2323	2398	2473	2550	2628	2707	2787	2869	2951	4
6	1895	1965	2035	2107	2179	2252	2325	2400	2476	2553	2630	2710	2790	2871	2953	6
8	1898	1967	2038	2109	2181	2254	2328	2403	2478	2555	2633	2712	2792	2874	2956	8
10	1900	1970	2040	2111	2183	2256	2330	2405	2481	2558	2636	2715	2795	2877	2959	10
12	1902	1972	2042	2114	2186	2259	2333	2408	2483	2560	2638	2717	2798	2880	2962	12
14	1904	1974	2045	2116	2188	2261	2335	2410	2486	2563	2641	2720	2801	2882	2965	14
16	1907	1977	2047	2119	2191	2264	2338	2413	2489	2566	2644	2723	2803	2885	2968	16
18	1909	1979	2050	2121	2193	2266	2340	2415	2491	2568	2646	2725	2806	2888	2971	18
20	1911	1981	2052	2123	2196	2269	2343	2418	2494	2571	2649	2728	2809	2890	2974	20
22	1914	1984	2054	2126	2198	2271	2345	2420	2496	2573	2651	2731	2811	2893	2976	22
24	1916	1986	2057	2128	2200	2274	2348	2423	2499	2575	2654	2733	2814	2896	2979	24
26	1918	1988	2059	2131	2203	2276	2350	2425	2501	2578	2657	2736	2817	2899	2982	26
28	1921	1991	2061	2133	2205	2278	2353	2428	2504	2581	2660	2739	2820	2902	2985	28
30	1923	1993	2064	2135	2208	2281	2355	2430	2506	2584	2662	2741	2822	2904	2988	30
32	1925	1995	2066	2138	2210	2283	2358	2433	2509	2586	2665	2744	2825	2907	2990	32
34	1928	1998	2069	2140	2213	2286	2360	2435	2512	2589	2667	2747	2828	2910	2993	34
36	1930	2000	2071	2143	2215	2288	2363	2438	2514	2591	2670	2749	2830	2913	2996	36
38	1932	2002	2073	2145	2217	2291	2365	2440	2517	2594	2673	2752	2833	2915	2999	38
40	1935	2005	2076	2147	2220	2293	2368	2443	2519	2597	2675	2755	2836	2918	3002	40
42	1937	2007	2078	2150	2222	2296	2370	2445	2522	2599	2678	2758	2839	2921	3005	42
44	1939	2009	2080	2152	2225	2298	2373	2448	2524	2602	2680	2760	2841	2924	3007	44
46	1942	2012	2083	2155	2227	2301	2375	2450	2527	2604	2683	2763	2844	2926	3010	46
48	1944	2014	2085	2157	2230	2303	2378	2453	2529	2607	2686	2766	2847	2929	3013	48
50	1946	2017	2088	2159	2232	2306	2380	2456	2532	2610	2688	2768	2849	2932	3016	50
52	1948	2019	2090	2162	2234	2308	2383	2458	2535	2612	2691	2771	2852	2935	3019	52
54	1951	2021	2092	2164	2237	2311	2385	2461	2537	2615	2694	2774	2855	2937	3022	54
56	1953	2024	2095	2167	2239	2313	2388	2463	2540	2617	2696	2776	2858	2940	3024	56
58	1956	2026	2097	2169	2241	2315	2390	2466	2542	2620	2699	2779	2860	2943	3027	58
60	1958	2028	2100	2171	2244	2318	2393	2468	2545	2622	2702	2782	2863	2946	3030	60

MERIDIONAL PARTS on a CHART, corresponding with the Earth's Convex Surface of a Spheroid. Diameters 266 to 265.

Degrees of LATITUDE.																
	30°	31°	32°	33°	34°	35°	36°	37°	38°	39°	40°	41°	42°	43°	44°	
0	1875	1945	2015	2085	2157	2229	2303	2377	2452	2529	2606	2685	2764	2845	2928	0
2	1878	1947	2017	2088	2159	2232	2305	2380	2455	2531	2609	2687	2767	2848	2931	2
4	1880	1949	2019	2090	2162	2234	2308	2382	2457	2533	2611	2690	2770	2851	2933	4
6	1882	1952	2022	2093	2164	2237	2310	2385	2460	2536	2614	2693	2772	2854	2936	6
8	1885	1954	2024	2095	2167	2239	2313	2387	2462	2539	2616	2695	2775	2856	2939	8
10	1887	1956	2026	2097	2169	2242	2315	2390	2465	2541	2619	2698	2778	2859	2942	10
12	1889	1959	2029	2100	2171	2244	2318	2392	2468	2544	2622	2700	2780	2862	2944	12
14	1892	1961	2031	2102	2174	2246	2320	2395	2470	2547	2624	2703	2783	2865	2947	14
16	1894	1963	2033	2104	2176	2249	2322	2397	2473	2549	2627	2706	2786	2867	2950	16
18	1896	1966	2036	2107	2179	2251	2325	2400	2475	2552	2629	2708	2789	2870	2953	18
20	1898	1968	2038	2109	2181	2254	2327	2402	2478	2554	2632	2711	2791	2873	2956	20
22	1901	1970	2041	2112	2183	2256	2330	2405	2480	2557	2635	2714	2794	2875	2958	22
24	1903	1973	2043	2114	2186	2259	2332	2407	2483	2559	2637	2716	2797	2878	2961	24
26	1905	1975	2045	2116	2188	2261	2335	2410	2486	2562	2640	2719	2799	2881	2964	26
28	1908	1977	2048	2119	2191	2264	2337	2412	2488	2564	2643	2722	2802	2884	2967	28
30	1910	1980	2050	2121	2193	2266	2340	2415	2490	2567	2645	2724	2805	2886	2970	30
32	1912	1982	2052	2123	2196	2268	2342	2417	2493	2570	2648	2727	2807	2889	2972	32
34	1915	1984	2055	2126	2198	2271	2345	2420	2495	2572	2650	2730	2810	2892	2975	34
36	1917	1987	2057	2128	2200	2273	2347	2422	2498	2575	2653	2732	2813	2895	2978	36
38	1919	1989	2059	2131	2203	2276	2350	2425	2501	2578	2656	2735	2816	2897	2981	38
40	1922	1991	2062	2133	2205	2278	2352	2427	2503	2580	2658	2738	2818	2900	2984	40
42	1924	1994	2064	2135	2208	2281	2355	2430	2506	2583	2661	2740	2821	2903	2986	42
44	1926	1996	2066	2138	2210	2283	2357	2432	2508	2585	2664	2743	2824	2906	2989	44
46	1928	1998	2069	2140	2212	2286	2360	2435	2511	2588	2666	2746	2826	2908	2992	46
48	1931	2001	2071	2143	2215	2288	2362	2437	2513	2590	2669	2748	2829	2911	2995	48
50	1933	2003	2074	2145	2217	2290	2365	2440	2516	2593	2671	2751	2832	2914	2998	50
52	1935	2005	2076	2147	2220	2293	2367	2442	2518	2596	2674	2754	2835	2917	3000	52
54	1938	2008	2078	2150	2222	2295	2370	2445	2521	2598	2677	2757	2837	2920	3003	54
56	1940	2010	2081	2152	2225	2298	2372	2447	2524	2601	2679	2759	2840	2922	3006	56
58	1942	2012	2083	2155	2227	2300	2375	2450	2526	2603	2682	2762	2842	2925	3009	58
60	1945	2015	2085	2157	2229	2303	2377	2452	2529	2606	2685	2764	2845	2928	3012	60

According to the Radius of Curvature = 3994 Miles, found for the Middle of a Degree of Latitude, measured in Lapland, Lat. 66° 20', and that of 3958.4 Miles found for the Middle of a Degree measured in France, Lat. 49° 22', the Ratio of the Equator's Semi-Diameter to the Earth's Semi-Axis is determined (by the Rules, p. 295.) as 1313.22 to 309.72, or their Squares, as 981.1 to 959.3; when the Focal Distance from the Center of the Ellipsis = 149 very nearly. Therefore to find the Meridional Parts to this Spheroid, Lat. 60°?

Product } Comp. Half } Log. Cotan. } Const. N°. } Foc. Dist. }
 Sine 60° = .8660254 } = .1290378 } 7° 25' } 82° 35' } 41° 17½' } .056375 by 9715.7 X .149 }
 By Foc. Dist. X d, 149 } Nat. Sine. } } } } } = 66.49 Dif. Merid. Parts. } Merid. Parts reqd 4461

MERIDIONAL

MERIDIONAL PARTS on Mercator's CHART, corresponding with the Earth's Convex Surface of a Sphere.

Degrees of LATITUDE.

	45°	46°	47°	48°	49°	50°	51°	52°	53°	54°	55°	56°	57°	58°	59°	
0	3030	3116	3203	3292	3382	3475	3569	3665	3764	3865	3968	4074	4183	4294	4409	0
2	3033	3118	3206	3295	3385	3478	3572	3669	3768	3868	3972	4078	4187	4298	4413	2
4	3036	3121	3209	3298	3388	3481	3575	3672	3771	3872	3975	4081	4190	4302	4417	4
6	3038	3124	3212	3301	3391	3484	3578	3675	3774	3875	3979	4085	4194	4306	4421	6
8	3041	3127	3214	3304	3394	3487	3582	3678	3777	3878	3982	4088	4197	4310	4425	8
10	3044	3130	3217	3307	3397	3490	3585	3682	3780	3882	3986	4092	4201	4313	4429	10
12	3047	3133	3220	3310	3400	3493	3588	3685	3784	3885	3989	4096	4205	4317	4433	12
14	3050	3136	3223	3313	3403	3496	3591	3688	3787	3889	3992	4100	4209	4321	4437	14
16	3053	3139	3226	3316	3407	3499	3594	3691	3791	3892	3996	4103	4212	4325	4441	16
18	3055	3142	3229	3319	3410	3503	3597	3695	3794	3895	4000	4106	4216	4329	4444	18
20	3058	3144	3232	3322	3413	3506	3600	3698	3797	3899	4003	4110	4220	4332	4448	20
22	3061	3147	3235	3325	3416	3509	3604	3701	3801	3902	4007	4114	4223	4336	4452	22
24	3064	3150	3238	3328	3419	3512	3607	3705	3804	3906	4010	4117	4227	4340	4456	24
26	3067	3153	3241	3331	3422	3515	3610	3708	3807	3909	4014	4121	4231	4344	4460	26
28	3070	3156	3244	3334	3425	3518	3614	3711	3811	3913	4017	4124	4235	4348	4464	28
30	3073	3159	3247	3337	3428	3521	3617	3714	3814	3916	4021	4128	4238	4352	4468	30
32	3075	3162	3250	3340	3431	3525	3620	3718	3817	3920	4024	4132	4242	4355	4472	32
34	3078	3165	3253	3343	3434	3528	3623	3721	3821	3923	4028	4135	4246	4359	4476	34
36	3081	3168	3256	3346	3437	3531	3626	3724	3824	3926	4031	4139	4249	4363	4480	36
38	3084	3171	3259	3349	3440	3534	3630	3728	3827	3930	4035	4143	4253	4367	4484	38
40	3087	3173	3262	3352	3443	3537	3633	3731	3831	3933	4038	4146	4257	4371	4488	40
42	3090	3176	3265	3355	3446	3540	3636	3734	3834	3937	4042	4150	4261	4375	4492	42
44	3093	3179	3268	3358	3449	3543	3639	3737	3838	3940	4046	4154	4264	4378	4496	44
46	3095	3182	3271	3361	3453	3547	3643	3741	3841	3944	4049	4157	4268	4382	4500	46
48	3098	3185	3274	3364	3456	3550	3646	3744	3844	3947	4053	4161	4272	4386	4504	48
50	3101	3188	3277	3367	3459	3553	3649	3747	3848	3951	4056	4164	4276	4390	4508	50
52	3104	3191	3280	3370	3462	3556	3652	3751	3851	3954	4060	4168	4279	4394	4512	52
54	3107	3194	3283	3373	3465	3559	3656	3754	3855	3958	4063	4172	4283	4398	4516	54
56	3110	3197	3286	3376	3468	3562	3659	3757	3858	3961	4067	4175	4287	4402	4520	56
58	3113	3200	3289	3379	3471	3566	3662	3761	3861	3965	4070	4179	4291	4405	4523	58
60	3116	3203	3292	3382	3475	3569	3665	3764	3865	3968	4074	4183	4294	4409	4527	60

MERIDIONAL PARTS on a CHART, corresponding with the Earth's Convex Surface of a Spheroid. Diameters 266 to 265.

Degrees of LATITUDE.

	45°	46°	47°	48°	49°	50°	51°	52°	53°	54°	55°	56°	57°	58°	59°	
0	3012	3097	3184	3272	3363	3455	3549	3645	3743	3844	3947	4052	4161	4272	4387	0
2	3014	3100	3187	3275	3366	3458	3552	3648	3746	3847	3950	4056	4165	4276	4391	2
4	3017	3103	3190	3278	3369	3461	3555	3651	3750	3850	3954	4060	4168	4280	4395	4
6	3020	3106	3193	3281	3372	3464	3558	3655	3753	3854	3957	4063	4172	4284	4399	6
8	3023	3108	3196	3284	3375	3467	3561	3658	3756	3857	3961	4067	4176	4287	4402	8
10	3026	3111	3198	3287	3378	3470	3565	3661	3760	3861	3964	4070	4179	4291	4406	10
12	3029	3114	3201	3290	3381	3473	3568	3664	3763	3864	3968	4074	4183	4295	4410	12
14	3031	3117	3204	3293	3384	3476	3571	3668	3766	3868	3971	4077	4187	4299	4414	14
16	3034	3120	3207	3296	3387	3480	3574	3671	3770	3871	3975	4081	4190	4303	4418	16
18	3037	3123	3210	3299	3390	3483	3577	3674	3773	3874	3978	4085	4194	4306	4422	18
20	3040	3126	3213	3302	3393	3486	3580	3677	3776	3878	3982	4088	4198	4310	4426	20
22	3043	3129	3216	3305	3396	3489	3584	3681	3780	3881	3985	4092	4201	4314	4430	22
24	3046	3131	3219	3308	3399	3492	3587	3684	3783	3885	3989	4095	4205	4318	4434	24
26	3048	3134	3222	3311	3402	3495	3590	3687	3787	3888	3992	4099	4209	4322	4438	26
28	3051	3137	3225	3314	3405	3498	3593	3690	3790	3891	3996	4103	4213	4325	4442	28
30	3054	3140	3228	3317	3408	3501	3596	3694	3793	3895	3999	4106	4216	4329	4445	30
32	3057	3143	3231	3320	3411	3505	3599	3697	3796	3898	4003	4110	4220	4333	4449	32
34	3060	3146	3234	3323	3414	3508	3603	3700	3800	3902	4006	4114	4224	4337	4453	34
36	3063	3149	3237	3326	3418	3511	3606	3703	3803	3905	4010	4117	4227	4341	4457	36
38	3065	3152	3240	3329	3421	3514	3609	3707	3807	3909	4013	4121	4231	4345	4461	38
40	3068	3155	3243	3332	3424	3517	3613	3710	3810	3912	4017	4124	4235	4348	4465	40
42	3071	3158	3246	3335	3427	3520	3616	3713	3813	3916	4020	4128	4239	4352	4469	42
44	3074	3160	3249	3338	3430	3524	3619	3717	3817	3919	4024	4132	4242	4356	4473	44
46	3077	3163	3251	3341	3433	3527	3622	3720	3820	3922	4028	4135	4246	4360	4477	46
48	3080	3166	3254	3344	3436	3530	3625	3722	3823	3926	4031	4139	4250	4364	4481	48
50	3083	3169	3257	3347	3439	3533	3629	3727	3827	3929	4035	4143	4254	4368	4485	50
52	3085	3172	3260	3350	3442	3536	3632	3730	3830	3933	4038	4146	4257	4371	4489	52
54	3088	3175	3263	3353	3445	3539	3635	3733	3834	3936	4042	4150	4261	4375	4493	54
56	3091	3178	3266	3356	3448	3542	3638	3735	3837	3940	4045	4154	4265	4379	4497	56
58	3094	3181	3269	3359	3452	3546	3642	3740	3840	3943	4049	4157	4269	4383	4501	58
60	3097	3184	3272	3363	3455	3549	3645	3743	3844	3947	4052	4161	4272	4387	4505	60

The Focal Distance by Sir Isaac Newton's Theory is .093, or thereabouts, as Mr. Emerson observes at p. 150 of his *Fluxions*—Hence the Meridional Parts by this New Theory, from Observations made at Quito in South America, making the Figure of the Earth nearer to a Sphere than by Sir Isaac Newton's Theory, are greater than those by that Theory—Therefore, until the exact Form and Dimensions of the Earth be determined, Tables of Meridional Parts for the Earth's supposed Form and Dimensions, as a Spheroid of some particular kind, will be a Refinement of no Certainty or Utility in the Practice of Navigation.

MERIDIONAL PARTS on Mercator's CHART, corresponding with the Earth's Convex Surface of a Sphere.

Degrees of LATITUDE.																
	60°	61°	62°	63°	64°	65°	66°	67°	68°	69°	70°	71°	72°	73°	74°	
0	4527	4649	4775	4905	5040	5179	5324	5474	5631	5795	5966	6146	6335	6534	6746	0
2	4531	4654	4780	4910	5044	5184	5329	5480	5637	5801	5971	6152	6341	6541	6753	2
4	4535	4658	4784	4914	5049	5189	5334	5485	5642	5806	5977	6158	6348	6548	6760	4
6	4539	4662	4788	4919	5053	5193	5339	5490	5647	5812	5983	6164	6354	6555	6768	6
8	4543	4666	4792	4923	5058	5198	5344	5495	5653	5818	5989	6170	6361	6562	6775	8
10	4547	4670	4797	4927	5063	5203	5349	5500	5658	5823	5995	6176	6367	6569	6782	10
12	4551	4674	4801	4932	5067	5208	5354	5505	5664	5829	6001	6182	6374	6576	6790	12
14	4555	4678	4805	4936	5072	5212	5359	5511	5669	5834	6007	6189	6380	6583	6797	14
16	4559	4683	4810	4941	5076	5217	5364	5516	5674	5840	6013	6195	6387	6589	6804	16
18	4564	4687	4814	4945	5081	5222	5368	5521	5680	5846	6019	6201	6393	6596	6812	18
20	4568	4691	4818	4950	5086	5227	5373	5526	5685	5851	6025	6207	6400	6603	6819	20
22	4572	4695	4822	4954	5090	5232	5378	5531	5691	5857	6031	6214	6407	6610	6826	22
24	4576	4699	4827	4959	5095	5236	5383	5537	5696	5863	6037	6220	6413	6617	6834	24
26	4580	4704	4831	4963	5100	5241	5388	5542	5701	5868	6042	6226	6420	6624	6841	26
28	4584	4708	4835	4967	5104	5246	5393	5547	5707	5874	6048	6233	6426	6631	6849	28
30	4588	4712	4840	4972	5109	5251	5398	5552	5712	5880	6054	6239	6433	6638	6856	30
32	4592	4716	4844	4976	5114	5256	5403	5557	5718	5886	6060	6245	6440	6646	6864	32
34	4596	4720	4848	4981	5118	5261	5409	5563	5723	5891	6066	6251	6446	6653	6871	34
36	4600	4724	4853	4985	5123	5265	5414	5568	5729	5897	6072	6258	6453	6660	6879	36
38	4604	4729	4857	4990	5127	5270	5419	5573	5734	5903	6078	6264	6460	6667	6886	38
40	4608	4733	4861	4994	5132	5275	5424	5578	5740	5909	6084	6270	6467	6674	6894	40
42	4613	4737	4866	4999	5137	5280	5429	5584	5745	5914	6089	6277	6473	6681	6902	42
44	4617	4741	4870	5003	5142	5285	5434	5589	5751	5919	6097	6283	6480	6688	6909	44
46	4621	4746	4875	5008	5146	5290	5439	5594	5756	5925	6103	6290	6487	6695	6917	46
48	4625	4750	4879	5012	5151	5295	5444	5599	5762	5931	6109	6296	6493	6702	6924	48
50	4629	4754	4883	5017	5156	5299	5449	5605	5767	5936	6115	6302	6500	6710	6932	50
52	4633	4758	4888	5022	5160	5304	5454	5610	5773	5942	6121	6309	6507	6717	6940	52
54	4637	4763	4892	5026	5165	5309	5459	5615	5778	5948	6127	6315	6514	6724	6947	54
56	4641	4767	4896	5031	5170	5314	5464	5621	5784	5954	6133	6322	6521	6731	6955	56
58	4645	4771	4901	5035	5174	5319	5469	5626	5790	5960	6139	6328	6527	6738	6966	58
60	4649	4775	4905	5040	5179	5324	5474	5631	5795	5966	6146	6335	6534	6746	6970	60

MERIDIONAL PARTS on a CHART, corresponding with the Earth's Convex Surface of a Spheroid. Diameters 266 to 265.

Degrees of LATITUDE.																
	60°	61°	62°	63°	64°	65°	66°	67°	68°	69°	70°	71°	72°	73°	74°	
0	4505	4627	4752	4882	5016	5155	5300	5450	5607	5773	5942	6121	6310	6510	6721	0
2	4509	4631	4756	4886	5021	5160	5305	5455	5612	5776	5947	6127	6317	6516	6728	2
4	4513	4635	4761	4891	5025	5165	5310	5460	5617	5782	5953	6134	6323	6523	6735	4
6	4517	4639	4765	4895	5030	5170	5315	5466	5623	5787	5959	6140	6330	6530	6743	6
8	4521	4643	4769	4899	5034	5174	5320	5471	5628	5793	5965	6146	6336	6537	6750	8
10	4525	4647	4774	4904	5039	5179	5324	5476	5634	5798	5971	6152	6343	6544	6757	10
12	4529	4651	4778	4908	5044	5184	5329	5481	5639	5804	5977	6158	6349	6551	6765	12
14	4533	4656	4782	4913	5048	5189	5334	5486	5644	5810	5983	6164	6356	6558	6772	14
16	4537	4660	4786	4917	5053	5193	5339	5491	5650	5815	5989	6171	6362	6565	6779	16
18	4541	4664	4791	4922	5057	5198	5344	5496	5655	5821	5995	6177	6369	6572	6787	18
20	4545	4668	4795	4926	5062	5203	5349	5502	5661	5827	6000	6183	6375	6579	6794	20
22	4549	4672	4799	4931	5067	5208	5354	5507	5666	5832	6006	6189	6382	6586	6802	22
24	4553	4676	4803	4935	5071	5212	5359	5512	5671	5838	6012	6196	6389	6593	6809	24
26	4557	4680	4808	4939	5076	5217	5364	5517	5677	5844	6018	6202	6395	6600	6816	26
28	4561	4685	4812	4944	5080	5222	5369	5522	5682	5849	6024	6208	6402	6607	6824	28
30	4565	4689	4816	4948	5085	5227	5374	5528	5688	5855	6030	6214	6409	6614	6831	30
32	4569	4693	4821	4953	5090	5232	5379	5533	5693	5861	6036	6221	6415	6621	6839	32
34	4573	4697	4825	4957	5094	5237	5384	5538	5699	5866	6042	6227	6422	6628	6846	34
36	4577	4701	4829	4962	5099	5241	5389	5543	5704	5872	6048	6233	6428	6635	6854	36
38	4582	4706	4834	4966	5104	5246	5394	5549	5710	5878	6054	6240	6436	6642	6861	38
40	4586	4710	4838	4971	5108	5251	5399	5554	5715	5884	6060	6246	6442	6649	6869	40
42	4590	4714	4842	4975	5113	5256	5404	5559	5721	5890	6066	6252	6449	6656	6876	42
44	4594	4718	4847	4980	5118	5261	5409	5564	5726	5895	6072	6259	6455	6663	6884	44
46	4598	4722	4851	4984	5122	5266	5415	5570	5732	5901	6079	6265	6462	6670	6892	46
48	4602	4727	4856	4989	5127	5270	5420	5575	5737	5907	6085	6272	6469	6678	6899	48
50	4606	4731	4860	4993	5132	5275	5425	5580	5743	5912	6091	6278	6476	6685	6907	50
52	4610	4735	4864	4998	5136	5280	5430	5586	5748	5918	6097	6284	6482	6692	6915	52
54	4614	4739	4869	5004	5141	5285	5435	5591	5754	5924	6103	6291	6489	6699	6922	54
56	4618	4744	4873	5007	5146	5290	5440	5596	5759	5930	6109	6297	6496	6706	6930	56
58	4622	4748	4877	5012	5151	5295	5445	5601	5765	5936	6115	6304	6503	6714	6938	58
60	4627	4752	4882	5016	5155	5300	5450	5607	5770	5942	6121	6310	6510	6721	6945	60

If you put l and g = the Logarithm Tangents of the Half Complements of the less and greater Latitude of two Places, (without their Indices) c the natural Tangent of the $\angle$ of the Ship's Course betwixt the two Latitudes, t the nat. Tangent of $51^\circ 38' 9''$, (whose Logarithm Tangent = 10.1015104) and d the Diff. of Long. in Miles. Then $\overline{l-g} \times c = t \times d$. Hence 1. $c = \frac{t \times d}{\overline{l-g}}$; 2. $d = \frac{\overline{l-g} \times c}{t}$; 3. $l = \frac{t \times d}{c} + g$.

$\overline{l-g} = l - \frac{t \times d}{c}$. The Half Complements of Lat. must be reckoned from the same Pole. The 3 last Figures of Log. Tangents of 7, are Decimals.

By these four Rules all Questions in Mercator Sailing, where Longitude is concerned, may be readily answered with Logarithms.

MERIDIONAL PARTS on Mercator's CHART, corresponding with the Earth's Convex Surface of a Sphere.

Degrees of LATITUDE.

	75°	76°	77°	78°	79°	80°	81°	82°	83°	84°	85°	86°	87°	88°	89°	
0	6970	7210	7467	7745	8046	8375	8739	9146	9606	10137	10765	11533	12522	13916	16300	0
2	6978	7218	7476	7755	8057	8386	8752	9161	9622	10156	10788	11562	12560	13974	16416	2
4	6986	7227	7484	7764	8068	8398	8765	9175	9639	10175	10811	11591	12589	14033	16537	4
6	6994	7235	7493	7774	8078	8410	8778	9190	9656	10194	10834	11620	12639	14093	16661	6
8	7002	7244	7502	7784	8089	8422	8791	9204	9672	10214	10858	11650	12679	14154	16792	8
10	7010	7252	7511	7794	8099	8433	8804	9219	9689	10234	10881	11680	12719	14216	16926	10
12	7018	7261	7520	7803	8110	8445	8817	9234	9706	10254	10905	11710	12760	14279	17067	12
14	7026	7269	7529	7813	8121	8457	8830	9248	9723	10274	10929	11740	12801	14343	17213	14
16	7034	7278	7538	7823	8131	8469	8843	9263	9740	10294	10953	11770	12842	14408	17366	16
18	7042	7286	7548	7833	8142	8481	8857	9278	9757	10314	10978	11801	12884	14475	17526	18
20	7050	7295	7557	7843	8153	8492	8870	9293	9774	10334	11002	11832	12927	14543	17693	20
22	7058	7304	7566	7853	8164	8504	8883	9308	9792	10354	11027	11863	12970	14612	17870	22
24	7066	7312	7575	7863	8175	8516	8896	9323	9809	10374	11052	11895	13014	14682	18056	24
26	7074	7321	7584	7873	8186	8528	8910	9338	9827	10395	11077	11927	13059	14756	18252	26
28	7082	7329	7594	7883	8196	8540	8923	9353	9844	10416	11102	11960	13104	14830	18461	28
30	7090	7338	7604	7893	8206	8553	8937	9368	9862	10437	11127	11992	13150	14906	18682	30
32	7098	7347	7613	7903	8217	8565	8950	9383	9879	10457	11152	12025	13196	14983	18920	32
34	7106	7355	7622	7913	8229	8577	8964	9399	9897	10478	11178	12059	13243	15062	19174	34
36	7114	7364	7631	7923	8240	8589	8978	9414	9915	10499	11204	12092	13290	15143	19450	36
38	7122	7372	7641	7933	8251	8601	8991	9430	9933	10521	11230	12126	13338	15226	19749	38
40	7130	7381	7650	7943	8262	8614	9005	9446	9951	10542	11257	12160	13386	15311	20076	40
42	7138	7390	7660	7953	8273	8626	9019	9461	9970	10564	11283	12194	13435	15398	20439	42
44	7146	7398	7669	7964	8284	8639	9033	9477	9989	10585	11310	12229	13485	15487	20843	44
46	7154	7407	7678	7974	8295	8651	9047	9493	10006	10607	11337	12263	13536	15579	21303	46
48	7162	7415	7688	7984	8307	8663	9061	9509	10025	10629	11365	12300	13588	15673	21832	48
50	7170	7424	7697	7994	8318	8676	9075	9525	10043	10652	11392	12336	13641	15770	22459	50
52	7178	7433	7707	8005	8329	8689	9089	9541	10061	10674	11420	12372	13694	15870	23226	52
54	7186	7441	7716	8015	8341	8701	9103	9557	10080	10696	11448	12408	13748	15973	24215	54
56	7194	7450	7720	8026	8352	8714	9118	9573	10099	10719	11476	12445	13803	16079	25609	56
58	7202	7458	7736	8036	8364	8727	9132	9590	10118	10742	11504	12483	13859	16188	27992	58
60	7210	7467	7745	8046	8375	8739	9146	9606	10137	10765	11533	12522	13916	16300	Infinite	60

MERIDIONAL PARTS on a CHART, corresponding with the Earth's Convex Surface of a Spheroid. Diameters 266 to 265.

Degrees of LATITUDE.

	75°	76°	77°	78°	79°	80°	81°	82°	83°	84°	85°	86°	87°	88°	89°	
0	6945	7185	7442	7719	8020	8350	8713	9120	9580	10111	10739	11507	12496	13891	16274	0
2	6953	7193	7451	7729	8031	8361	8726	9134	9597	10130	10762	11535	12535	13948	16390	2
4	6961	7202	7460	7739	8041	8373	8739	9149	9613	10150	10785	11564	12574	14007	16511	4
6	6969	7210	7469	7748	8053	8384	8752	9163	9630	10169	10808	11594	12613	14067	16636	6
8	6976	7218	7478	7758	8063	8396	8765	9178	9646	10188	10832	11623	12653	14128	16766	8
10	6984	7226	7487	7768	8073	8408	8778	9192	9663	10208	10855	11653	12693	14190	16900	10
12	6992	7235	7496	7777	8084	8419	8791	9207	9680	10228	10879	11683	12733	14253	17041	12
14	7000	7243	7505	7787	8094	8431	8804	9222	9697	10248	10903	11713	12775	14317	17187	14
16	7008	7252	7514	7797	8105	8443	8817	9237	9714	10268	10927	11744	12816	14383	17340	16
18	7016	7260	7523	7807	8116	8455	8830	9252	9731	10288	10952	11775	12859	14449	17500	18
20	7023	7269	7532	7817	8127	8467	8844	9267	9748	10308	10976	11806	12901	14511	17668	20
22	7031	7277	7541	7827	8137	8479	8857	9282	9765	10328	11001	11837	12945	14587	17844	22
24	7039	7286	7550	7837	8148	8491	8870	9297	9783	10349	11026	11869	12988	14658	18030	24
26	7047	7294	7559	7847	8159	8503	8884	9312	9800	10369	11051	11901	13033	14730	18226	26
28	7055	7303	7569	7856	8170	8515	8897	9327	9818	10390	11076	11933	13078	14804	18435	28
30	7063	7311	7578	7867	8181	8527	8911	9342	9835	10411	11101	11967	13123	14880	18657	30
32	7071	7320	7587	7877	8192	8539	8924	9358	9853	10432	11127	11999	13169	14957	18894	32
34	7079	7328	7596	7887	8203	8551	8938	9373	9871	10453	11153	12032	13216	15036	19149	34
36	7087	7337	7606	7897	8214	8563	8951	9389	9889	10474	11179	12066	13264	15117	19424	36
38	7095	7346	7615	7907	8225	8576	8965	9405	9907	10495	11205	12100	13312	15200	19723	38
40	7103	7354	7624	7917	8236	8588	8979	9420	9924	10517	11231	12134	13360	15285	20050	40
42	7111	7363	7634	7927	8248	8600	8993	9435	9943	10538	11258	12168	13410	15372	20413	42
44	7119	7372	7643	7937	8259	8613	9007	9451	9961	10560	11285	12203	13460	15461	20818	44
46	7128	7380	7653	7948	8270	8625	9021	9467	9980	10582	11312	12239	13511	15553	21277	46
48	7136	7389	7662	7958	8281	8638	9035	9483	9998	10604	11339	12274	13563	15647	21807	48
50	7144	7398	7671	7968	8293	8650	9049	9499	10017	10626	11366	12310	13615	15744	22453	50
52	7152	7407	7681	7979	8304	8663	9063	9515	10035	10648	11394	12347	13669	15843	23200	52
54	7160	7415	7691	7989	8315	8675	9077	9531	10054	10671	11422	12383	13723	15940	24189	54
56	7168	7424	7700	7999	8327	8688	9091	9547	10073	10693	11450	12421	13778	16052	25583	56
58	7177	7433	7710	8010	8338	8701	9105	9564	10092	10716	11478	12458	13834	16161	27966	58
60	7185	7442	7719	8020	8350	8713	9120	9580	10111	10739	11507	12496	13891	16274	...	60

LET f = Natural Secant of the Course, D = Miles Dist. run, L = Minutes Diff. Lat. Rad. = r ; Then $f \times L = r \times D$: Whence,
 1. $f = \frac{r \times D}{L}$. 2. $L = \frac{r \times D}{f}$. 3. $D = \frac{f \times L}{r}$; by which RULES, and the former, with Logarithms, all Questions in practical Navigation are readily answered.

ANOTHER RULE to find the Meridional Parts for a CHART corresponding to the Convexity of a SPHEROID. Multiply the Product of the Sine of the given Lat. (to Radius 1) and the constant Number 3437.87, by the Quotient, of the Square of the focal Distance from the Center of the Ellipsis in Parts of 1, the Semi-conjugate or Earth's Semi-Axis, divided by that Square more 1, and the last Product will be the meridional Parts less than those of the Sphere for the Spheroid of the same Latitude.

The

The LENGTH of any DEGREE, and also any Number of DEGREES, being elliptic Arcs of the EARTH'S MERIDIAN, in Minutes of the EQUATOR and Hundredth Parts. For any Latitude. According to Don Juan.

Lat.	Length of a Degree.	Arches of the Meridian.	Lat.	Length of a Degree.	Arches of the Meridian.
°	' H.	' H.	°	' H.	' Hun.
1	59.55	59.55	46	59.89	2745.17
2	59.55	119.10	47	59.90	2805.07
3	59.55	178.65	48	59.92	2864.99
4	59.56	238.21	49	59.93	2924.92
5	59.55	297.76	50	59.94	2984.86
6	59.55	357.31	51	59.95	3044.81
7	59.56	416.87	52	59.96	3104.77
8	59.57	476.44	53	59.97	3164.74
9	59.56	536.00	54	59.99	3224.73
10	59.57	595.57	55	60.00	3284.73
11	59.57	655.14	56	60.01	3344.74
12	59.58	714.72	57	60.02	3404.76
13	59.58	774.30	58	60.03	3464.79
14	59.59	833.89	59	60.04	3524.83
15	59.59	893.48	60	60.05	3584.88
16	59.60	953.08	61	60.06	3644.94
17	59.61	1012.69	62	60.07	3705.01
18	59.61	1072.30	63	60.08	3765.09
19	59.61	1131.91	64	60.09	3825.18
20	59.63	1191.54	65	60.10	3885.28
21	59.63	1251.17	66	60.11	3945.39
22	59.64	1310.81	67	60.11	4005.50
23	59.65	1370.46	68	60.13	4065.63
24	59.66	1430.12	69	60.13	4125.76
25	59.67	1489.79	70	60.14	4185.90
26	59.67	1549.46	71	60.15	4246.05
27	59.69	1609.15	72	60.16	4306.21
28	59.69	1668.84	73	60.16	4366.37
29	59.70	1728.54	74	60.17	4426.54
30	59.72	1788.26	75	60.18	4486.72
31	59.72	1847.98	76	60.18	4546.90
32	59.74	1907.72	77	60.19	4607.09
33	59.74	1967.46	78	60.19	4667.28
34	59.76	2027.22	79	60.20	4727.48
35	59.77	2086.99	80	60.20	4787.68
36	59.77	2146.76	81	60.21	4847.89
37	59.79	2206.55	82	60.21	4908.10
38	59.80	2266.35	83	60.21	4968.31
39	59.81	2326.16	84	60.22	5028.53
40	59.82	2385.98	85	60.22	5088.75
41	59.84	2445.82	86	60.22	5148.97
42	59.85	2505.67	87	60.22	5209.19
43	59.86	2565.53	88	60.22	5269.41
44	59.87	2625.40	89	60.22	5329.63
45	59.88	2685.28	90	60.22	5389.85

In the above TABLE the Length of any Arc, from the EQUATOR, is equal to Length of all the Degrees of Latitude as far as the given Latitude.

LENGTH of the Meridian Arcs of the SPHERE, by the Reverend Mr. Murdoch, confronted with the Arches of the SPHEROID, on the Left, by the Dons JUAN and ULLOA.

Lat.	Arches of the Sphere.	Arches Spheroid. Mr. Murdoch.	Diff. Sphre and Sphd.	Diff. of Sphd. Arches.	Lat.	Arches of the Sphere.	Arches Spheroid. Mr. Murdoch.	Diff. Sphre and Sphd.	Diff. of Sphd. Arches.
°	'	'	'	'	°	'	'	'	'
1	60	58.7	1.3	0.8	46	2760	2716.4	43.6	28.7
2	120	117.3	2.7	1.8	47	2820	2776.2	43.8	28.8
3	180	176.0	4.0	2.6	48	2880	2835.9	44.1	29.0
4	240	234.7	5.3	3.5	49	2940	2895.5	44.5	29.4
5	300	293.4	6.6	4.3	50	3000	2955.3	44.7	29.5
6	360	352.1	7.9	5.2	51	3060	3015.2	44.8	29.6
7	420	410.8	9.2	6.0	52	3120	3075.0	44.0	29.7
8	480	469.6	10.4	6.8	53	3180	3135.0	45.0	29.7
9	540	528.3	11.7	7.7	54	3240	3194.9	45.1	29.8
10	600	587.0	13.0	8.5	55	3300	3254.9	45.1	29.8
11	660	645.8	14.2	9.3	56	3360	3314.9	45.1	29.8
12	720	704.5	15.5	10.2	57	3420	3375.0	45.0	29.7
13	780	763.3	16.7	11.0	58	3480	3435.1	44.9	29.6
14	840	822.1	17.9	11.7	59	3540	3495.2	44.8	29.6
15	900	880.9	19.1	12.5	60	3600	3555.3	44.7	29.5
16	960	939.7	20.3	13.3	61	3660	3615.5	44.5	29.4
17	1020	998.5	21.5	14.1	62	3720	3675.7	44.3	29.3
18	1080	1057.4	22.6	14.9	63	3780	3736.0	44.0	29.1
19	1140	1116.3	23.7	15.6	64	3840	3796.2	43.8	28.9
20	1200	1175.2	24.8	16.3	65	3900	3856.5	43.5	28.7
21	1260	1234.1	25.9	17.0	66	3960	3916.8	43.2	28.5
22	1320	1293.0	27.0	17.8	67	4020	3977.2	42.8	28.3
23	1380	1352.0	28.0	18.4	68	4080	4037.5	42.5	28.1
24	1440	1411.0	29.0	19.1	69	4140	4097.9	42.1	27.8
25	1500	1470.0	30.0	19.7	70	4200	4158.4	41.6	27.5
26	1560	1529.0	31.0	20.4	71	4260	4218.8	41.2	27.2
27	1620	1588.1	31.9	21.0	72	4320	4279.3	40.7	26.9
28	1680	1647.2	32.8	21.6	73	4380	4339.8	40.2	26.5
29	1740	1706.3	33.7	22.2	74	4440	4400.3	39.7	26.2
30	1800	1765.5	34.5	22.7	75	4500	4460.8	39.2	25.9
31	1860	1824.7	35.3	23.2	76	4560	4521.3	38.7	25.6
32	1920	1883.9	36.1	23.8	77	4620	4581.9	38.1	25.1
33	1980	1943.1	36.9	24.3	78	4680	4642.5	37.5	24.7
34	2040	2002.4	37.6	24.8	79	4740	4703.1	36.9	24.3
35	2100	2061.7	38.3	25.2	80	4800	4763.7	36.3	23.9
36	2160	2121.0	39.0	25.7	81	4860	4824.3	35.7	23.5
37	2220	2180.4	39.6	26.1	82	4920	4884.9	35.1	23.2
38	2280	2239.8	40.2	26.5	83	4980	4945.5	34.5	22.8
39	2340	2299.2	40.8	26.9	84	5040	5006.2	33.8	22.3
40	2400	2358.7	41.3	27.2	85	5100	5066.8	33.2	21.9
41	2460	2418.2	41.8	27.6	86	5160	5127.5	32.5	21.4
42	2520	2477.7	42.3	27.9	87	5220	5188.2	31.8	21.0
43	2580	2537.3	42.7	28.2	88	5280	5148.8	31.2	20.6
44	2640	2596.8	43.2	28.6	89	5340	5209.5	30.5	20.1
45	2700	2656.6	43.4	28.6	90	5400	5270.2	29.8	19.6

By the ABOVE COMPARISON of Mr. Murdoch's Arcs of the Spheroid with those of the Sphere, and also with the Arcs of the Spheroid by Don Juan, you may see that Juan's Arcs are about two thirds as much greater than Mr. Murdoch's as his are less than those of the Sphere; and consequently are nearer to those of the Sphere by about 2 Parts of the 3 than the Arcs of Mr. Murdoch, by whose meridional Parts Mr. Martin's New Sea-Chart is constructed.

IN the above Table we have made Comparison of the Difference of Arcs, to give the better Idea of their Defects: as of the Defects of the meridional Parts by comparing them together.

The best Way of setting Falshood and Error in a proper Light being to compare and confront them with Truth, when it can be had.

Of the DIMENSIONS of the EARTH as a SPHEROID.

RULES for determining the DIMENSIONS of the EARTH as a Spheroid, or a flattened SPHERE, next the P O L E S.

PUT z = the Semidiameter of the Earth's Equator } The Semi-transverse.
 u = the Semidiameter of the Earth's Axis } The Semi-conjugate.
 x = the Part of the Semi-diameter of the Earth's Equator from the Center thereof, to where a Perpendicular falls on that Semidiameter from the Place of Latitude on the Earth's elliptic Meridian.
 y = that Perpendicular, or Semi-ordinate, let fall from the Place of Latitude, on the said Radius of the Equator, or Semi-Transverse.
 r = the Radius of Curvature, or of a Circle touching the elliptic Meridian of the Earth in the Place of Latitude.

i, t, s , Radius, Tangent, and Secant of the Angle of Latitude.

$\frac{u^2}{z^2} = a$. WHENCE, by the Properties of the Ellipsis and of Triangles, we have the following Rules, for determining the Requisites of the Earth's Dimensions.

RULE I. $\frac{u^2}{z} = az$ = the Latus rectum or Parameter of the elliptic Meridian.

$$\text{II. } x = \frac{y}{at} = \frac{z}{\sqrt{1+att}}; \text{ whence } y = atx = \frac{atz}{\sqrt{1+att}}.$$

$$\text{III. } r = \frac{s^3 y}{t+at^3} = \frac{s^3 ax}{1+att} = \frac{s^3 az}{1+att|^{3/2}}.$$

$$\text{IV. } z = \frac{r \times \sqrt{1+att|^{3/2}}}{s^3 a}, \text{ and } z^2 = \frac{r^2 \times 1+att^3}{s^6 a^2} = \frac{r^2 \times 1+att^3}{s^6 a^2}, \text{ for ANOTHER LATITUDE. Whence,}$$

$$\text{V. } a = \frac{r^2 \sigma^2 - \rho^2 s^2}{\rho^3 s^2 \tau \tau - r^3 \sigma^2 t t} = \frac{u^2}{z^2}.$$

CONSTRUCTION of Meridional PARTS continued.

THUS, Sine Lat. $60^\circ = .8660254$ } $\times d$ by Conf. N^o. 3437,87 } 2997,282742 } $\left. \begin{array}{l} \text{Produit.} \\ \text{fo. Dif.} \end{array} \right\} \text{ by ,00750466 } \left. \begin{array}{l} \text{fo. Dif.} \\ \text{fo. Dif. + 1} \end{array} \right\} \left. \begin{array}{l} \text{Last Produit.} \\ \text{fo. Dif.} \end{array} \right\} \left. \begin{array}{l} 4527 \text{ Mer. Parts Sphere.} \\ -22 \\ \hline 4505 \text{ Mer. Parts Sphd.} \end{array} \right\} \text{Lat. } 60^\circ.$

RULE for nearly determining the same. Multiply a few Figures of the natural Sine of the Latitude (to Radius 1) by 3 and the Product of that 3, and the first Figure, with what is carried, will be the Difference that the meridional Parts of the Spheroid are less than those of the Sphere, for the same Latitude, when the Difference of the Diameters of the Spheroid is small: otherwise this Rule is of no Use.

Example. Sine Lat. $60^\circ = .866$, &c. by 3 (as directed) = 25.98 Diff. Merid. Parts less, nearly as above: Agreeing with Don Juan's Tables for Semi-Diameters of the Spheroid as 266 to 265, compared with the meridional Parts of the Sphere, foregoing. See P. 292.

By M. Maupertuis the Ratio of the Earth's Diameters is as 313,22 to 309,72; from whence the two correct general Rules (before given) find the Difference of meridional Parts 66,49 for the same Latitude, viz. 60° .

By the other Ratio of the Diameters 266 to 265, we have only 22 for Difference.

Don John George Juan and Don Antoine de Ulloa, measured a Degree of the Meridian near the Equator in South America, in the Kingdom of Quito, as their Book sets forth, (first published in Spanish, and then translated into French) who first undertook the Work in the Year 1736, and afterwards found 56767,788 Toises in a Degree: which is greater than the spheroidal Theory gives it, by about 268 Toises, from a Computation depending on the Measure of a Deg. of the Merid. in Lapland, and another in France. From which last Mensuration in America, Don Juan determined the Ratio of the spheroidal Diameters of the Earth to be (as before observed) 266 to 265; a Ratio much nearer than that determined by Sir Isaac Newton (from the Theory of Gravity and Earth's Density) than was before determined from the Measure of a Degree in Lapland and another in France.

The meridional Parts for Latitude 89° to the Spheroid by Don Juan = 16273,7. Nat. Sine 89° is .9998 and $\times d$ by 3 (as directed) = 29,99, &c. = 30 the greatest Diff. the meridional Parts of the Spheroid are less than those for the Sphere. Whence from 16300 meridional Parts for Sphere, deduct 30, and there remains the meridional Parts for the same Latitude, viz. 16270.

Mr. Murdoch's meridional Parts (by which Mr. Martin's New Chart is constructed) and Don Juan's for the Spheroid, differ from the meridional Parts of the Sphere, for the same Latitude of 60° , by 66 and 22; and these from one another by 44, therefore the new Chart and Don Juan's Tables (published for equal Advantage to Navigation) differ twice as much from one another as Juan's Tables differ from the Sphere. That if Juan's Tables are near the Truth for the Spheroid, (making the Figure of the Earth almost as near to a Sphere as Sir Isaac Newton made it) then Martin's new Chart (from Murdoch's or Maupertuis's Principles) is doubly erroneous, (in the Instance before us, of the Proportion of merid. Parts for Lat. 60°) differing twice as much under Juan, as Juan differs under the Sphere, or Chart of Mercator.

OBSERVATION

From

Of the DIMENSIONS of the EARTH as a SPHEROID.

From above $z^{\frac{2}{3}} = \frac{r^{\frac{2}{3}} + r^{\frac{2}{3}} att}{s^2 a^{\frac{2}{3}}} = \frac{r^{\frac{2}{3}} + r^{\frac{2}{3}} att}{a^{\frac{2}{3}} + a^{\frac{2}{3}} tt}$, by Substitution of $s^2 + 1$ for s^2 ; because $s^2 - t^2 = 1$.

Whence,

$$\text{VI. } tt = \frac{r^{\frac{2}{3}} - a^{\frac{2}{3}} z^{\frac{2}{3}}}{a^{\frac{2}{3}} z^{\frac{2}{3}} - ar^{\frac{2}{3}}} = \frac{1 - a^{\frac{2}{3}}}{a^{\frac{2}{3}} - a}, \text{ when } r = z.$$

N. B. At the Equator (when $s = 1$, and $t = 0$) $r = az$.

At the Pole (when s and t are infinite and equal, and a finite being added to an infinite Quantity makes no Difference)

$$r = \left(\frac{s^2 az}{1 + att^{\frac{2}{3}}} = \frac{t^2 az}{att^{\frac{2}{3}}} = \frac{az}{a^{\frac{2}{3}}} \right) = \frac{z}{a^{\frac{1}{3}}}.$$

* * The above RULES (of genuine EXTRACTION) are correct and useful when the Radius of Curvature r can be first very nearly, or exactly, found. But Mr. MARTIN has applied them to no Purpose, Truth, or Advantage, by finding an incorrect Radius of Curvature as follows.

PROPOSITION I. To find the Radius of Curvature of the Earth in Miles, correspondent to the Middle of a Degree measured on the elliptic Meridian of the Earth's Surface.

Example. In Latitude $66^\circ 20'$, a Degree of the Earth's Meridian has been measured and found to be 57437.9 French Toises, whence the Radius of Curvature r , will be first found in Degrees, for the Middle of that Degree, by this PROPORTION, according to Mr. Martin. As 3,141593, &c. : 1 :: 360 : $2r$:: 180 : $r =$

	Logarithms.
180°	2.2552725
3,141593 : :	0.4971499
$r = 57^\circ, 29578$	4.7581226
Toises in 1° are 57437.9	4.7591985
Log. of Toises in r	6.5173211
Toises in an English Mile 823,97	2.9159114
English Miles in $r = 3994,115$	3.6014097

* * In the same Manner, from the Length of a Degree of the Earth's Meridian, near Paris, being measured, and found to be 56925.7 Toises, the Length of the Radius of Curvature of the Earth will be found 3958.4 English Miles, for the Middle of that Degree. A Degree by Don Juan = 56767.79 Toises, at the Equator gives Radius of Curvature 3947.42 English Miles. But this Curvature is according to the supposed spherical and not spheroidal Form of the Earth, and therefore incorrect.

OBSERVATION on the latter Part of the INTRODUCTION to the New Principles of Geography and Navigation, by B. Martin; pointing at Mr. EMERSON, and misrepresenting his Judgment and the TRUTH.

SAILING by the Spheroid instead of by the Sphere, (till the larger Errors of the Course and Distance can be truly corrected) is similar to correcting the small Equations of the Moon's Orbit, while the larger Ones, or even those of the mean Motions, are left largely defective.

To what Purpose then are the different Principles of Murdoch and Juan introduced to the Practice of Navigation, and called New Principles of Geography and Navigation, when neither (of those widely different Principles) are certain?

And for the Author of these Principles to suppose Mr. EMERSON depreciated what he did not, (who gave his Reason for what he said, and was unanswered) is the same as if Mr. Emerson were to suppose his Depreciator to be what he is not, and so press and degrade him with the Consequences of the erroneous Supposition.

RE-REVIEWER.

To the AUTHOR of the ROYAL ASTRONOMER and NAVIGATOR.

REMARKS on a BOOK intitled New Principles of Geography and Navigation, pretending to more Exactness than TRUTH and PRACTICABILITY.

AS to these New Principles of Navigation they are taken all from Murdoch. When I read them I found the Reasoning not Sterling. The Table of meridional Parts, the very Examples and the Remarks are all Murdoch's, making up the first Part of this new Navigation.

Murdoch's Book is a Quarto Pamphlet of 38 Pages, besides an Introduction of 32 Pages, printed in 1741, intitled Mercator's Sailing applied to the true Figure of the Earth; a Book exceeding the New Principles of Navigation; but the wiser Author (I have heard) has proposed to retract his Errors.

The principal Things of Note in the new Principles of Navigation are these.

In Page 12. the Author is pleased to tell us, from Maupertuis and Picard, that the Earth's Semi-axis to the Radius of the Equator is as 309,72 to 313,22; from whence it follows, that the Excess of the Height at Equator is $\frac{1}{50}$ of the Radius. Sir Isaac Newton made it $\frac{1}{230}$ only, by Computation.

Page 19. The Author of the New Principles calls the Distance of the Place (a Ship arrives at) from the Meridian of the Place sailed from, by the Name of Departure; though Departure and meridional Distance are of a different Nature: Which shews that this Author knows Nothing of what Departure is; and therefore a very fit Person to write a Book of Navigation!

Page

PROPOSITION

Of the DIMENSIONS of the EARTH as a SPHEROID.

PROPOSITION II. To determine the Ratio of the Semidiameter of the EARTH's EQUATOR, to the Semidiameter of the EARTH's AXIS, from the Radius of Curvature, and Latitude, for two Places, on the Earth's Surface being given?

To perform which, the Value of a must be first determined by RULE V. foregoing.

		Com. Logarithms.	By Lapland and France.	
Latitude.	$r=3994,115$ Miles	3.6014097	$a = \frac{r^{\frac{2}{3}} \sigma^2 - \rho^{\frac{2}{3}} s^2}{\rho^{\frac{2}{3}} r r s s - r^{\frac{2}{3}} t t \sigma^2}$	$r^{\frac{2}{3}} \sigma^2 = 593.604$
Lapland $66^\circ 20'$	$t=2,281,6693$	0.3582527	$= \frac{959.352}{981.345} = \frac{u^2}{x^2}$	$\rho^{\frac{2}{3}} s^2 = 1552.956$
	$s=2,491,1874$	0.3964064		$\rho^{\frac{2}{3}} r r s s = 2108.967$
				$r^{\frac{2}{3}} t t \sigma^2 = 3090.312$
			By Lapland and Equator.	
France $49^\circ 22'$	$\rho = 3958,4$ Miles.	3.5975195	$a = \frac{1298.347}{1310.543} = \frac{u^2}{x^2}$	$r^{\frac{2}{3}} \sigma^2 = 251.737$
	$r = 1,165,3472$	0.0664554		$\rho^{\frac{2}{3}} s^2 = 1550.084$
	$\sigma = 1,535,5892$	0.1862750		$\rho^{\frac{2}{3}} r r s s = 0.$
			By France and Equator.	$r^{\frac{2}{3}} t t \sigma^2 = 1310.543$
Equator 0°	$\rho = 3947,42$ Miles.	3.5963132	$a = \frac{338.7345}{339.8269} = \frac{u^2}{x^2}$	$r^{\frac{2}{3}} \sigma^2 = 250.234$
	$r = 0.$	0.0000000		$\rho^{\frac{2}{3}} s^2 = 588.9685$
	$\sigma = 1.$			$\rho^{\frac{2}{3}} r r s s = 0.$
				$r^{\frac{2}{3}} t t \sigma^2 = 339.8269$
				Here putting Lapland Letters for those of France.

Page 23. This Author of the *same Principles*, projects the Arch of the Meridian upon the *Tangent*, by an Eye at the Earth's Center, perhaps with Design to find the *meridional Parts*. If this was his Meaning, then the meridional Line for any Latitude is no more than the *Tangent* of that Latitude: because any Arch is projected into its *Tangent* by an Eye at the Center.

This Author likewise represents sailing by *middle Latitude* as not deserving his Notice or Regard; though every Seaman knows, in short Distances, as far as any Ship can run in 24 Hours, that the Difference by this Method from that of *Mercator*, in keeping a Ship's Reckoning of Longitude and Latitude, is not sensible; which Method of *Middle-Latitude-Sailing* is therefore a very useful *Approximation*.

Page 32. He speaks of a *Compass*, which is to be more than 100 Times stronger than those of the common Sort, by which, I suppose, that he intends to sail 100 Times more correctly than by the common Methods in Practice.

Page 35. He does not reckon *Departure* any Element of Navigation, or as belonging to the *Art of Sailing*, because it cannot be known from Observation; nor is it thought (by this Author) of any real Use, being determined by *Computation*; but leads (as he thinks) to Error. But does not every Body of Experience in Navigation know, that all the *Easings* and *Westings* are so many *Departures*; and that the Sum of them lying one Way, or Difference the other, is the total *Departure*, made in 24 Hours-Run; and extremely near the Truth.

In Page 36. He makes use of *Departure* in Calculations, though he had said before it was *useless*; who has here forgot himself. He resolves all his Cases *mechanically*, by Scale and Compass, and instead of a Scale uses *Paper-Schemes*, from which no Certainty can be expected; which, however, I submit to the Trial of others.

Page 43. He says the Way of Working on his Chart is the same as on any other; he only proposes to be more exact.—Why then (in the Name of Wisdom) does he call them *New Principles*?

He gives us a Table of the Magnitude of the Earth's Parallels, as *Murdoch* has done; nor seems to suppose them any Way different from those of the Sphere, for any Thing he has shewn to the contrary in his Book of *New Principles*. If so, he might as well suppose all the Degrees of the Meridian to be equal.

There is no such Thing in this whole Book as any Correction of Latitude or Longitude; as if they were but Trifles not worthy his Regard: while every experienced Navigator is sensible, that there is no Sailing without such Correction. But the Use of his new Compass and Dromometer may make that Correction needless, as a Means for the Defect. *amends is made*

In the second Part of this new Performance, the Author has given another Table of Meridional Parts from Don Juan's Book, quite different from the former he took from *Murdoch*.

Don Juan makes the Radius of the Equator to the Earth's Axis, as 266 to 265, and therefore the Excess of the Height of the Equator is $\frac{1}{265}$ Part of the Radius; which is not half so much as the other from *Maupertuis* and *Murdoch*, and is not far different from the Ratio by Sir Isaac Newton.

But, from his finding the great Difference, he softens a little, and says, that either the Earth is not a perfect Spheroid, or there are some Errors committed in the Mensuration: And is forced to say, that it is difficult to determine the true Ratio. What then shall we trust to? And what is come of this Bray of representing the Spheroidal Surface of the Earth and Sea, without any Error at all, (See Page 27).

This is to be remarked, that since his first Way from *Maupertuis* and *Murdoch* makes the Earth differ twice as much from a Sphere, as his second Way from *Juan* and *Newton*. That if *Juan* and *Newton* are right, (and who can shew the contrary) then it will follow that the Sphere

Of the DIMENSIONS of the EARTH as a SPHEROID.

By the three Ratios found of the Squares of the Earth's Diameters,

$$\frac{u^2}{z^2} = a = \frac{959.352}{981.345} = \frac{.977589}{1} = .977589; \frac{u}{z} = \frac{.988731}{1}$$

$$a = \frac{1298.347}{1310.543} = \frac{.9906937}{1} = .9906937; \frac{u}{z} = \frac{.995336}{1}$$

$$a = \frac{338.7345}{339.8269} = \frac{.9967855}{1} = .9967855; \frac{u}{z} = \frac{.9983917}{1}$$

Here you may see how this same Method, by Mr. Martin's mistaken Radius of the Earth's Curvature, gives the Ratios of the Earth's Diameters, very different.

The Ratio of the Earth's Diameters according to Don JUAN, $\frac{u}{z} = \frac{265}{266} = \frac{.996240}{1}$.

HENCE is shown the ERROR, of Mr. MARTIN's METHOD of determining the Ratio of the Earth's Diameters, by an incorrect Radius of Curvature.

OTHERWISE: According to M. MAUPERTUIS' METHOD of determining the said RATIO; Mr. Martin not chusing to depend on his RULES given, at P. 10. 11. in his *New Principles of NAVIGATION*; who has therefore followed THIS METHOD.

LET the Radius of the Equator to the Semi-axis be as 1 to n ; and A and B the Length of two different Degrees, measured in different Latitudes; and a and b the respective Sines of those Latitudes; then, from the Properties of an Ellipsis;

$$A \times 1 + \frac{1}{2} \times n^2 - 1 \times a^2 = B \times 1 + \frac{1}{2} \times n^2 - 1 \times b^2,$$

$$\text{Or, } 1 - n^2 = \frac{2 \times A - B}{3 \times Aa^2 - Bb^2}, \text{ where } n = \sqrt{1 - \frac{2 \times A - B}{3 \times Aa^2 - Bb^2}}.$$

Sphere is as near the true Figure of the Earth as his first Figure, drawn from Maupertuis. And therefore those that use the meridional Parts for the Sphere will come as nigh the Truth as those who follow his first Method from Murdoch and Maupertuis.

What then can be said of this Boasting Refiner? Can this Refined Navigation give the true Figure of the Earth so as to found any certain Computations upon it. If not, all that he has said imports Nothing to the Purpose. And who would not sooner follow the simple Method by the Sphere, than follow an uncertain Method, only for the Sake of its Novelty and Difficulty, when the Difference is insensible and useless?

When he comes to compare the different Measures of a Degree on the Earth's Surface, this Author is in the utmost Distress, and tries to accommodate them, by taking a Mean; but what Right has he to do that? Or, who will trust to the Truth merely upon his Word, when Sir Isaac Newton's would not be taken without Demonstration?

I wonder how he got it into his Conception, that Mr. Emerson depreciated that Discovery of the Spheroidal Figure of the Earth which is truly a very noble one in natural Philosophy. He only called it an idle Refinement when applied by Pedants to the Practice of Navigation, where it can make no sensible Difference; and must create an endless Deal of Trouble for no useful Purpose. Mr. Emerson gave his Reasons for calling it an idle Refinement, towards the latter End of his Preface, without Design (he says) of offending any Body. And what Right has any Man to depreciate another's Judgment, for what he advances, unless he properly answers him; which the Author of the *New Principles* (long since discovered) never once attempted.

And Mr. Emerson still continues to affirm, that the trifling Difference it makes is not worth the immense Labour (of making meridional Parts and new Arches) that it causes. That as we know nothing what Species of a Spheroid the Earth is, or whether it is a Spheroid at all, or not? I hope you will not follow this refined Author's Steps of inserting a Table of meridional Parts for the Spheroid, except it be to confront and compare with those of the Sphere, and confute the pretended Advantage of Spheroidal above Spherical Navigation by Examples given of both, to show the useless Difference.

The Reason why Simpson's Difference of the Meridional Parts is less than the Spheroidal Author's, is because Simpson uses Sir Isaac Newton's Numbers 229 to 230. I have found that the Difference = 34389s, where $s = \text{Nat. Sine Lat. } q = 1 - aa$, $a = \text{Semi-Axis of the Earth}$.

As to Robertson's Navigation you mention, it is rather a Course of Mathematics; for it contains almost all Branches of practical Mathematics. Barrow is much the same, and follows Murdoch in the Spheroid.

In the Voyages you have made you observe, that neither Course nor Distance can be truly ascertained. And therefore who would go about to take an immense Deal of Trouble to avoid an Error which is involved with another, ten or twenty Times as great; and which is unavoidable; and yet these Errors, as well one as another, may be all corrected from taking an Observation.

I know no Way of correcting the Departure but by finding the Longitude to every particular Course; yet this is troublesome, and in 24 Hours is insensible; and therefore in Practice every Body makes use of the whole Easting or Westing, instead of it.

YOUR METHOD of finding the Departure, Difference of Longitude, Latitude, &c. by the Seaman's COMPUTER, &c. is pretty enough, and will be of Use in Practical Navigation.

I know of nobody that has taken the Trouble to compute the Meridional Parts to the Spheroid, but Murdoch (which is only in whole Degrees) and what the Author of the *New Principles* gives us from Juan.

As to the Tables of reputed Longitude and Latitude of Places, there is a great Difference among Authors.

The best Method to ascertain them nearest (since exact Truth cannot be had) is by comparing several Books together, and where they differ, to apply to the best Maps.

The Longitudes are best reckoned from the westernmost Part of Africa, as being long since established by Custom, and ought not to be altered. Thus, the celestial Longitudes are reckoned from the first Point of Aries, and ought not to be altered; though some whimsical Persons, in Contradiction to a settled Custom, might reckon them from Libra, or any other Point of the Ecliptic.

NEWTONIENSIS.

Of the DIMENSIONS of the EARTH as a SPHEROID.

Or, (more simply) putting $m = 1 - n$, then $n = 1 - m$, and $n^2 = 1 - 2m + m^2$, where m^2 being very small, in Respect of the other Terms, may be rejected; and $n^2 = 1 - 2m$, and $1 - n^2 = 2m = \frac{2 \times A - B}{3 \times Aa^2 - Bb^2}$ above.

Whence, $m = \frac{A - B}{3 \times Aa^2 - Bb^2}$; or when B nearly A,

Then, $m = \frac{A - B}{3A \times a^2 - b^2}$, as is evident.

Now, if the Measure of one of the Degrees be taken near the Equator (when $b = 0$) then $m = \frac{A - B}{3Aa^2}$, as is also evident.

Whence, we have $B = A - 3Aa^2m$, the Length of a Degree of the Meridian at the Equator.

EXAMPLE. LET $A = 57437,9$ the Toises in a Degree in Lapland, and $B = 57050$ the Toises in a Degree at the Parallel of 45° in France.

Then, by the Equation foregoing, $m = 0,006646 = 1 - n$,

Whence, $1 - 0,006646 = 0,993354 = n$.

Therefore, $1 : n = ,993354 :: 266 : 264,3$ the Ratio of the Earth's Diameters.

CONSEQUENTLY, To find the Length of a Degree at the Equator?

LET $A = 57437,9$ Toises in a Degree measured in Lapland, and we have by the Equation $B = A - 3Ama^2 = 56501\frac{1}{2}$ Toises at the Equator: Or, if A be taken $= 57050$ Toises in a Degree of Latitude in France, under the Parallel of 45° , Then $B = A - 3Ama^2 = 56497,2$ the Toises in a Degree at the Equator.

From both which Methods, of M. Maupertuis, by a Degree measured in Lapland, and another measured in France, a Degree at the Equator should be nearly 56500 Toises; being less than the Degree measured there by Don Juan, at the Equator, by about 268 Toises.

BUT, admitting the Truth of Don Juan's Measure of a Degree at the Equator, and also of the Measure of a Degree before in France, Latitude 45° , Then $m = 0,003297$; Whence $1 : n = ,9967 :: 266 : 265,12$; according to which Ratio, viz. 266 to 265, Don Juan has fixed the Ratio of the Earth's equatorial to the polar Diameter; as on which Hypothesis he has computed his Table of meridional Parts of the Earth, as a Spheroid, which meridional Parts we have given with those of the Sphere for the Navigator's Trial and Curiosity.

If the two extreme Degrees, measured in Lapland and at the Equator, be taken as true, Then $A = 57437,9$ and $B = 56767,788$, whence $m = \frac{A - B}{3Aa^2} = 0,004625$.

Consequently, $1 : n = 0,99537 :: 266 : 264,73$, being a mean Quantity between 264,3 and 265,12 the Numbers before found, as Members of the Ratio, from the Measure of a Degree in Lapland and France, and France and at the Equator.

By M. Maupertuis' Method.	According to the New Principles of Navigat.	
NOW, 1 to ,99354	By Mr. Martin.	From a Degree measured in
1 to ,9967	1 to ,98873	Lapland and France,
1 to ,99537	1 to ,99839	France and at the Equator,
	1 to ,99533	Lapland and Equator.

WHENCE, it evidently appears, if M. Maupertuis' Method of determining the Ratio of the Earth's Diameters, by two different and distant Degrees measured on the Earth's Surface, be nearly true, as it appears to be consistent in the near Agreement of the different Answers, it also appears as evident, that the Method, by the New Principles of Geography and Navigation, for determining the Ratio of the same Earth's Diameters, is erroneous; from the different Answers to the same Requisites, deduced as above. For it supposes (what is not) an elliptic Arc to be circular, for a Degree; and the Middle thereof, of circular Curvature, to coincide with the elliptic Curvature of the Earth's Meridian. The Radius of Curvature likewise of the measured Degree considered as a circular Arc, to be the Radius of Curvature to the Middle of that Degree of the elliptic Meridian. Which to suppose, is against the Hypothesis admitted, and Dimensions of the Earth sought, as a Spheroid, formed by the Rotation of the Semi-elliptis about its shorter or polar Diameter; not "about its longer Diameter," as is asserted in the Introduction to the New Principles of Navigation.

HENCE, it necessarily follows, that the Solutions to the XI Problems, P. 12, 13, 14, of this New Navigation, are from erroneous Principles, differing from the Truth, by other Methods; as in the Scholium, P. 14, is signified.

Of the DIMENSIONS of the EARTH as a SPHEROID.

The MEASURE of the EARTH at different Times by different PERSONS.

Years since Christ.	PERSONS NAMES measuring	PLACES NAMES measured.	Geographical Miles.
800	Almamoon, Caliph of Babylon	Babylon	56, or 57.
1525	Fernelius	France	68.
	Snellius, a Dutch Mathematician	between Alcmæer and Bergen- op-Zoom, to $1^{\circ} 11' 30''$. . . Alcmæer and Leyden, a Mean of both	French Toises. 55021 in a Degree.
			Miles.
1635	Norwood, an Englishman	between London and York	$69\frac{1}{2}$ in a Degree. = 367196 Feet.
1654	Ricciolus, by several Methods found the Earth's Circumference		64363 Bologna Paces. or 61650 Toises.
LEWIS XIV. . . .	M. Picard measured between Malvoisine and Amiens		Toises.
	Less than Norwood in a Deg. by 240 Toises	788500 Toises to $1^{\circ} 22' 55''$. . .	57060 to 1° .
1718	M. Cassini measured the whole Meridian of France.		Toises.
		{ From Paris to Collioure Paris to Dunkirk Dunkirk to Collioure	{ 57097 56960 57060 } to 1° .
	Same as Picard		57033 to 1° .
	Mr. Muschenbroek, correcting Snellius, measured the same Arch, and found		57033 to 1° .
*** The above Measurements had been sufficient for determining the Earth's Dimensions as a Sphere, but Disputes arising about its Figure, according to Sir Isaac Newton, as a Spheroid, further Measure was made, to determine this important Truth, as near as possible.			
Began 1736	Messieurs Clairaut, Camus, Le Monnier, Maupertuis, the Abbe Outhier, of France, M. Celsus, Professor of Astronomy at Upsal	measured a Degree in } Lapland, near the } Arctic Circle . . . }	Toises. 57437,9 to 1° .
ended Dec. following.		Picard's less by	377.9
	Picard's corrected, allowing for Aberration of Light, of the observed Star, the Precession of Equinoxes, Refraction of Light, (before neglected)		56925,7 to 1° .
		Less than at the Polar Circle : . . .	512,2
Doubts arising about the Difference of a Degree in France, at the Return of the Academicians from the North, the Arch was remeasured by Cassini de Thury (Grandson to the former) and De la Caille, the Work was intitled, La Merid. de Paris verifiée, 45° Parallel. Toises 57050 to 1° .			
The Difference of a Degree in Lapland and France proved the Figure of the Earth to be no Sphere, but rather a Spheroid; the Ratio of which Diameters (the less Polar to the greater equatorial) were now to be correctly determined. On which Ratio were now to depend the Improvement of Geography and Astronomy, and also Navigation, so far as the Earth's Figure can be applied to the Practice of that Art. To which End a further Measure was applied near the Earth's Equator.			
Set out 1735	Don George Juan, Don Antoine de Ulloa, attending the Academicians		
See Juan's Book.	from France, by whom two Observatories were erected in the Plain of Yaruqui, one at Pueblo Viejo, and the other at Cuenca, by which, with a great Number of Signals erected, they formed and measured about 30 Triangles, and obtained a Base Line of 195734,574 Toises, between the Parallels of Pueblo Viejo and Cuenca		Pueblo Viejo to Cuenca. Lat. $0^{\circ} 32' 45''$ N. $2^{\circ} 54' 7\frac{1}{2}''$ S. 32 45

Diff. Latitude Sum $3^{\circ} 26' 52\frac{3}{4}''$.As $3^{\circ} 26' 52\frac{3}{4}''$: 195734,547 :: 1° at the Equator : 56767,788.

Taking 57437,9 } in a Degree { at the Arctic Circle,
57050, } under Parallel of 45° N. or France } by Rules before given { Ratio of

The Earth's Semi-Axis to the Semidiameter of the Equator, as 264,3 to 266.

Nearer to Sir Isaac Newton . . . Don Juan determines it to be 265 to 266.

It is known, that $1^3 : n^3$ (1 to n , being the Ratio of the Equatorial to the Polar Diameter) :: Length of a Degree at the Equator : to the Length of a Degree at the Pole, by the elliptic Theory. Or, $1 : n :: 266 : 265$. Therefore, $265^3 : 266^3 :: 56768$ Toises in a Degree, at the Equator : 57438 Toises in a Degree at the Pole. But, they are the Toises in a Degree at Lapland near the Arctic Circle, and much less than in a Degree at the Pole. Hence, the Number 265 is a little too large (as it is said by the Refiner) in the Ratio of 266 : 265, the Equatorial to the Polar Diameters of the Earth. And therefore the Matter is still in Dispute as to the true Figure of the Earth, whether it is an exact Spheroid, or not; and the Doubt is likely to remain, till we can measure the Length of a Degree at the Pole (non est habitabilis Niue). That, upon the whole (notwithstanding the late Author's pretended Improvements by the elliptic Chart, constructed from Meridional Parts on wrong Principles) the Mercator's Chart,

Of the DIMENSIONS of the EARTH as a SPHEROID.

for *practical Sailing*, will prove the Correcter of the two. Whatever might be expected from a *Chart* constructed on *Juan's* (instead of one constructed from *Murdoch's* or *Maupertuis's*) Principles.

That *unless* the Dimensions of the Earth be *exactly*, or very nearly, determined, and what Sort of a Figure it is, the Errors in Navigation from a wrong Figure of the Earth admitted will be greater than those from supposing it an *exact Sphere*, exclusive of the larger Errors in the *Course* and *Distance*, that are impracticable to correct, but by *celestial Observation*. Nor will the *incorrect* Determination of the Figure of the Earth have a better Effect, in the true Determination of the *Parallaxes* of the Moon, *Beginning*, *Middle*, and *End* of the *Solar Eclipses*, their Quantity of Duration, the Sun's *Parallax* by the *Transit* of *Venus* over his Disk, 1761, which will all be a little precarious.

COMPARISON of different SPHEROIDAL with the same SPHERICAL NAVIGATION.

EXAMPLE. Mr. *Murdoch* (and after him Mr. *Martin*) supposes a Ship sails from a Place in 25° Latitude South to another in 30° Latitude N. (the Difference of Latitude 55°) on a Course of 43° from the Meridian (being a Case that never happens in the Practice of Navigation) required the Difference of Longitude and Distance run?

The Difference of Longitude, by *Mercator*, is 3206'

The Difference of Longitude, by *Murdoch's Spheroid*, or Figure of the Earth 3141

Difference, less 65

Distance sailed according to the Sphere 4512'

Distance sailed by *Murdoch's Spheroid* 4423

Difference, less . . . 89

Difference of Longitude according to *Juan's Spheroid* 3148

Distance sailed, according to *Juan* 4482

Difference of *Juan*, from *Murdoch*, or *Martin*, in Longitude . . . 7+

Distance sailed 59+

Notwithstanding Mr. *Martin's* pretended Refinement of Navigation from Error, he, in one Part of his Introduction, speaks thus, (as if he had forgot himself) "It is too well known, how many Errors and Mistakes our most skilful Seamen are liable to in the Direction of their Course, their Distance run, &c." — Therefore to agree with M. *Maupertuis*, when all the Elements of Navigation are brought to Perfection, it will then be seen to what a Degree of Perfection this Element of Spheroidal Navigation (being applied to Practice) will have among the rest, the Errors of the Course and Distance still remaining.

TO determine (more correctly) the Ratio of the Diameters of the Earth, from the Measure of two different Degrees of Latitude on its Surface, supposing the Form of the Earth to be that of an oblate Spheroid?

PUT c = Semi-Axis.

t = Semidiameter of the Equator.

M, m the Measures of a Degree of the Meridian at the given Latitudes of the Places P, p .

S, s the Sines } of those Latitudes.

S', s' the Cosines }

τ' the Cotangent of the Latitude P .

r = tabular Radius.

$$A = M^{\frac{2}{3}}, a = m^{\frac{2}{3}}, B = \frac{ass}{ASS}, b = \frac{as's'}{AS'S'}, N =$$

$$\frac{A-a}{AS'S' - as's'} rr, n = \frac{A-a}{ass - ASS} rr.$$

$$\text{Whence, } \frac{c}{t} = \sqrt{1+n} \text{ or } = \sqrt{\frac{1}{1-N}},$$

$$\text{or } = \frac{\tau'}{r} \sqrt{\frac{b-1}{1-B}}, \text{ or } = \frac{\tau'}{r} \sqrt{\frac{1-b}{B-1}}.$$

of a Degree at different Places on its Surface are not true. — Hence, the immense Expence and Trouble the French Nation have been at in taking the several Mensurations of Arches on the Earth's Surface, are intirely thrown away. And we shall at last be forced to depend more on the Computations of Sir Isaac Newton, or Mr. Emerson, by the Laws of Gravity. While the different Densities of the Earth's Substance, next its Surface, and towards its Center, (on which that Computation depends) remain uncertain.

AND HENCE IS PROVED THE ABSURDITY OR UNCERTAINTY OF SPHEROIDAL NAVIGATION! AND CONSEQUENTLY OF ITS BEING AN IDLE OR USELESS REFINEMENT.

For were the true Figure of the EARTH and SEA determined (which may vary by the Effect of the Moon's Gravity more or less at different Times on the Sea) its Discovery would be a greater Advantage to Geography than to the PRACTICE of NAVIGATION: since the Spherical Surface of the Earth only considered, is sufficient to answer the End of all which Purposes.

Therefore Mrs. MOUNT and PAGE on Tower-Hill, may continue the Sale of their OLD MERCATOR CHARTS with superior Advantage to any NEW SPHEROIDAL ONES, constructed (from *Murdoch's*) doubly erroneous, when compared with the Sphere, and *Juan's* supposed truer Spheroid. — For the Meridional Parts of the New Spheroidal Chart (prematurely or inadvertently made by B. *Martin*) differ twice as much from those of the supposed truer Spheroid (last recommended for Practice in Navigation, by B. *Martin*) as those truer differ from the Meridional Parts of the Sphere.

Latitude.	Toises in Deg. measured.	Roots of which De. Toises.
Ex. <i>Lapland</i> $66^{\circ} 30'$	57437,9 = M	1488,672 = $M^{\frac{2}{3}}$
<i>France</i> $49^{\circ} 22'$	56925,7 = m	1479,811 = $m^{\frac{2}{3}}$
<i>Equator</i> 0°	56767,8 = m	1477,071 = $m^{\frac{2}{3}}$
By <i>Lapland</i> } $\frac{c}{t} = \sqrt{1+n} = \sqrt{1 - ,0221728} = \frac{,98885}{1}$		
<i>France</i> } $\frac{c}{t} = \sqrt{1+n} = \sqrt{1 - ,00926618} = \frac{,99535}{1}$		
<i>Lapland</i> } $\frac{c}{t} = \sqrt{1+n} = \sqrt{1 - ,00926618} = \frac{,99535}{1}$		
<i>Equator</i> } $\frac{c}{t} = \sqrt{1+n} = \sqrt{1 - ,00321503} = \frac{,99839}{1}$		
<i>France</i> } $\frac{c}{t} = \sqrt{1+n} = \sqrt{1 - ,00321503} = \frac{,99839}{1}$		
<i>Equator</i> } $\frac{c}{t} = \sqrt{1+n} = \sqrt{1 - ,00321503} = \frac{,99839}{1}$		

* From the above variable Ratios of the Earth's Diameters, deduced from unerring Rules (supposing the Earth to be in the Form of a Spheroid) it is evident, that either the Earth is not a Spheroid, or else the Mensurations taken

The PROPERTIES of a Right-angled PLANE TRIANGLE.

IF each Side of a right angled plane Triangle be made Radius, and Circles be described thereto from each End of the Hypotenuse, at the acute Angle, as a Center, then the two Sides, lying contiguous, and third Side lying remote to the same Center, will be the Natural Secant, Cofine, and Sine; Cotangent, Secant, and Tangent, respectively, of the Angle at the Center, included by the Hypotenuse and one Leg; each of those two Sides being made Radius.

Or, The natural Secant, Cofine, and Sine; Cotangent, Secant, and Tangent, of one Angle (included by the Hypotenuse and one Leg) under 45° , will be the natural Cofecant, Sine, and Cofine; Tangent, Co-secant, and Cotangent, respectively, of the other Angle, above 45° , and the contrary. And therefore these will be proportional to the Secants, Sines, Tangents, to the Radius in the Table.

On this Property of right-angled plane Triangles, the Seaman's Ready Computers No. I. II. III. are founded, by making each Side of a right angled Triangle Radius, = 10000, and taking out of the Tables for the 3 varied Sides, made Radius, 1, 2, 3, 4, 5, 6, 7, 8, 9 Times Radius, Cofine, and Sine; Radius, Secant, and Tangent; Radius, Cofecant, and Cotangent, correspondent, for all the Angles from 1 to 45° Degrees; whose Complements, having the same Numbers, from 45° to 90° Degrees, serve for 1, 2, 3, 4, 5, 6, 7, 8, 9 Times the same Complements, or for all the Angles from 45° to 90° Degrees.

LET 10000, 20000, 30000, 40000, 50000, 60000, 70000, 80000, or 90000 = Radius.		Computer, No. I.	
		Radius.	Other Sides.
Correspondent to which the other two Sides are set down in Computers, No. I. II. III. Being found by Multiplication of 1, 2, 3, 4, 5, 6, 7, 8, 9 into Natural	Cofine and Sine Tangent and Secant Cotangent and Cofecant	Hypotenuse.	Distance. Dif. Latitude. Departure
			Distance. Departure. Dif. Latitude.
			Computer, No. II.
		Longer Leg.	Radius. Other Sides.
			Dif. Lat. Departure Distance.
			Depart. Dif. Lat. Distance.
		Shorter Leg.	Computer, No. III.
			Radius. Other Sides.
			Depart. Dif. Latitude Distance.
			Dif. Lat. Departure Distance.

Some PROPERTIES of SINES, TANGENTS, &c.

THE Cofine, Tangent, Cotangent, and Secant are common to two Arches, which are Supplements to each other. Yet in an Arc ABOVE A QUADRANT they are Negative: being drawn the contrary Way to those of an Arc less than a QUADRANT.

In General	$\left\{ \begin{array}{l} \text{Sine} \\ \text{Co-versed Sine} \\ \text{Cofecant} \end{array} \right\}$	$\left\{ \begin{array}{l} \text{Affirmative} \\ \text{Negative} \end{array} \right\}$	For all Arcs	$\left\{ \begin{array}{l} \text{under} \\ \text{above} \end{array} \right\}$	180° .	Cofines	$\left\{ \begin{array}{l} \text{Affirmative} \\ \text{Negative} \end{array} \right\}$	fr. 0° to 90° , fr. 270° to 360° . fr. 90° to 270° .	
Tangents.	$\left\{ \begin{array}{l} \text{Affirmative} \\ \text{Negative} \end{array} \right\}$	$\left\{ \begin{array}{l} \text{from } 0^\circ \text{ to } 90^\circ, \text{ from } 180^\circ \text{ to } 270^\circ. \\ \text{from } 90^\circ \text{ to } 180^\circ, \text{ from } 270^\circ \text{ to } 360^\circ. \end{array} \right\}$				Cotangents	$\left\{ \begin{array}{l} \text{Affirmative} \\ \text{Negative} \end{array} \right\}$	$\left\{ \begin{array}{l} \text{from } 0^\circ \text{ to } 90^\circ, \text{ from } 180^\circ \text{ to } 270^\circ. \\ \text{from } 90^\circ \text{ to } 180^\circ, \text{ from } 270^\circ \text{ to } 360^\circ. \end{array} \right\}$	
Secants.	$\left\{ \begin{array}{l} \text{Affirmative} \\ \text{Negative} \end{array} \right\}$	$\left\{ \begin{array}{l} \text{from } 0^\circ \text{ to } 90^\circ, \text{ from } 270^\circ \text{ to } 360^\circ. \\ \text{from } 90^\circ \text{ to } 270^\circ. \end{array} \right\}$				Cofecants	$\left\{ \begin{array}{l} \text{Affirmative} \\ \text{Negative} \end{array} \right\}$	$\left\{ \begin{array}{l} \text{from } 0^\circ \text{ to } 180^\circ. \\ \text{from } 180^\circ \text{ to } 360^\circ. \end{array} \right\}$	
Versed Sines . . Affirmative, in general.						Sines Tangents &c.	$\left\{ \begin{array}{l} \text{of Negative} \\ \text{Arcs} \end{array} \right\}$	$\left\{ \begin{array}{l} \text{Have} \\ \text{contrary} \\ \text{Signs} \end{array} \right\}$	$\left\{ \begin{array}{l} \text{to those of} \\ \text{Affirmative} \\ \text{Arcs.} \end{array} \right\}$

For other infinite Properties of Sines, Tangents, &c. consult Mr. EMERSON'S TRIGONOMETRY, who has there exhausted the Subject, and left no Room for any Person to make Improvement after him.

SOME PROPERTIES of the SIDES and ANGLES of PLANE TRIANGLES.

- As Sum of Hyp. and $\frac{1}{2}$ longer LEG of a right angled Triangle is to 86, so is the shorter LEG to its opposite Angle, nearly.
- a being the Degrees of the less Angle, it will be,

$$\text{As } \frac{57.3}{a} + \frac{3a}{1000} : 1 :: \text{Hypotenuse} : \text{Opposite Side to that } \angle, \text{ nearly.}$$

- a being the Degrees of the less Angle, it will be,

$$\text{As } 1 : 1 - \frac{1\frac{1}{2}aa}{10000} :: \text{Hypotenuse} : \text{adjoining Side to that } \angle, \text{ nearly.}$$

- In any right-lined Triangle, As the Base : Sum Sides :: Diff. Sides : Diff. Segments Base by a Perpendicular.

By these 4 Properties, and 47 Proposition I. Euclid, all the Cases of right-lined Triangles may be solved without Tables: letting fall a Perpendicular from the End of given Side, and opposite to a given Angle.

- In any right-lined Triangle, As Sum of any two Sides : to their Diff. :: Tangent $\frac{1}{2}$ Sum of their opposite $\angle$ s : Tangent $\frac{1}{2}$ their Difference.
- In any right-lined Triangle, As twice the Rectangle of the Legs, including an $\angle$: Sum Squares of the Legs — Square Base :: Radius : Cof. vertical $\angle$ included by the Legs.

ANSWERS to all the CASES of PLANE TRIANGLES.



CASES of Right-angled PLANE TRIANGLES.

Cases.	GIVEN	SOUGHT	ANSWER.
			By PROPORTION.
1	Hyp. & $\angle$ BC, B C	Op. Leg CA BA	Rad. : Hyp. :: s. given $\angle$: opp. Leg. R : BC :: s. B : CA. s. C : BA. Or, Sec. $\angle$: Hyp. :: t. $\angle$: opp. Leg.
2	Hyp. & Leg BC, CA BA	Op. $\angle$ B C	Hyp. : Rad. :: Leg : s. opp. $\angle$. BC : R :: CA : s. B. BA : s. C. Or Leg : Rad. :: Hyp. : Sec. op. $\angle$ to Leg.
3	Hyp. & Leg BC, CA BA	Other Leg BA CA	Find the $\angle$ s by Case 2. And then the Leg by Case 1.
4	Leg & $\angle$ CA, B BA, C	Hyp. BC	s. $\angle$: Leg opp. : Rad. : Hyp. s. B : CA :: R : BC. s. C : BA. Or Rad. : Leg :: sec. adj. $\angle$: Hyp.
5	Leg & $\angle$ CA, B BA, C	Other Leg BA CA	s. $\angle$: opp. Leg. :: s. other $\angle$: op. Leg. s. B : CA :: s. C : BA. s. C : BA :: s. B : CA. Or Rad. : Leg :: t. adj. $\angle$: other Leg.
6	Legs BA, CA	$\angle$ B C	One Leg : Rad. :: other : t. op. $\angle$. BA : CA :: R : t. B } Complements of CA : BA :: R : t. C } one another.
7	Legs BA, CA	Hyp. BC	Find the $\angle$ s by Case 6. Then the Hyp. by Case 4.

EACH 4 of the following Terms, in the same Rectangle are proportional, read from Left to Right, or from Right to Left: Shewing the Variations of answering trigonometrical Cases by the Tables.

Radius : Sine : Cofine } N. B. Any two of these Proportional
Secant : Tangent : Radius } may be put in the Room of
Cofecant : Radius : Cotangent } any other two like Proportionals.
Thus, As Rad. : s. given $\angle$::
Or, As Sec. : t. given $\angle$:: Hyp. : opp. Leg to the given $\angle$.
Cofec. : Radius ::

The above like Proportionals may be also varied in the Solution of spherical Triangles further on, by Lord Napier's Proposition.

UNIVERSALLY.

IN any right-lined Triangle, whose Sides are represented by A, B, C, and opposite Angles by a, b, c, any three of the six Terms being given, (except the 3 Angles) the Rest are thus determined.

CASE I. When there is a Side and $\angle$ op. given.

Then $\frac{A}{s. a} = \frac{B}{s. b} = \frac{C}{s. c}$, whence the

Term reqd. is had.

When two Sides A, B, and an $\angle$ opposite to one of them A is given, if A is less than B, then b, c, and C have each two Values. If A is greater than B, then b, c, and C have but one Value.

CASE II. Two Sides (A the greater, B the less) and an Angle c included are given.

Put $D = \frac{180^\circ - c}{2}$, and $\frac{A-B}{A+B} \times t. D = t. E$.



Then $a = D + E$, and $b = D - E$.

Likewise, Put $L. 2A + L. 2B + 2L. s. \frac{1}{2}c - 20.000000 = L. F$. and $2L. A - B = L. G$: Then $\frac{1}{2}L. F + G = L. C$, the 3d Side, reqd. N. B. If the included $\angle$, c, is 90° ,

Then, $t. a = \frac{A}{B} \times r$; $t. b = \frac{B}{A} \times r$; and $C = \frac{s. a}{r} \times A$,

or $= \frac{s. b}{r} \times B$.

N. B. s, t, r, and L, denote Sine, Tangent, Radius, and Logarithm.



CASES of Oblique PLANE TRIANGLES.

Cases.	GIVEN	SOUGHT	ANSWER.
			By PROPORTION.
1	$\angle$ s & a Side A, B, BCA [BC, CA]	Other Sides BA CA	s. $\angle$: op. Side :: s. other $\angle$: op. Side. s. A : BC :: s. BCA : BA. s. B : CA.
2	2 Sides & $\angle$ opposite to one BC, CA, A	Other $\angle$ s B A BCA	Side : s. op. $\angle$:: Side : op. $\angle$. BC : s. A :: CA : s. B. CA : s. B :: BC : s. A. Now, two $\angle$ s known, their Sum fr. 180° leaves the 3d $\angle$ BCA.
3	2 Sides & $\angle$ opposite to one BC, CA, A	Other Side AB B	Find $\angle$ opposite Side required by Case 2. Then the required Side by Case 1. s. A : BC :: s. BCA : AB. s. B : CA :: s. BCA : AB.
4	2 Sides and $\angle$ included BA, BC, B	Other $\angle$ s BCA, A	Sum Sides : their Dif. :: Tan. $\frac{1}{2}$ Sum op. $\angle$ s : Tang. $\frac{1}{2}$ their Difference. Which added to and subtracted from the $\frac{1}{2}$ Sum gives the 2 Angles required.
5	2 Sides and $\angle$ included BA, BC B	The other Side. CA	Find the $\angle$ s by Case 4, and then the sought Side CA by Case 1.
6	3 Sides BA, BC, CA	$\angle$ opposite to one Side. A.	Let fall Perp. CD opposite $\angle$ required. Then, Base : Sum Legs :: Dif. Legs : Dif. Segments Base, BD - DA, where the Perpendicular CD falls, $\frac{1}{2}$ which Diff. = mD, added to and subtracted from half the Base gives the Segments BD, and DA. Whence $\angle$ A is found by Case 2d of Right-angled Triangles.

CASE III.

FROM the 3 SIDES given, A, B, (including a required $\angle$, c) and C the Side opposite to a required Angle, this Angle may be found thus.

RULE. Add the 3 Sides together, and take the $\frac{1}{2}$ Sum thereof, and set down its co. Logarithm. Under which set down the co. Logarithm of the Difference between which half Sum and the Side opposite to the required Angle, and then set down the Logarithms of the Differences between which Half Sum and the other two Sides; the Half Sum of which four Numbers will be the Logarithm Tangent of Half Angle, required.

Thus, If $\frac{A+B+C}{2} = H$ co. Log.

$H - C$ Side opposite $\angle$ required . . . co. Log.

$H - A$ Log.

$H - B$ Log.

$\frac{1}{2} \angle$ reqd. Log. Tang. $\frac{1}{2}$ Sum.

Doubled the $\angle$. See Page 116 for Example.

For, $H \times H - C : H - A \times H - B :: 1$ (Rad. sq.) t^2 , $\frac{1}{2} \angle$.
Also, $AB : H - A \times H - B :: 1$ (Rad. sq.) s^2 , $\frac{1}{2} \angle$.

HENCE this RULE. To the Log. Complements of the Sides including the sought Angle, add the Logarithms of the Difference between the Half Sum of all the 3 Sides and each of those including Sides, the half Sum of those 4 Numbers will be the Log. Sine of half the sought Angle.

Thus, A, including Side co. Log.
B, including Side co. Log.
 $H - A$ Log.
 $H - B$ Log.

$\frac{1}{2}$ sought $\angle$ Log. Sine $\frac{1}{2}$ Sum.
Doubled the $\angle$.

TABULAR RULES for answering all the CASES of SAILING.

A TABLE of the 6 PLAIN SAILING CASES.				In WORDS.		
Cases.	Given.	Sought.	ANSWER.	Given.	Sought.	ANSWER.
1	cd	δ	Rad. : cof. c :: d : δ . . by 2 Proportion. Rad. : sin. c :: d : e 1	Course Distance	Dif. Lat. Departure	As Rad. to Cof. Course so Dif. to Dif. Lat. Rad. to sin. Course so Dif. to Depart.
2	δe	c	δ : e :: Rad. : tan. c . . . 3 Sin. c : e :: Rad. : d 1	Dif. Lat. Departure	Course Distance	Dif. Lat. to Depart. so Rad. to Tan. Course. Sin. Course to Depart. so Radius to Dif.
3	ce	δ	Tan. c : e :: Rad. : δ . . . 3 Sin. c : e :: Rad. : d 1	Course Departure	Dif. Lat. Distance	Tan. Course to Depart. so Rad. to Dif. Lat. Sine Course to Depart. so Rad. to Dif.
4	δd	c	d : Rad. :: δ : cof. c 2 Rad. : Sin. c :: d : e 1	Dif. Lat. Distance	Course Departure	Distance to Rad. so Dif. Lat. to Cof. Course. Rad. to Sine Course so Dif. to Depart.
5	cd	e	Cof. c : Rad. :: δ : d 2 Rad. : tan. c :: δ : e 3	Course Dif. Lat.	Distance Departure	Cof. Course to Rad. so Dif. Lat. to Dif. Rad. to Tan. Course so Dif. Lat. to Depart.
6	de	c	d : Rad. :: e : sin. c 1 Rad. : d :: cof. c : δ 2	Distance Departure	Course Dif. Lat.	Dif. to Radius so Depart. to Sine Course. Rad. to Dif. so Cof. Course to Dif. Lat.
c denotes Course, d Distance run, δ Difference of Latitude, e Departure.						
A TABLE of the 3 PARALLEL SAILING CASES.				In WORDS.		
Cases.	Given.	Sought.	ANSWER.	Given.	Sought.	ANSWER.
1	al	d	Rad. : cof. a :: l : d . . by 4	Latitude Dif. Long.	Distance.	Rad. to Cof. Lat. so Dif. Long. to Dif.
2	ad	l	Cof. a : Rad. :: d : l . . . 4	Latitude Distance	Longit.	Cof. Lat. to Rad. so Dif. to Dif. Long.
3	dl	a	l : d :: Rad. : cof. a . . . 4	Distance Dif. Long.	Latitude.	Dif. Long. to Dif. so Rad. to Cof. Lat.
a denotes the Parallel of Latitude, l Difference of Longitude, and d the Distance of two Places in that Parallel.						
A TABLE of the 7 CASES of Middle LATITUDE and MERCATOR SAILING, concerning LONGITUDE.				In WORDS.		
Cases.	Given.	Sought.	ANSWER.	Given.	Sought.	ANSWER.
1	a, b, l	c	δ : l :: cof. μ : tan. c 7 λ : Rad. :: l : tan. c 9 m : Tan. $51^\circ 38' 9''$:: l : tan. c . . 10 $T = M' + L + N' + R$; $D = \Delta + C' + R$. By Equations 1 and 2. Cof. c : Rad. :: δ : d : Case 5 plain Sailing.	Two Lats. Dif. Long.	Course	Dif. Lat. to Dif. Lon. so cof. mid. Lat. to tan. Crse Mer. Dif. Lat. to Rad. so Dif. Lon. to Tan. Course Dif. Log. Tangents $\frac{1}{2}$ Comps. Lats. : Tan. $51^\circ 38' 9''$:: Dif. Long. : Tan. Course. Cof. Course to Rad. so Dif. Lat. to Dif.
2	a, b, c	l	Cof. μ : tan. c :: δ : l 7 Rad. : tan. c :: λ : l 9 Tan. $51^\circ 38' 9''$: tan. c :: m : l 10 $L = M + N + T - R$. By Equation 1. d as above.	Two Lats. Course	Dif. Long.	Cof. mid. Lat. to tan. Course so Dif. Lat. to Dif. Lon. Rad. to Tan. Course so mer. Dif. Lat. to Dif. Lon. Tan. $51^\circ 38' 9''$ to Tan. Course so Dif. Log. Tangents $\frac{1}{2}$ Comps. Lat. to Dif. Long. The Dist. is found as above, by Case 5, pl. Sail.
3	a, b, d	c	c found by Case 4. plain Sailing. l found as by 2. Case, Mercator.	Two Lats. Distance	Course Dif. Long.	Course found as above by Case 4. plain Sailing. Then Dif. Long. found by 2 Case, Mercator.
4	a, c, l	b	Tan. c : cof. a (for mid. Lat.) :: l : δ ; by which find mid. Lat. and repeat the Operation for the other Latitude. Tan. c : Rad. :: l : λ , subtract or add which Parts from or to those of other Latitude for Latitude correspondent. Tan. c : l :: tan. $51^\circ 38' 9''$: m . . . 10 Which added to Tan. $\frac{1}{2}$ Comp. given Lat. gives the other Latitude. $M = L + N' + T' + R$. Here m and b found. d found by Case 5 plain Sailing.	One Lat. Course Dif. Long.	Other Lat.	Tan. Course to Rad. so Dif. Lon. to mer. Dif. Lat. by this find mid. Lat. and repeat the Operation to get the other Latitude. Tan. Course to Rad. so Dif. Lon. to mer. Dif. Lat. subtract or add which Parts from or to those of the other Lat. and find the Latitude corresponding. Tan. Course to Dif. Lon. so Tan. $51^\circ 38' 9''$ to Dif. Log. Tangents $\frac{1}{2}$ Comps. Latitudes, which added to the tan. $\frac{1}{2}$ Comp. given Lat. gives the other. Dist. is now had by Case 5. plain Sailing.
5	a, c, d	b	δ is found and b by Case 1. plain Sailing. l is found now by 2 Case Mercator.	One Lat. Course, Dif.	Other Lat. Dif. Long.	Dif. Lat. and other Lat. find by Case 2. pl. Sail. Now, Dif. Long. is found by 2 Case Mercator.
6	a, d, l	c	Assume c , by which find δ , b , and e , till (by Trial and Error) δ : e :: λ : l . If meridional Parts of Dif. of Latitudes are greater than those for Lat. given, deduct the less from greater for merid. Parts to Latitude on contrary Side of Equator.	One Lat. Distance Dif. Long.	Course and other Lat.	Assume a Course, by which find Dif. Lat. other Lat. and Dep. till (by Trial and Error) Dif. Lat. to Depart. so Mer. Dif. Lat. to Dif. Longitude. The direct Method of solving this Case by one Rubes or Rebus, P. 64. Pal. 1759, is not true.
7	a, d, l	b	Assume a , and thence find b , till (by Trial and Error) δ : e :: λ : l . The above are all easily answered by Computers I. II. III.	Course Distance Dif. Long.	Both the Latitudes	Assume one Latitude, and find thence the other, till (by Trial and Error) Dif. Lat. to Dep. so Mer. Dif. Lat. to Dif. Longitude.
μ denotes Middle Latitude, λ the Meridional Difference of Lat.						
* The 6th and 7th Cases above are not taken Notice of before by any Writer on Navigation. No Solutions were ever given to them.						
Mr. B. Martin has represented the 6th as impossible, in his Spheroidal Navigation.						
N. B. The Cases of the Departure in Mercator, here omitted, may be answered by meridional Difference of Latitudes, and the 8th Proportion foregoing.						

EXAMPLES of answering the PRINCIPAL CASES of MERCATOR-SAILING. According to different METHODS.

1. GIVEN a, b, l ; To find c, d ? Here $T = M' + N' + L + R$; And $D = \Delta + C + R$.

By LOGARITHMIC EQUATIONS.

EXAMPLE. Required the Bearing and Distance of the Island of St. Vincent, (one of the Cape de Verd Islands) in Latitude $17^{\circ} 4' N$. Longitude $24^{\circ} 44' W$. of Greenwich, from the Lizard Point, in Latitude $49^{\circ} 55' N$. Longitude $5^{\circ} 19' W$. of Greenwich.

$a = 17^{\circ} 4'$	$a' = 72^{\circ} 56'$	$\frac{1}{2} a' = 36^{\circ} 28' 0''$	A = 9.8686804
$b = 49^{\circ} 55'$	$b' = 40^{\circ} 5'$	$\frac{1}{2} b' = 20^{\circ} 2' 30''$	B = 9.5624477
$d = 32^{\circ} 51' = 1971'$ or Miles,			$m = 0.3066327$
$l = 19^{\circ} 25' = 1165$			$M' = 0.5133816$
$\Delta = 3.2946166$			$N' = 6.1015104$
$C' = 0.0450185$			$L = 3.0663259$
$D = 3.3397051$			$R = 10.0000000$
$d = 2186,277$ Miles; $c = 25^{\circ} 38' 23'' W = T = 9.6812179$			

By DIFFERENCE of LOGARITHM TANGENTS.

As Dif. Log. Tangents $\frac{1}{2}$ Complements of Latitudes $36^{\circ} 28'$ and $20^{\circ} 2' 30''$; 3066,327	Logarithms.
To Dif. Long. $32^{\circ} 51' = 1971'$ Miles	co. 6.5133816
So Tang. const. $\angle 51^{\circ} 38' 9''$	3.0663259
	10.1015104
To Tang. Course $25^{\circ} 38' 23'' W$.	9.6812179
d is found as before.	

By COMPUTER, Number II.

Course.	Dif. Lat.	Dist.
$25^{\circ} 38'$	1000	1109.2
	900	998.29
	70	77.64
	1	1.00
1971		2186.13 Distance.

But taking to the nearest Course in whole Degrees will be sufficient.

By MERIDIONAL PARTS.

Lat.	Merid. Parts.	As Mer. Dif. Lat.
$49^{\circ} 55'$	3466,5	2427,5
$17^{\circ} 4'$	1039,	To Dif. Long. 1165' . . . 3.0663259
		So Radius . . . 10.0000000
$32^{\circ} 51'$	2427,5	To Tan. Course $25^{\circ} 38' . . 9.6812179$
Dif. Lat.	Mer. Dif. Lat.	

By COMPUTER, Number II.

2427,5) 1165(.4800 nearly, Quote, against which is Course $25^{\circ} 38'$; according to the Mercator-Triangle.

OR, By COMPUTER, Number II.

Against 26° Course, and 1971 Dif. Lat. you take out 2192 Miles Distance; being but 6 Miles too much in $32^{\circ} 5'$ Dif. Latitude.

2. GIVEN a, b, c ; To find l, d ? Here $L = M + N + T - R$; And $D = \Delta + C' + R = D$.

By LOGARITHMIC EQUATIONS.

EXAMPLE. From the Latitude $49^{\circ} 55' N$. Longitude $5^{\circ} 19' W$. a Ship sails on a Course S. $25^{\circ} 38' 23'' W$. until, by Observation, she finds herself in Latitude $17^{\circ} 4' N$. Required her Latitude and Distance run.

$a = 17^{\circ} 4'$	$a' = 72^{\circ} 56'$	$\frac{1}{2} a' = 36^{\circ} 28' 0''$	A = 9.8686804
$b = 49^{\circ} 55'$	$b' = 40^{\circ} 5'$	$\frac{1}{2} b' = 20^{\circ} 2' 30''$	B = 9.5624477
$d = 32^{\circ} 51' = 1971'$	$\Delta = 3.2946866$		$m = 0.3066327$
	$R + C' = 0.0450185$		$M = 9.4866184$
Miles.			$N = 3.8984896$
$d = 2186,277$	$D = 3.3397051$		$T = 9.6812179$
Long. went fr. $5^{\circ} 19'$			$L = 3.0663259$
Dif. Long. $19^{\circ} 25' = l = 1165$ Miles			
Ship's Long. $24^{\circ} 44'$			

By DIFFERENCE of LOGARITHM TANGENTS.

As Tan. const. $\angle 51^{\circ} 38' 9''$	co. 89.8984896
To Tan. Course $25^{\circ} 38' 23''$	9.6812179
So Dif. Log. Tangents $\frac{1}{2}$ Comps. Lats. $36^{\circ} 28'$ and $20^{\circ} 2' 30''$; 3066,327	3.4866184
To Dif. Long. $32^{\circ} 51' = 1971$ Miles	3.0663259
d is found as before.	

OR, By COMPUTER, Number II.

Against 26° Course, and 2427 Mer. Dif. Lat. you take out 1183' or $19^{\circ} 43'$ Depart. for Dif. Long. but 18' too much in about 33° Dif. Lat. Sailing.

By MERIDIONAL PARTS.

Lat.	Mer. Parts.	As Radius
$49^{\circ} 55'$	3466,5	10.0000000
$17^{\circ} 4'$	1039,	To Mer. Dif. Lat. 2427,5
		Tang. Course $25^{\circ} 38' 23''$ 9.6812179
$32^{\circ} 51'$	2427,5	To Dif. Long. 1165' . . . 3.0663771
Dif. Lat.	M. Dif. Lat.	
		By COMPUTER, Number II.
Course.	Mer. D. Lat.	Dep. for Dif. Long.
Against $25^{\circ} 38'$	2000	959.77
	400	191.94
	20	9.60
	7	3.43
	2427	1164.74 Dif. Long.

3. GIVEN b, c, d ; To find a, l ? Here $\Delta = D + C - R$, whence a and m are found; And $L = M + N + T - R$.

By LOGARITHM EQUATIONS.

EXAMPLE. From the Latitude $49^{\circ} 51' N.$ and $5^{\circ} 19' W.$ from Greenwich, a Ship runs on a Course $S. 25^{\circ} 31' 23'' W.$ 2186,277 Miles: Required from thence the Latitude and Longitude arrived in?

Lat. sailed from $49^{\circ} 55' N.$ $d = 2186,277.$ $D = 3.3397051$
 $c = S. 25^{\circ} 31' 23'' W.$ $C = 9.9549815$

Diff. Latitudes $32 \quad 51 \quad N = d \quad 1971 \text{ Miles, } \dots \Delta = 3.2946866$

Ship's Latitude $17 \quad 4 \quad N.$

$a = 17^{\circ} 4'; a' = 72^{\circ} 56'; \frac{1}{2} a' = 36^{\circ} 28' 0''$ $A = 9.8686804$

$b = 49 \quad 55; b' = 40 \quad 5; \frac{1}{2} b' = 20 \quad 2 \quad 30$ $B = 9.5620477$

$m = 0.3066327$

$M = 9.4866184$

$N = 3.8984896$

$T = 9.6812179$

By DIFFERENCE of LOGARITHM TANGENTS.

Find the Difference of Latitude $= 32^{\circ} 51' N.$ and Ship's Latitude $17^{\circ} 4' N.$ arrived in, by Plain-Sailing, as on the Right-Hand.

Then find the Difference of Longitude as in the former Example, by proportioning the Difference of Logarithm - Tangents of the Half Complements of Latitudes.

Long. sailed from $5^{\circ} 19'$

Diff. Long. $19 \quad 25 = l = 1165 \text{ Miles; } L = 3.0663259$

Ship's Long. $24 \quad 44$ arrived in.

The Course and meridional Difference of Latitude being found as in the former Example, the Difference of Longitude follows as in that Example, by proportioning the meridional Parts, and also by the COMPUTER.

All other Cases occurring in Practice, (except the 6th and 7th) are reducible to the foregoing; by which the Dead-Reckoning of a Ship (as it is called) may be duly kept, and corrected as follows by frequent Observation of the Latitude.

N. B. By the daily Course and meridional Difference of Latitude, (allowing for Lee-Way, Variation, Setting of Currents, &c.) you may take out the daily Difference of Longitude, from Computers Number II. or III. without taking Departure into the Reckoning, except you please: Which Departure may be readily taken out from the proper Difference of Latitude and Course given.

Of correcting the SHIP'S COURSE steered by the COMPASS.

CORRECTION of the Ship's Course by the Magnetical VARIATION.

THE Difference between the Sun's magnetical Amplitude, taken by the Compass, at his Rising (or Setting) and his apparent Amplitude at Rising (or Setting) by Tab. p. 204 to 208, is the magnetical Variation, E. or W. to be allowed for the same Way, or to the Right or Left of the Course steered by the Compass, for the Ship's true Course.

The magnetic Variation is likewise found by two magnetical Azimuths, taken at equal Altitudes of the Sun, on the same Day. This whole magnetical Arch being bisected, the Middle thereof will give you the true South or Meridian very nearly; which will shew the Difference, or Variation, E. or W. from the true Meridian, or South Point of the Compass: which is to the Right or Left thereof, respectively.

The magnetical Variation is otherwise determined by the Azimuth Compass, and a Plumb Line with some Star coming to the Meridian at a given Time of the Night, which will shew the Difference between the true and magnetical Meridian.

This Variation, at London, has altered as follows.

At	Year	Variation.	Now (in 1759) it proceeds westerly at London, about 1 Degree in 7 Years. In some Places it moves Eastward; in others it is without Motion. The Changes of Variation is also different in different Places. See Mr. Emerson's Table of Variation, Page 49. of his Navigation.
London in	1580	$11^{\circ} \frac{1}{4} E.$	
	1657	Northerly.	
	1723	$12^{\circ} W.$	
	1759	$15^{\circ} W.$	

There are Tables of magnetical Variation in *Philos. Transactions*, and elsewhere, not to be depended on; because there is continual Change of Variation in most Places.

FURTHER CORRECTION of the SHIP'S COURSE by allowing for LEE-WAY.

THE Ship falling off from the Course steered by the Compass, by several Causes and Impediments to continue that Course, especially the Impediment of Lee-Way, her Course by the Compass must therefore be rectified by allowing for Lee-Way, as follows.

Rule 1. Set the Ship's Wake by the Compass, and the Angle betwixt it and the Ship's Keel (or right Fore and Aft) is her Lee-Way, if there is no Current: for so much she falls off from her Course (steered) to the Leeward, or from the Wind.

Rule 2. Being within Sight of Land, the Angle taken betwixt the Ship's Keel (or right Fore and Aft) and that Point of Land always bearing on the same Point of the Compass, will be the Lee-way made by the Ship, or what she falls off from her Parallel, while she steers the same Course by the Compass.

REMARK. Different Ships make different Lee-Way, caused by the Difference of their Hulls in Building. Nor, as Ship-Builders allow, can any two Ships be built to fall alike for Swiftness: And Experience of Sailing in any one

Of correcting the SHIP'S COURSE steered by the COMPASS.

Ship will best determine her *Lee-Way* under certain Circumstances of *Wind, Sails set, Seas, and Weather*. A high Sea, or great Swell a-crofs the Course steered by the Compass, produce large *Lee-Way*; requiring sometimes much Skill and Experience to determine its Quantity: A Ship being out of *Trim*, or drawing more or less Water, affect her *Lee-Way*.

GENERAL RULES of LEE-WAY from the SAILS set, or furled.

1. A Ship upon a Wind with all Sails set, 1 Point *Lee-Way*.
2. Top-Sails reefed, 1 Point $\frac{1}{2}$.
3. Hard Wind, one Topsail in, 2 Points.
4. High Sea, hard Gales, both Topsails furled, 3 Points.
5. Only Mainsail and Mizzen set, 4 Points.
6. Only Mainsail out, 5 Points.
7. Only Mizzen set, 6 Points.
8. All Sails furled, 7 Points.

SOME Seamen make different Allowances, but these are the most common, when, in a dark Night, or great Sea, a Ship's Wake cannot be set. How fruitless then is *spheroidal Navigation*, when so great Difficulty, Doubts, and Difference frequently arise in adjusting the true Course of the Ship.

N. B. Hard Gales and high Seas setting with or against the Ship's Motion, varies the *Lee-Way*, more or less, of which Experience can only judge.

FURTHER CORRECTION of the Ship's COURSE, and also DISTANCE, by trying the SWIFTNESS and Setting of a CURRENT.

WHEN there is little Wind and a smooth Sea, a Ship's Current is tried by the known Methods in Practice, as follows.

You sink (out of a Boat, at a small Distance from the Ship, where the Current is) a large triangular Board, leaded on one Side, an Iron Pot, or a Kettle, below the Current, with a Line tied properly to either 60, 80, 100, or more Fathoms in Length, to be veered out into the Sea, at one End, to give the Sinker Depth enough below the Current; and then the End of the Line in Hand is fastened to the Boat's Stern, to bring her up, as if she were at Anchor; and keep her as steady as possible upon the Top of the Current.

NOW, heaving your Log, and turning your Half-Minute-Glass, you set the Drift of the Log, upon the Current, by the Compass, when the Length of the Log-Line run off the Reel, while the Half-Minute-Glass runs out, will give you the Swiftmess of the Current, or Number of Miles it runs per Hour; if you add about $\frac{1}{4}$, or more or less, for the Drift of the Boat with the Current, according to the Depth, Weight, or Capacity of the Sinker to keep the Boat more or less steady; which must be judged. And the Direction of the Line being set by the Compass, will shew you to what Point thereof the Current sets.

OR, the leaded Board, aforesaid, being hove into the Ship's Wake, as she sails, and suffered to sink below the Depth of the Current, the Line fastened thereto will shew you the Current's Direction. And if the Half-Minute Glass be turned, when sufficient Length of Line is veered out for sinking the Board below the Current, then, as many Knots as the more runs off, from a Mark in the loose-Line last come in Sight, while the Glass runs out, (measuring the Plus-line afterwards) so many Miles the Current and Ship together move in an Hour. With which Motion, if you compare the Ship's Motion, measured by the Log, upon the Top of the Current, you will have the Velocity of the Current, separately, as it is with, or thwarts, the Ship's Motion. But having the Direction and Velocity of both, the other Direction and Velocity is not wanted. This Way you may some Times lose your Board or Sinker, and Part of your Line.

The Quantity of a Current's Motion is determined by keeping the Reckoning of a Ship outward and homeward bound, as exact as possible; the Difference of which two Reckonings being the Current's Motion: supposing it constant.

In a furrowed Sea, a Current's Direction is a-crofs the Ridges of the Waves raised by the Wind.

REMARK. In several Straits there is a Current above, and another below. In the Straits of Gibraltar the Atlantic Ocean continually runs into the Mediterranean, while an under Current, from the Mediterranean, runs the contrary Way into the Atlantic.

There is a Current in the Baltic Sound, 4 or 5 Fathoms deep, and another under it.

A Current from the Euxine Sea sets through the Straits of Constantinople and Propontis.

And in all other Straits, where there are rapid upper Currents, it is probable there are likewise under Currents setting the contrary Way.

PROPOSITION. If a Ship be considered at Rest in the Sea, while that Sea has a Current setting towards some one Point of the Compass, it is the same Thing in Effect of Computation, as if there was no Current, while the Ship sailed with the Current's Motion towards that Point.

HENCE, in correcting your Course steered by the Compass, you may consider your Ship, sailing upon any Course, as if she sailed upon a new or altered Course, the same as is that of the Current's Motion, instead of considering that Motion.

And, the Distance on your Course, steered by the Compass, is corrected by the Current's Motion in the Ship's Direction (computed from its Motion per Hour in the Angle with the Ship's Direction) by adding that Motion to the Distance sailed, or subtracting it therefrom, as it sets towards, or contrary to, the Ship's Direction.

OF MEASURING the SHIP'S WAY.

Of the *Seaman's* COMPASS, LOG-LINE, LOG, HALF-MINUTE GLASS, and LOG-BOARD.

THE common *Mariner's* Compass is well known, and needs no Description. But the *Mariner's* Compass improved, and recommended for its greatest Advantage to Navigation (by its *Inventor*, Mr. *Emerson*) is as follows.

The Needle is fixed to a moveable plain Card, under the *Compass* Card, which going stiff and being set, as a *Rectifier*, to the present *Variation*, in the *Compass*-Card above it, the *Fleur de Luce* of that Card will always point to the true North and South Points of the Horizon, and likewise each Point of this Compass to the true Course steered by the Ship. By which Means frequent Trouble and Mistakes are prevented in correcting the several Courses steered by the Compass, to the true Course.

The *Log-line*, *Log*, and *Half-Minute Glass*, are well known, by *Seamen*, for their great Utility in steering the Ship, and measuring her Way through the Ocean.

When the *Log* is heaved, as many *Knots* and *Tenths* of Line as runs off the *Reel*, while the *Half-Minute Glass* runs out, so many Miles and Tenths of a Mile the Ship runs in an Hour.

The *Half-Minute Glasses* may be proved by a Thread, and a leaden Ball at the End thereof, hung up $39\frac{2}{8}$ Inches long, from the Point of Suspension of the Thread, to the Center of the Ball. Which Ball being put in Motion, will vibrate exactly in a Second of Time; and if the *Glass* runs out while 30 Swings or Vibrations are made, past the Perpendicular, then that *Glass* is true, otherwise not. Another Proof of *Half-Minute Glasses*, is by one already proved.

THE *Knots* of the *Log-line* are judged best to be 45 or 46 Feet asunder, betwixt Knot and Knot; though 50 Feet asunder agree nearest with the 120th Part (the Number of Half-Minutes in an Hour) of the 60th Part of a Degree of a great Circle of the Earth, or a Mile, by which the Knot-Divisions are computed. But the *Log* dragging somewhat after the Ship, measures her Distance this Way less than it is; and therefore, the less Divisions of 45 or 56 Feet, between Knot and Knot, are found to agree best with Practice. Some Navigators, in our Time of using the Sea, divided their *Log-line* by 42 Feet only between the Knots; which yet, by a due Correction of the *Dead-Reckoning*, by frequent Observations, have some Times very nearly agreed with making the Land, after a long Voyage. And, in the *Sea-Phrase*, it is always better for the *Reckoning* to be a-head of the Ship, than for the Ship to be a-head of the *Reckoning*.

But with Respect to measuring a Ship's Distance by the *Log*, if her *Reckoning* was kept by *Degrees and Hundredth Parts*, instead of by *Degrees and Sixtieth Parts*, as at present, there would be no Trouble of Reduction (as Mr. *Emerson* observes) of Degrees to Minutes, and of Minutes to Degrees. And the Knots of such a *Log-line*, measuring Hundredths of a Degree in Half-Minutes, would be 27 Feet asunder. In which Case of keeping a Ship's *Reckoning*, multiplying these *Cents* above Degrees by 6, the Product of the 1st Figure with what is carried from the 2d, would give the Miles above Degrees mentally. And cutting off 2 Figures to the Right, from any Number of *Cents*, would give the Degrees and *Cents* over, at once.

Some Masters of Ships, and other experienced *Seamen*, estimate the Ship's Distance sailed in an Hour, or other Interval, by throwing a Chip over Board, by the Motion of which along the Ship's Side they judge of the Distance. And a Ship's Way, in the Day-Time, may be nearly estimated by observing the Motion of the Bubbles, or Sea-Foam, passing by the Ship's Side from Stem to Stern.

But, for more Exactness and Certainty, in measuring the Ship's Way over the Ocean, the *Log* is generally hove every two Hours (and may be heaved every Hour if Navigators please) when the Distance, that Way measured, and Course steered by the Compass, are carefully chalked into the *Log-Board*, contrived with Partitions for that Purpose, as well as for entering all Remarks and Occurrences, that happen in 24 Hours. At the End of each of which Intervals, the Ship's *Reckoning* for the Day is made up: but not till after an Observation of the Sun, at Noon, is taken (if possible) by which the *Dead-Reckoning* (as it is called) is daily and duly corrected; or as often as an Observation of the Sun, at Noon, can be had.

MAY the 15th, 1760.


The LOG-BOARD.

Hrs.	Courses.	Knots.	Winds.	REMARKS and OCCURRENCES.
2	S. W.	6.5	N.W.b.N	Variation } from S. to E. } Left 2 Points. } or N. to W. } Hand.
4		7.6	NW.b.W	Squally with Rain.
6	S. S. E.	8.1	W. b. S.	Hard Gales.
8		7.0		Came up with the Triton.
10	N.W. $\frac{1}{2}$ W.	6.0	S. W. b. S.	Joined the Centurion, Anson, and De-
12		6.7		fiance.
2		6.5	W.S.W.	Lee-Way, } Took a Reef in our Top- 2 Points. } sails. [in Company.
4	S. W. b. S.	6.3	W. b. N.	2. The Thunder and Lightning Bombs Saw a Sail to Windward.
6		5.5		2. A Current sets W. b. N. 4 Hours, at $2\frac{1}{2}$ Miles an Hour.
8	S. E. b. E.	6.1	N. W. b. N.	0. Got ready for engaging. [Nymph.
10	S. W. b. W.	5.1		1. Lee-W. At Noon spoke with the Sea-
12	S. W.	4.3	W. N. W.	

Former 24 Hours, May 14, at Noon. } Latitude { $40^{\circ} 19' N.$ } By Observation
Longitude { $21^{\circ} 35' W.$ } corrected.

Last 24 Hours, May 15, 1759, at Noon. } Latitude { $39^{\circ} 2' N.$ } By Dead-Rec-
Longitude { $21^{\circ} 3' W.$ } koning.

The SEAMAN'S COMPASS.



Of correcting a SHIP'S DAILY RECKONING.

N. B. The Mean of the Knots set down every two Hours in the Log, being taken for the intermediate Hours, is a *Refinement of no Use in Practice*: since if it were, the Log might be heaved every Hour. And till a *Method* can be found of measuring the Distance and of steering the Ship's Course, with less *Doubt* and *Error*, heaving the Log every 2 Hours is sufficient. For, if a Ship's Way through the Water could be measured with more *Exactness*, yet as there will always be *Uncertainty in the Course* made good, especially in *high Seas, cross Winds, strong Tides, Currents, &c.* one general *Method* for correcting all Ship's Reckonings, (*except by observing the Longitude*) must suffice as follows.

RULES for CORRECTING a SHIP'S RECKONING.

A Ship's Reckoning is corrected by an Observation of the *Sun*, at *Noon*, as often as it can be taken; by which, finding your Latitude by Observation, you discover the *Truth* or *Error* of your Latitude by *Dead-Reckoning*. If these two agree, then your *Departure* or *Longitude* are supposed to be right, and require no Correction. But when they differ, you proportion the *Correction* of *Departure* and *Longitude*, by the *Error* of *Latitude*, found in your *Reckoning*.

FIRST, you may examine if you have rightly allowed for *Variation* and *Lee Way*, and made due Correction of your *Courses*, and especially by allowing for *Currents*, and if after rectifying any Mistakes you have committed, and again going over the *Operations*, your Latitude by Reckoning and Observation will not agree, you may suspect a *Current* to be the Cause of the Difference; as not knowing its Quantity of *Motion*. But, if you know its *Direction*, or *Course*, use the following

RULE I. For CORRECTION of DEPARTURE. — As Radius : to Tan. Angle of the Current's Direction with the Meridian :: so Error in Latitude : to Correction in Departure.

If the *Latitudes* will not yet agree, and your *Course* failed be *nearer* the *Meridian* than to the *East* or *West*, the *Error* is likely to be in the *Distance*. Here find the *Difference* of the *Latitude* and *Departure* since the last *Observation*, and say by

RULE II. For CORRECTION of DEPARTURE. — As *Diff. Lat: : to Error in Latitude :: so Departure : to Correction in Departure.* To be added or subtracted, according as the Ship is before or behind the Reckoning.

But if your *Course* is nearer to the East or West Points than the Meridian, the *Error* is likely to be in the *Course*. Here take the Difference of *Latitude* and *Departure* since the last Observation, and say by

RULE III: For CORRECTION of DEPARTURE. — As *Departure* : to *Error in Latitude* :: *so Difference of Latitude* : to *Correction in DEPARTURE*.

☞ If the *Course*, in this *Case*, be very near the *East* or *West*, this *Correction* then, so very small, will be *needless*.

WHEN you steer about the *Middle-Way*, betwixt the *East*, or *West*, and the *Meridian*, you may correct either *Course* or *Distance*, wherein you conceive the *Error* mostly lies: but, if you are not convinced of one being more erroneous than the other, you may divide the *Error* betwixt both and so correct each.

RULE IV. For CORRECTION of the LONGITUDE. — As *Error in Latitude : to meridional Difference of Latitude between the computed and observed Latitude :: so Correction in Departure : to Correction in Longitude.*
To be added or subtracted, according as the Ship is a-head of the Reckoning, or Reckoning a-head of the Ship.

ANOTHER RULE. To know if the ERROR of the *Ship's Reckoning* be in the COURSE or DISTANCE?

In sailing betwixt the Meridian and E. or W. } If the { *Dif. Latitude* } be the *greater*, the Error is probably in the { *Distance* }
 { *Departure* } { *Course* }

N. B. When the Difference of Latitude is greater than the Departure, and the Error of *Reckoning* be in the *Distance*, it may be thus *corrected*.

As *Dif. Lat. by Reckoning* : *true Dif. of Lat.* :: *Departure by Reckoning* : *to true Departure*
 (By similar Δ s included in one another) :: *Distance by Reckoning* : *to true Distance.*

Then, As *Corrected Dif. Lat.* : *corrected Departure* :: *Radius* : *Tangent true Course.*

☞ When the *Ship* has out-run the *Reckoning*, it is too little, and you must *add* the *Difference*; but the *Reckoning* having out-run the *Ship*, it is too much, and you must *subtract* the *Difference*.

CASE I.

Lat.	Ship's Mot.	Lat.	Ship's Mot.	} If the observed Lat. be	{ greater less }	} than Lat. by Reckoning.	{ Ship out run the Reckoning. Reckoning out-run the Ship. }	} + -
N.	towards N.	S.	towards S.					

CASE II.

Lat.	Ship's Mot.	Lat.	Ship's Mot.	} If the observed Lat. be	{ greater less }	} than Lat. by Reckoning.	{ Reckoning out-run the Ship. Ship out-run the Reckoning. }	} - +
N.	towards S.	S.	towards N.					

IF the Error in the Distance proceed from the Log, or the incorrect Way of measuring it, as in general the Ship's Distance is greater than is measured, then you may take 1 Mile in 10 ^{lefs}, if a great Sea follows the Ship; but less, when a less Sea follows. And when the Sea's Motion is contrary to or athwart the Ship's Motion, her Way is less than the Distance measured by the Log. And the Time of Tides and their Setting will alter the true Measure of the Ship's Way, which must be observed.

REMARK on the foregoing METHODS of CORRECTING a SHIP'S RECKONING.

ALL the METHODS of Correction here given, are but precarious, while there are no other to depend on: Except what we have shown by observing the Longitude, as near as is practicable.

The Agreement of several Ships' Reckonings, proceeding together in the same Voyage, is an Argument of their Truth; as their Difference may serve to correct one another.

The Reckoning (in Longitude and Latitude) of a Ship you meet at Sea, will sometimes be of Use to correct your own.

Here you suppose an Error in the Course or Distance only, while it may be in both. But if one Error ballances the other, so as to give the computed Latitude near that by Observation, then no Error in either Course or Distance will appear; and yet an Error may be in the Longitude, though you are not able to distinguish it. And if the Error in both Course and Distance lie one Way, so as to make great Difference in the Latitude, there may be but little Error in Longitude; which yet must be corrected in Proportion to the Error of Latitude, by the preceding Rules.

REMARKS on B. MARTIN'S NEW NAVIGATION, in a Second LETTER from a Correspondent, dated June the 16th, 1759.

TO the AUTHOR of the ROYAL ASTRONOMER and NAVIGATOR.

SIR,

I perceive, by the Sheets you sent me, that you have taken more Pains with B. Martin's Book than I think was worth the Time bestowed; for I am persuaded that Nobody will ever follow his Steps (or Method) by Sea, whatever they do by Land.

Could one measure the Sea as exactly as a Man can measure the Land, and could he take the Angle of the Course as near as one can take an Angle on the Ground, with an Instrument, this Refinement, of spheroidal Navigation, might be of some Consequence. But since neither the Course nor Distance can be truly known to a 20th Part, and often more, it must be a very idle Thing to introduce difficult Calculations to correct an Error, which is no Way sensible when found out. For what such a trifling Error as this amounts to, in a Day's Run, is swallowed up in the other grand Errors, which are unavoidable. And this Error, as well as those greater ones, are all corrected as soon as an Observation can be made.

The only Question that remains to be answered is, Whether this trifling and insignificant Error is worth all that laborious Calculation bestowed on it by B. M. in his Folio Book? I think not; and if he, or Murdoch, thinks otherwise, I have Nothing to do with that, or they with me. And also (which is an unanswerable Argument against it) he cannot tell what Species of a Spheroid the Earth is; the Parts of this Book are so contradicting, as to make the Eccentricity in one Page more than double that in another; which is therefore a hopeful Theory for a NAVIGATOR to depend on.

But is not the Motive of introducing (or rather lugging in) this Theory by the Head and Shoulders, Vanity more than Use, that the Proprietor might not miss the Opportunity of letting the Right Honourable the Lords of the Admiralty know how well he is acquainted with the Figure of the Earth? Who seems so fond of the Invention, and it so tickles his Fancy, that he would apply it to every Thing, whether it suits or not.

But, when he has contrived his Compass, which is to be 100 Times more exact than the common Sort, and his New Nautical Dromometer, by which he can get the Course to a Minute, and the Distance to the utmost Exactness, it will be Time enough then to take in the minuter Inequalities, such as proceed from the spheroidal Figure of the Earth; and get an Inch or two more that Way. So that it is to be hoped, from all these Improvements, there will be no need, afterwards, to correct either the Latitude or Longitude.

I shall only further take Notice, that, in the Preface to his Second Part, he says "he has added the Mechanic Principles of Navigation, as he has not seen them in any other Treatise of this Kind," which is a little ambiguous; for he has, or might have, seen them in EMERSON'S NAVIGATION; except he was resolved to see Nothing but what he did not like. But he is such a modest Proprietor of original Invention and Improvement, that he is not ashamed of any Thing he says, nor from whence he borrows his select Materials.

NEWTONIENSIS.

DIMENSIONS

DIMENSIONS and WEIGHT of SHOT.

TO determine the Weight of an Iron-Ball from its Diameter being given, and the Diameter from the Weight?

FIRST, As 3.1416 : to 1 :: so is the Measure round any Ball : to its Diameter.

NOW, As 100 : to the Cube of an Iron Ball's Diameter in Inches :: so is 14 to the Weight of that Ball in Pounds.

EXAMPLE. If a Ball measures 14 Inches round, required its Weight?

3.1416) 14.0000 (4.456 Inches,
Cubed = 88.478

Multiplied by 14 = 1238.692 lb. oz.

Divided by 100 = 12.387 = 12 6,192 Wt.

[reqd.]

If the Weight of a Ball be 42 Pounds, required its Diameter?

42 by 100 = 4200

Divided by 14 = 300 Its Log. = 2.4771213

3.1416 $\times$ 6.6948 = 20.177 } One Third = .8257071

Inches round the Ball. } Its No. 6.6948 In-
ches Diam. required.

The Method of directing a Ship through Sands, into a Harbour, or River's Mouth, by Sea and Land Marks, (such as Buoys, Light Houses, Steeples, Windmills, or other remarkable Objects, contrived or fixed on Purpose) is known to every experienced Navigator. But those who desire Satisfaction herein may consult Mr. EMERSON'S NAVIGATION, P. 62 and 63. See also his Mechanics (comprehending all other Books on the Subject) for computing the Force of a Ship's Tackle, which he has partly exemplified in his Navigation, p. 63. making the Force of the Hand to the Weight as 1 to 4. in a particular Instance. To which distinguished AUTHOR, and his WORKS, we have sometimes been beholden for Judgment and Improvement in our Subject: Who comprises all his Subjects of Science, in the fewest Words and shortest Method.

* * * THOSE who chuse to sail as near as they can by the Arch of a great Circle, or to use Great-Circle-Sailing, as preferable to spheroidal Sailing, (suppose from the Capes of Virginia to the Cape of Good Hope, or from the Cape of Good Hope to Cape Horn) may compute the Angles of Position, Length of the Arch betwixt two Places, and Latitudes of the Arch, to every 5 or 10 Degrees of Longitude, by Spheric Trigonometry; and then the Course and Distance, from one Latitude to the other on that Arch, by Mercator Sailing; placing the several constant Courses and Distances thereon, in a Table for putting in Practice, as near as possible. For the Answers to all the Cases of right and oblique angled spherical Triangles, see further on; being the Principles and Grounds of Practical Astronomy in the foregoing Part of this Work. To which Cases are added numerous Affections and Properties of Triangles, in general.

PLACE of TURNING to WINDWARD, and Distance of Head-LAND.

TO compute the PLACE of a Ship's Turning to Windward so as to reach her intended Port, or Object, by once Tacking?

WHEN a Ship or Vessel is close hauled, she lies within about $5\frac{1}{2}$ Points of the Wind, or $61^{\circ} 52'$, on either Side of it.

LET D be the Distance of the Port, or Object, at the Place sailed from;

A $^{\circ}$ the $\angle$ of the Course sailed, with the Port or Object, on the same Side of the Wind;

Then, $61^{\circ} 52' - A^{\circ} = \angle$ of the Wind and Port, at the Place sailed from, on the same Side of the Wind
[with the Course;

And $2 \times 61^{\circ} 52' - A^{\circ} = \angle$ of the Course and Port on the contrary Tack, or Side of the Wind = $\angle$ at
[the Port or Object, between the Place sailed from, and that of turning to Windward, or Tacking;

Also, $180^{\circ} - 2 \times 61^{\circ} 52' = 56^{\circ} 16' = \angle$ between the Place sailed from and intended Port, at the Place
[of Tacking.

Therefore, As Sine $56^{\circ} 16'$: to D, the Distance of the Port from the Place of Departure, :: so Sine $123^{\circ} 44'$ —
[A $^{\circ}$: to Distance between Place sailed from, and that of Tacking, or Turning to Windward, required.

The Place of Turning to Windward in a Current may be likewise determined trigonometrically.

To compute the Distance of a HEAD-LAND, or CAPE, at SEA?

SET the Angle of the Head-Land, and your Ship's Way, by the Compass, and after Sailing for some Time on that Course, set the Angle of the Head Land and your Ship's Way a second Time: Then, Say by PROPORTION, from the Tables of natural Sines, or by Logarithms.

As the Sine of the Remainder after the Sum of those Angles is taken from 180 Degrees : to Distance run between taking the first and second Angle, :: so is Sine of either Angle taken : to the Ship's Distance from the Head-Land, when the other Angle, with the Ship's Way, was taken.

The Connexion in this Page, and another or two, is as the Copy could come in.

TO measure the TONNAGE, or BURTHEN of a SHIP.

THE Rules laid down, by several Authors, are as follow.

RULE I. Divide the Product of her Breadth, half Breadth, and Length, in Feet, by 94, and the Quotient will give the Tuns of her Burthen.

RULE II. Divide the Product of her Length, Breadth, and Depth, in Feet, by 100 for Ships of War, or 95 for Merchant-Men, that allow Nothing for Guns, and the Quotient will give the Burthen in Tuns.

RULE III. Subtract two Thirds of her Breadth from her Length (measured from the Stern-Post to the upper Part of her Stem) and the Product of that Remainder, the Breadth, and Half-Breadth, in Feet, being divided by 100, or 94, will give the *King's*, or *Merchant's* Tonnage, respectively.

RULE IV. A Ship's Tonnage, or Burthen, is equal to Half the Weight of Water she will hold.

N. B. *There is such a Variety in the Forms of Ships, that no one Rule can measure their Burthens exactly; but the above Rules are the most general in Practice.*

POSITION of the SAILS and RUDDER, in Working a SHIP.

Observation 1. Placing the Sails according to different Positions in Respect of the Wind and Ship's Way, have more or less Force to draw the Ship along, or turn her about.

2. The Head-sails keep the Ship steady and serve to make her ware.
3. Main-sails move the Center of Gravity of the Ship.
4. Mizzen-sails serve to keep her from sheering backward and forward, and to force her Stern to Leeward.
5. Sails have more or less Force the tighter or slacker they are hoisted: a bagging Sail drawing less than one that is well hoisted.
6. The higher a Sail is hoisted the more Wind it receives to draw the Ship along, or turn her about.
7. When a Ship goes directly before the Wind, any one Sail placed at right Angles with her Way draws with the greatest Force. But since one Sail hinders the Wind from acting against another, being all placed in that Position, if they all receive the Wind at an Angle of about 60 Degrees they will draw together with the greatest Force.
8. Wet Sails draw more than dry Ones.
9. The nearer to, or further from, a right Angle any Ship's Sail is placed with the Way of her Course, the less or more Lee-Way she makes.
10. The nearer the Wind is with, or further from, the Ship's Way, the less or more Lee-Way she makes.
11. By the Principles of Mechanics, the Angle between the Wind and Ship's Sail must be about twice as much as the Angle between the Sail and the Ship's Way, to sail near the Wind with the greatest Motion.
12. If the Wind is almost upon the Beam, the Angle between the Wind and the Sail must be about once and half the Angle between the Sail and Ship's Way.
13. If a Ship sails almost before the Wind, the Angle between the Wind and Ship's Sail should be about equal to the Angle between the Sail and her Way. All these Positions of the Sails are necessary to give the Ship the greatest Motion possible.
14. The nearest a Ship can lie to the Wind being about $5\frac{1}{2}$ Points or 60 Degrees, the Angle between the Wind and Sail should be about 38° , and the Angle between the Sail and the Ship's Way (or Keel) 22° , or nearly in that Ratio, to make her go fastest to Windward.
15. If the Wind is nearly upon the Beam, the sharper the Head-sails are set they have the more Power to turn the Ship's Head about.
16. If the Wind's Way be nearly the same as the Ship's Way, the Sail ought to make an Angle with the Ship's Way of 54° , to turn her soonest about.
17. The most powerful Position of the Rudder to turn the Ship about has been computed, from Principles of Mechanics and Fluxions, at an Angle with the Keel of $54\frac{1}{2}$ Degrees.
18. The faster a Ship sails the better she answers her Helm.
19. If she sails slow she will scarcely steer. And the more she heels the less she answers her Helm.

See Mr. Emerson's Navigation, for further Satisfaction.

The PROPERTIES of MAGNETIC VARIATION.

- Property 1. Variation in *Quality* is either E. or W. Variation of the *magnetic* from the *true* North Point.
2. Variation in *Quantity* is the Number of Degrees (E. or W.) of the *magnetic* from the *true* North Point.
3. When } the *magnetic* { N. } varies fr. the *true* { N. } towards the { E. } Both to the *Right* of the same *tr.* Point.
Then } { S. } { S. } { W. }
4. When } the *magnetic* { N. } varies fr. the *true* { N. } towards the { W. } Both to the *Left* of the same *true* Point.
Then } { S. } { S. } { E. }
5. All the *magnetic* Points are carried as far as to the { Right } of the same *true* Point as is the { E. } Variation.
Left { W. }

Therefore,

6. The *true* Point, *Azimuth*, or *Amplitude*, (to which the corresponding *magnetic* Point is always directed) being reckoned on the same *magnetic* Compass from N. or S. to the E. or W. lies the same Way with the Variation, to the Right } of the *magnetic* Point, *Azimuth*, or *Amplitude*, as much as the Variation lies to the { Left } or { E. } of the *tr.* N.
Left { W. }

These Properties are evident by a RECTIFIER, or a *moveable* Compass on a *fixed* one. Hence the following

RULES for determining the *Quality* and *Quantity* of magnetic VARIATION.

Rule 1. The Situation of the *true* Azim. or *Amplitude* { Right } of the *magnetic* Azim. or { E. } Variation.
(reckoned on the same *magnetic* Compass) being to the { Left } Amp. shews { W. }

2. The *Distance*, or Difference between the *true* and *magnetic* Azimuth or *Amplitude* is the *Quantity* of Variation,

i. e. The *true* and *magnetic* *Amplitude* { both N. or S. } their { Dif. } in Degrees is the *Quantity* of Variation.
{ one N. other S. } { Sum }

Or, if you always reckon the *true* and *magnetic* *Amplitudes* from their respective North Points, and then count them on the same *magnetic* Compass,

3. At Sun-Rising } True { Amplitude being the greater, the Var. is { E. } But { W. } At Sun-Setting.
Magnetic { { E. }

Or, if you always reckon the *true* and *magnetic* *Azimuths* from their respective N. Points, and then count them on the same *magnetic* Compass,

4. In the Forenoon } True { Azimuth being the greater, the Var. is { E. } But { W. } In the Afternoon,
Magnetic { { E. }

N. B. The first Rule is general, and answers all Cases that can be proposed.

Example 1. At Sun-Rising | Mag. Amp. | True Amp. | True to the Left of magnetic Amplitude.
By second Rule Dif. 14° . . . Variation W. by 1st Rule.
Or 75° N. E. . . . 61° N. E. Magn. Amp. greater, theref. Dif. 14° W. Variat. by 3. Rule.

Example 2. At Sun-Setting | 8° W. S. | 14° W. N. | True to the Right of magnetic Amplitude.
By second Rule Sum 22° . . . Variation E. by 1st Rule.
Or, 98° N. E. . . . 76° . . . Magn. Ampl. greater, theref. Dif. 22° E. Var. by 3. Rule.

Example 3. In the Forenoon. | Mag. Azim. | Tr. Azim. | True to the Left of magnetic Azimuth.
By second Rule Sum 21° . . . Variation W. by 1st Rule.
Or, 100° N. E. . . . 79° N. E. Mag. Azim. greater, theref. Dif. 21° W. Var. by 4. Rule.

Example 4. In the Afternoon. | 30° 40' W. S. | 26° 14' W. S. | True to the Right of the magnetic Azimuth.
By second Rule Dif. 4 26 . . . Variation E. by 1st Rule.
Or, 120° 40' N. W. 116° 14' N. W. . . . Magnetic Azimuth greater, therefore Dif. 4° 26' E.
[Variation, by 4. Rule.]

PRACTICAL RULES for finding the LATITUDE of the PLACE of OBSERVATION.

The Meridian ALTITUDE and Declination of the SUN or STAR, given.

CASE I. Where the Object (SUN or STAR) rises and sets in a diurnal Revolution.

Rule 1. The Difference. Meridian Altitude and Declination both N. or both S.

Zenith-Distance } The Latitude { same } Name of Declination { greater } than the Zenith
Declination } { contrary } { less } Distance.

Rule 2. The Sum. Meridian Altitude and Declination one N. the other S.

Zenith-Distance } The Latitude { same Name of Declination:
Declination }

N. B. Meridian Altitude, or Zenith-Distance S. or N. means to the S. or N. of the Observer. And Zenith-Distance is the Complement of the Meridian Altitude.

Example 1. On the 16th of May, 1760, at Sea.

To Sun's Zenith Distance $26^{\circ} 44' S.$
Add Sun's Declination $19^{\circ} 16' N.$

Sum (by Rule 2) the Lat. of Place of Observ. $46^{\circ} 0' N.$

Example 2. On the 12th of June, 1761, at Sea.

From Sun's Declination $23^{\circ} 12' N.$
Subtract Sun's Zenith Distance $8^{\circ} 35' N.$

Diff. (by Rule 1) Lat. of the Place $14^{\circ} 37' N.$

CASE II. Where the OBJECT (SUN or STAR) neither rises or sets in a diurnal Revolution; but comes to the MERIDIAN, above the Horizon, on the contrary Sides of the elevated Pole, twice in 24 Hours.

Rule 3. The Sum. The SUN or STAR on the Meridian below the POLE.

Meridian Altitude } The Latitude { of the same Name with the Declination.
Comp. of Declinat. }

Rule 4. The Difference. The STAR on the Meridian above the POLE.

Meridian Altitude } The Latitude { of the same Name with the Declination.
Comp. of Declinat. }

Example 1. On the 12th of June, 1764, near Spitsbergen, Greenland.

To Sun's meridian Altitude below the Pole . . . $13^{\circ} 9' N.$
Add Complement of Sun's Declination $66^{\circ} 47' N.$

Sum (by Rule 3) the Place's Latitude $79^{\circ} 56' N.$

Example 2. On the 10th of July, 1761, at Sea.

To Sun's meridian Altitude below the Pole . . . $6^{\circ} 12' N.$
Add Comp. of Sun's Declination $64^{\circ} 47' N.$

Sum (by Rule 3) the Place's Latitude $70^{\circ} 59' N.$

See other Rules, in practical Astronomy, P. 190, 191.

Example 3. On the 28th of December, 1763, at Sea.

From Declination of Syrius $16^{\circ} 23' S.$
Subtract Zenith-Distance of * $10^{\circ} 9' S.$

Diff. (by Rule 1) Lat. of the Place $6^{\circ} 14' S.$

Example 4. On the 31st of August, 1764, at Sea.

To Zenith Distance of Fomalhaut, or brightest }
of the Southern Fishes $11^{\circ} 19' N.$
Add Star's Declination $31^{\circ} 1' S.$

Sum (by Rule 2) Lat. of the Place $42^{\circ} 20' S.$

REMARKS on II. CASE.

1. WHEN the Sun comes to the Meridian below, and a Star comes to the Meridian of the same Place, both above and below the same Pole, twice in 24 Hours, the Latitude of that Place is of the same Name with the Declination of the Sun or Star. And the Declination of the Sun or Star, in such a Circumstance must be equal to, or greater than the Complement of the Pole's Height. And therefore, as the Sun's greatest Declination is $23^{\circ} 28' \frac{1}{2}$, its Complement is $66^{\circ} 31' \frac{1}{2}$, which is the least Latitude of the same Name with the Sun's Declination, where the Sun can come to the Meridian, above the Horizon and below the same Poles twice in 24 Hours. And the least Latitude where a Star can come to the Meridian above the Horizon, and above and below the same Poles, twice in 24 Hours, is therefore the Complement of the Star's Declination of the same Name with the elevated Pole.

2. The Sun, or a Star, coming to the Meridian below the Pole, betwixt it and the Horizon, is then at its least Meridian Altitude, and coming to the Meridian on the contrary Side of the Pole, is then at its greatest.

3. Some Stars (whose Comp. of Declin. of the same Name with the elevated Pole, is less than the Comp. of the Lat.) come to the Meridian below the Pole, betwixt it and the Horizon, and above it, betwixt the Pole and Zenith (where the Sun never comes) at their greatest meridian Altitudes.

4. Every Star coming to the Meridian below and above the Pole, at its least and greatest Meridian Altitudes, respectively, is always to the Northward where the North Pole is elevated, or always to the Southward, where the South Pole is elevated.

5. Some Stars (whose Complement of Declination, of the same Name with the elevated Pole, is greater than the Complement of Latitude) come to the Meridian below the Pole and on the contrary Side thereof, beyond the Zenith betwixt the Pole and the Equinoctial, (where the Sun comes) at their greatest Meridian Altitudes.

For those Stars coming to the Meridian betwixt the Zenith and Equinoctial (coming also under the Pole, but never between the Pole and Zenith) the 1st and 2d Rules (in Case I) serve to find the Place's Latitude.

Geographical RULES for finding the tabular Difference of LONGITUDE.

To find the DIFFERENCE of LONGITUDE of two Places, according to the old Way of reckoning E. Longitude, from the Island of FERRO, round the GLOBE.

RULE I. Bound fr. less to greater Long. Sub. less from greater Bound fr. greater to less Long. Remainder } less } more } than 180° } Difference of Longitude . . . } E. } W. Take from 360° rem. Dif. Long. }

II. Sub. less from greater Remainder } less } more } than 180° } Difference of Longitude . . . } W. } E. Take from 360° rem. Dif. Long. }

To find the DIFFERENCE of LONGITUDE of two Places according to the new Way of Reckoning E. or W. Longitude from the OBSERVATORY or Metropolis of any NATION.

III. Both E. Longitude Sub. less from greater Remainder } Dif. Longitude } E. } W. bound to } greater } less } Longitude Both W. Longitude

IV. Sub. less from greater Remainder } Dif. Longitude } W. } E. bound to } greater } less } Longitude One E. other W. Long.

V. Sum of both } less } more } than 180° } Difference of Longitude } same } contrary } Name of Long. failed to. Take from 360° rem. Dif. Long. }

N. B. Mr. Atkinson, an Author in Vogue, has given Rules to find the Diff. of Lat. N. or S. from Latitudes given, both N. or S. or one N. and the other S. Also to find the other Lat. N. or S. from one Lat. N. or S. and Dif. of Lat. N. or S. given. But Reason finds these, with the least Attention. Whereas there is Trouble to the Attention, when the Mind is engaged, to find by the Table which Way the Longitudes lie, when wanted, without the above easy Rules, to prevent Mistake.

EXAMPLES of finding the Difference of Longitude by the OLD WAY of RECKONING the LONGITUDES E. from FERRO.

Longitudes
EXAM. I. From Havana 294° 40'
To Jamaica 300 6

Rule 1. Diff. Longitude 5 26 E.

EXAM. II. From London 18 10
To Martinico 319 45

301 35
360 00

Rule 1. Diff. Longitude 58 25 W.

Longitudes
EXAM. III. From Cape of good Hope 35° 35'
To London . . . 18 10

Rule 2. Diff. Longitude . . . 17 25 W.

EXAM. IV. From Cape Horn . . . 298° 15'
To Cape of good Hope 35 35

262 40
360 00

Rule 2. Diff. Longitude 97 20 E.

Longitudes
EXAM. V. From Jamaica 300° 6'
To Barbadoes 3 0 7

Rule 1. Diff. Longitude 20 1 E.

EXAM. VI. From London 18 10
To Lizard . . . 12 56

Rule 2. Diff. Longitude . . . 5 14 W.

EXAM. VII. From Constantinople 47 30
To Gibraltar 12 27

Rule 2. Diff. Long. . . 35 3 W.

The above Answers are according to Mr. Emerson's Navigation, and Tables of E. LONGITUDES reckoned from Ferro.

EXAMPLES of finding the Dif. of Long. by the NEW WAY of RECKONING the LONGITUDES E. or W. from Greenwich OBSERVATORY.

Longitudes
EXAM. I. From Havana 83° 0' W.
To Jamaica 76 37 W.

Rule 4. Dif. Long. 6 23 E.

EXAM. II. From London 0 5 W.
To Martinico 60 59 W.

Rule 4. Dif. Longitude 60 54 W.

Longitudes
EXAM. III. From Cape of Good Hope 19° 30' E.
To London . . . 0 5 W.

Rule 5. Dif. Longitude . . . 19 35 W.

EXAM. IV. From Cape Horn . . . 80 0 W.
To Cape of Good Hope . . . 19 30 E.

Rule 5. Dif. Longitude . . . 99 30 E.

Longitudes
EXAM. V. From Jamaica 76° 37' W.
To Barbadoes 58 55 W.

Rule 4. Dif. Longitude 17 42 E.

EXAM. VI. From London 0 5 W.
To Lizard . . . 5 19 W.

Rule 4. Dif. Longitude 5 14 W.

EX. VII. From Constantinople 28 51 E.
To Gibraltar . . . 4 58 W.

Rule 5. Dif. Longitude 33 49 W.

More EXAMPLES of finding the Difference of Longitude by the NEW WAY of RECKONING the Longitudes from Greenwich.

Longitude
EXAM. VIII. From Cape Farewell 46° 50' W.
To Admiralty Isle 59 45 E.

Rule 5. Dif. Longitude . . . 106 35 E.

Longitude
EXAM. IX. From Bantam 105° 5' E.
To Island Juan Fernandes 88 23 W.

Rule 5. Dif. Longitude 193 28 W.
166 32 E.

Longitude
EXAM. X. From Antigua 61° 50' W.
To Porto Bello 80 20 W.

Rule 4. Dif. Longitude 18 30 W.

REMARKS on the different BEGINNINGS of the terrestrial LONGITUDES.

1. IT has been observed, at p. 298, by Newtonianis, that the terrestrial Longitudes ought to be reckoned from Ferro, one of the Canary Isles, according to ancient Custom: and that for the same Reason that the celestial Longitudes are reckoned from the Beginning of Aries, according to ancient Custom. In which Computation of the terrestrial Longitudes, if all Nations agreed, the Confusion would be avoided now arising from the different Beginnings of Longitude among different Navigators in the same and different Nations.

2. The Latitude of each Place, on the Globe, begins at the Equinoctial on the Place's Meridian, and terminates on that Meridian at the Place whose Latitude is sought; by which, with Reference to the Equinoctial, the Quantity of each particular Latitude (of Places) is ascertained.

3. The Longitude of each Place, on the Globe, begins at some distant Meridian on the Equinoctial, where it is intersected by that Meridian, and terminates on the Equinoctial, where it is intersected by the Meridian, of the Place, whose Longitude is sought; by which, with Reference to some other Meridian, the Quantity of each particular Longitude (of Places) is ascertained.

4. Though reckoning the terrestrial Longitudes round the Globe easterly, from the Meridian of an Observatory, in any one Nation, would be equally easy, with reckoning from Ferro, for determining the Difference of Longitude betwixt any two Places; yet, by reckoning the Longitudes from Ferro only, according to original and ancient Custom established, or from some other Meridian universally agreed on, we should be able to compare the absolute Longitudes ascertained by different Nations, as we now compare the Latitudes, and so arrive at a still nearer and nearer Improvement in both. And each Nation would be able to make the same Comparison and Improvement for themselves from what is done by other Nations.

5. The Difference of Latitude of two Places is counted on the Meridian of one Place, betwixt the Parallels of both Places.

6. The Difference of Longitude of two Places is counted on the Equinoctial, betwixt the Meridians of both Places.

7. On whatever Meridian betwixt two Parallels of Latitude, the Difference of Latitude be reckoned, it is still the same in Quantity.

8. By whatever Method the Distance on the Equinoctial, or Difference of Longitude is reckoned betwixt two Meridians, it is still the same in Quantity, which is the Affection of Longitude chiefly sought, and what is chiefly wanted, from whatever Beginning Longitude is reckoned.

9. The celestial Longitudes are only wanted from the Intersections of the Ecliptic and Equinoctial, at Aries or Libra, to which ever they happen to be nearest, for the Purpose of computing the celestial Declinations in Astronomy, to which the terrestrial Latitudes in Geography are correspondent and similar.

A JOURNAL of a VOYAGE

IN THE

DREAD-NOUGHT MAN of WAR,

Captain CONDUCT, Commander.

From the *Lizard* { Latitude $49^{\circ} 55' N.$ } To the *Island of MARTI-* { Lat. $14^{\circ} 43' N.$
 { Longitude $5^{\circ} 19' W.$ } NICO in the *West-Indies* { Lon. $61^{\circ} 00' W.$ from *Greenwich* Observat.

Difference of LATITUDE $35^{\circ} 12' S$; Difference of LONGITUDE $55^{\circ} 41' W$.

The constant COURSE S. $52^{\circ} 23' \frac{1}{2} W.$ } DISTANCE on that Course, 3461 Miles.
 Or S. W. $7^{\circ} 23' \frac{1}{2} W.$ }

Began our Voyage on Thursday, the 8th of May, 1760.

Friday, May 9, 1760.

	Ho.	Courses steered.	Knots	Wind.	Courses corrected.	Points	Dist.
Afternoon.	2	W b. S	5.3	N	Sw.b.w $\frac{1}{2}$ W	$5 \frac{1}{2}$	5.3
	4	SW	5.0	NNW	SSW $\frac{1}{2}$ W	$2 \frac{1}{2}$	15.9
	6		5.4	N			
	8		5.5				
	10	SW b. S	5.7	NW b. W	Sb. W $\frac{1}{2}$ W	$1 \frac{1}{2}$	5.7
	12	SW	6.0	NNW	SSW $\frac{1}{2}$ W	$2 \frac{1}{2}$	28.0
After Midnight.	2		7.1	Nb. W			
	4		7.6				
	6		7.3	Nb. E			
	8	SW b. S	7.4		Sb. W $\frac{1}{2}$ W	$1 \frac{1}{2}$	7.4
	10	SW	6.5		SSW $\frac{1}{2}$ W	$2 \frac{1}{2}$	11.9
	12		5.4	NE			

On Board the DREADNOUGHT.

REMARKS and OCCURRENCES.

A Point $\frac{1}{2}$ Variation W.

Thursday, May 8, 1760, at Noon, the *Lizard*, by the Compass, bore NNE, Distance about 16 Miles. — Hence the Course steered from thence SSW, Distance 16 Miles. : Variation $16^{\circ} W$. Therefore, May 8, at Noon,

Tr. Course | Dist. | Dif. Lat. S. | Dep. W. | Lat. N. | Lon. W.
 S. $60 \frac{1}{2} W.$ | 16' | $15^{\circ} 89'$ | $1^{\circ} 74'$ | $49^{\circ} 39'$ | $5^{\circ} 22'$

At 6 lost Sight of the *Lizard*, bearing NE b. N $\frac{1}{2} N$.
 At 10 Scilly LIGHT-HOUSE bore N $\frac{1}{2} E$, Distance about 10 Leagues.
 Heard a Firing of Guns to the Westward.

Blows hard. Took a Reef in our Top-sails.

Fell in with a French Privateer, at 5 this Morning, *la Rochelle*, of 30 Guns, and took her.

The *Experiment*, *Defiance*, *Intrepid*, *Enterprise*, and *Success*, in Company.

Fair Weather.

Passed by the *China* Indiaman, bound for London.

Courses.	Pnts.	Dist.	N.	S.	E.	W.
Sw.b.w $\frac{1}{2}$ W	$5 \frac{1}{2}$	5.3	.	2'.50	...	4'.67
SSW $\frac{1}{2}$ W	$2 \frac{1}{2}$	55.8	.	49.21	...	26.30
S b. W $\frac{1}{2}$ W	$1 \frac{1}{2}$	13.1	.	13.39	...	4.06
Half { Sum			.	65.10	...	35.03
Dif			.			
Total, doubled			.	130.20		70.06

Twenty-four Hours RECKONING.

Course.	Dist.	Dif. Lat.		Departure.		SHIP'S PLACE.	
		N.	S.	E.	W.	Lat. N.	Lo. W.
S. $28^{\circ} 17' W.$	147'.85	...	$130^{\circ} 20'$	...	$70^{\circ} 06'$	$47^{\circ} 29'$	$7^{\circ} 8'$
Corrected by Observn		...	...	...	$-3^{\circ} 76'$	$+7'$	$-6'$
Reck'd Head of Ship		...	...	...	66.30	$47^{\circ} 36'$	$7^{\circ} 2'$
Corrected.		...	123.20	...			
Sun's Zenith Dist.	$30^{\circ} 21' N.$	Sun's Magnet. Ampl.		$12^{\circ} 17' E. N.$			
Sun's Declination	$17^{\circ} 34' N.$	Sun's True Ampl.		$28^{\circ} 17' E. N.$			
Lat. by Observation	$47^{\circ} 36' N.$	Needle's Variation		$16^{\circ} W.$			

This Day, at Noon, *Martinico* bore S. $53^{\circ} 53' W.$ Dist. 3347 Miles.

N. B. If your magnetical Variation is to the Right or Left of the true Point or Azimuth, (supposing yourself at the Center of the Compass) you allow for it in the true Course, the same Way, to the Right or Left of the Course steered by the Compass; for the Course made good by the Ship, when there is no Lee-Way. But when there is Lee-Way, you also allow for that in your Course corrected by Variation, as aforesaid, by taking (from the Wind) as many Points distant from your Course corrected by Variation, as is equal to the Lee-Way; for the Course then made good by the Ship.

HENCE also, if the Variation be to the Right or Left of the true Point of the Compass, by a certain Quantity, you must steer as much contrariwise, viz. to Left or Right, of your intended Course, to make that Course good by the Ship, independent of Lee-Way; or so much more to be allowed for towards the Wind, if the Distance of the Ship's Course from the Wind will admit.

10. The celestial Latitudes have no correspondent terrestrial Affection, or Property. And therefore the supposed Causes for the Difference in (Reckoning) the Beginnings of celestial and terrestrial Longitudes, by Astronomers and Navigators, respectively, in Use, are these two: 1. From the Equinoctial and Ecliptic Intersection, requiring that Beginning for Computation in Astronomical Purposes. 2. For the Purpose of Navigation; requiring the Beginning of Longitude at each Country's Metropolis, or Observatory, for adjusting the celestial Longitudes R. A. and Declinations correspondent thereto, to those under different Meridians, having the Diff. of Long. at once given, for the Purpose of Navigation. Whereas to find the Diff. of Long. had been a Problem, when the Long. is reckoned according to ancient Custom, and what would embarrass the Navigator; being attended with more Inconveniency to Astronomy than Reckoning the Longitude from an Observatory (to which the celestial Longitudes, Declination, and Right Ascensions are given) in any particular Nation. Reckoning Longitude from the *Lizard* is a Problem concerned only in a Ship's Reckoning, for comparing the Ship's Distance from her Port, and not concerned in her Longitude from *Greenwich*, which Geographical Longitude is wanted in all foreign Places, or Places on the Sea Coasts, remote from *Greenwich*, to compare with the geographical Longitude obtained from celestial Observation. Therefore the Longitudes from *Greenwich* in our Sea-Coast Table are given.

A JOURNAL of a VOYAGE from England to MARTINICO in the *West-Indies*.

Saturday, May 10, 1760.

On Board the DREADNOUGHT.

	Ho.	Courses steered.	Knots	Wind	Courses corrected.	Points	Dift.	REMARKS and OCCURRENCES.	
Afternoon.	2	S. W.	6.6	N. N. E.	SSW $\frac{1}{2}$ W	2 $\frac{1}{4}$	6'.6	A great Sea setting after us S. 1 $\frac{1}{2}$ Point Variation W, Saw a Sail to the N. E, Spoke with the <i>Mermaid</i> bound for <i>Leghorn</i> from <i>London</i> . Squally, with Rain. Rain. Spoke with the <i>Two-Sisters</i> bound for <i>Tarmouth</i> , from <i>Gibraltar</i> .	
	4	S. S. W.	6.2		S. $\frac{1}{2}$ W.	$\frac{1}{2}$	12.5		
	6		6.3						
	8	SW b. S	7.0	N. E. b. N.	Sb. W $\frac{1}{2}$ W	1 $\frac{1}{2}$	7.0		
	10	S. W.	7.1	N. E.	SSW $\frac{1}{2}$ W	2 $\frac{1}{2}$	21.4		
	12		7.3	N. b. E.					
After Midnight.	2		7.0	N N E				Squally. Gave Chace at 4 to a large Sail upon our <i>Starboard</i> Bow, proving to be the <i>Dragon</i> Man of War on a Cruise. A fresh Gale, and fair Weather. Spoke with the <i>Guardalupe</i> , bound for <i>London</i> , from <i>Antigua</i> . A Current setting S. E. 5 Hours, 2 Miles an Hour. Count for the Half Time only, where the Log. is doubled.	
	4	SW b. S	7.5	N E b. N	Sb. W $\frac{1}{2}$ W	1 $\frac{1}{2}$			
	6	SW	7.1	N N E	SSW $\frac{1}{2}$ W	2 $\frac{1}{2}$	26.2		
	8		7.0	N E					
	10		6.1						
	12		6.0	E N E					

							Twenty-four Hours RECKONING.							
Courses	Pnts	Dift.	N.	S.	E.	W.	Course	Dift.	Dif. Lat.		Departure		SHIP'S PLACE	
									N.	S.	E.	W.	Lat. N.	L. W.
SSW $\frac{1}{2}$ W	2 $\frac{1}{4}$	54'.2	. . .	47'.80	. . .	25'.55								
S $\frac{1}{2}$ W	$\frac{1}{2}$	12.5	. . .	12.44	. . .	1.23								
Sb. W $\frac{1}{2}$ W	1 $\frac{1}{2}$	7.0		6.70	. . .	2.03								
Current.														
ESE $\frac{1}{2}$ S	5 $\frac{1}{2}$	5.0		2.36	4.41	. . .	S. 19°. 24' W	146'.92	. . .	138'.60	. . .	48'.80	45° 17'	80 12'
2 $\frac{1}{2}$ Hours							Correction by Observ.		. . .	+15.00	. . .	+ 5.28	- 15	+ 8'
							Ship Head of Reckg.		. . .					
							Corrected.		. . .	153'.60	. . .	54'.08	45° 2'	80 20'
							⊙ Zenith Distance	27° 13' N		⊙ Magnet. Ampl.	100° 30' E. N.			
							⊙ Declination	17 49 N		⊙ True Ampl.	26 20 E. N.			
Half	{	Sum		69.30	4.41	28.81	Lat. by Observation	45 2 N		Needle's Variation	15 50 W.			
		Dif.				24.40								
Total, Doubled				138'.60		48'.80	This Day, at Noon, Martinico bore S. 55° 54' W. Distance 3044' Miles.							

Sunday, May 11, 1760.

On Board the DREADNOUGHT.

	Ho.	Courses steered.	Knots	Wind	Courses corrected.	Points	Dift.	REMARKS and OCCURRENCES.	
Afternoon.	2	S b. W	5.8	W.	Sb. E $\frac{1}{4}$ E	1 $\frac{1}{4}$	11'.4	1 $\frac{1}{4}$ Point Variation W. 1 Point Lee-Way. A great Sea following us, S. Spoke with the <i>Turkey-Merchant</i> , bound for <i>London</i> , from <i>Naples</i> . Cloudy. Lightning and Thunder from the S. W. Rain.	
	4		5.6	Wb. S					
	6	SSE	5.5	S. W.	SE $\frac{1}{4}$ E	4 $\frac{1}{4}$	11.4		
	8		5.9						
	10	NW b. W	6.6		NW b. W $\frac{1}{4}$ N	5 $\frac{1}{4}$	20.7		
	12		6.8						
After Midnight.	2		7.3					Rain. Passed by the <i>Jamaica</i> Frigate, bound for <i>London</i> from <i>Jamaica</i> . Squally. Spoke with the <i>Nightingale</i> from <i>Virginia</i> for <i>London</i> . Came up with the <i>London-Merchant</i> , an <i>English Privateer</i> . Fair Weather. A Current setting N. W. by W. 4 Hours, 3 Miles an Hour. Count for the Half Time only; the Log. being here doubled.	
	4	SSW	5.5	W	S $\frac{1}{4}$ E	$\frac{1}{4}$	10.6		
	6		5.1						
	8	ESE	4.8	Sb. W	E $\frac{1}{4}$ N	7 $\frac{3}{4}$	4.8		
	10	SSW	5.0	S. E.	SSW $\frac{1}{4}$ S	1 $\frac{3}{4}$	5.0		
	12	E b. S	5.3	Sb. E	E b. N $\frac{1}{4}$ N	6 $\frac{3}{4}$	5.3		

Twenty-four Hours RECKONING.							
Courses	Pnts	Dift.	N.	S.	E.	W.	
S b. E $\frac{1}{4}$ E	1 $\frac{1}{4}$	11'.4		11'.06	2'.77	. . .	
SE $\frac{1}{4}$ E	4 $\frac{1}{4}$	11.4	10'.64	7.65	8.45	. . .	
SW b. W $\frac{1}{4}$ W	5 $\frac{1}{4}$	20.7				17.75	
S $\frac{1}{4}$ E	$\frac{1}{2}$	10.6		10.58	0.52	. . .	
E $\frac{1}{4}$ N	7 $\frac{3}{4}$	4.8	0.23		4.79	. . .	
SSW $\frac{1}{4}$ S	1 $\frac{3}{4}$	5.0		4.70		1.68	
E b. N $\frac{1}{4}$ N	6 $\frac{3}{4}$	5.3	1.29		5.14	. . .	
W. N W	6	6.0	2.29			5.54	
Half } Dif.			14.45	33.99	21.67	24.97	
Sum				19.54		3.30	
Total, Doubled				39.08		6.60	
Course	Dift.	Dif. Lat.		Departure.		SHIP'S PLACE	
		N.	S.	E.	W.	Lat. N.	Lo. W.
S. 9°. 35' W	39'.64	. . .	39'.08	. . .	6.60	44° 23'	80 29'
Corrected by Observn		. . .	9.00	. . .	1.52	+ 9	- 2'
Reckg Head of Ship							
Corrected.		. . .	30'.08	. . .	5'.08	44 32	80 27'
⊙ Zenith Dist. 26° 27' N				⊙ Magnet. Ampl. 40° 47' W N			
⊙ Declination 18 5 N				⊙ True Ampl. 26 27 W N			
Lat. by Observation 44.32 N				Needle's Variation 14 20			
This Day, at Noon, Martinico bore S. 56° 22' W. Dist. 3229' Miles.							

Nearne's.

Nearness of SAILING to the WIND, and Resolving a TRAVERSE geometrically.

☞ If a Ship plies to Windward, and is close hauled, it is usual to allow 1 Point, from the Wind, for Lee-Way; when she lies within $5\frac{1}{2}$ Points of the Wind, viz. on either Side that Point of the Compass from whence the Wind blows; making a Course good $6\frac{1}{2}$ Points from the Wind. But when there is no Head Sea to make her fall off from her Course, the nearest she can lie to the Wind is reckoned about 5 Points, on either Side of it; then making her Course good about 6 Points from the Wind: (Currents, Hard Gales, and High Seas excepted) Otherwise, she may sail large (as 'tis called) with the Wind more or less abaft the Beam.

When the Wind blows directly aft, a Ship is then said to go right before the Wind, as to every experienced Seaman is known.

IN Imitation of the Common Form or Custom of Keeping a Ship's Journal, the Log is here heaved but every 2 Hours; though on Board most of our Men of War and East India Ships, it is generally heaved every Hour, for ascertaining the Distances run on the several Courses as near as possible. And therefore one Side of a Page in your Journal Book, being ruled for every Hour, instead of every two Hours, to contain one, instead of 2 Days Reckoning in each Page, your Ship's Place will be deduced more circumstantially, though perhaps not more exactly: Since its Place derived either from every Hour, or 2 Hours heaving the Log, must be duly corrected by Observation.

In flying to Windward, you may lay down your several Courses and Distances of the Traverse on Paper, by Scale and Compasses, and so determine, by the same Scale and Line of Chords, the whole Course and Distance, as well as whole Difference of Latitude and Departure, from the several Courses and Distances given, as follows.

RULES for laying down and Resolving a SHIP's TRAVERSE by the Line of CHORDS, with SCALE and COMPASSES.

Cases.	Northing and Southing, Easting and Westing.	Condition of Courses.	∠ between each 2 Courses.
1	2 Meridians, 2 Parallels	Subtract less from greater Course.	For the Points or Angle, between the Distances on each two succeeding Courses.
2	2 Meridians, 1 Parallel	Add both Courses.	
3	1 Meridian, 2 Parallels.	Subtract the Sum of both Courses from 16 Points.	
4	1 Meridian, 1 Parallel	Add 8 Points, the less Course, and Supplement of greater.	

☞ N. and S. are called Meridians, and E. and W. are called Parallels, in the Courses.

Supplement is what a Course wants of 8 Points or 180° .


To know the Meridians, and Parallels, the First and Second, Second and Third, Third and Fourth, &c. Courses, are compared together.

E X A M P L E.					
Compared	No.	Course	Points fr. Merid.	Points for plans	Dist. doubled
1 and 2	1	S. by E. $\frac{1}{4}$ E.	1 $\frac{1}{4}$	1 $\frac{1}{4}$	22,8
2 and 3	2	S. E. $\frac{1}{4}$ E.	4 $\frac{1}{4}$	13 by Case 4	22,8
3 and 4	3	N. W. by W. $\frac{1}{4}$ W.	5 $\frac{1}{4}$	1 . . . 1	41,4
4 and 5	4	S. $\frac{1}{4}$ E.	5 $\frac{1}{4}$	5 . . . 1	21,2
5 and 6	5	E. $\frac{1}{4}$ N.	7 $\frac{1}{4}$	8 . . . 2	9,6
6 and 7	6	S. S. W. $\frac{1}{4}$ S.	1 $\frac{1}{4}$	6 . . . 1	10,0
7 and 8	7	E. by N. $\frac{1}{4}$ N.	6 $\frac{1}{4}$	5 . . . 1	10,6
	8	W. N. W.	6	3 $\frac{1}{4}$. . . 3	12,0

These several Distances being laid down, or planned, according to the first $\angle$ 1 $\frac{1}{4}$ Point with the Meridian, and the $\angle$ of each succeeding Distance with the former, as in the 2 last Columns, you will find the whole Course and Dist. and whole Dif. of Lat. and Departure as in the foregoing Page.

An exact Geometrical PROJECTION of the DREADNOUGHT's TRAVERSE, 11th of May, 1760, according to the foregoing RULES.

Distances	Points
AB = 22.8	$\angle$ SAB = 1 $\frac{1}{4}$
BC = 22.8	$\angle$ ABC = 13
CD = 41.4	$\angle$ BCD = 1
DE = 21.2	$\angle$ CDE = 5
EF = 9.6	$\angle$ DEF = 8
FG = 10.0	$\angle$ EFG = 6
GH = 10.6	$\angle$ FGH = 5
Current, IH = 12.0	$\angle$ GHI = 3 $\frac{1}{4}$
Whence, IP = 66 = Departure W.	
PA = 39.1 = Dif. Lat. S.	
AI = 39.6 = Dif. run.	
$\angle$ SAI = $9^\circ 35'$ W. = the Course.	



N. B. AS is a North and South Line; N. at Top; S. at Bottom. E. to the Right; W. to the Left.

PLACES where the WINDS are partly *constant*. . . TIMES and PERIODS of blowing. . . DIRECTIONS of blowing.
In open Seas where there is no Interruption from interposing High-Lands, the *Trade-Winds* and *Monsoons* blow as underneath.

1. *Between the Tropics to Lat. 30° N. and S. . . . All the Year constantly. . . . Easterly Wind blows.*
Which Wind, but for Interruptions of *Mountains* and *High Land* on the opposing *Continents*, would blow quite round the *Globe*. And without these *Limits*, which are narrow, this Wind changes *instantaneously*.

- | | | |
|--|---|--|
| 1. <i>Between the Tropics to Lat. 30° N. and S.</i> | <i>All the Year constantly.</i> | <i>Easterly Wind blows.</i> |
| <i>Which Wind, but for Interruptions of Mountains and High Land on the opposing Continents, would blow quite</i> | | |
| <i>und the Globe. And without these Limits, which are narrow, this Wind changes instantaneously.</i> | | |
| 2. <i>From 30° to 35° Lat.</i> | <i>All the Year</i> | <i>Westerly Wind blows.</i> |
| 3. <i>Near the Tropic of Cancer</i> | <i>All the Year</i> | <i>N. Easterly.</i> |
| 4. <i>Near the Tropic of Capricorn</i> | <i>All the Year</i> | <i>S Easterly.</i> |
| 5. <i>Sun near the Tropic of Cancer</i> | <i>All Summer</i> | <i>Wind a Point or two more Southerly.</i> |
| 6. <i>Sun near the Tropic of Capricorn</i> | <i>All Winter</i> | <i>Wind a Point or two more Northerly.</i> |

6. *Sun near the Tropic of Capricorn* : : : : The Winds : : : : and a 2000 or 3000 more Northerly.
 ➡ The Sea generally follows the Direction of these Winds. In the *Pacific Ocean*, the Winds are very fresh, and regular, so as to require little Attendance on the Ship's Sails; and Storms and Tempests are seldom here known.

- | | | | |
|---|---|------------------------|--------------------------------------|
| 7. <i>On the Coast of Peru, the West of Africa,</i> | } | All the Year | <i>Southerly Wind blows.</i> |
| <i>Lat. S. and 100 Leagues from Guinea Coast</i> | | | |
| <i>In the Atlantic and Æthiopic Ocean,</i> | } | All the Year | <i>S. Easterly Wind blows.</i> |
| <i>Coast of Africa to 28° N. and S. Lat. And</i> | | | |
| <i>On the Americ. Coast fr. 31° N. to 4° N. Lat.</i> | } | All the Year | <i>Wind more Easterly, than</i> |
| <i>Near the Coast of Brasil, in America</i> | | | |
| <i>Near the Coast of Africa, in same Latitude, where,</i> | | All the Year | <i>The Wind more Southerly, tha.</i> |

8. On S. Coast of Guinea, fr. Sierre de Leon } Perpetual Wind Southerly and S. Westerly.
to St. Thomas's Isle }

Where are frequent *Calms* and *Tornados*.

- | | | |
|--|------------------------|---|
| 9. Latitude 10° on Guinea Coast | All the Year | West. |
| 10. Latitude 20 or 30° | All the Year | North West. |
| 11. Between 4 and 10° N. Lat. a Tract of Sea,
called the Rains . . . 100 or 150 Leagues
fr. Guinea-Coast | Perpetual | Calms, attended with Thunder and
[Lightning, and frequent Rains. |

Near the Caribbe-Islands . . Hurricanes, in August . . Wind more Easterly, sometimes E. or E. by S. but common'y a Point or These Winds gradually decrease their Strength, failing to the Westward. [two to Northward.]

- | | | | |
|-----|--|--|-------------------------|
| 12. | On the Brazil Coast | { From April
{ From September | South-West Wind blows. |
| | To South of Brazil | | North-East. |
| | | All the Year | Wind more Westerly. |
| 13. | On the West Coast of America | | Westerly. |
| 14. | In the Indian Ocean | Partly general, as in the Æthiopic and Atlantic Sea. | |
| | From Lat. 10° to 30° Lat. S. | Partly periodical . . All the Year | S. E. by E. Wind blows. |

15. From 20° to 10° Latitude S. As far as the *Molucco* Islands. A Degree more, near *Java*, and a Degree less near *Madagascar*. } **MONSOON** { From June till November . . South-East Wind blows.
From December till May . . North West.

16. From 3° S. Latitude Northwards { From October till April . . . Clear Breeze North-East.
 { From April till October . . . South-West or S. S. West, at-
 tended with Rains, and stronger of the two.

But these Winds are less constant in the *Gulf of Bengal* than in the *Indian Sea*.

17. *Between Madagascar and African Coast, as far as the Equinoctial* } *From April to October* . . . *S. S. West Wind blows, but more (Westerly near the Line.*
Rest of the Year *Easterly.*

18. *To the East of Sumatra, the Coast of Cambaia, and China, as far as the Philippine Islands and Japan* } *From November till May . . North Wind blows.*
 } *From May till Oct. or Nov. . . South Wind, varying a Point or*
 [two at Times.

19. *Between New Guinea and Sumatra, South of the Equinoctial, North Wind changes North-westerly, South to South-easterly, varying sometimes 5 or 6 Points.*

☞ The Time of these Winds changing is a Month or 6 Weeks later than their Times of blowing on the forementioned Coasts.

- | | | |
|--|---------------------------------|------------------------|
| 20. <i>Between the S. End of Madagafcar and the Shore</i> | } From October till May : . . . | South-East Wind blows. |
| | | From May |
| 21. <i>Beyond St. Lawrence, (otherwise called Madagascar) into the Sea</i> : | | South Wind. |

Of the TRADE WINDS, and MONSOONS.

REMARK. These *contrary* Winds shift not suddenly. In some Places they change to *Calms*; in others to *variable* Winds.

THE End of the *Westerly Monsoon* on the Coast of *Coromandel*, and the two *last Months* of the *Southerly Monsoon*, in the *Chinese Sea*, are generally *tempestuous*; so as to make the Navigation in these Seas very hazardous, at the *breaking up* of these Monsoons; for so these *Tempests* are called.

☞ In the *temperate Zones*, at all Times of the Year, the *variable Winds* blow from all Points of the Compass. And near the *Sea-Coasts*, the Winds blow with different Forces and Directions, by the Land differently reflecting Heat, and interrupting the Course of those Winds.

THE Winds, that constantly blow from, or nearly from, some certain Point of the Compass, are called *Trade Winds*: and such are the Winds Ships get into bound to the *West Indies*.

If these *Trade Winds* blowing, for a certain Time, in a constant Direction, or near it, shift their Direction, periodically, at a certain Time of the Year, and then blow *contrary*, they are called *Monsoons*: and such are the Winds Ships get into bound to the *East Indies*.

Experienced Navigators, bound to the *West Indies*, even to the Coast of *America*, are more solicitous to run into the Latitudes where these *Trade Winds* blow to favour their Passage through those Seas, than to attempt a *directer Course*, and be retarded in their Voyage by *contrary* or *adverse Winds*, as they may happen to blow. And by getting into these *Trade Winds*, to command what Course they please, having the Wind always more or less *abast* the Beam, they find it *commodious* to run down their Latitude, before the Ship comes near the intended Port or Island; and then they steer upon a *Parallel* (directly E. or W.) by which they are sure to hit their Port designed. Whereas the Island of *Barbadoes*, as well as other Places lying in the open Ocean, have sometimes been *mist*, or over run, by the *Inch* or two *Marksmen*, the *hair splitting* and *direct* Navigators; who, being homeward bound, sometimes run their Ship on the *Rocks of Scilly*, instead of *hitting the Lizard*.

For sailing into the *English Channel*, in homeward bound Voyages, see Directions by Captain Robert Brown, of the Navy, further on, recited from our History of the Islands of *Scilly*, for the Navigator's Purpose.

This Art of conducting a Ship in the *Trade Winds*, by a remoter Course, to make the *quicker Passage*, is known to the experienced Seaman; but is a Secret to the *Land Navigator*, (for whose Sake we insert it) that never made a Voyage to prove the Use of his *refined Theories*, and put his *infallible Land-Rules* in Practice.

If Ships, bound to, or from, the *East Indies*, arrive at the Place of a *Monsoon* too late for their Passage, when that *Monsoon* is shifted the *contrary Way* of the Voyage, they are forced to return to the next *convenient Port*, and there wait, *some Months*, for the Return of the *contrary Monsoon*, before they can pursue their Voyage. Whereas, if Ships rightly time their coming to the Place of the *favourable Monsoon*, they hasten their Passage through those Seas, and accomplish their Voyages in the *shortest Time*.

Of these different Sorts of *Trade Winds*, you may observe, that some blow from E. to W. some from S. to N. others from W. to E. — Some are constant in *one Quarter* all the Year; some blow one Half the Year *one Way*, and the other six Months quite *contrary*. Others blow six Months *one Way*, and then shifting only eight or ten Points, continue there six Months more, and then return again to their former Stations, as all these *shifting Trade-Winds* do; and so, as the Year comes about, they alternately succeed each other, in their proper Seasons.

There are other Sorts, called *Sea Winds* and *Land Winds*, differing much from any of the former, the one blowing by Day, the other by Night, constantly and regularly succeeding each other.

Within the *Torrid Zone* also are *violent Storms*, as fierce, if not fiercer than any are in other Parts of the World: And as to the *Seasons* of the Year, they are distinguished in these Parts only by *Wet* and *Dry*, (there never being Frost or Snow) which *wet* and *dry* Seasons do as successively follow one another, as the *Winter* and *Summer* do with us in *England*.

Here are also *strong Currents*, sometimes setting one Way, sometimes another; which though it is difficult to describe, with *desirable Accuracy*, yet we have attempted to give a particular Account of these Matters, from our own Observations, and the Authority of others, further on.

VARIATION of the COMPASS observed at different Times and PLACES.

Places Names	Variation.	Place's Names	Variation
Ascension Isle	1° W.	London	16° W.
Azores Isle	8 W.	Madagascar, W. Coast	19 W.
Baffin's Bay	57 W.	Madagascar, S. fr. Lat. 40°	25 W.
Baltic	10 W.	Majorca Isle	16 W.
Barbary, N. W. Coast	7 W.	Maldavia Isles	10 W.
— N. E. Coast	5 W.	Malacco	2 E.
Bombay	8 W.	New England Coast	10 W.
Brazil Coast, Lat. 18° S.	6 E.	Newfoundland Coast	21 W.
— Lat. 20° S.	12 E.	New Guinea Coast	6 E.
British Channel	15 W.	New Holland, S. and W. Coast	4 W.
Canary Isles	7 W.	New Zealand	9 E.
Cape Comerin	5 W.	Portugal Coast	9 W.
— Frio	12 E.	— W. 8°	6 W.
— of Good Hope	16 W.	River Grand before	14 E.
Chili, West Coast	8 E.	— Plate before	19 E.
England, Westward at Sea	12 W.	St. Helena	2 W.
France, W. Coast	11 W.	Sardinia	13 W.
— Westward at Sea 10°	8 W.	Sicily	12 W.
German Ocean	14 W.	Straits of Gibraltar	6 W.
Guinea, S. Coast	5 W.	— Magellan, E. Entrance	15 E.
Gulph of Bothnia	6 W.	— W. Entrance	13 E.
Hudson's Bay	30 W.	Terra Magellanica, E. Coast	19 E.
— Straits	40 W.	Tristian de Cunha Isle	0
Java Coast	2 W.	Van Diemens, Land	1 E.
Indian Sea	14 W.	Virginia	5 W.

See Variation Table by Mr. W. Mountain, F. R. S. in *Philosophical Transactions*.

REMARKS on the Vulgar ERROR in Books of Navigation, of supposing MERIDIONAL DISTANCE and DEPARTURE to be alike, and of a FIXED QUANTITY.

IN keeping a Ship's Journal, according to *Atkinson's Epitomy*, or *Art of Navigation*, the whole Departure or Difference of Easting or Westing is all along, in the daily Reckonings, reduced into Degrees and Minutes, by dividing the Departure of each daily Reckoning by 60. And these Degrees of Departure are very erroneously called by Mr. Atkinson, the Meridional Distance; being of another Nature to what Departure is.

Mr. Martin, in his *Spheroidal Navigation*, falls into the same Error, by making Departure from the first Meridian to be the Meridional Distance from that Meridian.

And Mr. Mountain, F. R. S. has settled his Longitudes of Places in the *West-Indies* (See p. 242, *Seaman's Compass rectified*, 1759.) by the Meridional Distance from Barbadoes to the Lizard, according to the Journals of experienced Mariners; which Journals, however, of Departure, for Meridional Distance, in long Voyages, can produce no Certainty: Since there may be as many different Departures, betwixt any two remote Places, as there are different Courses steered betwixt them. And these Departures will differ more or less, according as the different Courses in sailing betwixt the two Places, be nearer to the Meridian, or to an E. or W. Parallel.

1. For, in Sailing towards the Pole, if a Ship directs her Course N. or S. till she arrives in the Parallel of Latitude of the Port designed, and then pursues an E. or W. Course, till she arrives at her Port, the Departure by these Courses will be the least possible.

2. And if she directs her Course E. or W. in the Parallel of Latitude of the Place departed from, till she arrives in the Difference of Longitude of the Port designed, and then pursues a N. or S. Course, she will make the greatest Departure possible.

3. On the contrary, if a Ship sails directly S. or N. (according as the Place sailed from is N. or S. Latitude) towards the Equinoctial on this Side, till she arrives in the Parallel of Latitude of the Port designed, and then pursues an E. or W. Course, she will make the greatest Departure possible.

4. But, if she sails E. or W. in the Parallel of Latitude of the Place departed from, towards the Equinoctial, on this Side of it, till she arrives in the Difference of Longitude of the Place designed, and then pursues a S. or N. Course, she will make the least Difference of Departure possible.

5. Again, in Sailing betwixt one Latitude N. and the other S. across the Equinoctial, a Course directly on the Meridian betwixt the two Latitudes, and then an E. or W. Course, the Difference of Longitude, will make the Departure the least, provided the Latitude sailed from be the less.

6. But if the Latitude sailed from is the greater, then a W. or E. Course, the Diff. of Longitude, and a Course on the Meridian to the Place designed will give the Departure the least.

7. And sailing from one Latitude to the Equinoctial, then the Dif. of Longitude on the Equinoctial, and then on the Meridian to the other Latitude, (on 3 different Courses) will make the Departure always the greatest possible.

INFERENCE. But one Course (if practicable) betwixt any two Places on the Globe, will make the same Departure, or a Middle Departure, betwixt the greatest and least Departures in sailing backward or forward by 2 or 3 different Courses. So that, from hence, it is clear, that Ships may make as many different Departures as they make different Courses betwixt these 2 or 3 Courses, that make the greatest and least Departures. Q. E. D.

The Nature of variable Departure between two Places being shown as above, the Meridian or Meridional Distance between two Places will be the fixed Distance of their Meridians, at the Place sailed to, on a direct E. and W. Parallel. Which Meridional Distance is always equal to the least Departure, in sailing towards the Pole, but to the greatest Departure in sailing from the Pole, towards the Equinoctial. See Remark p. 276 of this Work. See also pages 68 and 69 of Mr. Emerson's *Navigation*, for the Distinction of Meridional Distance and Departure.

INDEXED. The Sum of all the Meridional Distances, made in Sailing, will be equal to the Sum of all the Departures, in which Sense only Meridional Distance and Departure are synonymous, or import the same, which Departure or Meridional Distance, is, however, proved to be very different in long Voyages; Because the same Courses and Distances, all the Way, are never pursued.

A TABLE shewing the Miles or Minutes of the Equator, that make a Degree of Longitude in any Latitude.

Deg ^s of Lat.	Minutes.	Deg ^s of Lat.	Minutes.	Deg ^s of Lat.	Minutes.
1	59.99	31	51.43	61	29.09
2	59.97	32	50.88	62	28.17
3	59.92	33	50.32	63	27.24
4	59.86	34	49.74	64	26.30
5	59.77	35	49.15	65	25.36
6	59.67	36	48.54	66	24.41
7	59.56	37	47.92	67	23.45
8	59.42	38	47.28	68	22.48
9	59.26	39	46.62	69	21.50
10	59.08	40	45.95	70	20.52
11	58.89	41	45.28	71	19.54
12	58.68	42	44.95	72	18.55
13	58.46	43	43.88	73	17.54
14	58.22	44	43.16	74	16.53
15	57.95	45	42.43	75	15.52
16	57.67	46	41.68	76	14.51
17	57.37	47	40.92	77	13.50
18	57.06	48	40.15	78	12.48
19	56.73	49	39.36	79	11.45
20	56.38	50	38.57	80	10.42
21	56.01	51	37.76	81	9.38
22	55.63	52	36.94	82	8.35
23	55.23	53	36.11	83	7.32
24	54.81	54	35.26	84	6.28
25	54.38	55	34.41	85	5.23
26	53.93	56	33.55	86	4.18
27	53.46	57	32.68	87	3.14
28	52.97	58	31.79	88	2.09
29	52.47	59	30.90	89	1.05
30	51.96	60	30.00	90	0.00

DIVISION of the OCEAN.

EASTERN, or Oriental Ocean; WESTERN, or Occidental; NORTHERN, or Septentrional; SOUTHERN, or Meridional Ocean. Other Seas take their Names from the Continents or Countries they wash; or from the Nature of those Seas.

As Chinese Sea, from adjacent Country of China.

Indian Sea, washing the Indian Shore.

Persian Sea, washing the Persian Shore.

Arabian Sea, washing the Arabian Shore.

Æthiopian Sea, washing the Æthiopic Shore.

Atlantic Sea, from Atlas, a huge Mountain in Africa, against which lies Mar del Nort, lying to the N. of America.

Mar del Zur, lying to the S. of America, or Pacific Ocean, from its Calmness.

Oceanus Hispanicus, or Spanish Sea, washing the Coast of Spain; called also Bay of Biscay, by the English.

English Channel, between England and France.

Irish Sea, or St. George's Channel, between England and Ireland.

German or British Ocean, between England and Holland.

Caledonian Sea, Northward of Scotland.

Hyperborean, or Frozen Sea, beyond the North Cape.

Tartarian Sea, or Scythian Ocean, far Eastward, &c.

Baltic Sea, between Denmark, Sweden, and Germany, whose Entrance is called the Sound.

Mediterranean, or Straits; called Levant by the Spaniards.

The Entrance is called Straits of Gibraltar.

Pontus Euxinus, or Black Sea, joining to Palus Meotis, or Mar del Zabèche.

Caspian, or Hircanian Sea.

Arabian Gulph, or Red Sea.

Persian Gulph, or Gulph de Elcatif.

Division of the Earth and Ocean.

DIVISION of the EARTH.

Four Continents, EUROPE, ASIA, AFRICA, AMERICA.

EUROPE Bounded.

To the North, by the Northern Ocean, or frozen Sea; on the South by the Mediterranean Sea lying betwixt it and Africa; on the East by the River Tanais; on the West by the Western, or Atlantic Ocean.

Divided into Chief Regions.

Poland, Muscovia, Sweden, Norway, Denmark, Germany, France, Spain, Holland, Flanders, Portugal, Italy, Hungary, Sclawonia, Greece, Dalmatia, Romania, little Tartary.

Chief Islands.

Great-Britain, Ireland, Sardinia, Sicily, Corsica, Candia, Negropont, Cyprus.

ASIA Bounded.

On the North by the Northern and Tartarian Sea; on the South by the Adriatic Gulph, or Red Sea; on the East by the Indian Ocean; and on the West by the River Tanais.

Divided into Chief Provinces.

Turkey in Asia, Arabia, Tartary, Persia, Natolia, Mesopotamia, Assyria, Chaldea, Syria, Armenia, Palestine, Georgia, Media, Partbia, China, India.

Chief Islands.

Japan, Sumatra, Borneo, in the Oriental Ocean; Rhodes, in the Mediterranean Sea; Metelino, Scio, Samos, &c. in the Archipelago, and some Islands in the East-Indies.

AFRICA Bounded.

On the North by the Mediterranean; on the East by the Red Sea; on the South by the Æthiopic or Southern Ocean; and on the West by the Atlantic Ocean.

Divided into Chief Provinces.

Ægypt, Barbara, Monematapa, Desarts of Zava, Negroland, Biledulgerid, Nubia, Guinea, Æthiopia, or Abyssinia.

Chief Islands.

The Canary Isles, Cape De Verd Isles, Isles of Azores, Madeira Islands, St. Thomas, Madagascâr, on St. Lawrence, Malta, in Mediterranean, St. Helena.

AMERICA Bounded.

On the North by the Northern Ocean; on the East by the Atlantic Ocean; on the South by the Magellanic Sea; and on the West by the South-Sea, or Mar del Zur.

Divided into North America.

Chief Provinces.

New Spain, Florida, Carolina, Virginia, Maryland, Pennsylvania, New Jersey, New York, New England, New France, Greenland, whether Continent or Island yet unknown.

The Chief Islands of North America.

Iceland, California, Newfoundland, Bermudas.

Into South America.

Chief Provinces.

Magellan, Brazil, Chili, the Amazons, Guiana, Peru, Panama, Cartagena, Paraguay, or Rio de la Plata.

Chief Islands of South America.

Terra del Fuego, Hispaniola, Cuba, Jamaica, Porto Rico, Bahama, Bermudas, Barbadoes, and the Rest of the Caribbee Islands. Moluque, or Molucca Islands, Java the Greater, Java the Less, and other numerous Islands in the East Indies.

From the Division of America as above, it appears that great Part of South America lies in North Latitudes.

See Gordon's geographical Grammar, for further Satisfaction.

AFFECTIONS of Spherical TRIANGLES in general.

1. EVERY great Circle divides the Globe into two equal Hemispheres.
2. Any two great Circles cutting each other, make the opposite Angles equal.

OF ANGLES.

In ANY Spherical TRIANGLE,

1. The external Angle is less than the Sum of the interior opposite Angles.
2. The Sum of any two Angles is greater than the Supplement of the 3d Angle.
3. The greatest Angle is opposite to the greatest Side, and the least Angle to the least Side, and the contrary.
4. Equal Angles are opposite to equal Sides, and the contrary.
5. There may be 3 right, or 3 obtuse Angles.
6. If each of the 3 Angles be acute, each Side will be less than a Quadrant.
7. If there are 2 obtuse Angles, there must be one Side greater than 90° .
8. The 3 Angles being equal, the three Sides will be equal (the Triangle being equilateral) and the contrary.
9. If there be one acute Angle, one Side will be less than a Quadrant.
10. If two Angles be greater than 2 right Angles, the Sum of their opposite Sides will be greater than 180° ; and therefore, one of them is, at least, greater than 90° .
11. If the Angles at the Base be of like Affection, (both greater or less than 90°) the Perpendicular falling, from the vertical Angle, on the Base, will fall within the Triangle: but if of different Affection will fall without.
12. If two Angles be obtuse, and one Angle acute, the Sides will be of like Affection with their opposite Angles.
13. When 2 $\angle$ s are of $\left\{ \begin{array}{l} \text{like} \\ \text{diff.} \end{array} \right\}$ Affection, and the Side included be $\left\{ \begin{array}{l} \text{more} \\ \text{less} \end{array} \right\}$ than 90° , the 3d $\angle$ is $\left\{ \begin{array}{l} \text{obtuse} \\ \text{acute} \end{array} \right\}$.
14. Two unequal $\angle$ s $\left\{ \begin{array}{l} \text{less} \\ \text{greater} \end{array} \right\}$ than 90° , the Side opposite to the $\left\{ \begin{array}{l} \text{less} \\ \text{greater} \end{array} \right\}$ Angle, will be $\left\{ \begin{array}{l} \text{less} \\ \text{more} \end{array} \right\}$ than 90° .
15. Two equal $\angle$ s being $\left\{ \begin{array}{l} \text{less} \\ \text{more} \end{array} \right\}$ than 90° , their opposite Sides will be $\left\{ \begin{array}{l} \text{less} \\ \text{more} \end{array} \right\}$ than 90° .
16. The Sum of any two $\left\{ \begin{array}{l} \text{Angles} \\ \text{Sides} \end{array} \right\}$ is $\left\{ \begin{array}{l} \text{more} \\ \text{less} \end{array} \right\}$ than the Difference between the 3d $\left\{ \begin{array}{l} \text{Angle} \\ \text{Side} \end{array} \right\}$ and 180° .
17. The Sum of the 3 $\left\{ \begin{array}{l} \text{Angles} \\ \text{Sides} \end{array} \right\}$ is less than $\left\{ \begin{array}{l} 540 \\ 360 \end{array} \right\}$ and more than 180° .
18. The Sum of any two Angles — 3d Angle (or any Angle and the Difference of the other 2 Angles) is less than 90° .

OF SIDES.

In ANY Spherical TRIANGLE,

19. ONE Side being greater than 90° , that Triangle has one obtuse Angle.
20. If each of the 3 Sides be greater than a Quadrant, each of the 3 Angles will be obtuse.
21. The Sum of any two Sides is greater than the 3d Side.
22. The more Degrees there be in each Side the more will the Sum of the Angles exceed 180° ; the Spherical Triangle then differing the more from a right-lined Triangle.
23. One of the equal Sides of an Isosceles Triangle being equal to, less, or greater than, 90° , the Angle at the Base will be also equal to, less, or greater than, 90° .
24. Either Side of an Isosceles Triangle is of like Affection with its adjacent or opposite Angle.
25. Two Sides being less than a Quadrant, there will be an acute Angle in that Triangle.
26. When 2 Sides are of like Affection, and the included Angle be acute, the third Side will be less than a Quadrant.
27. When 2 Sides are of different Affection, and the included Angle be obtuse, the third Side will be greater than a Quadrant.
28. When the Sum of any 2 Sides be equal to, less, or greater than, a Semicircle, the Sum of their opposite Angles will be also equal to, less, or greater than a Semicircle.
29. When the Sum of any two Sides is equal to, less, or greater than, a Semicircle, the internal Angle, included by those Sides, will be equal to, less, or greater than, the external opposite Angle, included by one of those Sides, and the third Side continued.
30. If two Sides be less, and one greater than 90° , the Angles will be of like Affection with their opposite Sides.
31. If two Sides are of $\left\{ \begin{array}{l} \text{like} \\ \text{diff.} \end{array} \right\}$ Affection, and the $\angle$ included be $\left\{ \begin{array}{l} \text{acute} \\ \text{obtuse} \end{array} \right\}$ the 3d Side will be $\left\{ \begin{array}{l} \text{less} \\ \text{greater} \end{array} \right\}$ than 90° .

AFFECTIIONS of Right-angled Spherical TRIANGLES.

In a RIGHT-ANGLED Spherical TRIANGLE.

1. EACH oblique Angle and its opposite Leg is of like Affection.
2. If one Leg be a Quadrant, the Hypothenuse will be a Quadrant.
3. If both oblique $\angle$ s, or both Legs are of $\left\{ \begin{smallmatrix} \text{like} \\ \text{diff.} \end{smallmatrix} \right\}$ Affection, the Hypothenuse is $\left\{ \begin{smallmatrix} \text{less} \\ \text{more} \end{smallmatrix} \right\}$ than 90° , and the contrary.
4. If the Hypothenuse is $\left\{ \begin{smallmatrix} \text{less} \\ \text{greater} \end{smallmatrix} \right\}$ than 90° , then either Leg is $\left\{ \begin{smallmatrix} \text{like} \\ \text{unlike} \end{smallmatrix} \right\}$ the adjacent $\angle$, and the contrary.
5. If the Hyp. and one Leg are of $\left\{ \begin{smallmatrix} \text{like} \\ \text{diff.} \end{smallmatrix} \right\}$ Affection, the other Side and op. $\angle$ are $\left\{ \begin{smallmatrix} \text{less} \\ \text{more} \end{smallmatrix} \right\}$ than 90° , and the contrary.
6. The Sum of the 2 oblique Angles is greater than one, and less than 3 right Angles.

REMARK. The doubtful Cases seldom arise in astronomical Computations, wherein are commonly used right-angled Triangles, whose Arcs are less than 90° : For when there are greater Arcs occur, it is usual for the Computer to take their Supplements, and work by the contrary Triangle.

Quicquid in Astronomicis, Geographicis, vel Nauticis, efficiendum, Trigonometriæ attribuendum est.

Emerson's Navigation.

TO sail nearly by the ARCH of a GREAT CIRCLE, when a given COURSE can be pursued.

STEER, from the Place of your Departure to the Port you are bound for, all the Way, by the Angle of Position, from your last Place arrived in, each Day, at Noon; by computing the Difference of Latitude and Longitude for each Day's Reckoning at Noon, and thence compute each successive Angle of Position; by which you are to steer from Day to Day.

To determine the Angle of Position from one given Place to another.

RULE. Find the Distance of each of two Places from the same Pole. Then (by Case 7th oblique Spherical Trigonometry) Say,

By PROPORTION.

1. As Cos. $\frac{1}{2}$ Sum : to Cos. $\frac{1}{2}$ Diff. of the Distance from the same Pole, :: so Cotan. $\frac{1}{2}$ Diff. Long. between the two Places, : to Tan. of an Arch-D $\left\{ \begin{smallmatrix} \text{less} \\ \text{greater} \end{smallmatrix} \right\}$ than 90° $\left\{ \begin{smallmatrix} \text{if } \frac{1}{2} \text{ Sum Polar Dist. } \left\{ \begin{smallmatrix} \text{less} \\ \text{greater} \end{smallmatrix} \right\} \text{ than } 180^\circ. 90^\circ \end{smallmatrix} \right.$
2. As Sine $\frac{1}{2}$ Sum : to Sine $\frac{1}{2}$ Diff. of the Distances from the same Pole, :: so the Cotan. $\frac{1}{2}$ Diff. Longitude between the two Places, : to Tang. Arch E less than 90° .

NOW, $\frac{D+E}{D-E}$ the $\angle$ of Position at the Place of $\left\{ \begin{smallmatrix} \text{greater} \\ \text{less} \end{smallmatrix} \right\}$ Latitude $\left\{ \begin{smallmatrix} \text{nearest to} \\ \text{furthest from} \end{smallmatrix} \right\}$ the same Pole.

Some PROPERTIES of Sines, Tangents, &c.

IF $r = \text{Rad.}$ $t = \text{Tan.}$ $\sigma = \text{Cofec.}$
 $s = \text{Sine}$ $\tau = \text{Cotan.}$ $v = \text{ver. Sine}$
 $c = \text{Cos.}$ $f = \text{Secant}$ $V = \text{verf. S. Sup.}$
 $1-s = \text{Cover. S.}$

Or, If $r = \text{Log. Rad.} = 10$ $t = \text{Log. Tan.}$ $f = \text{Log. Sec.}$
 $s = \text{Log. Sine of an Arc.}$ $\tau = \text{Log. Cot.}$ $\sigma = \text{Log. Cofec.}$
 $c = \text{Log. Cos.}$

$$\text{Then, } 1. s = \sqrt{1 - cc} = \frac{t}{f} = \frac{t}{\sqrt{1 + tt}} = \frac{c}{\tau} = \frac{1}{\sigma} = \sqrt{Vv}$$

$$2. c = r - v = V - 1 = \sqrt{1 - ss} = \frac{s}{t} = \frac{1}{f} = \frac{\tau}{\sigma} = \frac{\tau}{\sqrt{1 + tt}}$$

$$3. t = sf = \frac{s}{c} = \frac{s}{\sqrt{1 - ss}} = \frac{1}{\tau} = \frac{1}{\sqrt{\sigma^2 - 1}} = \frac{\sigma}{\tau} = \frac{f}{\sigma}$$

$$4. \tau = \sqrt{\sigma\sigma - 1} = \frac{1}{t} = \frac{1}{\frac{s}{c}} = \frac{c}{s} = \frac{c}{\sqrt{1 - cc}}$$

$$5. f = t\sigma = \sqrt{1 + tt} = \frac{1}{c} = \frac{t}{s} = \frac{\sigma}{\tau} = \frac{1}{\sqrt{1 - ss}} = \frac{1}{1 - v} = \frac{1}{V - 1}$$

$$6. \sigma = \sqrt{1 + \tau\tau} = \frac{1}{s} = \frac{f}{t} = \frac{t}{c} = \frac{1}{ct} = \frac{1}{\sqrt{1 - cc}} = \frac{f}{\sqrt{ff - 1}}$$

$$= \frac{1}{\sqrt{2v - vv}}$$

$$7. v = 2 - V = 1 - c = 1 - \sqrt{1 - ss} = 1 - \frac{1}{\sqrt{1 + tt}}$$

$$8. V = 2 - v = 1 + c = 1 + \sqrt{1 - ss} = 1 + \frac{1}{\sqrt{1 + tt}}$$

$$\frac{s+t}{t} = 1 + \tau = 1 + \frac{1}{f} = \frac{\tau + \sigma}{\sigma}$$

$$9. 1 = ss + cc = ff - tt = \sigma\sigma - \tau\tau = \tau\tau = cf = s\sigma.$$

$$\text{Then, } s = c + t - r = t + r - f = c + r - \tau = 2r - \sigma.$$

$$c = s + r - t = 2r - f = s + \tau - r = \tau + r - \sigma.$$

$$t = s + r - c = 2r - \tau.$$

$$\tau = c + r - s = 2r - t.$$

$$f = t + r - c = 3r - s - \tau = \sigma + r - \tau = 2r - c.$$

$$\sigma = \tau + r - c = 3r - c - t = f + r - t = 2r - s.$$

If the natural Sines of two different Arcs be denoted by S and s , and their natural Cosines by C and c , Radius being Unity.

$$\left. \begin{array}{l} \text{Sine of their Sum} = S + sC \\ \text{Sine of their Diff.} = S - sC \\ \text{Cosine of their Sum} = C - Ss \\ \text{Cosine of their Diff.} = C + Ss \end{array} \right\} \text{Make } S = s, \text{ and } C = c.$$

Hence, the Sine of twice an Arc, or of 2 equal Arcs $= 2CS$.

And Cosine of a double Arc $= C^2 - 2S^2$.

Also, if S and C = Sine and Cosine of $\frac{1}{2}$ Sum of 2 Arcs,

And s , c = Sine and Cosine of $\frac{1}{2}$ their Difference,

Then, $S + Cs$ = Sine greater, $S - Cs$ = Sine less,

And $C + Ss$ = Cosine greater, $C - Ss$ = Cosine less.

Again, If T and t denote the Tangents of two diff. Arcs, Rad. = 1.

$$\text{Then, } \frac{T+t}{1-tT} = \text{Tan. Sum; and } \frac{T-t}{1+tT} = \text{Tan. Diff. But}$$

for Tangent, Secant, Cosine, and Cotangent of an Arc above 90° take for an Arc as much below 90° , writing a negative Sign.


As Sum of Tangents of 2 $\angle$ s : to their Diff. :: so Sine of the Sum of those $\angle$ s : to Sine of their Difference.

For infinite other Properties, and Proportions of Arches, Sines, Tangents, &c. consult Mr. Emerson's Trigonometry.


ANSWERS

ANSWERS to all the CASES of RIGHT-ANGLED Spherical TRIANGLES.

IN a RIGHT-ANGLED spherical TRIANGLE, If H represent the Hypotenuse, or Side opposite to the Right-Angle, B, and P, the other Sides, Base and Perpendicular, or LEGS; b and p their opposite Angles; then any two of these 5 Terms being given the Rest are found by the ANSWERS to the CASES in the following TABLE.



OR, changing the SIDES of a Right-angled Spherical TRIANGLE, to ANGLES, and the ANGLES to SIDES, as in the annexed Figure, the Cases and their ANSWERS, are as in the following TABLE.



N. B. In the PROPORTIONS, r represents Radius; s Sine; s' Cofine; t Tangent; t' Cotangent; —, less than 90°; +, more than 90°; because Arcs and their Supplements have the same Sines, Cofines, Tangents, and Cotangents.

But here, instead of H, you must take its Supplement. And, for like, put unlike, and the contrary, where a Side or Angle is determined from the Species of H; or where H, in Species, is to be found.

Cases	Given.	Reqd.	ANSWER.		Given.	Required.	ANSWER.	
			By PROPORTION.	in Symbols.			By PROPORTION.	in Words.
1	H, p	B	$r : s'p :: t.H : t.B$ If H { like } p.		Hypoth. and $\angle$.	Leg adjacent.	Rad. : Cof. given $\angle ::$ Tan. Hyp. : Tan. Leg adj.	
2		P	$r : s.H :: s.p : s.P$, like p.			Leg opposite.	Rad. : Sine Hyp. :: Sine given $\angle ::$ Sine op. Leg.	
3	I.	b	$r : s'.H :: t.p : t'.b$ If H { like } p.			Other $\angle$.	Rad. : Cof. Hyp. :: Tan. given $\angle ::$ Cot. other $\angle$.	
4	H, B	p	$r : t'.H :: t.B : s'.p$ If H { like } B.		Hypo. b. and one Leg.	Adjacent $\angle$.	Rad. : Cot. Hyp. :: Tan. given Leg. : Cof. $\angle$ adj.	
5		b	$s.H : r :: s.B : s.b$, like B.			Opposite $\angle$.	Sine Hyp. : Rad. :: Sine giv. Leg : Sine opp. $\angle$.	
6	II.	P	$s'.B : r :: s'.H : s'.p$ If H { like } B.			Other Leg.	Cof. Leg giv. : Cof. Hyp. :: Rad. : Cof. other Leg.	
7	B, p	H	$s'.p : r :: t.B : t.H$ If B { like } p.		Leg and $\angle$ adjacent.	Hypotenuse.	Cof. giv. $\angle ::$ Rad. :: Tan. adj. Leg : Tan. Hyp.	
8		b	$r : s.p :: s'.B : s'.b$, like B.			$\angle$ opposite.	Rad. : Sine adj. $\angle ::$ Cof. adj. Leg : Cof. op. $\angle$.	
9	III.	P	$r : s.B :: t.p : t.P$, like p.			Leg opposite.	Rad. : Sine giv. Leg :: Tan. adj. $\angle ::$ Tan. op. Leg	
10	E. b	H	$s.b : r :: s.B : s.H$ both true.		Leg and $\angle$ opposite.	Hypotenuse.	Sine opp. $\angle ::$ Rad. :: Sine giv. Leg : Sine Hyp.	
11		P	$r : t.B :: t'.b : s'.P$ both true.			Other Leg.	Rad. : Tan. giv. Leg :: Cot. op. $\angle ::$ Sine other Leg.	
12	IV.	p	$s'.B : r :: s'.b : s.p$ both true.			$\angle$ adjacent.	Cof. giv. Leg : Rad. :: Cof. op. $\angle ::$ Sine $\angle$ adj.	
13	B, P	H	$r : s'.B :: s'.P : s'.H$ If B { like } P.		Legs.	Hypotenuse.	Rad : Cof. one Leg :: Cof. other Leg : Cof. Hyp.	
14		p	$s.B : r :: t.P : t.p$, like P.			$\angle$ opp. 1 Leg.	Sine one Leg : Rad. :: Tan. other Leg : Tan. op.	
15		b	$s.P : r :: t.B : t.b$, like B.			$\angle$ opp. other Leg.	[$\angle$]	
16	b, p	H	$t.b : t'.p :: r : s'.H$ If b { like } p.		$\angle$.	Hypotenuse.	Tan. one $\angle ::$ Cot. other $\angle ::$ Rad. : Cof. Hyp.	
17		B	$s.p : r :: s'.b : s'.B$, like b.		One Leg.		Sine one $\angle ::$ Rad. :: Cof. other $\angle ::$ Cof. op. Leg.	
18		P	$s.b : r :: s'.p : s'.P$, like p.		Other Leg.			

UNIVERSAL PROPOSITION. In any right-angled spherical Tr. angle, the Sine of the middle Part and Radius are reciprocally proportional to the Tangents of the Extremes conjunct, and to the Cofines of the Extremes disjunct.

Viz. Rad. $\times$ Sine middle Part = { Rectangle Tangents of Extremes conjunct. } One of the 5 Parts of a right-angled Triangle (omitting Radius) being reckoned middle Part; and the two contiguous or remote Parts, Extremes conjunct, and disjunct.

The Hypotenuse, and two adjacent Angles, are considered, in the above Proportions, as Complements of their real Arcs.

One Term	Hyp. or $\angle$	Leg
Middle Part	Cofine	Sine
Extr. conjunct	Cotangent	Tangent
Extr. disjunct	Sine	Cofine

Like RATIOS in the same RECTANGLE.
Radius : Sine : Cofine
Secant : Tangent : Radius
Cofecant : Radius : Cotangent.

EXAMPLE I. If B and p are given to find P? Here (among the 5 Parts, B, p, H, b, P) Sine B is middle Part, and Cotang. p, and Tang. P, Extremes conjunct; that is $s.B \times r = t'.p \times t.P$. Hence, $t'.p : r :: s.B : t.P$; Or, $r : t'.p :: s.B : t.P$. That is, (Putting $r : t$ for $t' : r$) $r : s.B :: t.p : t.P$.





EXAMPLE II. If P and b are given to find p? Here $s'.p$ is Middle Part, and $s.b$ and $s'.P$ Extremes disjunct. Therefore, $s'.p \times r = s.b \times s'.P$; Whence, $r : s.b :: s'.P : s'.p$.

In a right-angled spherical Triangle, if the Legs (and therefore the $\angle$ s) { be like } Then the { — } than 90°, { unlike } Hypotenuse { + } and contra.

ANSWERS to all the CASES of OBLIQUE Spherical TRIANGLES.

vide
note 331

IN any oblique angled spherical
TRIANGLE, ABC, whose
Sides,
 $\left. \begin{array}{l} BC = A \\ CA = B \\ BA = C \end{array} \right\}$ and op. $\angle s \left\{ \begin{array}{l} = a \\ = b \\ = c \end{array} \right.$
 Any 3 of which 6 Terms being gi-
 ven, the Rest are found as below.



Let $r = \text{Radius}$, $s = \text{Sine}$,
 $s' = \text{Cofine}$, $t = \text{Tangent}$, $t' =$
 Cotangent .

$\angle$ less
 $\triangleright$ greater } than.

$-$ less
 $+$ more } than 90°.

$d = \angle BCD$; $e = \angle ACD$.
 $D = BD$; $E = AD$.

Cases.	Given.	Sought.	ANSWER. } in Symbols. By PROPORTION.	GIVEN.	SOUGHT.	ANSWER. } in Words. By PROPORTION.
1	A, B, b	a	$s.B : s.b :: s.A : s.a$ both true, or ambiguous.	2 Legs $\angle$ opposite one of them.	Other op. $\angle$. Included $\angle$.	Sine 1 Leg : S. its op. $\angle ::$ S. oth. Leg : S. op. $\angle$. Rad. : Cof. Leg adj. to $\angle ::$ Tan. $\angle$ given : Cot. vert. $\angle$ included by adj. Leg and Perp. Tan. op. Leg : Tan. adj. Leg :: Cof. adj. vert. $\angle$: Cof. oth. ver. $\angle$ incl. by oth. Leg. & Perp. Then, $\angle$ required = Sum or Dif. vert. $\angle$.
2	...	c	$r : s'.A :: t.b : t'.d$ like } A, if } b - unlike } b + $t.B : t.A :: s'.d : s'.e$ like } B, if } b - unlike } b + Then $c = d \pm e$, both true.	...	Base.	Rad. : Cof. $\angle$ given :: Tan. adj. Leg. : Tan. adj. Segment of Base. Cof. adj. Leg : Cof. other Leg :: Cof. adj. Seg- ment : Cof. other Segment. Then Base reqd. = Sum or Dif. of Segments.
3	...	C	$r : s'.b :: t.A : t.D$ like } A, if } b - unlike } b + $s'.A : s'.B :: s'.D : s'.E$ like } B, if } b - unlike } b + Then $C = D \pm E$, both true. But, if $B = A$, or $= 180^\circ - A$, or be betw. A & $180^\circ - A$, then a like A only. And If A } like } b, then $c = d \pm e$ only; unlike } b, then $c = d \pm e$ only; And $C = D \pm E$ only.	...	Base.	
4	a, b, A	B	$s.a : s.A :: s.b : s.B$ both true.	2 $\angle s$ at Base and Leg oppo- site one of them.	Other Leg. Base.	Sine op. $\angle$: S. Leg given :: S. oth. $\angle$: S. op. Side. Radius : Cof. $\angle$ adj. to Leg :: Tan. Leg : Tan. adjacent Segment of Base. Tan. op. $\angle$ to Leg : Tan. adj. $\angle ::$ Sine adj. Segment : Sine other Segment. Then, Base reqd. = Sum or Dif. Segments.
5	...	C	$r : s'.b :: t.A : t.D$ like } b if A } - unlike } b if A } + $t.a : t.b :: s.D : s.E$ both true. Then $C = D \pm E$, if a } like } b. unlike } b. $r : s'.A :: t.b : t'.d$ like } b, if A } - unlike } b, if A } + $s'.b : s'.a :: s.d : s.e$ both true. Then, $c = d \pm e$ if a } like } b. unlike } b. But, if $A = B$, or $180^\circ - B$, or is between B and $180^\circ - B$, then B unlike b; and E + e are impossible.	...	Third $\angle$.	
6	...	c	$r : s'.A :: t.b : t'.d$ like } b, if A } - unlike } b, if A } + $s'.b : s'.a :: s.d : s.e$ both true. Then, $c = d \pm e$ if a } like } b. unlike } b. But, if $A = B$, or $180^\circ - B$, or is between B and $180^\circ - B$, then B unlike b; and E + e are impossible.	...	Third Side.	
7	A, B, b	a	$s. \frac{1}{2} A - B : s. \frac{1}{2} A + B :: t. \frac{1}{2} A - B : t. \frac{1}{2} C$. $s. \frac{1}{2} A - B : s. \frac{1}{2} A + B :: t. \frac{1}{2} A - B : t. \frac{1}{2} c$. Then, $c = d \pm e$ if a } like } b. unlike } b. But, if $A = B$, or $180^\circ - B$, or is between B and $180^\circ - B$, then B unlike b; and E + e are impossible.	2 Sides and 2 opp. $\angle s$.	Third Side. Third $\angle$.	Sine $\frac{1}{2}$ Dif. op. $\angle s$: Sine $\frac{1}{2}$ their Sum :: Tan. $\frac{1}{2}$ Dif. Legs : Tan. $\frac{1}{2}$ third Side. Sine $\frac{1}{2}$ Dif. Legs : Sine $\frac{1}{2}$ their Sum :: Tan. $\frac{1}{2}$ Dif. op. $\angle s$: Cot. $\frac{1}{2}$ third $\angle$.
8	A, C, b	a	$r : s'.b :: t.A : t.D$ like } A, if } b - unlike } b, if } b + $s'.E : s'.D :: t.b : t.a$ like } b, if D } < E. unlike } b, if D } > E. Or $s. \frac{1}{2} A + C : s. \frac{1}{2} A - C :: t. \frac{1}{2} b : t.E$ - Again $s'. \frac{1}{2} A + C : s'. \frac{1}{2} A - C :: t. \frac{1}{2} b : t.D$ + If $A + C > 180^\circ$. Then, $a = D + E$. And $a = D - E$. $r : s'.b :: t.A : t.D$ like } A, if } b - unlike } b, if } b + $s'.D : s'.E :: s'.A : s'.B$ like } E, if } b - unlike } E, if } b +	2 Sides and included $\angle$.	Other $\angle s$. Or Again	Rad. : Cof. included $\angle ::$ Tan. Side op. reqd. $\angle$: Tan. adjacent Segm. Base. The other Seg. Base = Dif. Base and adj. Seg. Sine latter Segm. : Sine former found Segm. :: Tan. $\angle$ included : Tan. sought $\angle$. Sine $\frac{1}{2}$ Sum Sides : Sine $\frac{1}{2}$ their Dif. :: Cot. $\frac{1}{2}$ incl. $\angle$: Tan. remote Segment Base. Cof. $\frac{1}{2}$ Sum Sides : Cof. $\frac{1}{2}$ their Dif. :: Cot. $\frac{1}{2}$ incl. $\angle$: Tan. adjacent Segment of Base. Then, other $\angle$ at Base = Sum } Base's $\angle$ at Vertex = Dif. } Segments.
9	...	B	$r : s'.b :: t.A : t.D$ like } A, if } b - unlike } b, if } b + $s'.D : s'.E :: s'.A : s'.B$ like } E, if } b - unlike } E, if } b +	...	Third Side.	Rad. : Cof. included $\angle ::$ Tan. of either Side : Tan. of adjacent Segment of Base. Whence other Seg. = Dif. Base and that Seg. Cof. adj. Seg. : Cof. remote Seg. of Base, :: Cof. adj. Side : Cof. remote Side.

errata
331

ANSWERS to the CASES of OBLIQUE Spherical TRIANGLES continued.

ANSWER.				ANSWER.			
Cases.	Given.	Sought.	In Symbols.	GIVEN.	SOUGHT.	In Words.	
			By PROPORTION.			By PROPORTION.	
9	b, c, A	a	$r : s'.A :: t.b : t'.d$ like unlike } b, if A { $+$ $e = c$ Dif. d, Then $s.d : s.e :: s'.b : s'.a$ like unlike } b, if d { $<$ $>$ $r : s'.A :: t.b : t'.d$ like unlike } b, if A { $+$ $e = c$ Dif. d. $s'.e : s'.d :: t.A : t.B$ like unlike } e, if b { $+$ Or, $s. \frac{1}{2} b + c : s. \frac{1}{2} b - c :: t. \frac{1}{2} A : t.E -$ And, $s'. \frac{1}{2} b + c : s'. \frac{1}{2} b - c :: t. \frac{1}{2} A : t.D \mp$ if $b + c > 180^\circ$. Then, $B = D + E$; $C = D - E$.	2 $\angle$ s and included Side.	Third $\angle$.	Rad. : Cof. included Side :: Tan. adj. $\angle$ at Base : Cot. adj. vert. $\angle$. Whence the remote vert. $\angle$ is known = Dif. whole vert. $\angle$ and adj. vertical $\angle$. Sine adj. : Sine remote vert. $\angle$:: Cof. adj. $\angle$ at Base : Cof. required $\angle$. Rad. : Cof. included Side :: Tan. adj. $\angle$ at Base : Cot. adj. vertical $\angle$. Hence remote vertical $\angle$ = Difference whole and adjacent vertical $\angle$. Cof. remote $\angle$: Cof. adj. vert. $\angle$:: Tan. adj. vert. $\angle$: Tan. Side required. Sine $\frac{1}{2}$ Sum $\angle$ s : Sine $\frac{1}{2}$ Diff. $\angle$ s :: Tan. $\frac{1}{2}$ adj. Side : Tan. remote Segment. Cof. $\frac{1}{2}$ Sum : Cof. $\frac{1}{2}$ Diff. $\angle$ s :: Tan. $\frac{1}{2}$ adj. Side : Tan. adj. Segment at Base. Then required Sides = Sum and Difference of Segments at Base.	
10	...	B	$r : s'.A :: t.b : t'.d$ like unlike } b, if A { $+$ $e = c$ Dif. d. $s'.e : s'.d :: t.A : t.B$ like unlike } e, if b { $+$ Or, $s. \frac{1}{2} b + c : s. \frac{1}{2} b - c :: t. \frac{1}{2} A : t.E -$ And, $s'. \frac{1}{2} b + c : s'. \frac{1}{2} b - c :: t. \frac{1}{2} A : t.D \mp$ if $b + c > 180^\circ$. Then, $B = D + E$; $C = D - E$.	...	Sides.	Or,	
11	A, B, C	b	$r. \frac{1}{2} C : t. \frac{1}{2} A + B :: t. \frac{1}{2} A - B : t.mD =$ Dif. Perpend. from Middle of the Base. Now, if Perp. within, $Bm + mD = BD$ } greater } Segm. $Am - mD = DA$ } less } Or $2 \times mD = BD$, when Perp. without. Now, $r.A : Rad. :: t.BD : Cof. b$. Case 4. Right angled Δ s. Or, $s.C \times s.A$ (including $\angle$ required) : $r^2 :: s. \frac{1}{2}$ $B + C - A \times s. \frac{1}{2} B + A - C : r^2. \frac{1}{2} b$. If $A + B$ } $<$ } 180° , the Perpend. falls nearest the } less } Side. } greater }	3 Sides.	$\angle$ at the Base.	Tan. $\frac{1}{2}$ Base : Tan. $\frac{1}{2}$ Sum Legs :: Tan. $\frac{1}{2}$ Dif. Legs : Tan. $\frac{1}{2}$ Dif. or $\frac{1}{2}$ Sum Segments of Base. Now, if Perpendicular falls within, $\frac{1}{2}$ Base } add } $\frac{1}{2}$ Dif. for } greater } Segm. } subtr. } less } Or $\frac{1}{2}$ Sum doub'd = wh. Base; the Perp. without. Tan. Leg adj. $\angle$ required : Rad. :: Tan. adj. Segment : Cof. $\angle$ adj. required. Or, Rect. $\angle$ Sines Sides (including required $\angle$) : Rad. Sq. :: Rect. $\angle$ Sines, of two Differen- ces betw. $\frac{1}{2}$ Sum of 3 Sides, and each Side including the required $\angle$: Square of Sine $\frac{1}{2}$ required $\angle$.	
12	a, b, c	A	As $\angle$ s at Base are like unlike } Perp. falls } within. } without. $r. \frac{1}{2} a + b : t. \frac{1}{2} a$ Dif. $b :: t. \frac{1}{2} c : t. \angle$ in- cluded by Perpendicular and Arch bisecting the vertical $\angle$. Whence $\angle BCD$ is known. For, $\frac{1}{2} c +$ angular Distance from Perpendicu- lar = greater $\angle$. $\frac{1}{2} c$, Diff. angular Distance from Perpendicular = less $\angle$. If $a + b$ } $<$ } 180° , the Perpend. falls nearest } less } Side } greater } $\angle$. } greater } less } Now, $t.d : t'.b :: r : s'.A$. 15th Case Right- angled Δ s. Or, $s.b \times s.c : r^2 :: s'. \frac{1}{2} a + c + b \times s'. \frac{1}{2} c + b$ $- a : s^2. \frac{1}{2} A$. Always let fall your Perpendicular from the End of a given Side, and opposite to a given $\angle$. Or, change the 3 given $\angle$ s into Sides, and Sides required into $\angle$ s, but instead of the greatest $\angle$, take its Supplement for one Side, then proceed as in Case 11.	3 $\angle$ s.	Side.	Cotan. $\frac{1}{2}$ Sum $\angle$ s at Base : Tan. $\frac{1}{2}$ their Dif. :: Tan. $\frac{1}{2}$ vert. $\angle$: Tan. $\angle$ included by Perpendicular and bisecting Arch of the ver- tical Angle. Then to $\frac{1}{2}$ vertical $\angle$ add angular Distance from Perpendicular for greater $\angle$. And, Diff. between $\frac{1}{2}$ vertical $\angle$ and angular Distance from Perpendicular = less $\angle$. Tan. adj. vert. $\angle$: Cot. adj. $\angle$ at Base :: Rad. : Cof. adj. Side, being Hypotenuse. Or, Rect. $\angle$ Sines of the $\angle$ s, including Side re- quired : Rad. Square :: Rect. $\angle$ two Co- sines, of $\frac{1}{2}$ Sum of the 3 $\angle$ s, and of $\frac{1}{2}$ Dif. be- tween the two including $\angle$ s, and third $\angle$: Sine Square of $\frac{1}{2}$ included Side, required.	
7	A, C, b	a	$r : s'.b :: t.A : t.D$ like unlike } A if b { $+$ $E = C$ Dif. D. $s.E : s.D :: t.b : t.a$ like unlike } b, if D { $<$ $>$ Or $s. \frac{1}{2} A + C : s. \frac{1}{2} A - C :: t'. \frac{1}{2} b : t.E -$ Again $s'. \frac{1}{2} A + C : s'. \frac{1}{2} A - C :: t'. \frac{1}{2} b : t.D \mp$ If $A + C$ } $<$ } 180° , Then, $a = D + E$. And $e = D - E$. $r : s'.b :: t.A : t.D$ like unlike } A, if b { $+$ $E = C$ Dif. D. $s'.D : s'.E :: s'.A : s'.B$ like unlike } E, if b { $+$ $-$	2 Sides and inclu- ded $\angle$.	Other $\angle$ s.	Rad. : Cof. included $\angle$:: Tan. Side op. req'd. $\angle$: Tan. adjacent Segm. Base. The other Seg. Base = Dif. Base and adj. Seg. Sine latter Segm. : Sine former found Segm. :: Tan. $\angle$ included : Tan. sought $\angle$. Sine $\frac{1}{2}$ Sum Sides : Sine $\frac{1}{2}$ their Dif. :: Cot. $\frac{1}{2}$ incl. $\angle$: Tan. remote Segment Base. Cof. $\frac{1}{2}$ Sum Sides : Cof. $\frac{1}{2}$ their Dif. : Cot. $\frac{1}{2}$ incl. $\angle$: Tan. adjacent Segment of Base. Then, other $\angle$ at Base = Sum } Base's $\angle$ at Vertex = Dif. } Segments.	
8	...	B	$r : s'.b :: t.A : t.D$ like unlike } A, if b { $+$ $E = C$ Dif. D. $s'.D : s'.E :: s'.A : s'.B$ like unlike } E, if b { $+$ $-$	...	Third Side.	Rad. : Cof. included $\angle$:: Tan. of either Side : Tan. of adjacent Segment of Base. Whence other Seg. = Dif. Base and that Seg. Cof. adj. Seg. : Cof. remote Seg. of Base, :: Cof. adj. Side : Cof. remote Side.	

The PROPERTIES of Spherical TRIANGLES.

1. IN any spherical Triangle, The Sines of the Sides are in Proportion to the Sines of the opposite Angles.

2. Letting fall a Perpendicular on the Base, from the vertical Angle, The Sines of the Segments of the Base are reciprocally proportional to the Tangents of the adjacent Angles.

3. The Cosines of the same Segments are directly proportional to the Cosines of the adjacent Sides.

4. The Cosines of the vertical Angles are as the Cotangents of the respective Sides.

5. The Sines of the vertical Angles are as the Cosines of the respective Angles at the Base.

6. As the Sine of Sum of the Legs : to Sine of their Difference :: so Cotangent $\frac{1}{2}$ Sum of Angles at Vertex : to Tangent $\frac{1}{2}$ their Difference.

7. As Sine Sum of the Sides : to Sine of their Difference :: so Cotangent $\frac{1}{2}$ vertical Angle : to Tangent $\frac{1}{2}$ Difference, or $\frac{1}{2}$ Sum, of the vertical Angles, according as the Perpendicular falls within or without.

8. As Tangent $\frac{1}{2}$ Sum of Sides : to Tangent $\frac{1}{2}$ their Difference :: so Tangent $\frac{1}{2}$ Sum of the Angles at the Base, : to Tangent $\frac{1}{2}$ their Difference.

9. As Cotan. of $\frac{1}{2}$ Sum of 2 Sides : to Tan. $\frac{1}{2}$ their Difference :: so Cotan. $\frac{1}{2}$ Base : to Tan. Distance from middle Base to where the Perpendicular falls ; being Tan. $\frac{1}{2}$ Difference of Segments of Base.

10. As Cotangent $\frac{1}{2}$ Sum Angles at the Base, : to Tangent $\frac{1}{2}$ their Difference :: so Tangent $\frac{1}{2}$ vertical Angle, or $\frac{1}{2}$ Sum vertical Angles, : to Tangent $\frac{1}{2}$ the Difference, or $\frac{1}{2}$ the Sum of vertical Angles, as the Perpendicular falls within or without.

11. As Tangent $\frac{1}{2}$ Sum Segments of the Base : to Tangent $\frac{1}{2}$ Sum Sides, :: so Tangent $\frac{1}{2}$ Difference of Sides : to Tangent $\frac{1}{2}$ Difference Segments of Base.

12. As Tangent $\frac{1}{2}$ Base : to Tangent $\frac{1}{2}$ Sum Sides, :: so Tangent $\frac{1}{2}$ Diff. Sides : to Tangent $\frac{1}{2}$ alternate Base = $\frac{1}{2}$ Difference or $\frac{1}{2}$ Sum Segments of the Base, as the Perpendicular falls within or without.

In a right-angled Triangle, the Rectangle of Tangent $\frac{1}{2}$ the Hypotenuse + $\frac{1}{2}$ one Leg, and the Tangent $\frac{1}{2}$ the Hypotenuse — $\frac{1}{2}$ that Leg = Tangent Square of $\frac{1}{2}$ the other Leg.

13. Letting fall a Perpendicular on the Base of a spherical Triangle.

As Sine Sum Angles at the Base : to Sine Difference :: so Tangent $\frac{1}{2}$ Sum of the Segments of the Base : to Tangent $\frac{1}{2}$ Difference.

14. Sine Sum Angles at the Base : Sine Difference :: so Tangent $\frac{1}{2}$ Base : to Tangent $\frac{1}{2}$ alternate Base.

15. Sine Sum Segments of the Base : to Sine of their Difference :: so Sine Sum Angles at the Vertex : to Sine of their Difference.

16. As Sine Base : to Sine vertical Angle :: so Sine Diff. Segments at Base : to Sine Diff. of Angles at Vertex, when the Perpendicular falls within ; or Sine of the Sum of Segments of Base : to Sine Sum of vertical Angles, when the Perpendicular falls without.

17. As Cosine $\frac{1}{2}$ Sum of the Sides : to Cosine $\frac{1}{2}$ their Difference :: so Cotangent $\frac{1}{2}$ included Angle : to Tangent $\frac{1}{2}$ Sum of opposite Angles.

18. As Sine $\frac{1}{2}$ Sum of the Sides : to Sine $\frac{1}{2}$ their Difference :: so Cotangent $\frac{1}{2}$ included Angle : to Tangent $\frac{1}{2}$ Difference of opposite Angles.

19. As Cosine $\frac{1}{2}$ Sum of 2 Angles : to Cosine $\frac{1}{2}$ their Difference :: so Tangent $\frac{1}{2}$ included Side : to Tangent $\frac{1}{2}$ Sum of the opposite Sides.

20. Sine $\frac{1}{2}$ Sum of two Angles : to Sine $\frac{1}{2}$ their Difference :: so Tangent $\frac{1}{2}$ included Side : to Tangent $\frac{1}{2}$ Difference of opposite Sides.

21. Cosine $\frac{1}{2}$ Sum Angles at the Base : to Cosine $\frac{1}{2}$ their Difference :: so Tangent $\frac{1}{2}$ Difference of opposite Sides : to Tangent $\frac{1}{2}$ Difference or $\frac{1}{2}$ Sum, of the Segments of the Base, as the Perpendicular falls within or without.

22. In any spheric Triangle, The Rectangle of the Sines of any two Sides, $\times$ Cosine of included Angle, + Rectangle of their Cosines = Cosine of third Side. (Radius being 1.)

23. The Cosine of an Angle of a spherical Triangle, = Cosine opposite Side — Rectangle of Cosines of the including Sides, the whole divided by the Rectangle of the Sines of the including Sides. (Radius being 1.)

24. The Rectangle of Sines of 2 Angles $\times$ Cosine of the included Side — Rectangle of their Cosines = Cosine of third Angle, Radius being 1.

25. The Cosine of any Side = Rectangle of the Cosines of the including Angles + Cosine of opposite Angle, and the whole divided by Rectangle of Sines of the including Angles, Radius = 1.

26. In any spherical Triangle, As the Product of the Sines of two Sides, including a required Angle, : to the Product of the Sines of two Differences between the $\frac{1}{2}$ Sum of the 3 Sides, and each of the two Sides, including the required Angle, :: so the Square of Radius : to the Square of the Sine of $\frac{1}{2}$ the required Angle.

27. As the Product of the Sines of two Angles, including a required Side, : to the Product of the Sines of two Differences, between the half Supplement of the three given Angles, and the Supplement of each Angle, including the required Side, :: so the Square of the Radius : to the Square of the Cosine of $\frac{1}{2}$ the required Side.

Or, taking the Supplement of the greatest Angle only, you may proportion, as by Property 26, by changing the Sides into Angles.

Viz. As the Product of the Sines of two Angles, including the required Side, : to the Product of the Sines of two Differences between the $\frac{1}{2}$ Sum of the 3 given Angles (the Supplement of the greatest being one) and each of the Angles, including the required Side, :: so is the Square of Radius, : to Square of the Sine of $\frac{1}{2}$ required Side.

REMARK. In plane Trigonometry, right-angled Triangles may be considered as spherical Triangles, whose Sides being very small Arcs may be considered as Right-Lines ; and therefore become equal to their Sines or Tangents.

HENCE,

The PROPERTIES of Spherical TRIANGLES.

HENCE, to plane Trigonometry, may be applied, all the Proportions of spherical Trigonometry, wherein Cosines and Cotangents are not concerned, by using the Word Side, instead of Sine of a Side, or Tangent of a Side. For, if the 3 Sides of a plane Triangle were given, an Angle may be found, as follows, by Property 26.

As the Product of 2 Sides, including a required Angle, : to the Product of two Differences, between $\frac{1}{2}$ Sum of the three Sides, and each of the Sides including the required Angle, :: so the Square of Radius : to the Square of the Sine of $\frac{1}{2}$ required Angle.

28. In any spherical Triangle, As Rectangle of Sines of 2 Sides, including a required Angle, : to Radius Square :: so versed Sine Base — versed Sine Difference of Sides, : to versed Sine included Angle, required.

:: And so Cosine Difference of Sides — Cos. Base : to versed Sine said included Angle, required.

29. Also, As Rectangle of the Sines of 2 Angles, including a required Side, : to Radius Square, :: so versed Sine Sum of including Angles, — versed Sine Supplem. 3d Angle : to versed Sine included Side, required.

:: And so versed Sine third Angle — versed Sine Supplem. Sum of including Angles, : to versed Sine included Side, required.

:: And so Cos. Difference of the including Angles, + Cos. 3d $\angle$: to versed Sine Supplement of included Side, required.

:: And so versed Sine Supplem. 3d Angle — versed Sine Diff. including Angles : to versed Sine Supplem. included Side, required.

:: And so versed Sine Supplem. Diff. including Angles — versed Sine 3d Angle : to versed Sine Sup. included Side, required.

30. If an Arc be drawn from the Vertex to the Middle of the Base of a spherical Triangle,

Then, the Sines of the vertical Angles are reciprocally as the Sines of the Sides, and directly as the Sines of the Angles at the Base.

31. If the vertical Angle of a spherical Triangle is bisected, As twice Cotangent of the bisecting Arc : to the Sum of the Cotangents of the Sides :: so Radius : to Cosine $\frac{1}{2}$ vertical Angle.

32. As double the Radius : to Tang. bisecting Arc :: so Sum Cotangents of Sides : to Cosine $\frac{1}{2}$ vertical Angle.

33. As one right Angle : to the Angle between 2 great Circles :: so the Area of a great Circle : to the lunular Area contained between those great Circles.

34. As 4 right Angles : to Angle intercepted between 2 great Circles :: so Surface of the whole Sphere : to Surface intercepted between those two great Circles.

:: And so the Solidity of the Sphere, : to the Solidity contained between the 2 great Circles.

For the Solidities are as the Surfaces, because they may be reduced to Pyramids of equal Height.

35. In any spherical Triangle, As two right Angles : to the Excess of 3 Angles above two right Angles, :: so the Area of a great Circle of a Sphere : to the Area of that Triangle.

In any spherical Polygon, put n = Number of Sides, $A = 180^\circ \times n - 2$,

Then, As 2 right Angles, or 180° : To Sum of all the Angles of the Polygon — A :: So the Area of a great Circle of the Sphere : To Area of that Polygon.

CORRECTIONS to be made in the Tt Sheet, with the PEN, before it is used.

Page	Col.	Line	For	Write
✓ 323	1	3 fr. Top	Buffin's, &c.	Baffin's Bay.
✓ 324	2	24	some Islands, &c.	some other Islands.
✓ Same	2	3 fr. Bot.	Divisions	Division.
✓ 326		26 fr. Top	less } than 180°	greater } than 90° .
✓ 328	4	Under Case 7, End of Line 1	$c \begin{cases} - \\ + \end{cases}$	$b \begin{cases} - \\ + \end{cases}$
		3	$\angle E$	$\angle C$.
		4	$D \pm$	$D \mp$
		Beginning of Line 7	$C = D - E$	$c = D - E$.
✓ 7		Under Case 2, Line 6	Sum and Dif. ver. $\angle$	Sum or Dif. ver. $\angle s$.

N. B. $mD = \frac{1}{2} BD - DA$, or $\frac{1}{2} BD$, $= \frac{1}{2}$ Dif. or $\frac{1}{2}$ Sum, Segments of Base, according as the Perpendicular falls within or without the Triangle; which Difference added to, and subtracted from $\frac{1}{2}$ the Base, or $\frac{1}{2} BA$, gives the Segments BD , and DA , from where the Perpendicular falls, within; but which Sum doubled is the whole Base when the Perpendicular falls without. See Fig. 1 and 2. Page 328.

SOME GENERAL OBSERVATIONS on the TIDES.

1. IF you observe the Times of the Tides, in any Place, at New and Full Moon, and also their Times at the Quarter Moons, then if you proportion the intermediate Aspects of the Moon with the Sun, from these Times, you will have the Times of the Tides for each Day of the Moon's Age, in that Place, very nearly.
2. AT London the Tides, at the Quarter-Moons, happen (generally) above an Hour sooner than the middle Time, between the Tide Times, at New and Full Moon. Thus, at London, the Tide-Times, at New and Full, are at about Half an Hour after two; the middle Time betwixt which Times, is half an Hour after eight; but the Quarter Moon-Tides generally happen, at London, about, or before, half an Hour after Seven.
3. AT Dublin, it is said, the Quarter Moon Tides happen half an Hour later than the middle Time counted from Tide-Times at New and Full. Thus, on New and Full, at the Custom-house of Dublin, the Tides happen about 11; the middle Time between which Times is about 5, for the Time of the Quarter-Tides: But they happen about half an Hour after 5, 'tis said.
4. THE higher the Moon rises above (or farther she descends below) the Horizon of any Place, the higher the Tide rises in that Place.
Therefore, in North Latitudes, the Moon having North Declination, being above the Horizon, and past the Meridian, she raises higher Tides than when she is under the Horizon, and past the Meridian.
☞ The contrary happens in North Latitudes and South Declinations of the Moon. Observe also the same in South Latitudes and South Declinations of the Moon. And the contrary in South Latitudes and North Declinations of the Moon.
5. Consequently, in North Latitudes, in the Summer, (when the Ecliptic at Noon rises higher than in Winter) the Evening are higher than the Morning Tides. And, in Winter (when the Ecliptic, at Noon, is lower than in Summer) the Morning are higher than the Evening Tides.
6. The highest Spring-Tides are always a little before and after the Equinoxes, about the Beginning of March, and End of September, each Year; but the Quarter or Neap-Tides, are least at those Times, and greatest of all in June and December, about the Solstices.
7. The highest Tides make the lowest Ebbs.
8. Hence, the highest Tide, at New and Full Moon, is always followed by the lowest Tide at the Quarter Moons.
9. The highest monthly Tides are generally observed to be about the second Day after Full or Change, in England; but in some Places abroad, as at Tonquin, and New Holland, the highest Spring-Tides are observed to be 3 Days after New and Full.
10. Tides, in like Circumstances, are highest at the Equator; and higher in less than greater Latitudes.
11. Tides are higher in larger Seas, and lower in less Seas.
12. Tides are higher at the Shores of Continents, where there are Rivers and Indraughts, than in the Middle of the Sea, where they pass freely; or are higher at those Shores, than they are about Islands, far from the Continent.
13. Within Rivers Mouths, the Tides ebb longer than they flow.
14. The Motion of the Tide is greatest a little after half Flood, and a little before half Ebb.
15. A little before high Water in a River, the under Water of the Tide runs down, while the upper Water runs up.
16. At the Top of high or low Water, the Tide has no direct Motion.
17. Strong Winds blowing the same Way with the Running of the Tide, bring it in sooner and make it rise higher; but blowing contrarily, against the Tide's Motion, keep it back later, and make it run lower.

TO find the Time of HIGH-WATER at a given TIME and PLACE?

RULE. By the following Tide-Table, find the Time of High-Water, or Full Sea, on the Day of Full or Change of the Moon, for the Place given; adding to which the Time of the Moon's Southings on the given Day, and the Sum will be the Time of High-Water for the Time and Place required. EXAMPLE. To find the Time of high Water, at London-Bridge, on May 10, 1760?

Time of High-Water at New and Full	2 ^h 30 ^m	} So for the Rest.
On May 10, 1760, Moon souths	8 4 Morning	
High-Water at London-Bridge, then	10 34 Morning	
Add	5 0	
Low Water at Gravesend	3 34 Afternoon.	

An *Alphabetical* TIDE - TABLE, from the best AUTHORITIES. Shewing the TIME of HIGH-WATER at the most remarkable SEA PORTS and SEA-COASTS, when it is NEW or FULL MOON. For finding the TIME of High-Water in those Places at any AGE of the MOON.

Places NAMES.	Time H. Water. h m	Places NAMES.	Time H. Water. h m	Places NAMES.	Time H. Water. h m	Places NAMES.	Time H. Water. h m
A.		<i>Bristol Key</i>	6 45	<i>Dunnoje</i>	9 45	H.	
ABERDEEN, Scot-		<i>Brouage without</i>	3 45	<i>Dunwich</i>	9 45	HAGUE	8 15
land	0 45	<i>Buchanefs</i>	3 0			<i>Hamborough</i>	6 0
<i>Abermorith</i>	6 0	<i>Bulloign</i>	11 0	E.		<i>Hampton Key</i>	0 0
<i>Aberwark</i>	4 30	<i>Buoy of Nore</i>	1 0	EDAM	1 30	<i>Harborough</i>	6 0
<i>Abroth</i>	3 15			<i>Edinburgh</i>	4 30	<i>Harfleur</i>	8 0
<i>Africa, West Coast</i>	3 0	C.		<i>Egmon</i>	4 30	<i>Haerlem</i>	9 0
<i>Aldborough</i>	9 45	CAEN, Normandy	9 0	<i>Eider</i>	0 0	<i>Hartlepool</i>	3 0
<i>Amazon's R. Mouth</i>	6 0	<i>Caldy</i>	5 15	<i>Elbe</i>	0 0	<i>Harwich</i>	11 0
<i>Ambleteuse, Fr. Coast</i>	11 0	<i>Calais</i>	11 30	<i>Emden</i>	0 0	<i>Hastings</i>	10 30
<i>America, W. Coast</i>	3 0	<i>Calshot</i>	11 15	<i>Emes</i>	9 0	<i>Havre de Grace</i>	9 0
<i>— E. Coast</i>	4 30	<i>Camvere</i>	1 30	<i>Emes Entrance</i>	7 30	<i>St Helens</i>	10 30
<i>Amsterdam</i>	3 0	<i>Canary Isles</i>	3 0	<i>Enchusen</i>	0 0	<i>Hern</i>	0 0
<i>Andrews St.</i>	2 15	<i>Cancale</i>	6 0	<i>Engomonts</i>	9 0	<i>Hever</i>	0 0
<i>Antwerp</i>	6 0	<i>Cape Blanco</i>	9 45	<i>Eftaple</i>	11 0	<i>Holms</i>	6 0
<i>Apenars</i>	12 45	<i>— Cantin</i>	0 0	<i>Exmouth</i>	6 20	<i>Holy Head</i>	1 30
<i>Apenmark, Fr. Coast</i>	2 15	<i>— Clear</i>	4 30	<i>Exwater</i>	7 30	<i>Home Head</i>	9 0
<i>Archangel</i>	6 0	<i>— De Four</i>	2 45			<i>Honfleur</i>	9 0
<i>Armentieres</i>	3 0	<i>— of good Hope</i>	3 0	F.		<i>Horn</i>	1 30
<i>Army</i>	1 30	<i>— Sierre de Lion,</i>		FAIR ISLE	0 0	<i>Hull</i>	6 0
<i>Audiern, Fr. Coast</i>	2 15	<i>in Guinea</i>	8 15	<i>Fair Isle Roads</i>	11 15	<i>Humber Mouth</i>	5 15
<i>Auray, Fr. Coast</i>	3 45	<i>Carmarthen Bay</i>	5 15	<i>Falmouth</i>	5 30	<i>Huncliff-Foot</i>	3 45
B.		<i>Caskets without</i>	8 15	<i>Fen, in E. Channel</i>	1 30		
BAJADOR, in Bar-		<i>Caskets within</i>	9 45	<i>Fescan</i>	9 45	I.	
bary	0 0	<i>Catnefs</i>	9 0	<i>Finkmark Coast</i>	1 30	JOHN-DE-LUCE	10 30
<i>Baltimore, Ireland</i>	4 30	<i>Chambernefs</i>	9 45	<i>Flamborough-Head</i>	4 0	<i>Ireland S. Coast</i>	5 15
<i>Barfleur, Fr. Coast</i>	7 30	<i>Cherburgh</i>	7 30	<i>Flanders Banks</i>	1 0	<i>— W. Coast</i>	3 0
<i>Barneville, Fr. C.</i>	7 0	<i>Chili Coast</i>	3 0	<i>Florida</i>	7 30	<i>Jutland Ifle</i>	0 0
<i>Baf without</i>	3 45	<i>Concarneau, Fr. C.</i>	3 0	<i>Flushing</i>	1 0		
<i>Bayonne, Fr. Coast</i>	3 30	<i>Condado</i>	0 0	<i>The Fly</i>	7 30	K.	
<i>Beachy, Eng Coast</i>	0 0	<i>Conquet, Fr. Coast</i>	2 15	<i>Fontenay Race</i>	2 15	KENTISH KNOCK	0 0
<i>Beauvoir, Fr. Coast</i>	3 15	<i>Cork</i>	6 30	<i>Foreland, N. and S.</i>	9 45	<i>Kildive</i>	9 0
<i>Bell Ifle</i>	3 30	<i>Corpus Christi Point</i>	1 30	<i>Forn</i>	5 15	<i>Kilduyn</i>	7 30
<i>Berguer, Holland</i>	1 30	<i>Cowes</i>	10 30	<i>Foulnefs</i>	6 45	<i>Kiltiars</i>	3 0
<i>Bermudas</i>	7 0	<i>Croyl</i>	11 15	<i>Fowey</i>	5 30	<i>Kinsale</i>	5 15
<i>Berwick</i>	2 15	<i>Cromer</i>	7 0	<i>France, W. Coast</i>	3 0	L.	
<i>Biscay Coast</i>	3 0	D.		<i>Firth</i>	11 0	LAMBAY	8 15
<i>Blacknefs</i>	1 30	DARTMOUTH	6 30	<i>Friesland Coast</i>	7 30	<i>Landfend</i>	7 30
<i>Blackney</i>	6 0	<i>David's Head</i>	6 0			<i>Lanion</i>	6 45
<i>Blacktail Beacon</i>	0 15	<i>Deal</i>	10 30	G.		<i>Lawrenefs</i>	4 30
<i>Blanket Race</i>	0 0	<i>Denbigh</i>	2 15	GALLICIA	3 0	<i>Leith</i>	4 0
<i>Blaver</i>	3 0	<i>Defire Port, in A.</i>		<i>Garande</i>	3 0	<i>Lenow</i>	9 45
<i>Bloy</i>	4 30	<i>merica</i>	0 0	<i>Garonne Mouth</i>	3 0	<i>Leofloff</i>	9 45
<i>Bluet without</i>	2 15	<i>Dieppe</i>	10 30	<i>Gascoigne Coast</i>	3 0	<i>— Road</i>	10 30
<i>Bordeaux</i>	5 0	<i>Dort</i>	3 0	<i>Gibraltar Road</i>	0 0	<i>Lime</i>	7 0
<i>Brazil Coast</i>	4 30	<i>Dover</i>	11 30	<i>Gouries Gut</i>	2 30	<i>Lifbon</i>	2 15
<i>Bremen</i>	6 0	<i>Downs</i>	1 15	<i>Goree</i>	1 30	<i>Lizard</i>	7 30
<i>Bree Sound</i>	4 30	<i>Dublin Bar</i>	9 15	<i>Gorend</i>	11 15	<i>London</i>	2 30
<i>Brest</i>	3 15	<i>— Custom house</i>	11 0	<i>Granville</i>	6 45	<i>Longsand Head</i>	10 30
<i>Britain, S. Coast</i>	3 0	<i>Dunbar</i>	2 0	<i>Gravelling</i>	0 0	<i>Loo</i>	6 0
<i>Bridgewater</i>	7 40	<i>Dundee</i>	2 15	<i>Gravefend</i>	1 30	<i>Loir Mouth</i>	3 0
<i>Bridlington Pier</i>	4 0	<i>Dungarven</i>	4 30	<i>Groyne</i>	3 0	<i>Lundey</i>	5 15
<i>Brille</i>	1 30	<i>Dungenefs</i>	9 45	<i>Guernsey</i>	1 30	<i>Lynn, Norfolk</i>	6 0
<i>Bristol</i>	6 30	<i>Dunkirk</i>	12 0	<i>Gunfleet</i>	10 30	<i>Lynn, without</i>	5 15

N. B. 0^h 0^m signifies 12^h 0^m, at Noon, or Midnight.

Alphabetical

Alphabetical TIDE-TABLE continued.

Places NAMES.	Time H. Water. h m	Places NAMES.	Time H. Water. h m	Places NAMES.	Time H. Water. h m	Places NAMES.	Time H. Water. h m
M.		<i>Poictu, S. Coast</i>	3 0	<i>Sandwich</i>	11 30	W.	
<i>MACKWELL'S</i>		<i>Port-Blank</i>	4 15	<i>Scarborough</i>	3 45	<i>Wales</i>	5 15
<i>CASTLE</i>	8 15	<i>Portbus</i>	3 0	<i>Scilly Islands</i>	3 45	<i>Wash (Lincoln-</i>	
<i>Maze</i>	3 0	<i>Portland</i>	8 30	<i>Sedmouth</i>	6 45	<i>shire)</i>	6 30
<i>Magnes Sound</i>	8 15	<i>Portugal Coast</i>	3 45	<i>Senegal</i>	10 30	<i>Waterford</i>	6 30
<i>Malden</i>	12 45	<i>Portsmouth</i>	12 30	<i>Seven-Cliffs</i>	9 0	<i>Weilands</i>	1 30
<i>St. Maloes</i>	6 0	Q.		<i>Seven Isles</i>	4 30	<i>Wells</i>	6 0
<i>Man Isle</i>	9 0	<i>QUEBECK (Cana-</i>		<i>Severn's Mouth</i>	6 0	<i>Weymouth</i>	7 0
<i>Margaret Road</i>	11 15	<i>da)</i>	6 0	<i>Seine's Mouth</i>	9 0	<i>Key</i>	6 45
<i>St. Mark</i>	2 15	<i>Queenborough (In-</i>		<i>Sheerness</i>	0 0	<i>Whitby</i>	3 0
<i>St. Matthew's Point</i>	3 45	<i>Shippy)</i>	0 0	<i>Shelbergb</i>	9 0	<i>Wieringham</i>	7 0
<i>Memifan</i>	3 30	R.		<i>Shetland</i>	3 0	<i>Wight Isle</i>	0 0
<i>Milford, and Moonl's</i>	5 15	<i>RAMEKINS</i>	1 30	<i>Shoe-Bacon</i>	0 30	<i>Winchelsea</i>	12 45
<i>Milford Haven</i>	7 30	<i>Ramsey</i>	5 15	<i>Shoreham</i>	10 30	<i>Winterton</i>	8 0
<i>Morbibam</i>	3 0	<i>Rebdan</i>	12 45	<i>Sleeve</i>	0 0	Y.	
<i>Mount's Bay</i>	4 30	<i>Redsand</i>	0 30	<i>Somme Mouth</i>	11 0	<i>YARM</i>	6 45
<i>Moufe-hole</i>	4 30	<i>Rhee In.</i>	3 0	<i>Sound</i>	3 45	<i>Yarmouth, Norfolk</i>	10 0
N.		<i>Rhodes</i>	11 15	<i>Southampton</i>	0 0	<i>Roads</i>	10 30
<i>NANTZ RIVER</i>	3 0	<i>Robin Hood's Bay</i>	3 0	<i>Spain, W. Coast</i>	3 0	<i>Youghall (Ireland)</i>	4 30
<i>Naze</i>	11 15	<i>Rochel</i>	3 45	<i>Spithead</i>	11 30	Z.	
<i>Needles</i>	10 0	<i>Rochfort</i>	4 15	<i>Spits</i>	0 0	<i>ZELAND COAST</i>	1 30
<i>Newcastle</i>	3 15	<i>Rochester</i>	0 45	<i>Spurn</i>	5 15	<i>Zerick-Sea</i>	3 0
<i>Newport (I. Wight)</i>	0 0	<i>Rohan</i>	3 45	<i>Staples</i>	3 45	N. B. The above TIDE-	
<i>St. Nicholas, in</i>		<i>Rofs</i>	5 15	<i>Start</i>	6 45	TABLE is settled from the	
<i>Ruffia</i>	6 45	<i>Rotterdam</i>	3 0	<i>Stockton</i>	5 15	best Authorities we could	
<i>Nore, W. End</i>	0 0	<i>Rouen</i>	1 15	<i>Swin</i>	0 0	procure; yet it is not im-	
<i>Normandy Coast</i>	10 30	<i>Rumney</i>	1 30	T.		probable the Times of High-	
<i>North Cape, Mag-</i>		<i>Rye</i>	11 15	<i>TEES MOUTH</i>	3 0	<i>Water</i> in some Places are not	
<i>gero</i>	3 0	S.		<i>Tenerif</i>	3 0	exactly as are here set down	
O.		<i>SAINT ANDREWS</i>	2 15	<i>Tenet</i>	1 30	— We shall be obliged to	
<i>OLLONE</i>	3 15	<i>— Augustine (Flo-</i>		<i>Terweer within</i>	0 45	Persons residing at the Pla-	
<i>Orfordness</i>	10 30	<i>rida)</i>	7 30	<i>without</i>	1 30	ces to inform us of any	
<i>Orkneys</i>	6 0	<i>— David's Head</i>	6 0	<i>Tergon</i>	9 45	Mistake; there being no	
<i>Orwell</i>	9 0	<i>— Helen's</i>	10 30	<i>Texel</i>	7 0	Way to come at Certainty	
<i>Ostend</i>	0 30	<i>— John de Lutz</i>	3 30	<i>Texel Cliffs</i>	5 0	in these Matters, but from	
P.		<i>— Lucas</i>	2 15	<i>Thames Mouth</i>	1 30	living Authors, who have	
<i>PENNES, Fr. C.</i>	3 45	<i>— Malps</i>	5 30	<i>Tinmouth</i>	6 0	frequently observed the	
<i>Peru Coast</i>	3 0	<i>— Mark</i>	2 15	<i>Torbay</i>	6 0	Times of highest Tide, at	
<i>Peter Port</i>	8 15	<i>— Matthew's Point</i>	3 45	<i>Treport</i>	10 30	New and Full Moon, in	
<i>Picardy Coast</i>	10 30	<i>— Michael</i>	5 30	V.		different Places.	
<i>Plymouth</i>	6 0	<i>— Nicholas (Russia)</i>	6 45	<i>VANNES</i>	3 45		
<i>Podofemsk, (Russia)</i>	6 45	<i>— Pol de Leon</i>	4 0	<i>Voard</i>	4 30		
		<i>— Powls</i>	6 0	<i>Vreck</i>	0 0		
		<i>— Valleri</i>	9 45	<i>Use</i>	3 0		
		<i>Salcomb</i>	6 0	<i>Ushant within</i>	4 30		

OBSERVATIONS

OBSERVATIONS on the different BEGINNINGS of LONGITUDES.

1. THE Difference of Longitude between the Meridian of an Observatory and some distant Meridian, is chiefly wanted for determining the Longitude, Declination, or right Ascension of the Sun, Moon, or Planet under that second Meridian, from their Data at the first, on a given Day and Year, with also their given diurnal Increase or Decrease.

2. As there must be a Question answered to determine the Difference of Longitude betwixt any two Meridians, whose Longitudes, according to antient Custom, are given from Ferro, the same Question may be as readily answered from the Longitudes East or West, counted from another given Meridian. But if the Longitudes of Places are given at once from the Metropolis or Observatory of any particular Nation, they are ready (without determining them) for the Service of that Nation's Navigation, to ascertain the Longitude, Declination, or right Ascension of the celestial Bodies under a distant Meridian. And on which Account we have given an extensive SEA-COAST TABLE, following, with the Difference of Longitudes from Greenwich, according to modern Custom, as well as the corresponding absolute Longitudes, according to antient Custom, from Ferro, which modern Custom has almost superseded.

Examples of the equal Facility of both these Ways of finding the required Difference of Longitude of Places, are seen in Page 317.

3. Hence future Astronomers and Navigators may compare the Longitudes of Places observed with these we have given from Ferro, or Greenwich, as a near STANDARD of the Truth; and may thence make their future Improvements, or Confirmation of our following Numbers, P. 337, &c.

THE Reviser of the Seaman's Calendar newly rectified, at Page 243, having regretted the Want of an Order of Parliament for causing Agreement among those concerned in settling the Beginning of Longitudes of Places (now confounding the inexperienced Navigators by so many different Countings and Beginnings thereof) we hope our Endeavours here have come up to his Wishes, and answered his Objections.

ASTRONOMICAL QUESTION, of great USE in NAVIGATION.

AT a given Time of the Year, in a given Latitude, the Times being given by a Clock or Watch, when the Sun was observed to have his Forenoon and Afternoon Altitudes alike, to determine from thence, the Time by the same Clock, or Watch, when the Sun passed the Meridian?

The ANSWER.

Put L = the Latitude of the Place, where the two Observations were made.

D = Sun's Distance from the Pole, at Noon.

T = $\frac{1}{2}$ Time (τ) between the two Observations, in Degrees and Decimals.

V = $\frac{1}{2}$ the Variation of the Sun's Declination, in that Time (τ) in Seconds of Time.

M = Middle-Time between the Observations, or $\frac{1}{2} \tau$ added to the Time of Forenoon-Observation.

Let $\frac{s.T \times s.L}{r} = t, A$ like T ; and $B = D$ Dif. A .

Then $\frac{s.B \times s.T}{s.A \times s.D} \times V = N$, Seconds of Time, the Correction or Interval between M , and the true Time of Noon.

And, $M \pm N$ = Time by the Clock or Watch, when the SUN was on the Meridian, according as he is in $\left\{ \begin{array}{l} \overline{\sigma}, \Omega, \pi, \epsilon, \eta, \zeta, \end{array} \right\}$ descending $\left\{ \begin{array}{l} \varphi, \omega, \chi, \gamma, \delta, \Pi, \end{array} \right\}$ ascending $\left\{ \begin{array}{l} \end{array} \right\}$ Signs.

SPHERICAL QUESTIONS answered.

1. The Hypotenuse, and Sum or Diff. of the Legs of a right-angled spherical Triangle being given to find the Triangle?

ANSWER. From twice the Cof. of the Hypoth. subtract the Cofine of Sum or Diff. of Legs, the Remainder will be the Cof. of an Arc, which being added to the said Sum, or Diff. gives the Double of the greater Leg, required.

2. One Leg, and the Sum, or Difference of the Hypotenuse and the other Leg, to find the Hypotenuse?

ANSWER. As Cof. $\frac{1}{2}$ Leg given, : to its Tangent :: so the Cot. $\frac{1}{2}$ Sum of Hyp. and other Leg, : to Tan. $\frac{1}{2}$ their Difference.

3. The $\angle$ at the Base, and Sum or Diff. of the Hypotenuse and Base, given, to find the spherical Triangle?

ANSWER. As Cotangent $\frac{1}{2} \angle$ given, : to its Tangent :: so Sine Sum of the Hyp. and adj. Leg : to Sine of their Difference.

4. The Hyp. and Sum or Diff. of the two adjacent $\angle$ s given, to find the Angles.

ANSWER. As Cot. $\frac{1}{2}$ the Hypotenuse : to its Tangent :: so Cof. Diff. two $\angle$ s : to Sine of the Excess of their Sum above a right Angle.

OBSERVATIONS

OBSERVATIONS on the ANSWERS to MARTIN's or MURDOCH's EXAMPLES, and on the Remarks by *Newtonienfis*, on SPHEROIDAL SAILING, Page 283.
With the ONLY INSTANCE (in a long RUN) wherein SPHEROIDAL SAILING is approved: Which Sailing, in general, having no real ADVANTAGE, the different meridional Parts of the SPHEROID are inserted to prove it, and shew the Extent of its USE.

OUR Correspondent *Newtonienfis* observes, on his first Remarks, Page 296, on B. Martin's Navigation, his Meaning is, That all B. Martin's Examples are like Murdoch's, in respect of their being what never happen in Practice, and that some Examples are numerically the same as what are given by Murdoch. — What is become of this Bray, Page 297, Line 3. Bottom, should be this Bray. — *Newtonienfis* further observes. —

What we had to do with all those Examples was only to shew that the several Cases of Sailing might be differently answered, both according to the Sphere and Spheroid, by our Computers, No. I. II. III. more easily than by the common Methods in Practice; at the same Time we gave Instance, or made Comparison, of our correct Answers with the Answers given by B. Martin to the same Questions or Cases of Sailing. Whether all Mr. Martin's Examples of the Cases of Sailing are the same in all Respects with Mr. Murdoch's, we had no further Regard to, than to shew how much *spheroidal* Navigation differs from itself, according to the different *spheroidal* Figures of the Earth, assumed by B. Martin from *Maupeituis* or *Murdoch*, and from *Juan*: the Form thereof, by the last of which Authors, being considered as much the nearest to Truth, and Sir Isaac Newton's Theory.

The different *spheroidal* meridional Parts, according to *Murdoch* and *Juan*, are inserted in our Work for the Navigator to compare their Use in any Case of Sailing, he chuses to make Trial of, and also to try their Use, respectively, with the meridional Parts of the Sphere. By which he will find that *spheroidal* Navigation, according to *Martin* or *Murdoch*, differs more from *spheroidal* Navigation, according to *Juan*, than *Juan's* differs from the Navigation of the Sphere.

And the only Instance, wherein the *spheroidal* meridional Parts that we have inserted and contrasted with those of the Sphere, are designed to be applied to Practice, for determining the Ship's Place, is when you are sure that your Course and Distance, in a long Run, are both very correct. Which Ship's Place may then serve to compare with the Result of your several daily Reckonings on that Course, according to the common Navigation of the Sphere, and thereby to mark the Difference in Longitude, according to the most correct Form of the terrestrial Spheroid yet discovered. A bare Quotation of a few Numbers or meridional Parts of the Spheroid the most approved, though of equal Authority with the Truth of a greater Number given, could not have evinced their Utility or Inutility, in the Extent thereof, so well as by the 6 half Pages, to every other Minute of the Quadrant, and Length of the Arches to every Degree of Latitude, as we have inserted.

REMARK. Since our printing the 6th and 7th Cases of Mercator-Sailing, we have observed, that the 6th Case and its Variation is printed in the *Miscellanea Curiosa*, for the Months of January, February, and March, 1734, by Mr. John Turner of York, our deceased Friend and Correspondent; being the 7th Question there; though we proposed both those Cases without having observed them elsewhere; nor have we seen either of these Cases proposed in any Work of Navigation but our own, though they are chiefly speculative.

3

VARIATION of CASE 6th of MERCATOR SAILING. (See Page 305.)

A Ship sailed from a Port in Latitude 54° N. upon an unknown Course between the S. and the W. till her Departure was $44^{\circ}.45'$ Miles, and her Difference of Longitude $74^{\circ}.17'$ Miles, required, from thence, her Course, Distance run, and Difference of Latitude?

ANSWERED by the DIRECT METHOD of Solution.

PUT x = Length of the Arch of Latitude arrived in; then, according to Dr. Halley (in Vol. I. *Philos. Transf.*) the meridional Parts of that Latitude will be $= \frac{1}{a} \times x + \frac{1}{6} x^3 + \frac{1}{24} x^5 + \frac{1}{83} x^7 + \frac{1}{262} x^9$, &c. (where a = the Length of the Arch of 1 Min. = ,00029088,

and consequently $\frac{1}{a} = 3438$.) Now, to make all the Parts of the Mercator-Triangle alike, multiply the Departure and also Difference of Longitude by a . Let $44^{\circ}.45' \times a = c$; and $74^{\circ}.17' \times a = d$; and $54^{\circ} \times 60 \times a = b$; $n = 3864$, the merid. Parts for Lat. 54° .

Then, $n - x = \frac{1}{6} x^3 - \frac{1}{24} x^5 - \frac{1}{83} x^7 - \frac{1}{262} x^9$, &c. = merid. Diff. Latitude; the prop. Diff. Lat. = $b - x$; Say, $b - x : c :: n -$

$x - \frac{1}{6} x^3 - \frac{1}{24} x^5$, &c. : d ; whence (by multiplying Extreams and Means) we have, $\frac{bd - dx}{c} = n - x - \frac{1}{6} x^3 - \frac{1}{24} x^5 - \frac{1}{83} x^7 - \frac{1}{262} x^9$, &c.

Put $\frac{bd}{c} = m$, $\frac{-d}{c} = -p$; and $m - px = n - x - \frac{1}{6} x^3$, &c. Then, by Transposition, $n - m = 1 - px + \frac{1}{6} x^3 + \frac{1}{24} x^5 + \frac{1}{83} x^7$, &c.

Put $n - m = q$, $1 - p = f$, and Letters for the known Coefficients of the Powers of x . Then, $fx + rx^3 + sx^5 + tx^7 + gx^9$, &c. = q .

By Reversion of Series $x = \frac{q}{f} - \frac{rq^3}{f^4} + \frac{3r^2 - fs}{f^7} q^5 + \frac{8frs - f^2t - 12r^3}{f^{10}} q^7 + \frac{55r^4 - 55fr^2s + 10f^2rs + fs^2 - f^3g}{f^{13}} q^9$, &c. = ,9231.

Now, x , or the Length of the Arch being known, the corresponding Latitude in Degrees is thus found, by saying, As 1.5708 : 90° :: ,9231 : $52^{\circ}.89' = 52^{\circ}.53'$, the Latitude arrived in. — Whence the proper Dif. of Lat. is 67° : Dep. $44^{\circ}.45'$:: Mer. Lat. 112 : Dif. Long. 74 Miles; And Distance run 80 Miles; Course (by Computer, No. I.) $33^{\circ} 7' \frac{1}{2}$ S. Westerly.

The indirect Method of Solution, by Trial and Error, (see Page 305) is preferable.

3

3

CASE 6th of MERCATOR-SAILING. (See Page 305.)

A Ship sails from a Place in Latitude 54° N. Distance 80 Miles, and finds her Dif. of Long. to be $74^{\circ}.17'$ Miles, required her Course, Departure, and Difference of Latitude?

Answered by the DIRECT METHOD of Solution.

In this Case, by 47. e. 1. (taking the same Letters as in the former Case) As $b - x$ proper Dif. Lat. : $\sqrt{c^2 - b^2 + 2bx - x^2}$ the Departure :: $n - x - \frac{1}{6} x^3 - \frac{1}{24} x^5$, &c. Mer. Dif. Lat. : d , Dif. Longitude. Whence, we have an infinitesimal Equation to be squared, and then

the Value of x is had by Reversion of infinite Series, if any Body chuses to take the Time and Trouble.

The indirect Method of Solution, by Trial and Error (Page 305) is preferable.

3

The MARINER'S SEA-COAST TABLE.

Of the principal Ports, Harbours, Head-Lands, Bays, Rivers-Mouths, and Islands, lying contiguous round the WORLD. With their Latitudes, North or South, and Longitudes East from Ferro Island; also their Difference of Longitude East or West, from GREENWICH OBSERVATORY. According to the best Authorities, and some* according to the correctest Observations.

* The Latitudes and Longitudes of Places marked with a * are from correct and repeated Observations. The Rest are improveable, or want Confirmation.

- | | | |
|------------------------------|-----------------------|-----------------------|
| I. The Coast of ENGLAND | } EUROPE. | |
| II. ——— SCOTLAND | | |
| III. ——— IRELAND | | |
| IV. ——— HOLLAND and FLANDERS | | |
| V. ——— FRANCE and PORTUGAL | | |
| VI. ——— The MAIN CONTINENT | | |
| VII. ——— The Islands | | } within the Straits, |
| VIII. ——— BARBARY and GUINEA | | |
| IX. ——— The Western | | } AFRICA. |
| X. ——— The Canary | | |
| XI. ——— The Cape de Verd | | |
| XII. ——— The Southern | | |
| XIII. ——— The Main CONTINENT | | |
| XIV. ——— The Islands | } in the East } ASIA. | |
| | | |

- | | | |
|---|------------|-----------|
| XV. The Coast of AMERICA in the South-Sea, from California to Cape Horn | } AMERICA. | |
| XVI. ——— BRAZIL in South America, from Cape Horn to Cape Roque | | |
| XVII. ——— The Main CONTINENT of the West Indies | | |
| XVIII. ——— The Caribbee | | } ISLANDS |
| XIX. ——— The Bahama | | |
| XX. ——— CAROLINA, VIRGINIA, MARYLAND, PENNSYLVANIA, NEW ENGLAND, and NEWFOUNDLAND | | |
| XXI. ——— HUDSON'S BAY and the Straits | | |
| XXII. ——— ICELAND, GREENLAND, NOVA ZEMBLA, and the Northern ISLES | | |
| XXIII. ——— The SOUND and BALTIC, EUROPE. | | |

Places Names.	Lat. N.	Long. fr. Ferro.	D. Long. fr. Gr. E. or W.	Places Names.	Lat. N.	Long. fr. Ferro.	D. Long. fr. Gr. E. or W.	Places Names.	Lat. N.	Long. fr. Ferro.	D. Long. fr. Gr. W.
I. COAST of ENGLAND.								II. COAST of SCOTLAND.			
1. BERWICK	55 48	15 55	1 45 W	36. Rye	51 2	18 20	0 40 E	72. Lancaster	54 42	12 59	4 41 W
2. Newcastle	55 12	16 5	1 35 W	37. Beachy	50 46	18 0	0 20 E	73. White-Haven	54 17	14 5	3 35 W
3. Shields	55 2	16 12	1 28 W	38. Shoreham	50 55	17 18	0 42 W	74. Isle of Man, W. End . . .	53 45	12 35	5 5 W
4. Sunderland	54 55	16 15	1 25 W	39. Portsmouth	50 49	16 35	1 5 W	75. Holy Head	53 24	12 45	4 55 W
5. Hartlepool	54 43	16 39	1 1 W	40. I. Wight, Newport . . .	50 42	16 10	1 30 W	76. Carlisle	54 47	14 30	3 10 W
6. Stockton	54 33	16 10	1 30 W	41. Pool	50 56	15 41	1 59 W				
7. Whitby	54 26	16 45	0 55 W	42. Weymouth	50 42	14 55	2 45 W	1. SKY ISLAND, N. End . . .	57 45	11 50	5 50 W
8. Scarborough	54 20	16 55	0 45 W	43. Portland	50 30	14 51	2 49 W	2. St. Kilday	57 52	7 50	9 50 W
9. Humble-Head	54 8	17 46	0 6 E	44. Chiddock	50 46	14 35	3 5 W	3. Island of Lewis, N. End . . .	58 20	10 35	7 5 W
10. Burlington	54 0	17 43	0 3 E	45. Lime	50 45	14 20	3 20 W	4. Farro-Head, or C. Wrath . . .	58 34	12 25	5 15 W
11. Spurn	53 45	17 48	0 8 E	46. Topham	50 38	14 7	3 33 W	5. Sbetland, S. End	60 4	15 35	2 5 W
12. Hull	53 52	17 15	0 25 W	47. The Berry	50 25	13 46	3 54 W	6. Fair-Isle	59 30	15 15	2 25 W
13. Grimsby	53 30	18 31	0 51 E	48. Torbay	50 32	13 56	3 44 W	7. Isles of Orkney	59 10	14 13	3 27 W
14. Boston	53 10	18 0	0 20 E	49. Dartmouth	50 26	13 57	3 43 W	8. Cathness Point	58 40	14 59	2 41 W
15. Lynn	52 55	18 7	0 27 E	50. The Start	50 7	13 48	3 52 W	9. Buchanefs	57 45	16 17	1 23 W
16. Wells	53 10	18 35	0 55 E	51. The Eddystone	50 10	13 11	4 29 W	10. Aberdeen	57 22	15 45	1 45 W
17. Blakeny	53 8	18 30	0 50 E	52. Plymouth	50 31	13 18	4 22 W	11. Dundee	56 28	14 45	2 45 W
18. Cromer	53 10	18 40	1 0 E	53. Ramhead	50 16	13 0	4 40 W	12. Leith	56 0	14 39	3 1 W
19. Winterton	52 57	18 57	1 17 E	54. Foy	50 24	13 0	4 40 W	13. EDINBURGH*	55 58	14 35	3 5 W
20. Yarmouth	52 42	19 13	1 33 E	55. Falmouth	50 11	12 25	5 15 W	14. Glasgow	55 52	13 30	4 10 W
21. Aldborough	52 16	19 3	1 23 E	56. LIZARD*	49 55	12 55	4 45 W				
22. Orfordness	52 12	18 51	1 11 E	57. Land's End	50 0	11 35	6 5 W	III. COAST of IRELAND.			
23. Ipswich	52 10	18 40	1 0 E	58. Gulf-Rock	49 56	11 49	6 11 W	1. FAIR FORE-LAND . . .	55 5	11 5	6 35 W
24. Harwich	52 5	18 53	1 13 E	59. SCILLY Islands, St. Mary's	50 0	10 50	6 50 W	2. Londonderry	55 0	9 45	7 55 W
25. Colchester	51 58	18 31	0 51 E	60. Seven Stones	50 10	10 55	6 45 W	3. Belfast	54 39	11 5	6 35 W
26. LONDON*	51 31	17 35	0 5 W	61. Hartland Point	51 10	13 0	4 40 W	4. Island-Torr	55 9	9 5	8 35 W
27. Rochester	51 28	18 1	0 21 E	62. Lundy Isle	51 20	12 55	4 45 W	5. Isles of Arran	54 8	8 36	9 4 W
28. Margate	51 27	18 49	1 9 E	63. Mort-Bay, or Point . . .	51 12	12 55	4 45 W	6. Stags of Broad Haven . . .	54 7	7 29	10 11 W
29. North Foreland	51 27	18 59	1 19 E	64. Bristol	51 33	13 0	4 40 W	7. Isles of Arrian	54 55	7 45	9 55 W
30. Sandwich	51 25	18 49	1 9 E	65. Swandsey	51 40	13 10	4 30 W	8. Galway	53 7	7 55	9 45 W
31. The Downs	51 23	19 0	1 20 E	66. Caldy-Island	51 33	12 21	5 19 W	9. Gall, or Doen's Head . . .	52 40	8 5	9 35 W
32. South Foreland	51 10	18 55	1 15 E	67. Milford	51 45	12 20	5 20 W				
33. Ripples-Sand	51 53	19 1	1 21 E	68. St. David's Head . . .	52 0	12 13	5 27 W				
34. Dover	51 6	18 53	1 13 E	69. Bardsey-Isle	52 44	12 35	5 5 W				
35. Dongennefs	50 57	18 26	0 46 E	70. Liverpool	53 20	14 35	3 5 W				
				71. Westchester	53 37	13 15	4 25 W				

The MARINER'S SEA-COAST TABLE.

Places Names.	Lat. N.	Long. fr. Ferro.	D. Long. fr. Gr. E. or W.	Places Names.	Lat. N.	Long. fr. Ferro.	D. Long. fr. Gr. E. or W.	Places Names.	Lat. N.	Long. fr. Ferro.	D. Long. fr. Gr. E. or W.
III. COAST of IRELAND continued.											
10. Lupis's Head	52 24	7 20	10 20 W	20. West Penmark	47 48	13 11	4 29 W	28. Cape St. Angelo, or Angulo	36 32	41 31	23 51 E
11. Limerick	52 23	8 0	9 40 W	21. Bell-Iyle	47 18	14 19	3 21 W	29. Athens*	38 5	41 32	23 52 E
12. Galway	53 7	7 55	9 45 W	22. Nantz*	47 13	15 56	1 44 W	30. Cape Martelo, S. Point of Negro-			
13. Sline Head	53 20	5 40	11 20 W	23. Island Dieu	46 34	15 22	2 18 W	pont	38 7	42 38	24 58 E
14. Blasques	52 0	5 39	12 1 W	24. Hey's Isle	46 24	15 21	2 19 W	31. Cape Colonne	37 45	42 17	24 37 E
15. Shilloks	51 30	5 40	12 0 W	25. Island de Rey, the Middle				32. Salonica	40 41	40 48	23 8 E
16. Cow and Calf	51 22	6 59	10 41 W	26. Isles of Oleron	46 10	16 5	1 35 W	33. Cape Monte Sancto . . .	49 26	42 37	24 57 E
17. Mizen-Head	51 16	6 5	11 35 W	27. Rochel*	46 10	16 24	1 16 W	34. Gallipoli	40 33	44 55	27 15 E
18. Old Head of Kinsale . . .	51 35	8 37	9 3 W	28. Bourdeaux*	44 50	17 6	0 34 W	35. Constantinople*	41 0	46 33	28 53 E
19. Cape Clear	51 17	6 25	11 15 W	29. St. Sebastian	43 24	16 8	1 32 W	36. Cape Barbador, or Baba . . .	39 30	44 5	26 25 E
20. Kinsale	51 50	7 55	9 45 W	30. Bilboa	43 29	14 37	3 3 W	37. Smyrna*	38 28	45 0	27 20 E
21. Cork	51 49	8 5	9 35 W	31. Cape Pinas	43 56	11 35	6 5 W	38. Cape Babernola, or Blanc . . .	38 9	44 7	26 27 E
22. Waterford	52 9	8 55	8 45 W	32. — Ortegual	44 4	9 47	7 53 W	39. Ephesus	38 1	45 28	27 48 E
23. Wexford	52 13	10 8	7 32 W	33. — Corumna or Groin	43 28	8 15	9 25 W	40. Antiochetta	36 30	50 21	32 41 E
24. DUBLIN	53 12	10 39	7 1 W	34. — Finifter	43 12	7 55	9 45 W	41. Scanderoon, or Alexandretta	36 34	54 5	36 25 E
25. Lambay-Island	53 24	12 5	7 35 W	35. Isles of Bajo- na	42 29	8 8	9 32 W	42. Antiocha	36 11	54 0	36 20 E
IV. COAST of HOLLAND and FLAN- DERS.				36. Oporto	41 10	8 10	9 30 W	43. Aleppo*	35 45	55 0	37 20 E
1. SCAM	57 30	27 55	10 15 E	37. Burlings	39 35	8 11	9 29 W	44. Tortosa	35 0	54 5	36 25 E
2. Holy Land, or Helighland Isle	54 24	26 10	8 30 E	38. Rock of Lif- bon	38 52	7 45	9 55 W	45. Tripoly	34 38	53 50	36 10 E
3. Hamburgh	53 41	28 15	10 35 E	39. Lisbon*	38 42	8 42	8 58 W	46. Joppa, or Jaf- fa	32 27	53 55	35 15 E
4. Biemen	53 30	26 35	8 55 E	40. Cape St. Vin- cent	36 53	8 29	9 11 W	47. Jerusalem*	31 55	53 0	35 20 E
5. Embden	53 5	25 10	7 30 E	41. Cape St. Maria	36 58	9 5	8 35 W	48. Alexandria*	31 7	47 57	30 17 E
6. Ameland-Island	53 30	23 55	6 15 E	42. Cadiz*	36 33	11 38	6 7 W	49. Cape Rufato	32 48	39 0	21 20 E
7. Scheling	53 27	23 33	5 53 E	43. Cape Trefalgar	36 10	11 34	6 6 W	50. Cape Mesura- to	32 21	33 52	16 12 E
8. The Fly	53 16	23 5	5 25 E	VI. COAST on the Main CONTINENT within the STRAITS.				51. Tripoly*	32 54	30 45	13 5 E
9. The Texel	53 15	22 45	5 5 E	1. GIBRALTAR	36 12	12 42	4 58 W	52. Sufa	35 53	28 9	10 29 E
10. AMSTERDAM*	52 23	22 39	4 59 E	2. Malaga	36 48	13 45	3 55 W	53. Cape Bona	37 3	28 39	10 59 E
11. Rotterdam	51 55	22 5	4 25 E	3. Cape de Gat	36 40	15 55	1 45 W	54. Tunis	36 50	27 52	10 12 E
12. Antwerp	51 10	21 55	4 15 E	4. Cape Paul	38 15	17 40	0 0 E	55. Bona	37 2	25 54	8 14 E
13. The Brill	52 0	21 35	3 55 E	5. Alicant.	38 35	17 50	0 10 E	56. Seven Capes	37 15	24 35	6 55 E
14. Middleburgh in Zealand	51 37	21 33	3 53 E	6. Cape St. Mar- tin	38 46	18 15	0 35 E	57. Gigeri	37 14	23 50	6 10 E
15. Sluys	51 14	21 18	3 38 E	7. Barcelona*	41 26	19 53	2 13 E	58. Cape Tidelles, or Dellys . .	37 15	21 53	4 13 E
16. Oflend*	51 14	20 36	2 56 E	8. Marseilles*	43 18	23 2	5 22 E	59. Algier*	36 50	20 51	3 11 E
17. Dunkirk*	51 2	20 2	2 22 E	9. Toulon*	43 7	23 37	5 57 E	60. Cape Tenes	36 50	19 25	1 45 E
V. COAST of FRANCE and PORTUGAL.				10. Genoa	44 25	26 18	8 38 E	61. Oran	36 2	18 1	0 21 E
1. CALAIS*	50 58	19 29	1 49 E	11. Leghorn	43 28	28 10	10 30 E	62. Cape de Tres Forcas . . .	35 30	15 31	2 9 W
2. Dieppe	49 56	18 44	1 4 E	12. Civita Vecchia	42 10	30 0	12 20 E	63. Ceuta	35 54	12 50	4 50 W
3. St. Valery	50 10	18 31	0 51 E	13. Rome*	41 54	30 9	12 29 E	64. Tangier	35 42	12 13	5 27 W
4. Sain - Head, or Cape de Antifer	49 44	18 9	0 29 E	14. Naples*	40 51	32 20	14 40 E	65. Tetuan	35 27	12 29	5 11 W
5. Rouen Mouth	49 34	18 5	0 25 E	15. Cape Spartaven- to	37 55	34 30	16 50 E	VII. ISLANDS within the STRAITS.			
6. Cape Barfleur	49 43	16 21	1 19 W	16. Cape Colonne	38 56	35 0	18 0 E	1. ALBORAN	35 54	15 6	2 34 W
7. CapedelaHogue	49 47	15 35	2 5 W	17. Gallipoli	39 56	36 16	18 36 E	2. Formentura	38 33	19 30	1 50 E
8. Alderney	49 50	15 23	2 17 W	18. Cape St. Maria or Lucia	39 45	36 35	18 55 E	3. Ivica	38 50	19 15	1 35 E
9. Caskets	49 50	15 15	2 25 W	19. Ancona	43 40	32 1	14 21 E	4. Majorca	39 30	20 38	2 58 E
10. Guernsey	49 33	14 55	2 45 W	20. Venice*	45 25	29 45	12 5 E	5. Minorca City	39 51	22 27	4 47 E
11. Jersey	49 15	15 15	2 25 W	21. Zara	44 30	34 10	16 30 E	6. C. To- lare } End of } Sar- parde, N. } dinia	38 46	26 47	9 7 E
12. St. Malou*	48 39	15 38	2 2 W	22. Ragusa	42 55	36 25	18 45 E	7. C.S. Re- parde, N. } dinia	41 10	27 25	9 45 E
13. Morlaix	48 33	13 46	3 54 W	23. Cattaro	42 47	36 52	19 12 E				
14. Island de Bas	48 50	13 35	4 5 W	24. La Valona	40 45	38 0	20 0 E				
15. Ushant	48 30	12 33	5 7 W	25. Point Palerma	40 0	37 50	20 10 E				
16. Conquet	48 27	12 35	5 5 W	26. Lepanto	38 10	40 27	22 47 E				
17. Brest*	48 23	13 9	4 31 W	27. Cape Matapan or Caliga . .	36 33	40 16	22 36 E				
18. Camarita Bay	48 25	13 7	4 33 W								
19. Seams	48 2	12 35	5 5 W								

The MARINER'S SEA-COAST TABLE.

Places Names.	Lat. N.	Long. fr. Ferro.	D. Long. fr. Gr. E.	Places Names.	Lat. N.	Long. fr. Ferro.	D. Long. fr. Gr. E. or W.	Places Names.	Lat. S.	Long. fr. Ferro.	D. Long. fr. Gr. E. or W.
VII. ISLANDS within the STRAITS continued.				54. } C. Sygre Mety-line. P. O-liva	39 15	43 40	26 0 E	29. Cape Lopar	0 55	27 30	9 50 E
8. Port Mahon, Minorca . .	39 42	21 47	4 7 E	55. } Sciatto, or Schate	38 57	44 15	26 55 E	30. Cape Negro	16 8	30 6	12 26 E
9. Galitia	37 41	26 19	8 39 E	56. Scio or Xio	38 22	43 47	26 7 E	31. River Congo	5 40	33 0	15 20 E
10. Sardinia, S. End	38 46	26 47	9 7 E	57. Patmos	37 20	44 30	26 50 E	32. Angola	8 57	33 31	15 51 E
11. Asinaria	41 3	26 43	9 3 E	58. Zino	37 35	43 11	25 31 E	33. Cape St. Thomas . . .	24 10	32 18	14 38 E
12. Corfica, S. End	41 24	27 21	9 41 E	59. Zio, or Sea	37 37	42 25	24 45 E	34. Cape Secos, Seco, or Seca . .	29 0	33 31	15 51 E
13. Corfica, N. End	41 24	27 21	9 41 E	60. Fermina	37 24	42 33	24 53 E	35. Cape of Good Hope	34 15	34 40	17 0 E
14. Cape Corfo	42 56	27 25	9 45 E	61. Sifanto, or Sorfante	36 58	42 55	25 15 E	36. Cape Aguilhas or Lagullas . .	34 54	38 55	21 15 E
15. Gorgona	43 34	27 13	9 33 E	62. Millo, Mila, or Melo	36 40	42 35	24 55 E				
16. Capraia, or Cap-tia	43 3	28 2	10 22 E	63. W of End C. St. John	35 15	41 35	23 55 E	IX. The WESTERN ISLANDS.			
17. Lilboa, or Elba	42 45	28 35	10 55 E	64. E Can. C. So-lomon	35 0	44 43	27 3 E		N.	0 /	0 /
18. Planofa	42 30	28 24	10 44 E	65. Rho- } N End	36 27	46 13	28 33 E	1. Corvo	39 54	346 40	31 0 W
19. M. Christo	42 17	28 35	10 55 E	66. } S End	36 1	46 3	28 23 E	2. Flores (Azores)	32 32	346 41	30 59 W
20. Palmarolla, or Palmeria . .	41 3	31 10	13 30 E	67. C. Tranquil	36 1	46 3	28 23 E	3. Fyal	38 53	349 20	28 20 W
21. Ponfa	41 0	31 12	13 32 E	68. } C. Baf-	34 57	49 58	32 18 E	4. Pico	38 40	350 15	27 25 W
22. Ifchia	40 54	32 5	14 25 E	69. } SW End of C. St. Andrea	35 31	52 35	34 55 E	5. St. George	38 52	351 32	26 8 W
23. Strombello	38 58	33 35	15 55 E	70. } S Cy-prus C. de Gatte	34 30	50 45	33 5 E	6. Tercera	38 57	352 1	25 39 W
24. Volcanello	38 27	33 8	15 28 E					7. Gratiofa	39 16	350 34	27 6 W
25. Fellicur	38 30	32 35	14 55 E					8. Whale Fish	39 37	352 24	25 16 W
26. Allicur	38 28	32 17	14 37 E					9. Vajo, or Vejia	38 43	353 47	23 53 W
27. Uffica	38 36	31 27	13 47 E					10. St. Michael	38 6	353 59	23 41 W
28. Trepano, W. End of Sicily	37 57	30 48	13 8 E	VIII. COAST of BARBARY and GUINEA.				11. Formigas, or Hormigas	37 40	354 48	22 52 W
29. Palermo in Sicily	38 6	31 25	13 45 E	1. CAPE SPAR-TEL	35 50	11 46	5 54 W	12. St. Mary's	36 59	353 57	23 43 W
30. C. Passaro, W. End of Sicily	36 38	33 15	15 35 E	2. Salle	33 51	10 10	6 30 W				
31. Messina, in Sicily	38 7	33 55	16 15 E	3. Cape Can'in	32 36	8 25	9 15 W	X. CANARY ISLANDS.			
32. Maritimo	38 12	34 45	17 5 E	4. Cape de Geer	30 27	7 29	10 11 W		N.	0 /	0 /
33. Syratufe, in Sicily	37 3	33 20	15 40 E	5. Cape de Non	28 15	6 31	11 9 W	1. SANCTO Porto	32 58	1 41	15 59 W
34. C. Molin, in Sicily . . .	37 42	33 31	15 51 E	6. Cape Bajadore	26 4	2 0	15 40 W	2. Maderia W. End	32 23	0 9	17 31 W
35. Cape Passaro	36 38	33 15	15 35 E	7. Cape Olerado	23 41	1 40	16 0 W	3. Salvages	30 0	0 45	16 55 W
36. Pantalania	36 48	29 55	12 15 E	8. Cape Blanco	20 32	0 0	17 40 W	4. Palma	28 40	359 59	17 41 W
37. Limofa	35 56	30 46	13 6 E	9. Senegal	15 28	1 5	16 35 W	5. FERRO*	27 54	0 0	17 40 W
38. Lampidofa	35 34	30 21	12 41 E	10. Cape de Verd*	14 43	0 30	17 10 W	6. Gomero	28 6	0 30	17 10 W
39. Malta*	35 54	32 9	14 29 E	11. Mouth of River Gambia . . .	13 8	2 4	15 36 W	7. Pico Tenerif	28 23	0 7	16 33 W
40. Ouore l'oeil, a Rock . . .	35 45	34 11	16 31 E	12. Sierra Leona	8 36	4 38	13 2 W	8. Grand Canary	27 52	1 25	15 15 W
41. Grande, or I. Longo . .	44 25	33 53	16 13 E	13. Cape de Monte or Mount . .	6 23	5 35	12 5 W	9. Allegranfa	29 20	5 2	12 38 W
42. Poma	43 10	33 42	16 2 E	14. Monferado	6 5	7 33	10 7 W	10. Lancerota	29 2	4 50	12 50 W
43. St. Andrea	43 12	34 1	16 21 E	15. River Sester or Sefos	5 24	8 59	8 41 W	11. Forteventura, S. W. End	28 5	3 59	13 41 W
44. Leffa	43 6	34 15	16 35 E	16. Cape de Palmas	4 13	10 50	6 50 W				
45. Agofa	42 58	34 55	17 15 E	17. River St. Andrea	5 5	13 20	4 20 W	XI. CAPE DE VERD ISLANDS.			
46. Corzola	43 3	35 7	17 27 E	18. Cape 3 Points	4 28	15 45	1 55 W		N.	0 /	0 /
47. Melada	42 52	35 41	18 1 E	19. Jaque Jaque	4 16	7 48	9 52 W	1. St. ANTONIO	17 24	352 33	25 7 W
48. Corphu	39 42	37 41	20 1 E	20. Affene	4 15	10 18	7 22 W	2. St. Vincent	17 4	352 56	24 44 W
49. Cephalonia	38 15	38 35	20 55 E					3. St. Lucia, or round Island	17 5	352 55	24 45 W
50. Zante	37 46	38 49	21 9 E	21. Cape Corfe	4 55	19 35	1 55 E	4. St. Nicholas	16 50	353 37	24 3 W
51. Sapiensa	36 40	39 20	21 40 E	22. River de Volta, or Actara . .	5 55	21 0	3 20 E	5. Isle de Sal	16 50	355 27	22 13 W
52. Rhodes City*	36 42	45 40	28 0 E	23. River Formosa	7 0	24 15	6 35 E				
In the ARCHIPELAGO.				24. Cape Formosa	4 15	24 15	6 35 E				
53. Taffo	40 40	42 52	25 12 E	25. New Calabar Ent.	4 42	26 8	8 28 E				
54. Lemnos	39 59	43 12	25 32 E	26. Old Calabar Ent.	4 10	27 20	9 40 E				
				27. River Camerones	3 25	27 45	10 5 E				
				28. River Angra	0 50	27 36	9 56 E				

The MARINER'S SEA-COAST TABLE.

Places Names.	Lat. N.	Long. fr. Ferro.	D. Long. fr. Gr. E. or W.	Places Names.	Lat. N.	Long. fr. Ferro.	D. Long. fr. Gr. E.	Places Names.	Lat. N.	Long. fr. Ferro.	D. Long. fr. Gr. E.
XI. CAPE DE VERD ISLANDS continued.				24. Cape Glado	0 1	0 1	0 1	78. Point Romania	0 1	0 1	0 1
6. Bonavista	16 5	355 28	22 12 W	25. River de Sinda	25 50	81 9	63 29 E	79. Point Cui	1 16	120 50	103 10 E
7. Mayo, or Island	15 14	355 33	22 7 W	26. Diu Head	24 55	85 10	67 30 E	80. Siam Entrance	12 10	116 37	98 57 E
8. St. Jago	15 8	354 39	23 1 W	27. Surat*	21 2	87 25	69 45 E	81. Camboida En-	14 18	118 30	100 50 E
9. Fuego, or Fulgo.	14 50	353 54	23 46 W	28. Deman	21 10	90 0	72 20 E	82. Cape Anarilla,	10 28	122 35	104 55 E
10. Bravas	14 28	353 33	24 7 W	29. Bombay Island	20 6	91 5	73 25 E	or Avarilla	13 25	125 38	107 58 E
11. St. Paul	1 20	352 5	25 35 W	30. St. John's	19 18	90 41	73 1 E	83. Cochín, or Chin-	14 5	125 31	107 51 E
XII. SOUTHERN ISLANDS.				31. Chaul, or	19 55	90 55	73 15 E	chen	22 13	131 26	113 46 E
	N.			Choale . . .	18 31	91 12	73 32 E	84. Macao*	20 50	123 15	105 35 E
1. FERNANDE POO	2 40	28 5	10 25 E	32. Dabul	18 3	91 35	73 55 E	85. Canton*	23 8	130 43	113 3 E
2. De Prince	1 40	26 50	9 10 E	33. Rajapour Isle	17 2	91 45	74 5 E	87. Amoy, or Que-	24 35	134 25	116 45 E
3. St. Thomas	0 0	25 55	8 15 E	34. Goa*	15 31	91 25	73 45 E	moy Island	26 30	135 55	118 15 E
4. St. Matthews	S.			35. Carwar	14 47	92 35	74 55 E	88. Hockfew	27 51	136 25	118 45 E
5. Annabona	2 10	25 2	7 22 E	36. Manqueloire	12 53	93 0	75 20 E	89. River Swadia	29 59	138 10	120 30 E
6. Ascension	7 20	4 30	13 10 W	37. Tellicherry	11 42	93 0	75 20 E	90. Liampo, Lingpo,	30 0	138 10	120 30 E
7. St. Helena Nova	16 c	23 49	6 9 E	38. Calicut	11 16	93 5	75 25 E	or Ningpo	33 45	137 36	119 56 E
8. St. Helena*	15 56	12 25	4 15 W	39. Cannanou	10 22	93 10	75 30 E	92. NANQUIN*	39 54	134 3	116 23 E
9. Tristia de A-	37 5	3 45	13 55 W	40. Cochín	9 54	93 30	75 50 E	93. PEKIN*			
cuntia . . .				41. Anjanga	8 29	94 0	76 20 E				
XIII. Main CONTINENT in the EAST INDIES.				42. Cape Comerin	7 50	95 0	77 20 E	XIV. ISLANDS in the EAST INDIES.			
	S.			43. Colombo, in	7 7	97 5	79 25 E		S.		
1. BAY de ALLO-	0 1	0 1	0 1	Zeylon . . .	6 10	97 45	80 5 E	1. St. Paul	38 20	93 0	75 20 E
GO or Dallagoa	25 50	48 40	31 0 E	44. Point de Galle,	6 2	98 28	80 40 E	2. Amsterdam Isle	38 40	90 15	72 35 E
2. River St. Lucia	28 20	49 52	32 12 E	or Gallo, on the	8 40	99 15	81 35 E	3. Romeras de Caf-	38 45	94 52	67 12 E
3. Cape St. Martin	22 40	52 40	35 0 E	same	6 23	99 20	81 40 E	4. St. Joan de Lif-	25 24	71 5	53 25 E
or Mary	23 40	53 52	36 12 E	45. Dunder-Head,	9 50	97 40	80 0 E	bon	19 50	79 5	61 25 E
4. Cape Corientes	15 4	58 45	41 5 E	on the same	11 1	97 30	79 50 E	5. Diego Roys	16 38	82 5	64 25 E
5. Mofambique	10 17	57 45	40 5 E	46. Trincomale	11 15	97 25	79 45 E	6. St. Brandon	20 10	70 30	52 50 E
6. P. de Aguada,	9 0	56 55	39 15 E	47. Great Bassas	11 45	97 19	79 39 E	7. Mauritius	11 15	78 5	60 25 E
or Del Gada	4 50	56 34	38 54 E	Shoals . . .	11 54	97 53	80 13 E	8. Malha	25 47	60 45	46 5 E
7. Cape de Falso	3 50	56 5	38 25 E	48. Jetrapatam	12 5	97 23	79 43 E	9. C. St. Ma- End	12 10	68 40	51 0 E
8. Tongon	2 58	57 10	39 30 E	49. Negrapatam	12 35	97 40	80 0 E	ry, S. of			
9. Mombafo	1 20	57 48	40 8 E	50. Trincumbar	13 11	98 7	80 27 E	10. C. D'Am- Ma-			
10. Molinde, or	0 41	58 50	41 10 E	51. Porta Nova	13 30	97 57	80 17 E	bre, N. da-			
Melinde . .				52. Pondicherry*	14 16	97 47	80 7 E	[gascar.			
11. River Lamos				53. Fort St. David,	16 6	98 17	80 37 E	11. Terra del Ma-	19 29	62 10	44 30 E
12. River de Fuego				or Tregapatam	16 8	99 7	81 27 E	dagas	17 21	60 55	43 15 E
13. Magadoxa	2 20	62 25	44 45 E	54. Congmere	16 22	98 45	81 5 E	12. St. John de No-	17 36	61 15	43 35 E
14. Cape de Bassas,	4 6	65 13	47 33 E	55. Fort St. George	16 30	99 32	81 52 E	va	13 10	63 13	45 33 E
or Boxos . .	11 44	68 55	51 15 E	or Madrasappa-	17 43	101 32	83 52 E	13. St. Christova	12 10	62 38	44 58 E
15. Cape Gardefoy	13 0	64 35	46 55 E	tam	18 49	102 50	85 10 E	14. Mayetto	12 5	61 58	44 18 E
16. Aden	14 10	62 25	44 45 E	56. Pullicat	19 51	104 17	86 37 E	15. Joanna	11 40	61 25	43 45 E
17. Mocha	15 23	69 45	52 5 E	57. Armegon	29 11	104 58	87 18 E	16. Mohilla	8 7	57 50	40 10 E
18. Cape Matriaca	17 0	73 10	55 30 E	58. Petapoly	20 42	105 27	87 47 E	17. Comero, or An-	6 48	57 32	39 52 E
19. Defar	22 41	77 20	59 40 E	59. Diu Point, or	21 16	105 23	87 43 E	gazecha	5 20	57 10	39 30 E
20. Cape Refulgat	23 32	77 20	59 40 E	Mecha . . .	21 25	105 32	87 52 E	18. Morfia or Mon-	10 30	62 14	44 34 E
21. Cape Muea, or	29 45	66 55	49 15 E	60. Masulapatam	23 9	106 40	89 0 E	sia	10 14	68 52	51 12 E
Mufcat . . .	27 20	74 15	56 35 E	61. Masipore	23 57	108 30	90 50 E	19. Zanzibar	9 30	70 16	52 36 E
22. Bufero				62. Visagapatam	25 6	106 20	88 40 E	20. Penda	9 55	71 25	53 45 E
23. Gomboroon				63. Birmilapatam	22 17	109 56	92 16 E	21. Comero			
				64. Pondy	20 10	111 15	93 35 E	22. Cosmedelo			
				65. Jacarnaut Pa-	18 35	113 55	96 15 E	23. Juan de Nova			
				god	1 16	119 45	102 5 E	24. Aftore Isle			
				66. Arsipare	2 5	119 5	101 25 E	25. Agalega, or Gal-			
				67. Point Palmiras				lega			
				68. Balafore Road				26. Setta Harna-			
				69. Fipley				nes			
				70. Hughly							
				71. Dacca							
				72. Cafimbazar							
				73. River Bengal							
				74. River Aracan							
				75. Pegu							
				76. Malacca*							
				77. Formosa							

The MARINER'S SEA-COAST TABLE.

Places Names.	Lat. S.	Long. fr. Ferro.	D. Long. fr. Gr. E.	Places Names.	Lat. N.	Long. fr. Ferro.	D. Long. fr. Gr. E. or W.	Places Names.	Lat. S.	Long. fr. Ferro.	D. Long. fr. Gr. W.
XIV. ISLANDS in the EAST INDIES Continued.				XV. COAST of AMERICA in the SOUTH-SEA, from California to CAPE-HORN.				XVII. COAST of the Main CONTINENT in the WEST INDIES.			
27. Quevelo, Que- bello	0 1	0 1	0 1	74. Celebes, N. End	1 40	139 55	121 15 E	13. River Grande Entrance	0 1	0 1	0 1
28. Bassas de Ban- has	3 53	70 11	52 31 E	75. Meridano, W. Part	6 40	136 50	119 10 E	14. Isl. St. Catherine	31 55	325 35	52 5W
29. Hermanos	3 32	72 20	54 40 E	76. Borneo, N. Part	7 40	130 40	113 0 E	15. Cape Frio	27 50	328 35	49 5W
30. Isle Gratio	6 10	80 36	62 56 E	77. Leuco- nia SW Part	12 30	137 45	120 5 E	16. Cape St. Thomas	23 0	335 15	42 25W
31. Padra Banhas	2 10	83 7	65 27 E	78. Formo- ja NE Part	18 55	137 40	120 0 E	17. Spirito Santo	22 10	335 20	42 20W
32. Bassas de Chages, or Island Chagos	6 55	86 20	68 40 E	79. Formo- ja S. Part	12 0	137 31	119 51 E	18. Cape de Abrolhos	19 59	335 25	42 15W
33. Diego Gratioja	8 40	86 0	68 20 E	80. Formo- ja N. Part	25 30	138 20	120 40 E	19. Port Segura	18 15	336 28	41 12W
34. Three Germans	4 30	80 18	62 38 E	81. Piscadore Islands	23 30	136 10	118 30 E	20. Bay Todos Sanc- tos	16 31	337 0	40 40W
35. Cross-Island	4 10	75 40	58 0 E					21. River St. Fran- cisco	12 46	336 35	41 5W
36. Yas de Amber	0 0	70 5	52 25 E					22. Olinda,* or Per- nambuco	10 50	339 45	37 55W
37. Sacatora, or Zucatora . . .	12 21	71 40	54 0 E	1. CAPE St. SE- BASTIAN . . .	12 45	249 40	128 0W	23. Cape St. Augustin	8 13	342 30	35 10W
38. Abdeleur Is.	12 4	70 29	52 49 E	2. Cape St. Lucas, or Lucia . . .	23 20	265 49	111 51W	24. Rio Grande	8 35	342 15	35 25W
39. Cubello . . .	8 10	89 20	71 40 E	3. Cape Corientes	18 50	267 5	110 35W	25. Cape Roque	5 20	341 38	36 2W
40. Malique	9 0	90 33	72 53 E	4. Aquapulco	17 5	273 17	104 23W	26. Island Ferdinan- do Lorenha	5 0	341 48	35 52W
41. Garipe, or Gripe	10 40	90 12	72 32 E	5. Aquatulco	15 27	275 32	102 8W	27. Para River	3 50	346 25	31 15W
42. Qualpera	10 0	91 5	73 25 E	6. Guatimala	14 25	276 35	101 5W	28. Island St. Paul	2 50	334 48	42 52W
43. Andomaon, or Antada	11 10	91 7	73 27 E	7. Panama	8 50	295 43	81 57W	29. Isl. Trinidad	1 20	352 5	25 35W
44. Ceylon, S. End, or C. Gallo	6 8	98 50	81 10 E	8. Bay Bonaven- ture	3 24	299 29	78 11W	30. Isl. des Picos	20 30	347 35	30 5W
45. Maldivas Part	0 25	93 57	76 17 E	9. Island Gallope- ga	0 0	287 25	90 15W	31. Isl. de Martin- vas	22 30	352 20	25 20W
46. vias Part	7 14	90 39	72 59 E					32. Isl. St. Mariade Agosta	19 30	350 40	27 0W
47. Yas de Diego Reys	0 20	89 35	71 55 E	10. Cape de Adjugo	6 30	292 45	84 55W	33. Isl. Ascension	19 40	348 25	29 15W
48. Manila	14 25	134 41	117 1 E	11. Lima*	12 1	300 51	76 49W		7 40	3 10	14 30W
49. Ay- nian Part	19 30	124 35	106 55 E	12. Arica	18 29	304 25	73 15W				
50. J- pan Part	19 55	127 30	109 50 E	13. La Serena	29 0	301 13	76 27W				
51. J- pan Part	35 30	158 5	140 25 E	14. Isl. Juan Fer- nandes	33 15	289 17	88 23W	1. RIVER AMA- zon's Mouth	0 0	327 39	50 1W
52. Verkin's Island	15 0	146 5	128 25 E	15. Baldivia	39 35	296 25	81 15W	2. Cape Orange	4 25	326 10	51 30W
53. Cocos	2 22	111 42	94 2 E	16. Port Steven	46 50	294 59	82 41W	3. Suranam	6 25	320 45	56 55W
54. Andaman . .	4 10	108 37	90 57 E	17. Cape Victory	52 0	294 25	83 15W	4. N. Cape, the Middle of Cao- pory Isl. . . .	2 5	327 39	50 1W
55. Borneo . . .	12 35	109 54	92 14 E	18. Cape Horn, S. Part Terra del Fuego	56 55	297 40	80 0W	5. Cape Nassau	8 5	319 40	58 0W
56. Nicobar . . .	4 50	128 30	110 50 E					6. Cape 3 Points	10 30	315 25	62 15W
57. Sumatra, N. W. End	7 11	111 15	93 35 E					7. Mouth of Oro- noque River	8 15	318 10	59 30W
	5 22	112 25	94 45 E					8. Isl. Trinidad	10 15	317 18	60 22W
59. Bencola . . .	3 55	121 43	104 3 E					9. Isl. Margarita	11 20	314 15	63 25W
60. Sumatra, S. E. End	5 22	122 45	105 5 E					10. Isl. Curacao, or Quicasoe . . .	12 10	309 40	68 20W
61. Jambe	1 19	120 30	102 50 E	1. Le Maire Straits	54 34	304 34	73 6W	11. Cape Conquin- baco	12 40	306 53	70 47W
62. Bantam	6 11	123 3	105 50 E	2. Cape Virgin Ma- ry of Magellan Straits	52 0	302 30	75 10W	12. Cartagena*	10 27	302 14	75 26W
63. Batavia	6 16	124 21	106 41 E	3. Point of River Julian	48 40	303 1	74 39W	13. Darien, or Scots Settlement	8 30	298 50	78 50W
64. Nassau Island	2 54	117 7	99 27 E	4. Peppy's Isles	47 20	310 55	66 45W	14. Porto Bello*	9 33	297 50	79 50W
65. Engano, or Trompeus . .	5 50	119 18	101 38 E	5. Cape Blanco	46 50	305 28	72 12W	15. Nicaragua En- trance	11 25	293 20	84 20W
66. Selam	8 20	119 48	102 8 E	6. River Camero- nes	45 30	304 35	73 5W	16. Cape Camaron	16 10	294 5	83 35W
67. Princes Island	6 30	121 37	103 57 E	7. Point de los Le- ones	44 0	306 55	70 45W	17. Cape Honduras	16 25	292 50	84 50W
68. Java, E. End	8 32	131 5	113 25 E	8. Bay Siffendo	42 35	319 35	58 5W	18. Salamancha	16 50	288 25	89 15W
69. Straits of Sun- dy	6 2	123 21	105 41 E	9. Cape St. An- dreas	38 40	314 30	63 10W	19. Cape Catocha	21 10	291 25	86 15W
70. Banca, S. End	3 20	124 20	106 40 E	10. River Plate	35 40	319 59	57 41W	20. Cape Condefe- do	20 40	287 55	89 45W
71. Borneo, South Point	3 54	131 12	113 32 E	11. Buenos Ayres* in River Plate	34 35	319 35	58 5W	21. Campecha	19 30	285 25	92 15W
72. Banda Isles	4 55	144 52	127 12 E	12. Cape St. Maria	34 30	320 55	56 45W	22. Trieste, or Triest Isles	18 10	286 15	91 25W
73. Celebes, S. End	5 10	136 42	119 2 E								

The MARINER'S SEA-COAST TABLE.

Places Names.	Lat. N.	Long. fr. Ferro.	D. Long. fr. Gr. W.	Places Names.	Lat. N.	Long. fr. Ferro.	D. Long. fr. Gr. W.	Places Names.	Lat. N.	Long. fr. Ferro.	D. Long. fr. Gr. W.
XVII. COAST of the Main CONTINENT in the WEST-INDIES continued.											
23. Sierra, or Cape de Martin . .	19 10	181 50	95 50W	38. Aruba	12 50	309 7	68 33W	32. St. Andrew	12 23	297 18	80 22W
24. La Vera Cruz	19 12	279 45	97 55W	39. E. End of His- paniola . .	18 18	308 20	69 20W	33. Caimanuback	19 8	297 24	80 16W
25. Mexico*	20 0	274 0	103 40W	40. St. Domingo, in Hispaniola . .	18 25	308 5	69 35W	34. Pedro Shoals, N. Side . .	17 10	299 39	78 1W
26. Tompeck	22 20	277 20	100 20W	41. W. End of His- paniola . .	18 26	302 59	74 41W	35. St. Milan	17 10	296 7	81 33W
27. River Spiritus Sanctus, or Mis- sissippi River's Mouth . . .	28 54	280 45	96 55W	42. East End of Ja- maica . .	18 0	301 45	75 55W	36. Guayana	16 53	289 5	88 35W
28. La Phillipina, or Apalachia . .	29 47	292 51	84 49W	43. Port Royal in Jamaica . .	17 40	301 3	76 37W	37. Cozumelli	19 30	288 30	89 10W
29. Cape Florida	24 57	297 5	80 35W	44. West End of Ja- maica . .	18 8	298 35	79 5W	38. Zuna Quita	17 1	287 47	89 53W
30. Escondido	30 20	288 5	89 35W	45. East End of Cu- ba . . .	20 15	303 40	74 0W	XX. COAST of CAROLINA, VIRGINIA, MARYLAND, PENNSYLVANIA, NEW ENGLAND, and NEWFOUNDLAND.			
31. Bay Hondy	22 45	293 55	83 45W	46. Havanna	22 40	294 40	83 0W	1. Bay of St. Au- gustine . .	30 10	296 52	80 48W
32. Cape St. Anto- nio	21 45	292 3	85 37W	47. West End of Cu- ba . . .	21 40	291 5	86 35W	2. Port Royal	32 0	297 52	79 48W
XVIII. CARIBBEE ISLANDS.				XIX. BAHAMA ISLANDS.				3. Charles Town, upon Ashley Ri- ver . . .	33 5	298 49	78 51W
1. TRINIDAD	10 15	317 18	60 22W	1. BERMUDAS	32 25	313 55	63 45W	4. Cape Roman or Cattit . .	33 27	299 45	77 55W
2. Tobago, West End . . .	11 10	318 25	59 15W	2. North Part of Bahama Bank	27 50	299 0	78 40W	5. Cape Fear	33 58	300 41	76 59W
3. Barbadoes, at Bridge-Town	12 58	318 45	58 55W	3. Bahama Island	26 50	298 19	79 21W	6. Cape Hatteras	35 15	303 15	74 25W
4. Granado	11 57	317 15	60 25W	4. Abaco, South End . . .	26 0	303 49	73 51W	7. Cape Henry	37 0	302 11	75 29W
5. Granadillos	12 20	317 40	60 0W	5. Harbour Island	25 37	300 48	76 52W	8. Cape Charles	37 16	303 19	74 21W
6. Boquia	12 50	317 42	59 58W	6. Andros, North End . . .	25 10	298 55	78 45W	9. Cape May	39 6	303 56	73 44W
7. St. Vincent	13 12	317 23	60 17W	7. Providence	25 0	300 15	77 25W	10. Cape Hinlopen, or Cape James	38 50	302 39	75 1W
8. St. Lucia	13 55	317 31	60 9W	8. Eleuthera, South End . . .	24 40	301 39	76 1W	11. Long Island, Middle . .	40 50	304 50	72 50W
9. Martinico*	14 43	316 41	60 59W	9. Cat Island, Mid- dle . . .	24 25	302 26	75 14W	12. Sundry-Hook	40 25	303 41	73 59W
10. Dominico	15 23	317 5	60 35W	10. Watling's Island	24 3	303 0	74 40W	13. New York	40 58	303 42	73 58W
11. Marigallante	15 58	317 15	60 25W	11. Rum-Key	23 45	302 45	74 55W	14. Fisher's Islands	41 20	306 55	70 45W
12. Guadalupe	16 10	316 20	61 20W	12. Exuma	23 22	307 40	76 0W	15. Montock Point	41 18	307 15	70 25W
13. Desada	16 20	316 25	60 15W	13. Crooked Isl. N. End . . .	22 56	303 23	74 17W	16. Block Island	41 15	307 37	70 3W
14. Antigua	17 5	315 50	61 50W	14. Long Isl. South End . . .	22 41	302 53	74 57W	17. Elizabeth's Is- land . . .	41 35	308 22	69 18W
15. Barbuda	17 56	316 55	60 45W	15. Atwood's Keys	23 10	304 0	73 40W	18. Martha's Vine- yard . . .	41 14	308 26	69 14W
16. Monserat	16 45	315 20	62 20W	16. Mayaguana	22 35	303 49	72 51W	19. Nantucket Is- land . . .	41 15	308 47	68 53W
17. Rodunda	16 55	316 20	61 20W	17. French Keys	22 40	303 55	73 45W	20. S. End of Nan- tucket Shoals	39 50	309 17	68 23W
18. Nevis	17 5	315 3	62 37W	18. Marapervouz	21 58	302 50	74 50W	21. South End of St. George's Bank	41 55	310 25	67 15W
19. St. Christopher's	17 17	314 55	62 45W	19. Hogsties	21 17	303 40	74 0W	22. Plymouth	42 2	308 45	68 55W
20. Fustatia	17 25	314 15	62 25W	20. Hineago, West End . . .	20 52	303 49	73 51W	23. Boston Entrance	42 30	308 12	69 28W
21. Saba	17 35	314 55	62 45W	21. West Caicos	21 38	305 37	72 3W	24. Cape Cod	42 12	308 40	69 0W
22. St. Bartholo- mew	17 52	315 29	62 11W	22. Caicos, North End . . .	20 50	306 15	71 25W	25. Cape Ann Island	42 46	307 50	69 50W
23. St. Martin's	18 6	315 25	62 15W	23. Turks Isl.	21 35	307 27	70 13W	26. Pascataway En- trance . .	43 26	307 25	70 15W
24. Anguilla	18 17	315 22	62 18W	24. Abrolho Bank, N. End . .	21 35	308 29	69 11W	27. North Yar- mouth . .	44 14	309 37	68 3W
25. Sambreiro	18 35	315 5	62 35W	25. Plate Wreck	20 10	309 20	68 20W	28. Panobscot River	44 40	310 25	67 15W
26. Aneguda	18 47	314 49	62 51W	26. Mucars	21 30	301 0	76 40W	29. Point, or Port Royal . .	44 45	311 55	65 45W
27. St. Cruise	17 52	314 5	63 35W	27. Verd	21 17	301 19	76 21W	30. Cape Sable	43 50	312 37	65 3W
28. Virgins	18 30	314 10	63 30W	28. Cajad Zal	23 10	297 11	79 29W	31. Island Sable	44 20	318 34	59 6W
29. St. Thomas	18 30	314 13	63 27W	29. Pinos	21 20	293 25	84 25W	32. Cape Breton	46 0	319 5	58 35W
30. St. John de Port Rico	18 30	311 58	65 42W	30. Great Camains	18 54	296 6	81 34W	33. Cape Charles, or Chr. Straits	52 10	323 35	55 5W
31. Isles and Rocks of Aves . . .	12 1	313 5	64 35W	31. Little Camains	19 30	297 11	80 29W	34. Anti Costi Isl. the Middle	49 40	316 50	60 50W
32. Tortuga	11 10	313 41	63 59W								
33. Margaritta	11 20	314 15	63 25W								
34. Blanco	11 50	313 55	63 45W								
35. Testigos	11 35	314 54	62 46W								
36. D'Orchila	11 45	313 5	64 35W								
37. Bonairy	12 12	311 7	66 33W								

The MARINER'S SEA-COAST TABLE.

Places Names.	Lat. N.	Long. fr. Ferro.	D. Long. fr. Gr. W.	Places Names.	Lat. N.	Long. fr. Ferro.	D. Long. fr. Gr. W. or E.	Places Names.	Lat. N.	Long. fr. Ferro.	D. Long. fr. Gr. E.			
XX. COAST of CAROLINA, VIRGINIA, MARYLAND, PENNSILVANIA, NEW ENGLAND, and NEWFOUNDLAND, continued.				20. Viner's Isl.	53	5	293 32	84 8W	25. Black Point, or South End of Charles Isle	78	0	28 5	10 25 E	
35. Quebec*	46	55	307 47	69 53W	21. Cape Henrietta Maria . . .	55	7	293 5	84 35W	26. Deer Sound	79	15	30 15	12 35 E
36. Tadoussack	49	0	310 30	67 10W	22. New Severn	56	0	289 15	88 25W	27. Fowl Sound	77	28	30 25	12 45 E
37. Bay of Brest	52	10	320 38	57 2W	23. Port Nelson, or York Fort . .	57	10	283 37	94 3W	28. Bell Sound	77	15	30 15	12 35 E
38. Bell-Island	52	7	322 0	55 40W	24. Cape Churchill	59	0	282 15	95 25W	29. Horn Sound	76	45	31 11	13 31 E
39. Cape St. John	50	25	324 47	52 53W	25. Sir Thomas Roe's Welcome	64	15	285 31	92 9W	30. Point Lookout	76	25	33 11	15 31 E
40. Cape Bonavista	49	15	325 23	52 17W	26. Cape Southamp-ton	61	55	290 47	86 53W	31. Helie's Sound	78	55	39 25	21 45 E
41. French Factory	50	10	316 25	61 15W	27. Shark Point	64	30	294 40	83 0W	32. Cape Barcan, or Barcam . . .	78	15	39 46	22 6 E
42. Virgin Rocks	46	6	319 55	57 45W	28. Nottingham Isl.	63	30	297 42	79 58W	33. Cape Blanco } Edge of S. M.	77	45	41 5	22 25 E
43. Cape Roye	48	0	319 55	57 45W	29. Mills Isles	64	26	297 17	80 23W	34. Duck's Cove }	77	35	40 45	23 5 E
44. Isl. St. Paul	47	10	319 17	58 23W	30. Salisbury Island	63	45	300 20	77 20W	35. Negre-Point }	76	55	41 5	23 25 E
45. Cape St. Lau-rensa . .	47	30	323 12	54 28W	31. Salvages Island	62	40	307 18	70 22W	36. Hope Island	76	18	41 20	23 40 E
46. Placentia Bay	47	45	323 37	54 3W	32. Qu. Ann's Fore-land	63	48	302 50	74 50W	37. Whale's Head	77	18	39 5	21 25 E
47. Cape St. Mary	47	10	324 12	53 28W	33. Resolution Isle	61	50	312 31	65 9W	38. Lee's Foreland	78	5	41 0	23 20 E
48. St. John's Har-bour . . .	48	0	325 56	51 44W	34. Cape Elizabeth	62	3	310 45	66 55W	39. Cherry, or Bear Island	74	30	35 43	18 3 E
49. Bay of Bulls	47	50	316 6	51 34W	35. Cape Farewell	59	45	330 50	46 50W	40. Ice Point, or Cape Desire	77	40	86 45	69 5 E
50. Cape Race	46	40	325 43	51 57W	XXII. COAST of ICELAND, GREEN- LAND, NOVA ZEMBLA, and the NOR- THERN ISLES.				41. Admiralty-Island	75	5	77 25	59 45 E	
51. Cape St. Francis	48	9	325 49	51 51W	1. SOUND ROYAL	66	22	353 2	24 38W	42. Langeneß	74	20	71 11	53 31 E
52. Conception Bay Entrance	48	20	325 27	52 13W	2. Bargarer's Point	66	20	1 0	16 40W	43. Crofs-Point	72	0	70 47	53 7 E
53. Barcaleau Isl.	48	40	325 39	52 1W	3. Whale's Bay	65	27	357 2	20 38W	44. Fretumborough	70	0	78 55	61 15 E
54. Trinity Bay En-trance . .	48	52	325 15	52 25W	4. Grime's Hole, or Guberman's Rocks	66	23	348 5	29 35W	45. Colgoyen Isl.	69	0	62 35	44 55 E
55. Penguin Isl.	50	0	324 45	52 55W	5. Gamar Isles, or Gille	65	48	350 41	26 59W	46. Cape Candynose	69	5	60 10	42 30 E
56. Grey's Isl.	50	35	324 10	53 30W	6. Westmania Isles	63	30	354 41	22 59W	47. Cape Barfo	66	30	55 35	37 55 E
XXI. COAST of HUDSON'S BAY, and the STRAITS.				7. Rook-Point	64	0	351 11	26 29W	48. Catnose	65	43	52 45	35 9 E	
1. BUTTON'S ISLES . . .	60	25	311 8	66 32W	8. Halliford . .	64	30	342 52	34 48W	49. Archangel	64	30	58 5	40 25 E
2. Cape Charles	62	10	302 0	75 40W	9. Snow-Hill	65	11	350 21	27 19W	50. Crofs-Island	66	31	54 8	36 28 E
3. Cape Walsing-ham . . .	62	35	299 40	78 10W	10. Fair Foreland	66	20	351 8	26 32W	51. Cape Gallant, or Sweet-Nose	68	10	52 20	34 40 E
4. Cape Jones	54	55	298 37	79 3W	11. Roge Point, or Orgel Bay . .	66	0	352 10	25 30W	52. Kilduyn Isl.	59	30	48 55	31 15 E
5. Mansfield Isl. the Middle	61	42	297 5	80 35W	12. Marza, or Lun-geneß	66	8	353 35	24 5W	53. River Kola, En-trance	69	10	48 40	31 0 E
6. Sleeper's Isles	60	10	296 5	81 35W	13. Grims Isl.	67	15	355 1	22 39W	54. Fisher's Island	70	0	44 13	26 33 E
7. Baker's Dozen Isles . . .	57	56	296 10	81 30W	14. Isles of Ferro	62	6	12 35	5 5W	55. North Cape	71	23	40 37	22 57 E
8. Bears-Isles	54	25	293 50	83 50W	15. Langeneß	66	46	4 45	12 55W	56. Surroy Island	71	5	34 15	16 35 E
9. Cubbs-Isle	54	10	294 55	82 45W	16. Silly, or Pappy Isl.	64	50	5 25	12 15W	57. Tromsund Isl.	70	25	33 5	15 25 E
10. Weston's Isl.	52	58	294 47	82 53W	17. Horn Bay	64	50	5 35	12 5W	58. Isl. Sanien, S. W. Point . .	69	35	30 35	12 55 E
11. Solomon's Tem-ple Island	53	5	296 35	81 5W	18. Merchant's Fore-land	63	25	0 30	17 10W	59. Loe-fort, West Point	68	15	26 35	8 55 E
12. Shepherd's Isl.	51	45	296 47	80 53W	19. Portland	64	2	356 30	21 10W	60. S. W. Point of fame . . .	68	15	27 5	9 25 E
13. Danby's Island	52	15	295 39	81 1W	20. Green's Island	66	50	352 55	24 45W	61. Werro, or Weroy Island	67	30	25 5	7 25 E
14. Charlton's Is-land	52	8	285 15	81 25W	21. Beerenbergh, or John Main's Is-land	71	45	22 5	4 25 E	62. Dronen, or Dron-them	63	40	27 50	10 10 E
15. Rupert's River	51	30	298 9	79 31W	22. Hacluit's Head Hand	79	55	28 35	10 55 E	63. Ransdel	63	15	24 5	6 25 E
16. Frenchman's River . . .	51	20	297 21	80 19W	23. Fair Foreland	79	20	28 27	10 47 E	64. North Point	62	20	23 1	5 21 E
17. Point Comfort	51	24	296 24	81 16W	24. Cape Cold, N. End of Charles Isl.	79	0	27 35	9 55 E	65. Standland	62	10	22 13	4 33 E
18. Moufe River's Mouth . .	51	18	294 22	83 18W						66. Kat's Nefs, or Scut's Nefs, S. Point	61	45	21 11	3 31 E
19. Albany Fort	52	26	292 45	84 55W						67. Hearle Isl. S. End	60	40	21 13	3 33 E
										68. North Bergen	60	10	23 15	5 35 E
										69. Bommel Isl. N. Part	69	25	23 15	5 35 E
										70. Jedder	58	5	23 45	6 5 E
										71. Naze of Nor-way	57	45	22 59	7 19 E
										72. Maesterland	57	53	29 20	11 40 E

The MARINER'S SEA-COAST TABLE.

Places Names.	Lat. N.	Long. fr. Ferro.	D. Long. fr. Gr. E.	Places Names.	Lat. N.	Long. fr. Ferro.	D. Long. fr. Gr. E.
XXIII. COAST of the SOUND and BALTIC.				26. Arensbergh, in Ofel Island . .	58 20	41 7	23 27 E
1. MAERDEN, or MARDEL . .	58 19	26 32	8 52 E	27. Parnaw	58 30	43 22	25 42 E
2. Caperwick	59 20	27 45	10 5 E	28. Runen Island	57 55	41 35	23 55 E
3. Larwick	58 54	26 55	9 15 E	29. Riga	57 4	42 50	25 10 E
4. Ansluo, or Christiana	59 40	27 35	9 55 E	30. Domenels	57 30	41 35	23 55 E
5. Gottenberg Gat	57 50	29 50	12 10 E	31. Der Winda	57 15	39 41	22 1 E
6. Cape Kol	56 30	29 48	12 8 E	32. Der Memel	55 48	39 11	21 31 E
7. Elsinore	56 22	31 17	13 37 E	33. Coningsburgh	54 43	39 10	21 30 E
8. Elfenbergh	56 10	30 5	12 25 E	34. Dantzic*	54 22	36 28	18 48 E
9. Valsterborn	55 28	30 35	12 55 E	35. Heel	54 40	36 43	19 3 E
10. Christianople	56 10	33 35	15 55 E	36. Wisby, in Gotland	57 30	36 5	18 25 E
11. Calmer	56 40	34 10	16 30 E	37. Gotland, N. End	58 0	37 50	20 10 E
12. {Oe- } S. End	56 17	33 5	15 25 E	38. Gotland, S. End	57 0	35 20	17 40 E
13. {land } N. End	57 20	34 35	16 55 E	39. Bornholm	55 15	32 20	14 40 E
14. Landfort	58 40	35 55	18 15 E	40. Carmin, or Hammin	54 10	32 35	14 55 E
15. Stockholm*	59 20	37 5	19 25 E	41. Jafmond or Rugen	54 30	31 35	13 55 E
16. Aboo	60 40	38 45	21 5 E	42. Rostock	54 37	36 15	18 35 E
17. Rastbergh	60 28	40 10	22 30 E	43. Straelfund	54 25	31 51	13 11 E
18. Borgo	60 40	43 35	25 55 E	44. Wismar	54 10	29 35	11 55 E
19. Pelting Sound	60 32	44 25	26 45 E	45. Lubeck	54 6	27 35	9 50 E
20. Wyburgb	60 52	46 51	29 11 E	46. Copenhagen*	55 41	30 25	12 45 E
21. Petersburg*	60 0	48 0	30 20 E	47. Moscow, Russia	55 56	58 0	40 20 E
22. Narva	59 27	46 0	28 20 E	48. Elfenore	56 22	30 17	12 37 E
23. Revel	59 35	42 26	24 46 E	49. Uraniburg*	55 54	30 33	12 53 E
24. Nargin Isl.	59 35	42 5	24 25 E	50. Anout, or Anbolt Isl. . . .	56 50	28 41	11 1 E
25. Sybrannest in Dagoo, or Dagerort	59 0	40 35	22 55 E	51. Lefon, or Lefnow Isl. . . .	57 5	28 5	10 25 E
				52. The Scaw	57 30	27 55	10 15 E

The Places in the foregoing Sea-Coast Table are given contiguously, under distinct Heads, and numbered for Reference, to show the Situation and Geography of Places, in Respect of each other, as well as their Latitudes and Longitudes. Whereas an alphabetical Table of the same Places (especially of the less remarkable) had required above double Room to explain their Situation. An Instance of which may be seen, of the Room taken to explain the Situation of alphabetical Places, in Robertson's Navigation; which yet gives no Idea of their Contiguity.

The Differences of Longitudes are given from Greenwich Observatory, for determining the Place, Declination, or Right Ascension, of the celestial Bodies observed under a distant Meridian: All Ephemerides, in Books of Navigation, being properly made for the principal Place of a Nation's Observatory.

Hence the Reason of each Nation beginning the Longitude, or giving the Difference of Longitude, from their Metropolis, or principal Place of Observation; for the Purposes for which it is principally wanted.

Therefore, the Longitudes from Ferro-Island, serve only to compare with what all other Nations have fixed, or shall fix, them at; by first discovering the Difference of Longitudes from each Metropolis or principal Observatory of each Nation, respectively; and then reducing that Difference of Longitude to the Longitude from Ferro; by which Means an Agreement or Difference in the Longitudes, among the different Discoverers or Navigators, will be immediately seen; and thence a Certainty, or Improvement therein, may the sooner be had.

And we are of Opinion that all Nations should publish their celestial Observations quarterly, or annually, as they are made, for the Use of the Navigator, as here mentioned, and the Good of Mankind in general. Instead of which, we find the celestial Observers of our own British Nation, by some unaccountable Disposition or Principle they are possessed of, more backward or averse, than those of any other Nation, of communicating their Observations (though they are paid for it with the public Money) for the Use of the Mathematicians, in perfecting the Theory of Motion, and making Astronomical Improvements: On Account of which Degeneracy, they may be deemed MERE OBSERVERS! far different from that Great Mathematical Genius, and Astronomer, (Second to Sir ISAAC NEWTON) the late famous and communicative Dr. HALLEY!

An alphabetical TABLE of the LATITUDES of *promiscuous* and remarkable PLACES, with their LONGITUDES from the Island of FERRO, and Difference of Longitude from Greenwich OBSERVATORY. According to the Authority of the Sieur DESPLACES, in his *Ephemerides*, from 1715 to 1725.

Places Names.	Lat. N. or S.	Long. fr. Ferro.	D. Long. fr. Gr. E. or W.	Places Names.	Lat. N. or S.	Long. fr. Ferro.	D. Long. fr. Gr. E. or W.
AGRA, in Mogul	26 43 N	94 24	76 30 E	Marseilles	43 19 N	23 7	5 17 E
Alexandria, Syria	36 32 N	54 0	36 10 E	Martinico	14 44 N	316 41	61 9 W
Aleppo, Syria	36 0 N	55 0	37 10 E	Messina	38 21 N	33 56	16 6 E
Alexandria, Ægypt	31 11 N	47 54	30 4 E	Mexico	20 0 N	274 0	103 50 W
Amsterdam	52 23 N	22 53	5 3 E	Milan	45 20 N	26 35	8 45 E
Antibes	43 34 N	24 48	6 58 E	Montpellier	43 37 N	21 32	3 42 E
Avignon	43 57 N	22 32	4 42 E	Moscow	55 30 N	57 0	39 10 E
				Munich, Bavaria	48 18 N	29 30	11 40 E
Barcelona	41 26 N	19 53	2 3 E				
Batavia, Indies	6 15 S	124 0	106 10 E	Nants	47 13 N	16 7	1 42 W
Berlin	52 33 N	31 7	13 17 E	Nancy	48 42 N	23 57	6 7 E
Boulogna, Italy	44 30 N	29 17	21 27 E	Naples	41 5 N	32 20	14 30 E
Bordeaux	44 50 N	16 55	0 55 W	Nice	43 41 N	25 4	7 14 E
Bourges	47 5 N	20 4	2 13 E				
Breslau, Silesia	51 3 N	34 47	16 57 E	Olinda, America	8 13 S	342 30	35 20 W
Brest	48 23 N	13 6	9 4 E	Ostend	51 10 N	20 31	2 41 E
Bruges	51 12 N	20 47	2 57 E	Oxford, England	51 45 N	16 26	1 24 W
Cadix	36 37 N	11 50	10 20 E	Padua	45 31 N	29 1	11 11 E
Cairo, Ægypt	30 2 N	49 35	31 45 E	PARIS OBSERVA-			
Cambray	50 10 N	20 54	3 4 E	TORY . . .	48 50 N	20 0	2 10 E
Calais	50 57 N	19 27	1 37 E	Parma, Italy	44 44 N	28 27	10 37 E
Cape Bonne Espe-				Pekin, China	39 54 N	134 16	116 26 E
rance	34 15 N	37 44	19 54 E	Perpignan	42 41 N	20 33	2 43 E
Cape de Verd	14 43 N	0 30	17 20 W	Porto Bello	9 33 N	297 50	80 0 W
Carthage	10 30 N	302 10	75 40 W	Pondicherry, East			
Cayenne, America	4 56 N	324 30	53 20 W	Indies . . .	11 55 N	97 30	79 40 E
Chester, England	53 13 N	14 51	2 58 W	Prague	50 4 N	32 22	14 32 E
Constantinople	41 6 N	46 33	28 43 E				
Copenhagen	55 41 N	30 25	12 35 E	Canton, China	23 8 N	130 43	112 53 E
Cracow, Poland	50 10 N	38 0	20 10 E	Quebec	46 55 N	307 47	70 3 W
Dunkirk	54 22 N	36 11	18 21 E	Rochelle	46 10 N	16 37	1 13 W
Dublin	53 11 N	10 30	7 20 W	Rodes	44 20 N	20 14	2 24 E
				Rome	41 54 N	30 20	12 30 E
Edinburgh	56 10 N	15 0	3 50 W	Rotterdam	51 50 N	22 30	4 40 E
				Rouen	40 41 N	40 48	22 58 E
Florence	43 47 N	28 59	11 9 E				
Frankfort	50 4 N	26 10	8 20 E	Siam, Indies	14 18 N	118 30	100 40 E
				Smyrna	38 28 N	45 0	27 10 E
Guadalupe, Ameri-				Stockholme	59 36 N	36 15	18 25 E
ca	16 20 N	316 11	61 39 W	Strasbourg	48 35 N	25 25	7 35 E
Geneva	46 12 N	24 0	6 10 E	Surat, Indies	21 10 N	90 0	72 10 E
Goa	15 31 N	91 25	73 35 E				
Goree, Cape Verd	14 39 N	0 35	17 15 W	Toulon	43 7 N	23 35	5 45 E
GREENWICH OB-				Tripoly	32 54 N	31 11	13 21 E
SERVATORY . .	51 29 N	17 50	0 0	Turin	44 50 N	25 10	7 20 E
Hamburg	53 41 N	28 15	10 25 E	Valparaiso, South			
				America . .	33 28 N	305 28	72 22 W
Jerusalem	31 50 N	53 30	35 40 E	Vienna	48 14 N	34 32	16 42 E
Isle of Ferro	28 5 N	0 0	17 50 W	Venice	45 35 N	30 10	12 20 E
Ispahan, Persia	32 30 N	70 30	52 40 E	Uraniburg	55 34 N	30 32	12 42 E
Lima, in Peru	12 20 S	296 45	81 5 W				
Lipfic	51 19 N	30 0	12 10 E				
Lisbon	38 45 N	9 15	8 35 W				
LONDON	51 31 N	17 45	0 5 W				
Macao, China	22 12 N	130 48	112 58 E				
Malacca, Indies	2 12 N	119 45	101 55 E				
Madrid	40 26 N	14 20	3 30 W				
Malta	35 43 N	32 9	14 18 E				

In the SEA-COAST TABLE, the Difference of Longitude from Greenwich to Paris Observatory, is taken according to Dr. Halley, 2° 20' E. which taken from 20°, East Longitude of Paris from Ferro Isle, leaves 17° 40' E. Longitude of Greenwich from Ferro.

But, in the foregoing alphabetical Table, the Difference of Longitude from Greenwich to Paris Observatory is taken 2° 10' E. only, according to the French Authority of De la Caille, by which the Differences of Longitudes E. or W. of Greenwich Observatory are adjusted from the given Longitudes (according to the Sieur DESPLACES) from Ferro.

The Difference of Longitude from London to Ferro in the New Sea-Calendars corrected, and Mariner's Compass newly rectified, is 17° 45' W. which is 17° 50' W. from Greenwich, and as much East from Ferro, agreeing with the French Authority; which we have corrected from Halley's Authority, being only 10' Difference.

Hence, if you add 10' to some of the Longitudes from Ferro, in the Sea-Coast Table, they will nearly agree with some of these alphabetical Longitudes from Ferro.

There is still another very large Variation-Table of Latitudes and Longitudes in Robertson's Navigation, as if the terrestrial Latitudes and Longitudes were never to be fixed, but shifted their Places with the Latitudes and Longitudes of the Stars. In our Sea-Coast Table the Latitudes and Longitudes of Places are according to the present best Authorities.

Captain BROWN's DIRECTIONS for the Safety of Homeward-bound SHIPS, coming into the ENGLISH CHANNEL.

FIRST, I recommend that all Ships and Vessels, coming off the Ocean, be steered in a *Parallel of Latitude*, not more nor less than $49^{\circ} 30'$; keeping your *Lead* going, endeavour to strike the Ground in 100 or 120 *Fathom* Water, which I call the outer *Edge* of the *British Bank*, or *Soundings*. Steer from thence E. b. S. $\frac{1}{2}$ S. in Order to keep your *Latitudes* by the *Compass*, till by your *Log* you have run 80 *Leagues* from the above *Sounding* of 100 or 120 *Fathom* Eastward. Then may you *haul* to the Northward and make Land at Pleasure. But, if interrupted by Clouds, or hazy Weather, so that you have missed an Observation for several Days, whereby you cannot, with Certainty, determine your *Latitude*, in such Cases, if you come into *Soundings* from the *Western Ocean*, observe, as before, to get Ground, if possible, in 100 or 120 *Fathoms*, which obtained, keep your *Log* and *Lead* going every Hour; steering E. b. S. $\frac{1}{2}$ S. till, by your *Distance*, you have run from the aforesaid *Soundings* 40 or 45 *Leagues*, and shoaled your Water; gradually lessening to 60 *Fathoms*. Then you may find it difficult to determine whether you are to the Northward, or the Southward of *Scilly*; for the *Soundings* on both Sides I have often found to be pretty near alike. Therefore, to resolve this Doubt, I recommend steering a Southerly Course from the aforesaid *Distance* run, and Depth of 60 *Fathom*; and as you sail to the Southward you will deepen your Water from 60 to 70 and 75 *Fathom*; which having done, you may depend on the *British Channel* being open, and clear from the Danger of either running ashore on *Scilly*, or into *St. George's Channel*, too often the Fate and Case of Ships, who, for Want of such Helps, are sometimes lost, with the Lives on Board, or dangerously bewildered.

When you have got the Depth of 70 or 75 *Fathom*, aforesaid, immediately alter the Southerly Course to E. or E. b. S. till, by your *Distance*, you are shot within *Scilly-Islands*; the said Places lying about 62 or 3 *Leagues* from the Western Edge of the *British Soundings*, then may you haul to the Northward, and make the Land, as you think proper.

But, if you come from the Southward, the Coast of Spain and Portugal, or Bay of Biscay, you must be likewise careful how you come in with the Channel, in thick Weather; for as you strike Ground with your *Lead*, you will often find coarse *Soundings*; and, if near *Ushant*, Gravel, with small Stones; which Ground is much steeper than the Edge of the *Western Bank*. For, if you come into your *Soundings* with the Channel open, steering to the Northward, to make the Land's End, Lizard, &c. in running eight or ten *Leagues*, you will go from 100, to 75, or 70 *Fathom*, in the said *Distance*. Whereas being to the Westward, you may run 20 or 30 *Leagues*, and not make more Difference in your *Sounding* than aforesaid. So that, from what I have advanced, it will be easy to determine whether you have the Channel open or not, a *Dispute* that has often puzzled the most experienced *Mariners*, and skilful *Navigators*, using these Seas.

However, I must observe, that some Times, in the aforesaid *Soundings*, I have met with a strong Northerly Current, at the Rate of about one Mile an Hour, which, and about 17 Degrees Variation West, at this Time, should be duly accounted for, in Order to keep the true *Parallel of Latitude*, afore-mentioned. As, likewise, in some *Sea-Charts*, the *Latitude* of the Land's-End, Lizard, &c. are laid down, ten Miles to the Northward of their true *Latitude*. So that if these *Impediments* are not all duly considered, and allowed for, I say, from what I have already observed, an Error may be easily contracted, greatly endangering the Loss of a Ship.

These Directions being carefully observed, your *Latitude* and *Distance* will be corrected, and a tolerable Knowledge of the Ship's Position may be inferred. And I would also advise all Ships to be careful how they deal with the French Coast. For let the Weather be as it will, after having run the *Distance* aforesaid, off the *Islands* of *Scilly*, then make bold with your own Coast, in Order to shun falling in with the *Islands* of *Guernsey*, *Jersey*, &c. which so often ends in frightful Circumstances of both Ships and Lives.

AND, lastly, observe that in sounding a Stream, and to the Westward of *Scilly*, you will find bluish, oozy Ground; so that when, by your *Lead*, you have such Ground, you may be assured where you are, and therefore shape your Course accordingly.

Note, That in coming up the Channel when you are abreast of the Lizard, you will have 50 *Fathom* Water, and of the Start Point 45 *Fathom*.

The practical Mariner finding it difficult to get Ground in a Storm, or hard blowing Weather at 100 or 120 *Fathom*, coming into Channel, he may follow the above Rules; getting Ground at 80 *Fathom* or less, by making proper Allowance, according to the Depth of Water.

COMMENCEMENT of the most FAMOUS ÆRAS in YEARS current and complete, before and since CHRIST, Julian Styl.
 The Want of Distinction before, between current and complete Years, has occasioned great Confusion in CHRONOLOGY.

Remarkable ÆRAS.	Week-Day.	Month Day. O. S.	CURRENT.				COMPLETE.				Days.	Olympiads.	Olympic Years.	Dif. com.	Cycles.		
			Years before Christ.	Dif. com.	Yrs. Jul. Per.	Dif. com.	Years before Christ.	Dif. com.	Years Julian Period.	Dif. com.					Y. O. S.	Y. O. S.	Indict.
Commencement of the Jul. Period	Monday	Jan. 1	4713		1		4712		1 bef. o fin. Per		364	Current	Current		1	1	1
The common Æra of Creation.				763		763		763		763							
Strauchius. Neglecting other traditional Æras of 5508, 5199, and 3955 before Christ	unknn.	unknn.	3950	1656	764	1656	3949	1656	763	1656	unknn.	...	...		8	4	14
The Æra of the Deluge	unknn.	unknn.	2294	1518	2420	1518	2293	1518	2419	1518	unknn.	...	...		12	7	5
— The Greek Olympiads . . .	Monday	Solft. July 8	776		3938		775		3937		23	1	1		18	5	8
— The Foundation of Rome	Monday	Apr. 21	753		3961		752		3960		254	4	6		23	13	9
— Nabonassar, King of Babylon	Wednes.	Feb. 26	747		3967		746		3966		308	2	8		6	15	7
— The Death of Alexander the Gr.	Sunday	No. 12	324		4390		323		4389		49	1	114		423	22	10
— The Syrians and Chaldeans	Monday	Oct. 1	312		4402		311		4401		12	1	117		12	6	13
— Jul. Cæsar, Corrn. of Calend.	Friday	Jan. 1	45		4669		44		4668		91	1	117		267	21	14
— Spain; or Spanish Æra	Sunday	Jan. 1	38		4676		37		4675		364	4	183		7	28	11
— CHRIST, or Christian Æra	Satur.	Jan. 1	1		4714		1 bef. o fin. Chr		4713		38	1	195		38	10	2
— The Æthiopians and Abyssines, called also Dioclesian, or Æra of Martyrs . . .	Friday	Aug. 29	284		4997		283		4996		283	4	265		283	13	19
— Hegira, or Turkish and Arabic Æra	Friday	July 16	622		5335		621		5334		254	4	265		338	13	19
— Jeldegird, or Persian Æra	Tuesday	Jun. 16	632		5345		631		5344		168	2	350		10	15	10
— 2d Persian, or Jallalean Æra	Thurs.	Mar. 14	1079		5792		1078		5791		198	4	352		447	24	6
— Pope Gregory's reforming the Calender	Friday	Oct. 5	1582		6295		1581		6294		292	3	464		503	24	16
											87	2	590		23	6	10

Chronological Dates of all Kinds are most truly ascertained by a Concurrence or Correspondence discovered, in the Week-Days, from the Years of different Æras, to the same Event.

N. B. The above Table correctly distinguishes between the current and complete Julian Years, before and since Christ, (confusedly given by Chronologers) and therefore the Æras here given are fitted for the Reduction of Chronology, or for reducing one Account of Time to another. Which is effected by Means of Tables, further on, estimating the Number of Years and Month-Days of any particular Kind, equivalent to a given Number of Julian Years and Month-Days, and the contrary. And these Tables are likewise fitted for determining the Number of the Ferie, or Week-Day, correspondent to a given Month-Day of different Kinds of Years. See p. 141, for the Form, Length, Beginning, and Ending, of the Jewish, Egyptian, Arabic, Turkish, and also antient Grecian Year. Any Number of which, and their Month-Days, are reducible to an equivalent Number of Julian Years and Month-Days; by the RULES and EXAMPLES hereafter exhibited.

The Egyptian or Persian Year consists of 365 Days without any Variation; and each Month of 30 Days, with 5 Days added at the End of the last Month. This Year is completed 6 Hours sooner than the mean Julian Year, and gains 1 Day of the Julian Account in 4 Years. And in the Space of 1460 Egyptian or Persian Years, the same Day of the Year and Month returns. For 1460 Years Julian make 1461 Years Egyptian; when the 1st Day of the Egyptian Month Thoth happens on the 1st of January Julian Style, 4 Years after which the 1st of Thoth happens on the 31st of December, and in 4 more on the 30th Day, &c. retreating 1 Day in every 4 Years Julian Style. Therefore, 1 Day retreated or retrograded every 4 Years Julian is 365 Days retreated in 1460 Julian Years, when the 1st of Thoth returns to January 1, Julian Style. This Egyptian or Persian Year, though fit for astronomical Computations, is improper for the Purposes of civil Life. The antient Persians, by a Correction of their Calender, at certain Periods, made their Year to consist of 365^d 5^h 48^m 53^s 20th, according to the French Authors.

The Arabians and Turks make their Years consist of 12 synodical Lunations, and in 30 Years, which is their Period, they reckon 19 Years of 354 Days, and 11 Years of 355 Days, viz. 2d, 5, 8, 10, 13, 16, 19, 20, 24, 27, and 30th, of 355 Days each; the Rest, 1st, 3, 4th, &c. of 364 Days; which Period of Years contain 10631 Days exactly. They make the Product of 30 Years, Periods, or Cycles in Succession, that is of 29^d 12^h 44^m to a Lunation. The Commencement of their Year is unfixed, and wanders through all the Seasons successively.

	d	h	m		d	h
Their Lunar Year is	354	8	48	} Julian Year 365 6 at a Mean		
Solar Year about	365	5	49			
Dif.	10	21	1			
				Therefore . . . 537 Turkish	} Or 235 Turkish Years	
				Make 521 Julian Years.		Make 228 Julian Years.

The Jewish and Grecian Years follow their particular Forms as described in Page 141.

The Year, represented by a Ring, or Snake with the Tail turned into its Mouth, has different Beginnings, and Successions of the same Parts in different Nations. Some begin their Year at the Solstices, others at the Equinoxes. And, in France, before the Year 1564, we find, by a French Ephemeris, that their Year commenced at Easter. But Charles IX. we find, in the same Ephemeris, ordered the Commencement to be on the 1st of January, which was confirmed and established by Order of the French Council and Parliament to begin the 1st of January, 1567.

The Astronomers begin their Year on Jan. 1, or 31st of December, at Noon.

REDUCTION of CHRONOLOGY.

A TABLE for reducing Julian Years into Days, and the contrary.

Jul. Yrs.	Days corresponding.	Jul. Yrs.	Days corresponding.
*1	365	600	219150
*2	730	700	255675
*3	1095	800	292200
4	1461	900	328725
8	2922	1000	365250
12	4383	2000	730500
16	5844	3000	1095750
20	7305	4000	1461000
24	8766	5000	1826250
28	10227	6000	2191500
32	11688		
36	13149	N. B. There being 1461 Days in every 4 Julian Years, when either 1st, 2d, or 3d Year (marked with *) is Bissextile, 1 Day must be added to the Number of Days in the right Hand Column, for the true No. of Days in Succession from the 1st Year, as appears from the 3 first Years of 365 Days each. See P. 129.	
40	14610		
44	16071		
48	17532		
52	18993		
56	20454		
60	21915		
64	23376		
68	24837		
72	26298		
76	27759		
80	29220		
84	30681		
88	32142		
92	33603		
96	35064		
100	36525		
200	73050		
300	109575		
400	146100		
500	182625		

TABLE of the Number of DAYS to the END of the Julian MONTHS.

No.	Months.	Years com.	Years Bissext.
1	January	Ds 31	Ds 31
2	February	59	60
3	March	90	91
4	April	120	121
5	May	151	152
6	June	181	182
7	July	212	213
8	August	243	244
9	September	273	274
10	October	304	305
11	November	334	335
12	December	365	366

N. B. 365^d 5^h 49^m Julian Yr.

A TABLE for reducing Turkish or Arabic Years into Days, and the contrary.

Turkish Yrs.	Days corresponding.	Turkish Yrs.	Days corresponding.
1	354	21	7442
2	709	22	7796
3	1063	23	8150
4	1417	24	8505
5	1772	25	8859
6	2126	26	9213
7	2480	27	9568
8	2835	28	9922
9	3189	29	10276
10	3544	30	10631
11	3898	60	21262
12	4252	90	31893
13	4607	120	42524
14	4961	150	53155
15	5315	180	63786
16	5670	210	74417
17	6024	240	85048
18	6378	270	95679
19	6733	300	106310
20	7088	600	212620

Years 2, 5, 8, 10, 13, 16, 19, 22, 24, 27, 30, contain 355 Days each.

Years 1, 3, 4, 6, 7, 9, 11, 12, 14, 15, 17, 18, 20, 22, 23, 25, 26, 28, 29, contain 354 Days each: Making 30 Years of 10631 Days.

N. B. 235 Turkish Years make 228 Julian Years, or 537 Turkish make 521 Julian Years.

TABLE of the Number of DAYS to the END of the Turkish or Arabic MONTHS.

Beginning.	No.	Turkish Months.	Days
16 July	1	Muh' Arram	30
15 Aug.	2	Sapbar	59
13 Sept.	3	Rabie I.	89
13 Octo.	4	Rabie II.	118
11 Nov.	5	Giumadi I.	148
11 Dec.	6	Giumadi II.	177
9 Jan.	7	Regeb	207
8 Feb.	8	Sabeen	236
9 March	9	Ramadhan	266
8 April	10	Scheual	295
7 May	11	Dulkadate	325
7 June	12	Dulbakate	354
		In Year intercalary	355

365^d 8^h 48^m Turk. Yr.

TABLE for reducing Persian or Egyptian YEARS into DAYS, and the contrary.

Yrs.	Days.	Yrs.	Days.
1	365	30	14950
2	730	40	14600
3	1095	50	18250
4	1460	60	21900
5	1825	70	25550
6	2190	80	29200
7	2555	90	32850
8	2920	100	36500
9	3285	200	73000
10	3650	300	109500
11	4015	400	146000
12	4380	500	182500
13	4745	600	219000
14	5110	700	255500
15	5475	800	292000
16	5840	900	328500
17	6205	1000	365000
18	6570		
19	6935		
20	7300		

Doctor Newton, in his *Cosmographia*, gives an Account of 245 Days to the End of the 8th Persian Month *Aban*, instead of 240, which Month he calls *Apenina Wabak*, and differs 5 Days in the following Months as he does in the Names of some other of his Months, as also does Mr. *Ferguson*, from some of our Persian and Turkish Names of Months from good Authority, which makes the 12th Persian, not 8th Month, intercalary of 5 Days; like the 12th Egyptian Month.

TABLE of the Number of DAYS to the END of each Persian MONTH.

No.	Persian Months.	Days.
1	<i>Pharavardin</i>	30
2	<i>Adar-pabascbt</i>	60
3	<i>Cbardad</i>	90
4	<i>Tbir</i>	120
5	<i>Mardad</i>	150
6	<i>Scheberiz</i>	180
7	<i>Mebar</i>	210
8	<i>Aban</i>	240
9	<i>Adar</i>	270
10	<i>Di</i>	300
11	<i>Bebemen</i>	330
12	<i>Asphardar, or Aspirer</i>	365

TABLE of the Number of DAYS to the END of each Egyptian MONTH.

Beginning.	No.	Egyptian Months.	Ds.
29 Aug.	1	<i>Tbeth</i>	30
28 Sept.	2	<i>Paopbi</i>	60
28 Oct.	3	<i>Aibir</i>	90
27 Nov.	4	<i>Cbojac</i>	120
27 Dec.	5	<i>Tibi</i>	150
26 Jan.	6	<i>Mebir</i>	180
25 Feb.	7	<i>Phamenoth</i>	210
27 March	8	<i>Parmutbi</i>	240
26 April	9	<i>Pachon</i>	270
26 May	10	<i>Payni</i>	300
25 June	11	<i>Epiphi</i>	330
25 July	12	<i>Meferi, or Epagomene, 5 Days added.</i>	365

A TABLE for finding the WEEK-DAYS. According to the Turkish Account of Time.

Sunday.	Monday.	Tuesday.	Wednesday.	Thursday.	Friday.	Saturday.
I.	II.	III.	IV.	V.	VI.	VII.
	180	Turkish	150	Cycles	120	
90		60		30		0
Turkish	2	Years	and	1	Months.	Mubarram
5	Sepbar	Rabie I.	4	Rabie II.		3
		7			6	
	10			9	Giumadi I.	8
13			12		11	
Giumadi II.	Regeb	15			14	
			Sabeen	17		16
21			20	Ramaden	19	Scheual
Dulkadati		23			22	
	26			25		24
29		Dulbakati	28		27	
				30		

Arabic or Turkish CYCLES, the Products of 210 Years.

210	420	630	840	1050	1260	1470
1680	1890	2100	2310	2520	2730	2940

365.5.49 is a Gregorian year

Construction.

REDUCTION of CHRONOLOGY.

Construction. The Numbers at the Top of the Table are the Products of 30 Years in a retrograde Order. All the other Numbers in the Table are the Cycles current from 1 to 30 Years. The Months are placed in the Intervals, (showing the Fera or Week-Day to the Beginning of each Month, for given Years, in the same Column) to find the Fera, or Week-day, as follows.
 The 210 Cycle of Years serve for 30 Weeks, and answers to the Cycle of the Sun 28.

EXAMPLE I. To find the Fera, or Week-Day, at the Death of the Emperor AMURAT II. in the 855th Year of the Turkish Hegira, on the 1st Day of the Month Muharram, current?

The Years of the Cycle nearest to 855 are 840.
 — 840

Rem. 15 Years.

Seek, from the Table, the Fera to these 15 Years, being 3, and the Fera to the Month Muburram being 7, add these together, making 10, to which add the 1st Day of Muburram, and the Sum will be 11, which divided by 7, there will remain 4, or the 4th Fera being Wednesday, required.

OTHERWISE, To find the Julian Date, O. S. corresponding to the said Turkish Date?

Turkish Years completed	Days
600	212620
240	85048
14	4961

Years 854 completed.

Muburram 1 Day current . . 0, or 1 Day less completed [than current.

Sum . . . 302629 Days completed.

Julian Years 800 completed 292200

28	10429
28	10227

Add, 828 Yrs. Jul. and 202 Days, both completed.

To July 16th, 622 Years current, when the Hegira commenced.

Sum, Jul. 16th, 1450 current, and 202 Days more completed;

Or, February 3, 1451, current, O. S. (by Table, P. 32.) When the Emperor Amurat died.

NOW, the Dominical Letter (by Tab. p. 148.) for 1451, O. S. is C, and February 3, (by Table, P. 149) is evidently on a Wednesday, required.

EXAMPLE II. To find the Turkish Date corresponding to the Julian Date of the 29th of May, 1452, O. S. (being on a Monday) when Mahomet II. begun his Reign?

Yrs cur. Chr. D.	Yrs cur. Ds cur.	Yrs	Days
1452, May 29 =	1451 . 515 =	1452 + 150,	Bisextile,
622, July 16 =	622 . 197	when Hegira begun.	

Julian Years Diff. 829 . 318 completed.

† To find the Week-Day?

856 Year current

— 840 Cycle

Rem. 16

Fera 7 }
 Giumadi I. 6 }
 Day 10 }

7)23(3

Monday . . 2 Rem.
 Fera.

† The above follows and refers to what is done at the End of the Operation in the next Column. See †.

Julian Years	Days
800	292200
28	10227
1	365
	+ 318
829	
	Sum, 303110
Turkish Years 600	212620
	90490
240	85048
	5442
15	5315

Sum, 855 Years, and . 127 Days completed.
 Rabie II. . . 118

Giumadi I. . . 9 completed.

Or, 856 Year of the Hegira, on the 10th Day of Giumadi I. current. †

Le Sieur Desplaces, by Mistake, (in the Preface to his Ephemerides, from 1715 to 1725, Page 30) brings out 856 of Hegira, completed on the 1st of Saphar current, who says the Turks reckon it 857 of Hegira, on the 1st of the Month Saphar, current, when Mahomet II. began his Reign, which cannot agree with the 29th of May, 1452, current, on a Monday: The 1st of Saphar, in the 857 of Hegira, current, being on a Sunday, as our Table foregoing shews, according to the Rule given. And the 1st of Saphar, in the 856 of Hegira current, by the same Rule, is on a Tuesday.

† The same Method serves for reducing the Julian into Persian or Egyptian Date, and the contrary.

Whence also the Persian may be reduced to the Turkish Account, and the contrary.

† Though subtracting Julian Years and Days current from other Julian Years and Days current, leaves the Difference in Years and Days complete, the same as if those Julian Years and Days current were first reduced to Years and Days complete, and their Difference then taken, both the Numbers of Years current being Bisextile, or both common; yet, as these Numbers may be of different Kinds, if the current Years and Days be always reduced to those which are complete, and then the Difference or Sum thereof be taken, for Reduction of one Date to another, as different Occasions shall require, the Hazard of mistaking a Day, in computing by Years current, will, by this Means, be avoided.

EXAMPLE

REDUCTION of CHRONOLOGY.

EXAMPLE III. To find the YEAR of the TURKISH HEGIRA corresponding with the 20th of May 1761, N. S. on a Wednesday; being May 9, 1761, O. S. current, or May 8, 1760 Years complete?

Jul. Years complete	Days
1000	365250
700	255675
60	21915
1760, Apr. complete . .	120
In May, Days complete . .	8
From Chr. to the present Julian Date . .	642968
From Chr. to the Hegira, . . Deduct . .	227016
Since the Hegira . .	415952 Days, complete,
Turkish Years 900 . . Deduct	318930
	97022
270 . . Deduct	95679
	1343
3 . . Deduct	1063
1173 Ys complete, & 280 Days, complete,	
Ramadhan, complete . .	266
Scheual . .	14 Days, complete.

The Turkish Hegira begun July the 16th, 622 current.

Jul. Yrs. comp.	Days
600	219150
20	7305
1	365
Comp. Years 621, June, complete . .	181
In July, Days complete . .	15
To the Hegira	227016 Ds.
In 1st Example	302629 Ds.
	since the Hegira.
Jul. Yrs. comp.	529645 Sum.
1000	365250 —
	164395
400	146100 —
	18295
48	17532 —
	763
2	730

HENCE to
prove the
Answer to
the First
EXAM.

1450 Yrs complete, and 33 Days, complete.
Viz. 1451, February 3, current, as before, the Date
of the Christian Æra, corresponding with the Date of the Hegira,
855, on the 1st Day of the Month Mubarram.

EXAMPLE IV. To reduce the Julian Years and Months current, suppose the 11th of September, 1781, O. S. since Christ, into the Years and Months current of the Persian Jeshdegird and the contrary?

Jul. Yrs. comp.	Days
1000	365250
700	255675
80	29220
1780 Yrs. Aug. complete	243
In September, Days complete . .	10
Since the Christian Æra . .	650398
To the Jeshdegird . .	230639
Since the Jeshdegird . .	419759
Persian Years 1000 Deduct	365000
	54759
100 Deduct	36500
	18259
50 Deduct	18250
1150 Years complete, and 9 Days	
complete of Pharawardin.	
Viz. in 1151 Year of Jeshdegird, on the 10th of Pha-	
rawardin, current.	

The Date of the Jeshdegird may be reduced to the Julian Account, and thence to the Hegira, or any other Account of Time you please, by the foregoing Rules.

The Persian Jeshdegird began June the 16th, 632 of Christ current, on a Tuesday.

Jul. Yrs. comp.	Days
600	219150
28	10227
3	1095
631 Ys comp. May comp. 152 Bissextile.	
In June, compl. . .	15
To the Jeshdegird, . .	230639
CONTRARILY, To find the Christian Æra, correspondent to the	
1151 Year of Jeshdegird, on the 10th of Pharawardin, current?	
Persian Years	Days
1000	365000
100	36500
50	18250
1150 Years complete.	
In Pharawardin	9
Æra of Jeshdegird . .	419759
To that Æra from Christ . .	230639
Years Julian 1000 Sum . .	650398
	Deduct 365250
	285148
700 Deduct	255675
	29473
80 Deduct	29220
1780 Years complete and 253 Days.	
Aug. complete . .	243
In Sept. Days 10 complete.	
Or Sept. 11, 1781, current, required.	

EXAMPLE

REDUCTION of CHRONOLOGY.

EXAMPLE V. To find the WEEK-DAY according to ANY ACCOUNT of TIME?

RULE. Divide the Number of the Days complete in the Account by 7, and add thereto the Number of the FERIA on which the Account or ÆRA began, then dividing the Sum by 7, the Remainder will be the Number of the FERIA corresponding to the Week-Day, required.

N. B. The complete Days are always 1 less than the Days current.

THUS, in Example I. The Number of Days in the 855th Year of Hegira, on the 1st of Mubarram, is 302629, which divided by 7, leaves 5, to which 6 being added, the Number of FERIA (answering to Friday) when the ÆRA began, the Sum is 11, which divided by 7, the Remainder 4 is the FERIA answering to Wednesday, required, as before found by other Methods.

Again, in Example II. To the 856th Year of Hegira, on the 10th Day of Gumadi I. current, are 303110 Days complete, which divided by 7 there remains 3, to which if 6 be added, the Number of FERIA (answering to Friday) when the ÆRA began, the Sum is 9, which divided by 7, the Remainder 2, is the FERIA answering to Monday, as before found by other Methods.

Again, in Example III. To the 20th of May, 1761, N. S. are 642968 Days complete, which divided by 7 there remains 4, to which adding 7, the Number of the FERIA when the Christian ÆRA began (answering to Saturday) the Sum is 11, which divided by 7, the Re-

mainder 4, is the Number of the FERIA answering to Wednesday, as before found by other Methods.

Again, in Example III. To 1774 Years of the Hegira, on the 15th Day of Sheual current, are 415952 Days complete, which divided by 7, there remain 5, to which adding 6, the Number of FERIA when the Hegira began (answering to Friday) the Sum is 11, which divided by 7, the Remainder 4 is the Number of FERIA, answering to Wednesday, as before found, correspondent to the Date of the Christian ÆRA, M. y 20, 1761, N. S.

Again, in Example IV. To 1151st Year of Jeshdegird, on the 10th of Pharwardin current, are 419759 Days complete, which divided by 7, there remains 4, to which adding 3, the Number of the FERIA when the Jeshdegird-ÆRA began, (viz. Tuesday) the Sum will be 7, the Number of the FERIA, or Saturday, on the Day of the Jeshdegird, required, corresponding to the 11th of September, 1781, current, O. S.

By the like Methods, the Reduction of all ÆRAS of Chronology are performed, and the Week-Day to the Month-Day in each ÆRA found.

The OLD is reduced to the NEW STYLE, and the contrary, by the Methods already shewn.

If the Week-Day comes out the same to the corresponding Dates of Chronology, by different Methods of computing the Week-Day, you may be sure the Reduction of the Chronology is made right.

CHRONOLOGICAL SCALE of an ÆRA beginning BEFORE CHRIST.

Succesive Years complete of the ÆRA before Chr.									Succesive Years complete of the ÆRA since Chr.								
Be g. Æra	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
9	8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8	9
Years current before Christ Julian Style.									Years current since Christ Julian Style.								

The Spaces of the under Column in which the Numbers are placed are for the Years current, from their Beginning, on the Left: The Ends of those Spaces, to the Right, are for the Years complete, before and since Christ.

CHRONOLOGICAL SCALE of an ÆRA beginning SINCE CHRIST.

Retrogressive Years complete of the ÆRA before Chr.									Succesive Years complete of the ÆRA since Chr.								
10	9	8	7	6	5	4	3	2	1	Be g. Æra	1	2	3	4	5	6	7
9	8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8	9
Years current before Christ Julian Style.									Years current since Christ Julian Style.								

The Spaces of the under Column in which the Numbers are placed are for the Years current, from their Beginning, on the Left: The Ends of those Spaces, to the Right, are for the Years complete, before and since Christ.

AXIOMS.

1. A Year current, is the Year running on, not yet completed.
2. Years current are always 1 more than Years complete, in all Dates, before and since Christ.
3. Days current are 1 more than Days complete, &c.
4. Years and Days complete added to, or subtracted from, Years and Days complete, the Remainder will be Years and Days complete.
5. Years and Days complete being added to, or subtracted from, Years and Days current, the Sum, or Remainder, will be Years and Days current.
6. Years and Days current being subtracted from Years and Days current, the Remainder will be Years and Days complete.

OF ÆRAS BEGINNING and DATES, both Before CHRIST.

PROPOSITION I. To determine the Number of Julian Years current of an Æra, from the Beginning to the present Date thereof, before Christ, being given, and thence to give other chronological Rules?

RULE. By the former of the two chronological Scales it is evident, that If you deduct the whole Julian Years current before Christ, of the present Date, from the whole Julian Years current, of the Date when the Æra before Christ, began, both in the lower Column, the Remainder

REDUCTION of CHRONOLOGY.

will be the Number of whole Years complete, in the upper Column, (and also Years and Days complete, which Years and Days are in both Dates) since the Commencement of that Æra.

Which whole Years Difference are complete Years, at each annual Return of the same Month-Day, on which the said Æra began. To which Difference of Dates current, or Number of complete Years, since the Æra, if 1 Year be added, it will give the Number of Years current of that Æra; because (as was before observed) the Number of Years current are always 1 more than the Number of Years complete.

Otherwise, If you subtract 1 Year from the present Year's Date current, before Christ, (to make that Date complete Years) and then deduct them from the Years of the current Date at the Æra's Beginning, before Christ, the Remainder will also be the Year of that Æra, since it began.

PUT Æ for the Years current at the Æra's Beginning before Christ.

P, for the present Year, current before Christ.

Y, for the Number of Years current of that Æra.

HENCE, we have these EQUATIONS.
$$\left. \begin{array}{l} 1. \text{Æ} - P + 1 = Y \\ 2. \text{Æ} - Y + 1 = P \\ 3. P + Y - 1 = \text{Æ} \end{array} \right\} \begin{array}{l} \text{Julian} \\ \text{Years} \\ \text{current.} \end{array}$$

EXAMPLE 1. To find the Number of the Julian Year current, Y, or Year of the Julian Period, beginning in the 4713th Year current before Christ, answering to the 324th Year, present Date, P, before Christ?

By the 1st Equation.

$$\begin{array}{r} \text{Æ} = 4713 \\ + 1 \\ \hline 4714 \\ P = 324 - \end{array}$$

The Year of the Julian Period, Y = 4390 required. current.

EXAMPLE 2. To find P, the present Date, current before Christ, from the Year, Y, of a given Æra, the 93d current of the Building of Rome, and the Beginning of that Æra, Æ, 753 Years current before Christ?

By the 2d Equation.

$$\begin{array}{r} \text{Æ} = 753 \\ + 1 \\ \hline 754 \\ Y = 93 - \end{array}$$

Present Date before Christ, P = 661 required. current.

EXAMPLE 3. To find the Date, Æ, before Christ, of the Beginning of an Æra, from Y, the Years of that Æra, 280 current since it began, and P, the present Date, in the 45th Year current before Christ, being given?

By the 3d Equation.

$$\begin{array}{r} P = 45 \\ Y = 280 \\ \hline 325 \\ - 1 \\ \hline \end{array}$$

The Æra, the Death of Alexander the Great 324 bef. Chr. current, reqd.

OF ÆRAS beginning BEFORE, and DATES SINCE CHRIST.

PROPOSITION II. To determine the Years, Y, of an Æra, Æ, beginning at a given Date, Julian Style, before Christ, answerable to a present Year's Date, P, Julian Style, since Christ; and thence to give other chronological Rules?

RULE. By the former of the two chronological Scales, it is evident, that If 1 less than the Years current since Christ (being the Years complete) be added to the Years current of the Commencement of an Æra, before Christ, the Sum thereof will be the Years complete of that Æra, since it began, at Return of the same Month-Day.

And therefore 1 being added to those complete Years (by Axiom 2) will be the Years current of that Æra.

Otherwise, the Number of the present Julian Year current, since Christ, being added to the Number of Years current, before Christ, at the Commencement of the Æra, the Sum thereof will be the Number of the Year current of that Æra, since it began.

HENCE, we have these EQUATIONS.
$$\left. \begin{array}{l} 1. \text{Æ} + P = Y \\ 2. Y - \text{Æ} = P \\ 3. Y - P = \text{Æ} \end{array} \right\} \begin{array}{l} \text{Julian} \\ \text{Years} \\ \text{current.} \end{array}$$

EXAMPLE 1. To find Y, the Number of the Julian Year of the Greek Olympiads, whose Æra, Æ, began in the 776th Julian Year current before Christ, answering to the present Year, P, 1760, since Christ, current?

By the 1st Equation.

$$\begin{array}{r} \text{Æ} = 776 \\ P = 1760 \end{array}$$

Julian Year of the Greek Olympiads . . Y = 2536 current, req.

EXAMPLE 2. To find P, the present Date, current since Christ, from the Number of the Year Y, of a given Æra, 2507 Years current, of Nabonassar, the Beginning of which Æra, Æ, being 747 Years before Christ?

By the 2d Equation.

$$\begin{array}{r} Y = 2507 \\ \text{Æ} = 747 \end{array}$$

Present Date since Christ, P = 1760 required. current.

EXAMPLE 3. To find the Date, Æ, before Christ, of the Beginning of an Æra, from Y, the Years of that Æra, 1815, current since it began; and P, the present Year's Date, in the 1770th Year current, since Christ, being given?

By the 3d Equation.

$$\begin{array}{r} Y = 1815 \\ P = 1770 \end{array}$$

The Æra, at Julius Caesar's reforming the Calendar 45 required. current bef. Chr.

OF ÆRAS beginning and DATES, BOTH SINCE CHRIST.

PROPOSITION III. To determine the Years, Y, of an Æra, Æ, beginning at a given Date Julian Style, since Christ, answerable to a given present Year's Date P, Julian Style, since Christ, and thence to give other chronological Rules.

RULE. By the latter of the two chronological Scales, it is evident, that, If you deduct the whole Years, of a Date at the Beginning of an Æra, from the whole Years of a present Date, the Difference in whole Years will be the whole Years complete since that Æra began, and consequently 1 Year added thereto, will be the whole Years current of that Æra; which are completed at the Return of the same Month-day, since Christ.

HENCE, we have these EQUATIONS.
$$\left. \begin{array}{l} 1. P - \text{Æ} + 1 = Y \\ 2. Y + \text{Æ} - 1 = P \\ 3. P - Y + 1 = \text{Æ} \end{array} \right\} \begin{array}{l} \text{Julian} \\ \text{Years} \\ \text{current.} \end{array}$$

In the Account of Time, by the Julian Period, all other Rules are reduced to these Equations.

EXAMPLE

EXAMPLE 1. To find the Number of the Julian Year current, Y, of the Turkish Hegira, whose Æra, Æ, began in the 622d Year current since Christ, answering to the present Date, P, 1765 current, since Christ?

$$\begin{array}{r} P = 1765 \\ + 1 \\ \hline 1766 \\ E = 622 \end{array}$$

EXAMPLE 2. To find P, the present Date current since Christ, from the Year, Y, of a given *Æra* 1129 Years Julian of the Persian *Jesdegrid*, current; the Beginning of which *Æra*, *Æ*, being in the 632d Year current, since Christ?

$$\begin{array}{r} Y = 1129 \\ \text{Æ} = 632 \\ \hline 1761 \\ \hline \text{I} \end{array}$$

EXAMPLE 3. To find the Date, $\mathcal{A}$, since Christ, of the Beginning of an $\mathcal{A}$ ra, from Y, the Years of that $\mathcal{A}$ ra, 1477, current since it began; and P, the present Year's Date, in the 1760 Year current, since Christ, being given?

$$\begin{array}{r} P = 1760 \\ Y = 1477 \\ \hline 283 \\ + 1 \end{array}$$

Those who desire greater Exactness than this near Way of answering these chronological Cases, by whole Years current, may apply the Years and Days complete, of one *Æra*, at the Commencement of other *Æras*.

RULE II. 'Take the *Sum* of the *Dates*, in complete Years and Days, being one *Date* before and one *Date* since Chrif.

EXAMPLE 1. *Alexander the Great* died }
in the 324th Year current, *before Christ*, on } Yrs. Ds.
the 12th of *November* } 323 . 315 complete.

Julius Caesar reformed the *Calendar*, in the }
the 45th Year current, *before Christ*, on the 1st }
of *January* } 44 . 0 complete.

Diff. 279 . 315 complete.

Therefore, when *Julius Cæſar* reformed the *Calendar*, it was in the 280th *Julian Year* current or 279th *Julian Year* and 315 Days complete, of the *Æra* of *Alexander's Death*.

By PROP. I. *Equat. 1.* $\begin{matrix} \text{Æ} = & \text{Yrs.} \\ & 324 \\ \text{P} = & -45 \end{matrix} \left. \vphantom{\begin{matrix} \text{Æ} \\ \text{P} \end{matrix}} \right\} \text{current before Christ,}$

279 + 1 = 280 Year current.

EXAMPLE 2. The Foundation of Rome, in the 753^d Year current, before Christ, on the 21st Day of April } Yrs. Ds.
Turkish Hegira, begun in the 622^d Year } 752. 254 complete.
current, since Christ, on July 16 } 621. 196 complete.

Sum 1374. 85 complete.

Therefore, the *Turkish Hegira* (see *Tab. p. 129.*) commenced on the 1375th Year current of the Building of Rome, or on the 1374th Year and 85th Day thereof, complete.

By PROP. II. *Equat. 1.* $\begin{matrix} \text{Æ} \\ \text{P} \end{matrix} = \begin{matrix} 753 \\ 622 \end{matrix} \left. \begin{matrix} \text{Yrs.} \\ \text{current before} \\ \text{current since} \end{matrix} \right\} \text{Christ.}$

EXAMPLE 3. The *Æra* of *Martyrs*, or }
Dioclesian, begun in the 284th *Julian Year* } Yrs. Ds.
current since Christ, on *August 29* . . . } 283 . 241 complete.
The *Æra* of the Reformation of the *Ju-* }
lian Calendar by Pope *Gregory* began in 1582 }
since Christ, on *October 5* } 1581 . 278 complete.

Dif. 1298. 37 complete.

Therefore, the *Grigorian Æra* commenced in the 1299th *Julian Year* current of the *Æra* of Martyrs, in the 1298th Year thereof and 37 Days complete.

By PROP. II. *Equat.* 1.
$$\begin{array}{r} \text{P} = 1582 \\ \text{Æ} = 284 \\ \hline 1298 + 1 = 1299 \text{ Years current} \\ \text{[since Christ.]} \end{array}$$

N. B. Hence the Truth of the foregoing Rules, for answering chronological Questions, in whole Years, are proved.

☞ When the Days complete, in any Julian Date, are less than those, in another Julian Date to be taken from them, you must subtract 1 Year from the Answers, in whole Years current, had from the Equations, for the true whole Number of Years current.

☞ By taking the Years and Days *Difference complete* betwixt any former and succeeding *Dates*, you may reduce the *Julian Account* of Time to any other Account of Time you please, in *Years and Days*, and the contrary: Especially by a Table of the Days in the Years and Months, according to the new Account; like what is done in Page 350 foregoing.

RULE. Subtract 1 from the current *Olympiads* for the complete Number thereof; multiply the *Remainder* by 4, adding thereto 1 less than the Number of the odd Olympic Years current for the Number of the Olympic Years complete.

EXAMPLE 1. *In the 3d Year of the 16th Olympiad.*

$15 \times 4 = 60$
 Add 1 less than the No. of odd Olymp. Yrs. $+ 2$

Complete Olympic Years 62 required.
Or current 63.

REDUCTION of CHRONOLOGY.

EXAMPLE 2. In the 1st Year of the 195th Olympiad current?

$$\begin{array}{r} - 1 \\ 194 \times 4 = 776 \text{ Years complete.} \\ \text{Or current 777.} \end{array}$$

Now, these Olympic Years, Y, being given since the Æra of the Olympiads, Æ, 776 Years current before Christ, began, are reduced into P, the present Date, before or since Christ, by the former Equations, as have been shewn, in the I. and II. PROPOSITIONS.

And P, Years current given before or since Christ, may be reduced into Y, Olympic Years, from the Olympic Æra, beginning 776 Years current before Christ; and then those Olympic Years, Y, may be reduced into Olympiads, and odd Years complete and current, by the Converse of the foregoing Rule for reducing Olympiads into Olympic Years, as follows.

RULE. For reducing Olympic Years complete into OLYMPIADS.

Divide the Olympic Years complete by 4, and the Quotient will be the Olympiads, complete; to which add 1, for the Olympiads current: Then add 1 to the Remainder for the odd Olympic Years current.

EXAMPLE 3. 63 Olympic Years current.

$$\begin{array}{r} \text{complete.} \\ 4 \overline{) 63} \begin{array}{l} 15 + 1 = 16 \text{ Olympiads current.} \\ 60 \\ \hline 2 + 1 = 3 \text{ Olympic Years current.} \end{array} \end{array}$$

Olympiads are reduced into Olympic Years, Y, and then into Years of the Julian Period, P, by considering the Æra of Olympiads, Æ, beginning its Æra in the 3938th Year of the Julian Period; coming under PROPOSITION III. and Equation 2.

EXAMPLE 4. To find the present Year, P, of the Julian Period, answering to the 2490th Year, Y, current of the Olympic Æra, Æ, that begun in the 3938th Year current of the Julian Period?

$$\begin{array}{r} \text{By Prop. III. Equat. 2.} \quad \begin{array}{r} Y = 2490 \\ \hline \text{Æ} = 3938 \\ \hline 6428 \\ - 1 \\ \hline \end{array} \end{array}$$

Current Years of the Julian Period, required P = 6427

CONTRARILY, P and Æ being given, Y the Olympic Years are thence found, and may be reduced to Olympiads and Years Olympic, complete and current.

PROPOSITION VI. To find the Cycles of the Sun, Moon, and Indiction, answering to the Olympiads?

RULE. Reduce the Olympiads to the Olympic Years, current, add thereto 17, 4, and 7, and divide by 28, 19, and 15, respectively, and the remaining Numbers will be those of the respective Cycles, required.

EXAMPLE. To find the respective Cycles to the 3d Year of the 623d Olympiad, or 2491 Olympic Years current.

$$28 \overline{) 2508} (89 \quad \quad 19 \overline{) 2495} (131$$

$$\text{Cycle } \odot . . 16 \quad \text{Cycle } \text{D} . . 6$$

$$15 \overline{) 2498} (166$$

Indiction . . 8 required.

Answering to the 1715th Julian Year current since Christ.

$$\left. \begin{array}{l} \text{For, } 2491 + 17 = 2508 \\ \quad \quad + 4 = 2495 \\ \quad \quad + 7 = 2498 \end{array} \right\}$$

The Olympic Year, commencing in July, the Moon after the Summer Solstice, has 2 Cycles, answering to two successive Years of the Julian Period, (each commencing with January) the former Cycle belonging to the first, and the latter Cycle to the last, Half of the same Olympic Year.

PROPOSITION VII. To find the Number of Egyptian Years current, of Nabonassar, answering to any Julian Year current, since Christ; O. S.?

RULE. Add 747, (the Date of the Julian Year current before Christ, when the Nabonassar-Æra began, on the 26th of February, being on the 1st Day of the Egyptian Month, Tbeth) to the present Date, since Christ; and the Sum thereof will give you the Number of Julian Years since that Æra; (by Prop. II. and Equat. 1.)

NOW, Subtract from which Sum 2 (because the 747th Year before Christ is the 2d Year after Bissextile) and divide the Remainder by 4, to discover the Number of Bissextiles happening since that Æra; then add the Number of Days expressed by the Quotient, (or 1 more if Nothing remains) to the preceding Number of Bissextile Years, and the Result thereof will give the Egyptian Years required; complete to the last Return of the same Month-Day, on which the Æra began; when those Years begin to be current.

But the Plus Days, above a Year, or Years, of 365 Days, must be reckoned back from the 25th of February, at Noon, to find the Month-day in the Julian Style, to which the Commencement of the Egyptian Year is now fallen back. This is easily performed by the Help of the Table, P. 129.

And, If the Number of the Egyptian Year and Month-day thereof be required for any Julian Month-day preceding, or following, that on which the Æra of Nabonassar began, you must deduct, or add, the preceding or succeeding Number of Days thereof, respectively, from and to the before found Days which the Æra's Commencement has fallen back, in this Julian Account, and you will have the Number of the Egyptian Years and Days current, required, correspondent to the Julian Month-day and Year given.

If Nothing remains, after the Julian Years, less 2, are divided by 4, then 1 must be added to the Quotient, for the true Number of Bissextiles

EXAMPLE. To find the Number of Egyptian Years, Y, of the Æra of Nabonassar, Æ, commencing February 26, in the 747th Year Julian current, before Christ, answering to P, the present Date of July 16, 1760, current since Christ, Julian, or O. S.?

$$\begin{array}{r} \text{By PROP. II. Equat. 1.} \quad \begin{array}{r} P = 1760 \\ \hline \text{Æ} = 747 \end{array} \end{array}$$

$$Y = 2507 \text{ current.}$$

$$\begin{array}{r} - 2 \\ \hline 4 \overline{) 2505} (626 \\ \text{Rem. 1} \end{array}$$

$$1 \text{ Year Egyptian} . . - 365$$

Days to be reckoned back fr. Feb. 26, the Æra's falling back, 261

From February 26, to July 16, forward . . Days 140

$$\begin{array}{r} \text{Years } 2507 \text{ Julian.} \quad 401 \\ \text{Years } + 2 \text{ Egyptian.} \quad 1 \text{ Egyptian Year} - 365 \end{array}$$

$$\begin{array}{r} \text{Years } 2509 \text{ Egyptian, reqd. and 36 Days} \quad \text{Days} . . 36 \\ \text{since that Year's Commence-} \\ \text{ment; being the 6th Day of the} \\ \text{Month Paophi, current.} \end{array}$$

REDUCTION of CHRONOLOGY.

But, (by Tab. P. 129) to February 25 . . . 56 —, in a com. Yr.
To be reckoned back 261 in both Yrs.

To be reckoned back in the preceding Year 205 —
365

Corresponding with the 9th of June, in the }
Year of Christ 1760, on which the E- }
gyptian Year now begins } 160 reckoning forward.
From whence reckoning . . . 140 Days more forward.

October 27, 1760, Julian, O. S. . . 300, by Tab. P. 129.
On which, July 16, in the 1st Egyptian Year of Nabonassar, now
happens, by falling back, in the Julian Account.

N. B. The Month-day of any Julian Year, before Christ, may be de-
termined in the same Manner, on which any Month-day of the 1st Egyp-
tian Year of Nabonassar happens; only, instead of adding 747 to the Date
(as directed) since Christ, you must now subtract the Date before Christ
from 748, to find the Julian Year current of the Nabonassar Æra; (by
PROP. I. Equat. 1.) and then subduct 2 from the Remainder, which
Result must then be divided by 4, to find the Number of Bissextiles; and so
proceeding to find the Egyptian Year current, and the Month-day of the
Julian Year, on which its Commencement falls back, as before.

OTHERWISE. To find the Egyptian Year and Month-
day current of the Æra of Nabonassar, commencing February
26, 747 Years current before Christ, answering to the Date
of the Christian Æra, of July 16, O. S. 1760, current, or
July 27, 1760, N. S.

DATE SINCE CHRIST.

Jul. Yrs. comp.	Days.
1000	365250
700	255675
56	20454
3	1095

Julian Years 1759 complete.

June, complete . . . 181
In July, Days complete . . . 15

Since Christ, to the present Julian Date . . . 642670
Before Christ, at the Æra of Nabonassar . . . 272785

Sum . . . 915455 Days
[complete.]

Egyptian Years 2000 . . . 730000 —

500 . . . 185455 —
182500 —

8 . . . 2955 —
2920 —

Egyptian Years 2508 complete, and . . . 35 Days

Thoth, complete 30 — comp.
[comp.]

Paophi 5 Day
Viz. in 2509 Egyptian Years current, on 6 Day of
Paophi. Current.

ÆRA BEFORE CHRIST.

Julian Yrs comp.	Days.
700	255675
44	16071
2	730

Julian Years 746 complete.

From the 25th of Feb. to the Years End 309 complete.

Before Christ, at the Beginning of the }
Æra of Nabonassar . . . } 272785 Days complete.

Mr. Ferguson, in his *Astronomy*, has given us August 29, O. S.
on which the first Day of the Egyptian Month Thoth happened, (and
which we have also observed) but he has not given us the Year of Christ,
or Year of the Julian Period, nor any Point of Time, when this Event
happened, which we have here † supplied below.

REMARK. The Beginning of the Æra of Alexander the Great,
on November 12, on the 324 Julian Year current, before Christ, was
on the 1st Day of the Egyptian Month Thoth; as also was the Begin-
ning of the famous Æra of Nabonassar, so well known among the
ANTIEN ASTRONOMERS; because King PTOLEMY, and the
Rest, adjusted all their *Astronomical Computations*, by the Years of that
Æra, according to the Egyptian Style, so very fit for their Purpose.

Likewise, the Beginning of the Æra of Martyrs, or of the Diocle-
sian Æra, on August 29, in the 284th † Julian Year, current, since
Christ, fell on the 1st of the Egyptian Month Thoth; which is the
Egyptian Æra given us by Mr. Ferguson, in his *Astronomy*; though he
has not given us the Julian Date, or Year, correspondent.

Those who would reduce any Date, or Year of an Æra, in the Ju-
lian Style, before or since Christ, to the Year and Month-day of any
other Æra, (and the contrary) according to the Difference between the
Julian and Egyptian Years, must follow the foregoing Rules of the VII.
PROPOSITION, (and the Converse thereof, Mutatis Mutandis).

THUS have we completed THE WHOLE BUSINESS OF
CHRONOLOGY, (as promised in former Pages) so far as relates to
the calculative or most useful Part; to which the historical Part is only
wanting, to perfect the Subject, with which our *Astronomy* is so closely
connected. See Sir Is. Newton's *Chronology*.

INTRODUCTION

TO THE

COMMENT

ON THE

MOON'S THEORY.

I. DEFECT of ASTRONOMICAL TABLES, *constructed from OBSERVATIONS.*

TO gain, from *Observation*, such a complete Knowledge of the *lunar Motions*, as to be able from thence to frame *Tables* accurately enough to represent the Moon's true *Place*, or *Longitude*, in every Point of her Orbit, during an entire Revolution of her *Apogee* and *Node*, through the Space of near 19 Years, has hitherto proved ineffectual Labour. For, though the *lunar Tables* framed, from Time to Time, from such *lax* and *sparing* Observations, as the Astronomers of the *last Century* contented themselves with making, agree sometimes, even in the *Oscillants* and *Quadratures*, surprizingly with the Heavens, yet, in *general*, it has happened much otherwise. And they have been found to deviate, so widely, from Observation, that no Use can be made of them for obtaining the *Longitude to the Degree of Exactness necessary for the Purpose of Navigation*: Not even the celebrated *Tables* of Dr. Halley, imperfectly founded on the *Theory* of the great Sir ISAAC NEWTON.

1. *Such Tables in vain.*

2. *Unfit for the Purpose of Navigation.*

II. DEFECT of TABLES *supplied by OBSERVATIONS.*

YET this Defect, as this great Astronomer, Dr. Halley, observes, is not the Fault of the *Art*, but of the *Artificer*. For, by observing the *Period* of the *lunar Inequalities*, which is performed in *eighteen* Years and *eleven* Days, or two Hundred and *twenty-three* LUNATIONS, it is found that the Return of ECLIPSES, and other *Phænomena* of the Moon's Motion, are very regularly performed. That whatever *Error* is found in a former *Period*, the same is again repeated in a succeeding one, under the like Circumstances, of the Distance of the Moon from the Sun, and of the Sun from the Moon's *Apogee*.

But, then to gain a certain and distinct Notion of all these *Irregularities*, which, from Observation, throughout this *Period*, disturb the *lunar Motions*, would not only exceed the Power of any one Person, in the short Duration of his Existence, but would necessarily be the Work of Ages to perform; it requiring no fewer than 194400 Observations to complete the *lunar Theory*; being $= 360 \times 60 \times 9$; that is to every Minute in the *Zodiac*, for one Revolution of the *Apogee*.

Dr. Halley, to correct the Computation from his *Tables*, has given an *Abacus* of the *Errors* observed, throughout one *Period*, joined to a correct Series of Observations from whence they are deduced.

III. Of TABLES CONSTRUCTED according to the THEORY.

ON the other Hand, to deduce the Theory of the Moon, *à priori*, or to shew, from the Laws of Gravity and Attraction, the Manner in which the Motion of a *Satellite*, revolving round its Primary, is disturbed by the SUN, the Quantity of that disturbing Force, and its Proportion to the *centripetal* Force, by which it is retained in its proper

INTRODUCTION to the COMMENT.

Orbit, with the Nature and Quantity of the Errors thence arising, both in *Longitude* and *Latitude*, is an Attempt, which neither the *Mathematicians* of remote Ages, or those of more modern *Date*, since the Revival of Learning in the *West*, have made the Object of their Studies and Inquiries. This wonderful *Exertion* of the Powers of the Mind, greatly beyond what the human Understanding has before been thought equal to, was reserved for *Newton*. For *Him* it was destined to raise the *English* Reputation in the *cælestial Sciences* above that of other Nations, and to teach the admiring World, the true *Causes* of the Order and Harmony observed throughout our PLANETARY SYSTEM.

“ *Discimus hinc tandem, quâ causâ argentea Phœbe*
 “ *Passibus haud æquis, graditur: cur subdita nulli*
 “ *Hactenus astronomo, numerorum fræna recusat:*
 “ *Cur remeant nodi; curque auges progrediuntur.*”

Halleius in *Newtonum*.

Every *Englishman* who values his Country, and who is emulous of having its Reputation eminently raised in the Opinion of other Nations, for the Progress made in every Branch of *Literature*, and of *Philosophy* in particular, must venerate the Name of *Newton*, for establishing that Reputation on the most solid Foundation. And there are few who will blame this Attempt of making him more universally known and understood upon a Subject, the most useful of any in the Science of *Astronomy*; though the most perplexing and difficult to understand! I mean the true Theory of the *lunar Motions*, deduced from the Theory of Gravity by our great *Mathematician*!

1. *Discovery and Honour of the Moon's Theory, Sir Isaac Newton's.*

IV. ACCOUNT of the SUBSEQUENT COMMENT.

IN the *Progress* of the subsequent *Comment*, the COMMENTATOR has endeavoured to be clear as well as concise: closely following Sir *Isaac Newton's* Arguments. He has traced out from thence the general *Principles* on which the several *Theorems* are established, and from whence the different *Problems* are deduced, without entering into the *fluxionary Method* in their Solution: giving only, in general, the *Result* of the Computations, as they are derived from the *Principles of Motion*. To have gone further, had been, at once, to copy all that our Author has wrote upon this Subject, who reasons too closely for his Reasoning to be abridged, and too clearly for giving the Sense of his Subject in fewer or fitter Words. What is here attempted, will, I hope, answer the Purpose for which it is given the Public. It is intended as a *Foundation* for better *lunar Tables* than yet have been published; such as may properly be called TABULÆ verè NEWTONIANÆ. And these Tables deviate only in very minute Particulars from the Theory, as Sir *Isaac* left it; having but three † additional Equations; except the Equation which corrects the Moon's Variation, according to her Anomaly, may be admitted as additional also; which cannot entirely be so esteemed, as our Author himself points it out to the Consideration of future Astronomers.

2. *Principles of the Moon's Theory.*

3. *Design of the Comment and Tables.*

4. *Three Equations only added to the Moon's Theory.*

S. C.

† These three additional Equations are reduced to One. For the VI and VII, are only different Parts of the same Equation, and the latter Part Supplemental of the First. And the IX, or Evection-Equation, is no other than the second Equation of the Apogee, and of the Eccentricity, transferred to the Moon's Anomaly, or, indeed, is but a Part of the whole central Equation, according to *Newton*; the Mean Central being the other Part.

A N

Explanatory Comment

ON THE

MOON'S THEORY.

According to which PRINCIPLES, the Astronomical TABLES at the BEGINNING of THIS WORK (preceding the SUPPLEMENTAL TABLES) are CONSTRUCTED.

I. Of the MOON'S MOTION.

IN establishing the *Theory of the Moon*, from the Laws of *centripetal Forces*, as laid down in the *first Book* of the *Principia*, our *Author* first considers the Moon as revolving round the Earth, in the same Manner as the Earth, or any other of the *primary Planets*, revolves round the Sun; and from the *Phænomena*, or observed Motion, or Distance of the Moon, demonstrates, that the Moon, in revolving round the Earth, preserves that established Law of Motion, in describing *Areas proportionable to the Times*, no less than the *primary Planets* in their respective Revolutions round the Sun: neglecting however those inconsiderable Errors which arise from the Action of the Sun upon the Moon in her Orbit.

1. Moon describes Areas round the Earth nearly proportionable to the Times.

2. Law of the Moon's Motion, and of her centripetal Force.

Our *Author* next observes, that the Force, by which the Moon is retained in her Orbit, tends towards the Earth, agreeable to the Law of *centripetal Forces*. (B. I.) And also, that the Force is, *reciprocally*, as the Squares of the Distance from the Earth's Center; as is the more evident from the very slow Motion of the Moon's *Apsides*, amounting but to $3^{\circ} 3'$ at a mean

Rate in one Revolution of the Moon.

He further demonstrates, that the Moon gravitates towards the Earth, and by that Force of Gravity is continually drawn off from a *rectilinear Motion*, and thereby retained in her Orbit. This Property of the Moon's Motion is partly to be deduced from the foregoing Proposition; but more extensively from the following Theorem, which shews, that the Force, by which she is retained in her Orbit, is no other than that which we call Gravity, and by which all Bodies are made to descend towards the Earth.

The correct mean Distance of the Moon from the Earth, according to Observation, is $60\frac{1}{2}$ of the Earth's Semidiameters. Let us, says our *Author*, assume the mean Distance of 60 Semidiameters in the Syzies; and suppose one Revolution of the Moon, in respect of the fixed Stars, to be completed in $27^d 7^h 43'$, as Astronomers have determined; and the Circumference of the Earth to be equal to 123249600 Paris Feet, as the French have found by Mensuration. And, now, if we

imagine the Moon, deprived of all revolving Motion, to be let go, so as to descend towards the Earth, with the Impulse of all that Force, by which she is retained in her Orbit, she will, in the Space of one Minute of Time, describe, in her Fall, $15\frac{1}{2}$ Paris Feet. For, the versed Sine of that Arc which the Moon, in the Space of one Minute of Time, would, by its mean progressive Motion, describe, at the Distance of 60 Semidiameters of the Earth, is accurately 15 Feet, 1 Inch, and 1 Line $\frac{2}{3}$. Wherefore, since that Force, in approaching the Earth, increases in the reciprocal duplicate Proportion of the Distance, and upon that Account, at the Surface of the

Earth,

COMMENT on the MOON'S THEORY.

Earth, is 60×60 Times greater than at the Moon; a *Body*, in our Regions, falling with that Force, ought, in the Space of one Minute of Time, to describe $60 \times 60 \times 15\frac{1}{2}$ Paris Feet; and in the Space of one Second of Time $15\frac{1}{2}$ Paris Feet: And, with this very Force, we actually find Bodies, here upon Earth, do really descend. For, a *Pendulum*, oscillating Seconds, in the Latitude of Paris, will be 3 Paris Feet and $8\frac{1}{2}$ Lines in Length, as *Huygens* has observed: And the Space which a heavy Body describes, by falling, in one Second of Time, is to Half the Length of the *Pendulum*, in the duplicate Ratio of the Circumference to the Diameter of a Circle, and is therefore 15 Feet, 1 Inch, and 1 Line $\frac{2}{3}$. And, therefore, the Force, by which the Moon is retained in her Orbit, becomes, at the very Surface of the Earth, equal to the Force of Gravity, which we observe in heavy Bodies there. And hence the Force, by which the Moon is retained in her Orbit, is that very same Force which we call Gravity, and no other. For, were Gravity another Force, different from that, then Bodies would descend to the Earth with the joint Impulse of both Forces, and descend with double Velocity, and in one Second of Time described $30\frac{1}{2}$ Paris Feet, altogether against Experience.

OUR AUTHOR thus demonstrating, that the Moon is retained in her Orbit by Gravity; and, by Analogy of Reasoning, explaining likewise to us, that the same Sort of Force retains the *Primaries*, and all their *Secondaries*, in their respective Orbits; he refers us, next of all, to those Principles already laid down, from the Theory of Gravity, in the first Book, as the Foundation on which the lunar Theory is built. And observes, that all the Motions of the Moon, her *Apogee*, and *Nodes*, are to be derived from thence, as well as the Inequalities of those Motions from the Action of the Sun upon the Earth, the Primary, as well as on the Moon, the Secondary Planet.

The several Inequalities of the Moon, from a mean Motion, shall be considered, when we have attended to that Proposition, (B. III. 25.) which assigns the Quantity of the solar Force on the Moon, by which her Motion round the Earth is accelerated or retarded. For, having the Quantity of that disturbing Force, and the Proportion it bears to that by which it is retained in its Orbit, we shall know better how to compute the lunar Inequalities, and to find the Agreement of the Computation with Observation.

II. To FIND the FORCES with which the SUN disturbs the MOTION of the MOON.

IN the annexed Figure (see Prop. 25. B. III.) let S represent the Sun; T, the Earth; P, the Moon; A, C, B, D, the Moon's Orbit: In SP continued take SK, equal to ST, and let SL be to SK, in the duplicate Ratio of SK to SP; draw LM parallel to PT; and if SK or ST represent the accelerative Force or Gravity of the Earth towards the Sun (for on Account of the Smallness of the Angle PST these Lines may be esteemed equal to each other) then SL will represent the accelerative Force of the Moon towards the Sun, when at P. But, that Force is compounded of the Parts SM, and LM; of which the Force LM, and that Part of SM, represented by TM, disturb the Motion of the Moon. For, as the Earth and Moon are revolved about their common Center




Fig. I.

of Gravity, the Motion of the Earth will be also disturbed by the like Forces. But we may consider the Sums of both the Forces and Motions, as in the Moon, and represent the Sum of the Forces by the Lines TM and LM. The Force LM in its mean Quantity, is to the centripetal Force, by which the Moon is retained in its Orbit, revolving round the Earth, supposed at Rest, at the Distance PT, in the duplicate Proportion of the periodical Time of the Moon about the Earth, to the periodical Time of the Earth about the Sun, that is, in the duplicate Proportion of $27^d 7^h 43'$ to $365^d 6^h 9'$: or as 1000 to 178725, or as 1 to 178 $\frac{2}{3}$. But, the mean Force LM, is to the Force of Gravity on the Surface of our Earth, (assuming the Moon's mean Distance to be $60\frac{1}{2}$ Semidiameters of the Earth) as $1 \times 60\frac{1}{2}$ to $60 \times 60 \times 60 \times 178\frac{2}{3}$, or as 1 to 6380926; whence, the Proportion of TM, LM, TM, is also given.

For, by Prop. IV. B. 3. If both the Earth and Moon were revolved about their common Center of Gravity, the mean Distance of one from the other would be nearly $60\frac{1}{2}$ mean Semidiameters of the Earth; and the Force by which the Moon may be kept revolving in its Orbit about the Earth Quiescent, at the Distance, PT, of $60\frac{1}{2}$ Semidiameters of the Earth, is to the Force by which it may be revolved at the same Time, at 60 Semidiameters, as $1 \times 60\frac{1}{2}$ to 60; and this Force is to the Force of Gravity with us very nearly, as 1 to 60×60 .

The Proportion of the solar Force to that which retains the Moon in her Orbit being thus given, the next Thing, to be inquired into, is the Manner in which this solar Force affects the Moon's Motion, and the Change it produces in the Form of the lunar Orbit; which are included in the following Propositions and Theorems of this Third Book.

COMMENT on the MOON'S THEORY.

III. To determine the SOLAR FORCE producing the CHANGES and INEQUALITIES in the LUNAR ORBIT.

SIR Isaac Newton has demonstrated (Prop. XXVI. B. 3.) that the Areas, which in equal Times are described by the Moon, in the Syzgies and Quadratures, are to each other, respectively, as 11965 to 11865, supposing the Earth quiescent, and the Moon performing one Revolution in $27^d 7^h 43'$. But the Synodical Month consists of $29^d 12^h 44'$, therefore, the Increments of the Motion must be enlarged in the same Proportion; and the Moment of the Area in the Syzgie will be to the Moment of the Area in the Quadrature, as 11073 to 10973.

1. A Deviation from equal Areas described in equal Times, by the Moon.

But, to understand more fully those Principles on which these Propositions are established, we must go back to the FIRST BOOK of the PRINCIPIA, where, in the second Corollary of the 66th PROPOSITION, Sir ISAAC NEWTON deduces, that in a SYSTEM of THREE BODIES, T, P, S, (see Fig. I.) If the accelerative Attractions of any two of them, towards a Third, be to each other reciprocally, as the Squares of the Distances, the Body P, revolving round T, will describe its Area swifter near the Conjunction A, and the Opposition B, than it will in the Quadrature C, or D.

2. Properties of Motion, in a System of 3 Bodies, explained.

For, every Force, with which the Body P, is acted on, and not the Body T, and which does not act in the Direction of the Line PT, does either accelerate, or retard the Description of the Area, according as it is directed in Consequentia, or Antecedentia: such is the Force TM. This Force, in the Passage of the Body P, from C to A, is directed in Consequentia of its Motion, and therefore accelerates it; then as far as D, in Antecedentia, retards it; then in Consequentia, as far as B, the Motion is again accelerated. And, lastly, from B to C, in Antecedentia, is again retarded. Therefore, *ceteris paribus*, the Planet P must move swifter in the Conjunctions than in the Quadratures.

Hence also the Orbit of the Body P, (Cor. 4. Prop. same) *ceteris paribus*, is more curved at the Quadratures than at the Conjunction and Opposition; for the swifter Bodies move the less they deflect from a straight Line. And, besides, the Force KL, or TM, is contrary, at the Conjunctions, to the Force with which the Body T attracts the Body P; and therefore diminishes that Force: But the Body P will be less deflected from its rectilinear Course the less it is impelled towards the Body T.

Hence, lastly, (see Cor. 5) *ceteris paribus*, supposing the Orbit to be circular, the Body P recedes farther from the Body T, in the Quadratures than in the Conjunctions: For, if the Orbit is eccentric, its Eccentricity will be greatest (see Cor. 9) when the Apfides are in the Syzgies; and then the Body P, at the higher Apfis, may go farther from the Body T, placed in or near the lower Focus, than at the Quadratures.

That the Decrease of the centripetal Force, at the Conjunctions, should be the Cause of the Body P, coming nearer to the Earth; and the Increase, of the same Force, at the Quadratures the Cause of the Body P receding then farthest from it, Dr. Rutherford observes, appears to be a Paradox; and to which he gives the following Solution.

3. An astronomical Paradox.

When the Moon, or Body P, is at A, the ablatitious Force is greatest of all; and the Moon, or Body P, is less impelled towards the Earth, than the Force of Gravity would impell it, and, consequently, it will run farther from the Earth than if the centripetal Force of the Earth, only, acted upon it, and therefore when it has reached the Quadrature D, will be at a greater Distance. When, on the other Hand, it has arrived there, the centripetal Force is increased by the additious Force, expressed by LM, or PT. By this Cause, the Course of the Body P, is more incurvated towards the Earth; and when it is arrived at B, approaches nearer to it, than if the solar Force had not disturbed its Motion.

4. Solution of the Paradox.

IV. To determine the RATIO of the DIAMETERS of the MOON'S ORBIT without ECCENTRICITY.

THE Curvature of the Moon's Orbit, or the Proportion of the Diameter AB to CD, is deduced from the 28th Prop. of the 3d Book of the Principia, wherein Sir Isaac considers the Orbit without Eccentricity; and lays down, as his fundamental Principle, by which he solves this Problem, that the Curvature of the Orbit, described by a Body, attracted in Lines perpendicular to it, is as the Force of Attraction, directly, and the Square of the Velocity, inversely. And estimates the Curvature of Lines compared with one another, according to the evanescent Proportion of the Sines or Tangents of their Angles of Contact to equal Radii, supposing these Radii to be infinitely diminished.

The Attraction of the Moon towards the Earth in the Syzgies, is the Excess of its Gravity towards the Earth, above the solar Force, 2. PK. by which the Moon gravitates more towards the Sun than the Earth; and, in the Quadratures the Attraction of the Moon, towards the Earth, is the Sum of the Gravity of the Moon towards the Earth, or of PT and of the solar Force KT. In the former Case, the Excess of the Gravity will be $178725 - 2000 = 176725$; in the latter, the Sum of the Gravities will be $178725 +$

1. Solution of the Problem.

AT+CT

1000 = to 179725. For, putting N = to ——— 17825, equal to the accelerative Gravity of the Moon towards

2

by which the
is also attracted
towards the Earth

the

COMMENT on the MOON'S THEORY.

the Earth; the mean Force ML in the Quadratures = to PT, or TK = 1000, and the mean Force, TM, in Syzigies, will be equal to 3000; from which, taking the mean Force ML, there remains 2000 the Force by which the Moon is drawn from the Earth, and which before was called 2. PK. Now, the Velocity to the Moon in Syzigies A and B, is to the Velocity in the Quadratures C and D, as CT to AT. And the Moment of the Area described by the Moon in the Syzigies, to that described in the Quadratures, as 11073 CT, to 10973 AT. Taking this Ratio twice, *inversely*, and the former *once*, directly, and the Curvature of the Orbit in Syzigies will be to that in Quadrature, as $2151969 \text{ AT} \times \text{CT} \times \text{N} - 24081 \text{ AT}^3$ to $2191371 \text{ AT} \times \text{CT} \times \text{N} + 12261 \text{ CT}^3$. Our Author, proceeding, finds, by a *biquadratic* Equation, the Ratio of the *shortest* Diameter CD, of the lunar Orbit, to be to the *longest* AB, exclusive of Eccentricity, as 99281 to 100719, being the Ratio of the Moon's Distance in Syzigies, to her Distance in Quadrature; that is, in round Numbers, as 69 to 70.

Mr. Machin, in his *Treatise on the Law of the Moon's Motion*, subjoined to the *English Translation of the Principia*, makes the Proportion of the shorter to the longer Axis of the Moon's Orbit to be as 59 to 60.

But this Ratio, as he observes, is somewhat larger than the Proportion given by Newton; though not so large as that given from Observation by Dr. Halley, which is that of $44\frac{1}{2}$ to $45\frac{1}{2}$, or as 89 to 91.

The Transverse to the conjugate Diameter of the *elliptic Epicycle* (so called) of Machin is as 2 to 1; whose Center he supposes is carried to the *Left* round a circular Orbit, while the *Track* of the Moon's elliptic Orbit is carried round, in the Epicycle's Periphery, to the *Right*; describing equal Areas in equal Times, about the Center thereof; with the Epicycle's Transverse Diameter always parallel to the *Line of the Moon's Conjunction and Opposition*. The Moon is always at the remoter End of the shorter Axis of the elliptic Epicycle when in the *Quarters*; and at the near End (from the Center of the Moon's Orbit) at *New* and *Full*. See the *Figure and Explanation* farther on.

Mr. MACHIN gives these following SHORT and READY RULES for deducing these PROPORTIONS.

Let L be the *periodical* Time of the Moon; S, the Period of the Sun from the *fixed Stars*; M, the *synodical* Revolution of the Moon from the Earth; D, the Difference of the Periods of the Sun and Moon. Then the Difference of the two Axes will be to their Sum, as $3 \text{ L} \times \frac{\text{M} + \text{L}}{2}$ to 4 DD

— SS, by the Principles of Sir Isaac Newton; but 3 LL to $2 \text{ SS} - \text{LL}$, according to Machin's Principles: Whence $\text{SS} - 2 \text{ LL}$ to $\text{SS} + \text{LL}$, is his Proportion of the *less* to the *greater* Axis of the Moon's Orbit.

From the foregoing Data, the Diameters of the Moon's Orbit, and the Inequality of the Moments of the Area, which the Moon, by a Radius drawn to the Earth, describes, our great Mathematician next deduces that Inequality called

V. The MOON'S FIRST EQUATION of VARIATION or REFLECTION.

IT was first discovered by Tycho Brahe; but its true Cause never assigned, till our Author undertook to settle it from the Principles of Gravity.

If the Moon (as has been supposed) actually revolved in an *Ellipsis*, round the Earth, *quiescent*, placed in its Center, whose Diameters being as 69 to 70, and the Areas, by Radii drawn from the Earth's Center, were all proportionable to the Time of Description, then the *Tangent* of any Angle of its Motion, in that Orbit, would be to the *Tangent* of its mean Motion from the Quadrature, as 69 to 70. But, as our AUTHOR observes, the Description of the Areas from the Quadrature to the Conjunction, or Opposition, is accelerated, and ought to be in such Proportion, that the Moment of the Area in Syzigies, may be to the Moment of the Area in Quadrature, as 11073 to 10973. Which is readily brought to Computation, by diminishing the *Tangent* of this Angle from the Quadrature, in the *subduplicate* Proportion of 10973 to 11073, that is, in the Ratio of 68,6877 to 69. Hence, the Angle of the true Motion, from the Quadratures, will be, in the *Orbants*, (when the Angle of mean Motion is 45°) = $44^\circ 27' 28''$; the Difference between which, and the Angle of mean Motion, 45° , gives, for the MEAN VARIATION, at a Maximum, $32' 32''$. But this is, as I said, supposing the Earth *quiescent*. But as the Earth moves, and the Sun thereby is, *apparently*, carried in *Consequentia*, through the *Ecliptic*, the Moon, before it overtakes the Sun, describes an Angle greater than a Right-Angle, in its Progress from the Quadrature; and That, in Proportion of the periodical to the synodical Revolution of the Moon, $27^d 7^h 43^m$ to $29^d 12^h 44^m$. Whence, it comes to pass, that all the Angles about the Center of Motion are dilated in that Ratio, and the GREATEST VARIATION, which, in the former Supposition, amounts but to $32' 32''$, becomes $35' 10''$.

According to Mr. Machin's Principle, before laid down, the Diameters of the elliptic Orbit being to each other as 59 to 60, the VARIATION-EQUATION, at a Maximum, would be but $28' 53''$, which increased in the duplicate Ratio of the periodical to the synodical Revolution, or in the Square of $27^d 7^h 43^m$ to $29^d 12^h 44^m$, gives for the GREATEST VARIATION but $33' 45''$.

2. Ratio of the Diameters of the lunar Orbit, without Eccentricity, determined, according to Sir Isaac Newton.

3. Ratio according to Mr. Machin.

4. Ratio according to Dr. Halley.

5. Machin's Rules for determining the same.

1. Method of determining the Quantity thereof.

2. Greatest Quantity of Variation determined according to Sir Isaac Newton.

3. According to Mr. Machin.

Emendation. P. 359, L. 12, fr. Bot. for 1,000 r. 1000; L. 9 fr. Bot. for 178725, r. 178,725. L. 5 fr. Bot. same P. also, for, at the same Time, r. in the same Time.

P. 360, L. 5 fr. Bot. for, more towards the Sun than the Earth, r. more or less than the Earth towards the Sun.

Line 3, fr. Bot. after KT, introduce and r. by which the Moon is also attracted towards the Earth.

COMMENT on the MOON'S THEORY.

4. According to Dr. Halley. If Dr. Halley's Ratio, from Observation, of the shorter to the longer *Axis*, be taken as 89 to 91, it will give the VARIATION at a Maximum, $38' 12''$.

5. According to Mr. Mayer. Mr. Mayer, the Author of the *Göttingen Tables*, asserts, that the VARIATION, at a Maximum, is equal to the Half-Angle of EVECTION-EQUATION; that is, to $40' 20''$. But he gives no Demonstration of this; and it appears, by the *Construction of his Tables*, that the Correction of Evection, according to Anomaly, and the Correction of Anomaly itself, according to the variable Eccentricity, are, by him, both flung into his Variation-Equation, to raise it to that Quantity.

This Method of his is well adapted for the Ease of Computation, but not to be admitted as Theory against the Principles of Newton, founded on the truest Geometry; which limits the Quantity, as above, to $35' 10''$.

6. Variation proportional to the Semi-Angle of Evection. By examining into this Question, and proving, for my own Satisfaction, the Truth of these Computations, I have been led to enquire what certain Proportion, the greatest mean VARIATION, $32' 32''$ (the Earth quiescent) as Sir Isaac has given it, bears to the Semi-Angle of EVECTION-EQUATION, at a Maximum, and have found it to be in the duplicate Ratio of the synodical Revolution of the Moon from the Sun, to the synodical Revolution of the Moon from her Apogee. And

7. Proportional to 45° . I farther found, that the Tangent of the Angle of Variation is to the Tangent of the Angle of 45° , in the Ratio of 69 to 70, and the Sub-duplicate of 10973 to 11073, conjunctly; so likewise the Tangent of the Arg. of EVECTION is to the Tangent of the Angle of 45° , in the former Ratio of 69 to 70, and the simple Ratio of 10973 to 11073, conjunctly. Which Arg. of Evection $\angle 44^\circ 19' 40'' 40'''$, being subducted from the Angle of 45° , leaves the Semi-Angle of Evection-Equation $40' 19'' 20'''$, nearly as before.

8. A Remark. It is well worth observing, that as there is a given Difference, or Ratio, between the Angle of Evection-Equation and Variation, both being at a Maximum, so the Sum of these Equations should be, when taken together, equal to the correct Quantity of Equation of the Earth's Orbit, at a Maximum, $1^\circ 55' 50''$.

For the Evection-Equation, $1^\circ 20' 40'' +$ Variation, $35' 10''$, is equal to $1^\circ 55' 50''$, the above Equation. So that the Force by which the Curvature of the Moon's Orbit is changed, and becomes more or less eccentric than in its mean State, and also by which the Areas described are rendered unequal to one another, generates an Equation of the Moon's Longitude proportionable to that generated, in the Sun's Longitude, by the Eccentricity of the Earth's Orbit.

OUR AUTHOR has here given the mean Quantity of the Variation only, neglecting, as he says, the Difference which may arise from the Curvature of the Orbit, or the stronger Action of the Sun upon the Moon, when horned and new, than when full and gibbous. In other Situations of the Earth, that is, when the Earth is near the Aphelion or Perihelion, the GREATEST VARIATION is compounded of the duplicate Proportion of the Time of the synodical Revolution, directly, and the triplicate Proportion of the Distance of the Earth from the Sun inversely: and therefore, according to these Proportions in the Aphelion, the GREATEST VARIATION is $33' 14''$: in the Perihelion, $37' 11''$.

9. Variation greatest at the mean Distance of Moon from the Earth. In these Computations, our Author has neglected to take into Consideration the Change which the variable Eccentricity of the Moon's Orbit must make in the VARIATION-EQUATION: leaving it for future Astronomers to determine the Quantity thereof from Observation: For, by the Moon's being more or less removed from the Earth, in her Orients, by the variable Eccentricity of her Orbit, the Variation-Equation will become more or less; the Ratio between the centripetal and the disturbing Force being less or greater in the one Case than the other. OBSERVATION shews that this Consideration is not wholly to be neglected. For, Tables formed from these Principles are found to err most where this Correction of the Variation, according to the Moon's Anomaly, is greatest; that is, when the Moon is in Orients, and near one of her Apfides.

10. Variation changeable at other Distances. It were much to be wished, that Sir Isaac had not contented himself to pass this by, as a Remark, for others to improve on; but had deduced the Quantity directly from the Theory of Gravity. However, this Equation of the Moon's mean Variation, according as she is at a greater or less Distance from the Earth, may be obtained by the following Rule.

The centripetal Force in the Quadratures, is to the accelerating Force in the Syzygies, as the Sum of the greatest and least centripetal Forces, to 3 Times their Difference.

Again, the accelerating Force, when the Moon is at a mean Distance from the Earth, is to the Difference of the greatest and least accelerating Force, when the Moon is in the Apfides of her Orbit, as the Radius is to the Sine of Half the mean elliptic Equation at a Maximum, or to the mean Eccentricity (Radius being 1) and the Subduplicate of the periodical to the synodical Revolution of the Moon from the Sun conjunctly.

Therefore, on the Principles of Machin's Equant, as the Sum of the greatest and least centripetal Forces, is to the Difference of the accelerating Forces at each Apfis, so is the Radius to the Sine of this other greatest Equation = $1' 39''$, viz. As 357,50 : 171 :: Radius : S. $1' 39''$.

COMMENT on the MOON'S THEORY.

VI. Of the MOON'S SECOND EQUATION of VARIATION.

BUT there is still a farther Change in the lunar Orbit, arising from the stronger Action of the Sun upon the Moon, when new and falcated, than when full and gibbous, from whence arises another Equation, which Sir Isaac calls the second Variation.

The centripetal Force in Opposition being to the same in Conjunction as 180 to 178 nearly, the Moon recedes somewhat farther from the Earth, and moves somewhat slower in the Conjunction than in the Opposition. This Force generates a new Equation of mean Motion, which adds to it from Full to the New Moon, and subtracts from New to the Full Moon. And the Sine of this Equation is to Radius as 2 to 357. Mr. Whiston, for explaining this Equation, erroneously distinguishes between an arithmetical Point of true, and geometrical Point of mean, Motion; by which he endeavours to give the Laws of this Equation; but as this Distinction of his two Points is without sensible Difference, his Reasoning is inconclusive. Sir Isaac Newton gives the Maximum of this Equation at 2', as it is retained in the Tables. But by the foregoing Analogy it comes out no more than 1' 56".

But then this second Equation of Variation has been observed, Sir Isaac says, to be liable to Change.

For, when the Sun is in Conjunction with the Apogee of the Moon, it is, according to our Author, about 1' greater, and as much less, when in Opposition; on Account of the variable Eccentricity of the lunar Orbit: The solar Force being greater in one Situation and less in the other.

Dr. Gregory, who first published this famous Theory to the World, gives us a very perplexed and confused Account of this Equation; nor had Sir Isaac Newton himself Observations sufficient to speak of this Change with Certainty. In his First Edition of the Principia, he says, that the changeable Quantity is to be determined by the Situations of the Apogees of the Sun and Moon; and is greatest when they are in Conjunction, and least when in Opposition: It is least also when in Quadrature with each other, and greatest also when the Moon's Apogee is in Conjunction with the Sun; but whether less in Opposition to the Sun, Dr. Gregory, Leadbetter, and others, tell us, that our Author could not determine, for Want of Observations. But this supposed Change, in this Equation, is not from the mutual Situation of the Apogees, or of the Sun, in Regard to the higher or lower Apogee of the Moon's Orbit; but of the Moon in respect to her Distance from the Earth; and is, in Truth, not an Increase or Decrease of this second Variation, but of the first only; and from the Principles deduced, in the foregoing Paragraph, to which the Reader is referred back.

VII. Of the MOTION of the MOON'S APOGEE, or APSIDES.

FROM the 7th and 8th Corollary of the 66th Proposition of the 1st Book. Our Author derives the Motion of the Moon's Apogee. The Inequalities of which Motion are, as he observes, very great, and generate the principal Equation of the Apogee.

From what follows, it will appear evident, that the Line of the Apogees must have, on the whole, a progressive Motion: but very irregular. For, when the Moon's Apogee is in Conjunction and Opposition with the Sun, the Apogee goes forward at the greatest Rate; but, when in Quadrature with the Sun, it even goes backward. For, in the Quadrature, the Force by which the Body P, (see Fig. 1.) is urged to the Body T, the Force MT, then vanishing, is compounded of the Force ML, and the centripetal Force with which the Body T attracts the Body P. The first Force, ML, is increased, nearly, as PT. The other decreases in a duplicate Proportion of that Distance. Therefore the Sum of these two Forces decrease in a less Proportion than the duplicate. Whence the Line of the Apogee, (by the 1. Coroll. of the 45th Prop. of the 1st Book) will recede. In the Conjunction and Opposition, the centripetal Force is as the Difference of the Force KL, and PT; which, because KL is nearly increased as the Distance PT, decreases in more than a duplicate Ratio of that Distance; and therefore causes (Cor. 1. Prop. 45, B. I.) the Apogees to go forward. But, since the Force KL, in the Syzigies, is nearly twice the Force LM, in the Quadratures, the Excess will be on the Side of the Force KL; and, by Consequence, the Line of the Apogees will, on the whole, be carried forward.

Sir Isaac Newton does not determine the Quantity of this progressive Motion of the Moon's Apogee, from the Theory of Gravity, the Solution he gives of the PROBLEM, "TO find the MOTION of the APSIDES (see Prop. 45. Probl. 31. B. 1.) in ORBITS nearly CIRCULAR," accounts, when applied to the Moon's Orbit, but for Half the Quantity of the progressive Motion of the Apogee in one Lunation, viz. 1° 31' 28", nor has Mr. Machin succeeded better. He tried what it would do, considering the Cause of this Motion as arising from the Variation of the Moon's Gravity to the Earth, as it revolves in the elliptic Epicycle, aforementioned, for ascertaining its true Quantity: the Difference of the Forces being, in that Case, double the former; but it gave not the Quantity of Motion with that Exactness as was requisite: Nor was there any Method, he thought, upon commonly received Principles, sufficient for this Purpose.

Nevertheless, this Problem has lately been very satisfactorily solved by the Rev. Mr. Patrick Murdoch, (see Phil. Transactions, V. 47, P. 62) by which Solution he determines the Quantity in a synodical Month to be 3° 17' 28", exceeding the Motion by the Tables but 4". And he observes, that the Reason, that other Authors have failed in giving the true Quantity of Motion of the Apogee, from Sir Isaac's Principles, is, that they all have considered the Earth as quiescent; although the Case, in Nature, and to which they mean to apply Sir Isaac's Rule, is

1. Change of the 2d Variation.

2. Controversies about the 2d Variation.

1. Irregular Motion of the Apogees explained.

2. Problem solved and Quantity of the Motion of the Apogees determined by the Rev. Mr. Murdoch.

COMMENT on the MOON'S THEORY.

different. For, as this learned Writer observes; "The Earth and Moon revolve round their common Center of Gravity, their Distance from which being inversely, as their Masses, and the Forces by which each is attracted by the other, as also the Forces of the Sun to disturb their Motion being in the same Ratio; It follows, that the Earth in its Motion round the common Center of Gravity, will suffer these Disturbances every Way similar to those of the Moon: and the whole Motion of the Apis, resulting from the two disturbing Forces, will be, nearly, double of what either of them could produce separately round a fixed Center."—So far this Author. And whoever considers the Solution of this difficult Problem, will not only be satisfied of this Gentleman's great Abilities, but join in his Conclusion from it: That in this, as in the other Phenomena of the celestial Motions, the Principles and Rules which Sir Isaac Newton lays down are fully confirmed and verified.

3. Quantity of D's Apogee, in a syderal Year, according to Mr. Machin's Rules.

4. According to Sir Isaac's Theory.

Mr. Machin, however, has deduced the Motion of the Apogee from other Principles, which he has not explained, but contented himself with giving Rules founded thereon, for obtaining the mean Motion of the Apogee, its Equation and the Change of Eccentricity: which are these that follow.

"That the mean Motion of the Moon, from its Apogee, is to its mean Motion, in the subduplicate Proportion of 8235 to 8376, or as 1877 to 1893. According to which Proportion the Motion of the Apogee, in a syderal Year, ought to be $40^{\circ} 40' 40'' \frac{1}{2}$. In Sir Isaac's Theory the Numbers stand at $40^{\circ} 40' 43''$. According to Tycho Brahe $40^{\circ} 40' 47''$.

VIII. To determine the CHANGE of ECCENTRICITY of the MOON'S ORBIT.

1. Eccentricity determined.

THE least is to the mean Eccentricity, in the duplicate Ratio of the periodical Time of the Sun's Revolution, to the mean synodical Time of its Revolution from the Moon's Apogee: which Ratio is nearly as 107 to 136.

2. Greatest 2d Equation of the Moon's Apogee.

LASTLY, for the SECOND EQUATION of the APOGEE — "The Sine of the greatest Equation is to the Radius, as the Difference of the greatest and least Eccentricities are to their Sum; that is, as 29 to 136. This determines the Maximum of this Equation at $12^{\circ} 18' 43'' 16'''$. Sir Isaac has determined it from Observation to be $12^{\circ} 18'$. And the Rule I have already given, for determining the Semi Equation of Erection, assigns it the same Quantity.

3. Mr. Machin's Rule for determining the Eccentricities of the Moon's Orbit, and correcting the lunar Theory.

Mr. Machin also makes Mention of another Rule, derived from a different Method, which presupposes the Knowledge of the mean Eccentricity, and which, he says, would not only determine the variable Eccentricity with greater Exactness; but would also correct the modern Theories of the Moon in a Particular, wherein their greatest Error, he says, seems to consist: that is, in placing the Earth in the lower Focus of the Ellipse, which the Moon describes in her Revolution round the Earth; whereas, it ought, HE SAYS, "to be in a certain Point more near the Perigeon." What these Principles were from, whence he derived this Correction, or how he founded this new Hypothesis, it is not possible to collect from this Sketch, or Abstract of a larger

Work intended; He having given no Data for our Guide, in his short Treatise on the Moon's Gravity.

IX. CHANGE of the INCLINATION of the MOON'S ORBIT; With the RETROGRADATION of her NODES, explained, and determined in a circular and elliptic ORBIT.

BUT, it is not only the Motion of the Moon, as to Longitude, which is affected by these disturbing Forces: The Inclination of her Orbit likewise suffers Change by them, and her Nodes, (or the Points, where the Orbit of the Moon intersects the Plane of the Earth's Orbit) are carried with a retrograde Motion, through every Point of the Ecliptic. The Causes of these Changes are explained in the 10th and 11th Corollaries of the 66th Prop. B. 1. Principia. The Summary of which Explanation is as follows.

1. Change of the Orbit's Inclination described.

LET the disturbing Forces be represented, as in Fig. I. by the Lines TM, ML. It is evident, that this last Force, acting in the Plane of the Moon's Orbit, never can disturb the Motion as to Latitude; and that the former Force, TM, when the Nodes are in Syzygies, acting also in the same Plane of the Orbit, cannot, at that Time, affect these Motions. But, when the Nodes are in Quadrature, it must disturb them very much. For attracting the Moon perpetually, out of her Orbit, it diminishes the Inclination of the Plane thereof in the Passage of the Moon from the Quadratures to the Syzygies; and again increases the same in her passing from the Syzygies to the Quadratures. Hence, the Inclination is least of all, when the Moon is in Syzygies, and returns to the first Quantity, nearly, when the Moon arrives at the next Node.

The Moon being thus, perpetually, drawn from the Plane of her Orbit, it is manifest as she continually recedes from that Plane, in her Passage from C, that she will intersect the Plane of the Earth's Orbit, not in the opposite Node D, but in some Point more near to S, which, from thence, becomes a new Place of the Node, in Antecedentia to the former Place. The Nodes therefore, when near the Quadratures, recede perpetually; but, in the Syzygies, where no disturbing Forces act in the same Plane, are quiescent; and in intermediate Places, recede with a quicker or more slow Motion, as they are nearer to the one Point than the other.

COMMENT on the MOON'S THEORY.

Our Author has computed the Quantities of these Irregularities, which take no less than FIVE (See from Prop. 36 PROPOSITIONS in the III. BOOK. It would be difficult to render such an *abstruse Subject* clearer to 35 inclusive.) than in the Words of the Author, or to explain the Grounds of his Computations so, as to give more Satisfaction. Let it suffice to trace out the Steps he proceeds in. And first, He begins with the LEAST DIFFICULT PROBLEM, To find the HORARY MOTION of the NODE in a CIRCULAR ORBIT?

2. Quantity of the horary Motion of the Moon's Node, in a circular Orbit determined.

3. The Quantity determined in an elliptic Orbit.

5. *Annual Motion of the Nodes determined.*

Fig. III.

1. Method of determining the 2d Equation of the Node.

2. The greatest semi-annual Equation of the Moon's Node determined.

COMMENT on the MOON'S THEORY.

XI. The QUANTITY of INCLINATION of the MOON'S ORBIT, with the ECLIPTIC, deduced from Her HORARY VARIATION.

1. The Horary Variation.

Our Author further computes, from the same Principles, the Horary Variation of the Orbit's Inclination to the Ecliptic, and gives a Method of computing the Quantity of Inclination at any given Time.

The Mean Variation of the Inclination of the Orbit is given at $16' 23'' \frac{1}{2}$; but this is abstracted from the Consideration of the Moon's Situation in her Orbit. For, if the Nodes are in Quadrature, the Inclination is more by $2' 43''$, the Sum of the Horary Increments, than when in Syzygie; and the whole Mean Variation $16' 23''$, diminished by half that Quantity, = $1' 21'' 30'''$, becomes $15' 1'' 30'''$, when the Moon is in Quadrature: And being increased, by the

2. The Greatest Variation of the Inclination of the Moon's Orbit, determined.

same becomes $17' 44'' 30'''$, when the Moon is in Syzygies. If therefore the Moon be in Syzygies, the WHOLE VARIATION, in the Passage of the Node from the Quadratures, to the Syzygies, will be $17' 44'' 30'''$, or rejecting thirds, $17' 45''$. And therefore, if the Inclination, when the Nodes are in Syzygies, be $5^\circ 17' 20''$, it will be $4^\circ 59' 35''$ when they are in Quadrature, and the Moon in Syzygies: Observation confirming the Accuracy of all these Computations.

3. Mr. Machin's Determination of the same Quantities nearly.

Mr. Machin derives the same Quantities, nearly, by a different Method, as Sir Isaac gives; but as the Proportions, for forming his Elliptic Equant, are the same with those Sir Isaac has given, it is unnecessary to say more of it here. Our Author thought it worthy of a Place in the Scholium to these Propositions, though the Method of computing from this Elliptic Equant was, at the same time, found out by Dr. Pemberton, who made use of it for the same Purpose: the Investigating the Motion of the Nodes. I shall refer the Reader for further Instruction, as to this Method, to Machin's Theory of Gravity, which is founded on the known Properties of the Figure he applies to that Purpose. That, where equal Areas are described in equal Times, about the Center of the Elliptic or any other Equant, the Rays, from the Center, are always reciprocally in the Subduplicate Ratio of the Velocity about the Center.

4. The Investigation of the Moon's Nodes according to Machin.

THE Foregoing are the PRINCIPLES, from which our AUTHOR has given the LUNAR THEORY, a Priori; shewing how the IRREGULARITIES, in the Motion of the Moon, may be derived from PHYSICAL CAUSES. What is farther wanted to compleat it, and reduce those IRREGULARITIES to Computation, is supplied in the SCHOLIUM to the FOREGOING PROBLEMS.

XII. Of the SOLAR FORCES, acting on the MOON, in her ORBIT.

1. The Sun produces the Inequalities of the Moon's Motions by Turns.

WE have been sufficiently instructed, already, that the Irregularities, in the Moon's Motion, proceed from the disturbing Forces of the Sun, which increase and diminish, by Turns, the Areas which she describes; and which, were that disturbing Force taken away, would, like the Areas, described by the primary Planets, round the Sun, be equal or proportional to the Time of Description.

2. The Force of the Sun on the Moon's Orbit according to the different Distances.

It is also evident, that the Action of the Sun upon the Moon's Orbit, must be increased or diminished, according to the less or greater Distance of the Sun from the Earth. And, as the Ablatitious Forces are double those of the Addititious, nearly, that the Orbit of the Moon must be more, in general, dilated, and its Periodical Revolution round the Earth more slowly performed, than if these Forces had not disturbed her Motion at all. These disturbing Forces increase and decrease in the Triplicate Ratio of the Sun's apparent Diameter. For they are, as the Square of the Distance of the Sun, inversely, and the Distance of the Moon from the Earth, directly. In this Proportion, therefore, is the Orbit of the Moon dilated in the Perihelion, and contracted in the Aphelion. The Orbit of the Moon being thus by Turns, dilated and contracted, she takes a longer Time to perform her Revolution in the dilated than in the contracted Orbit. The Quantity of this Irregularity, and of the Equation thence generated, is determined by Sir Isaac, from Corol. 6. Prop. 66. B. I. Wherein he demonstrates that the periodical Time of the Moon, will be increased and diminished, in a Ratio compounded of the sesquialterate Ratio of the increasing and decreasing Radii, and of the subduplicate of that Ratio in which the Earth's central Force is diminished, or increased, by the Increase or Decrease of the Sun's Action upon the Earth. From which

3. Proportional Change of the Solar Forces disturbing the Moon's Motion.

4. Change of the Lunar Orbit explain'd.

XIII. The Greatest Annual EQUATION of the MOON,

1. Greatest Annual Equation.

According to Sir Isaac Newton is $11' 49''$: The Eccentricity of the Orbis Magnus being to the Radius, as 1687 to 100000. But Observation shews this to be somewhat too large; so that if this Equation were farther diminished in the subduplicate Ratio also of the synodical to the periodical Times, its Maximum would be (to the above given Eccentricity 1687) = $11' 29''$ with which Observation more nearly agrees. If we assume a less Eccentricity for the Earth's Orbit, viz. 1685, or as much as shall correspond to $1^\circ 55' 50''$, for the Equation of the Sun's Center, the annual Equation of the Moon will be $11' 25''$ only; as by our Tables; somewhat greater than given by Mayer in the Gottingen Tables.

COMMENT on the MOON'S THEORY.

Mr. Machin, substituting an Equation which shall be to the Sun's Equation, as the mean Solar Force, to the Force of the Moon's Gravity, or as 47 to 8400; and increasing that Equation, in the Ratio of the Sun's Period, to the mean Synodical Period of the Moon from the Sun, or of 99 to 8, computes this Equation at $12' 5''$. Horrox gives its Maximum $11' 30''$.

But, the Solar Force, on the Moon's Orbit, which acts in a Triplicate Ratio of its Distance from the Earth, inversely, not only affects the mean Motion of the Moon, but generates Equations both in the Mean Motion of the Apogee, and Nodes: They being accelerated as the Mean Motion of the Moon is retarded, and retarded when that is accelerated. For, though the Equation of the Nodes is to be subtracted, when that of the Moon's mean Place is likewise subtracted, yet it is, in Fact, an Acceleration of the Motion of the Nodes, which is Retrograde in Itself. For, determining the Quantity of these Equations. The MOTION of the SUN is, in the reciprocal duplicate Ratio of its Distance from the Earth: This Motion generates, (the Eccentricity of the Orbit, being $16\frac{1}{2}$ to 2000 its M. Diam.) an Equation of the Center, which, at a Maximum is $= 1^{\circ} 56' 20''$. If the Motion of the Sun had been in a Triplicate Ratio, the Equation thence generated would have been $2^{\circ} 54' 30''$. Therefore, the greatest Equation of the Apogee and Node is to $2^{\circ} 54' 30''$, as the mean diurnal Motion of the Apogee, and of the Node, respectively, is to the mean diurnal Motion of the Sun, Hence

2. Mr. Machin's greatest Quantity of the Annual Equation of the Moon.

XIV. The GREATEST EQUATION of the MOON'S APOGEE,

Is $19' 43''$, and the GREATEST EQUATION OF THE NODE $9' 24''$. But the greatest EQUATION OF THE SUN being but $1^{\circ} 55' 50''$, the Annual Equations of the Moon's Apogee and Nodes will be reduced to $19' 38''$ and $9' 21''$, respectively, as in the Tables.

Mr. Machin's RULE, derived from his Elliptic Equant, is, that if the Sun's Force is in a Triplicate Ratio of the Sun's Distance, the Rays of the Equant shall be in the Sesquiplicate Ratio of the Distance. That, if the Revolution of the Motion, to be equated, was performed in the Time of the Sun's Revolution, this Equation would be to the Equation of the Sun's Center, nearly as 3 to 2. Or if the Force decreased as m , any other Power of the Sun's Distance, the Equation would be to that of the Sun's Center as m to 2. But, if the Motion be performed in any other Period, any small annual lunar Equation, arising from the different Distance of the Sun, at different Times of the Year, would then be more or less, in the Ratio of the synodical Revolution of the Sun from the Motion to be equated, to the periodical Revolution of that Motion to be equated. THIS RULE makes the greatest Equation of the Node $8' 56''$; of the Apogee, $21' 57''$.

1. Mr. Machin's Rule for determining the greatest Equation of the Apogee and Nodes.

2. Mr. Machin's greatest Equations thereof.

These three Annual Equations, according to Newton, bear such Proportion to the Central Equation of the Earth, that when that is greatest, the Rest are so; and when that is diminished, the Rest will be diminished proportionably. For the Sun's greatest Equation is to the greatest Equation of the Moon, Moon's Apogee, and Node, respectively, as the present Equation of the Sun to the present Equation of the Moon, Apogee, and Node, respectively.

XV. A SECOND EQUATION of the MOON'S ORBIT.

By the Theory of Gravity, Sir Isaac Newton found, that the Moon's Orbit was also somewhat more dilated when the Sun and Moon's Apfides were in Conjunction, than when at Right Angles with each other; which Difference arises from the Orbit being the most Eccentric and Elliptical in the Former Situation, and least of all in the Latter. From hence the Mean Motion requires a farther Correction, the Moon moving slower in a dilated Orbit, and swifter in the more contracted one. This Equation our Author determines, from Observation, to be $3' 45''$, at a mean, when the Sun is in Orients with the Moon's Apfides: at other Times, it is to that Quantity, as the Sine of the double Distance of the Apfides, from the nearest Syzygy or Quadrature, is to the Radius. Its least Quantity in the Aphelion of the Earth is $3' 34''$; and its greatest, in the Perihelion, $3' 56''$: being in the reciprocal triplicate Ratio of the Sun's Distance from the Earth.

1. The greatest Quantity of this Equation determined.

Though our Author deduces this from Observation, alone, yet the Quantity may be deduced from the following Analogy: That is, the Sine of this Equation is to the Sine of the greatest Second Equation of the Moon's Apogee, in the duplicate Ratio of the synodical Time of the Moon from the Sun, to the synodical Time of the Sun from the Moon's Apogee. By this RULE the Equation comes out $3' 46''$.

2. Whence this Equation derived.

The synodic Time of the Sun from the Moon's Apogee $= 412^d.786$; Moon from Sun $29^d.53$; Greatest second Equation of the Moon's Apogee $= 12^{\circ} 18'$.

XVI. A THIRD ANNUAL EQUATION of the MOON.

From the Laws of Gravity, our Author has likewise deduced a Third Annual Equation of the Moon's Motion from the Situation of the Sun with the Nodes. For the Solar Force being somewhat Greater on the Lunar Orbit, when they

COMMENT on the MOON'S THEORY.

1. *Its greatest Quantity determined.* are in Conjunction with the Sun, than when in Quadrature, there arises a *small Equation* from this different *Dilatation* of the Orbit, as the Sun and Nodes are situated in respect of each other. Which Equation he has computed, (for it is too small to be determined by Observation) to be at a Maximum, in the Orients, $47''$. At other Times, it is to this Quantity, as the Sine of the double Distance of the Sun from the Node, is to the Radius. At other Distances, in Orients, as Cube of Sun's Distance from the Earth; and therefore in Perigee $49''$, in Apogee $45''$.

2. *Otherwise determined.* The mean Inclination of the Moon's Orbit being $5^{\circ} 8' 28''$, and the half Difference of the greatest and least Inclination $8' 52''$, this Equation will be to the Sine of the half Difference as the Sine of the mean Inclination to the Radius. For the Force by which the Moon is drawn from the Earth being lessened in the Ratio of the Cosine of the Inclination of the Moon's Orbit to Radius, this Equation of mean Motion, generated by the Increase or Decrease of that Force, as the Sun is in Conjunction or Quadrature with the Moon's Nodes, will be as the Sines of that Difference to the Sines of the mean Inclination, making the Radius of the Circle which the Moon describes, according to her mean Motion, or while the Sun is in Orients with the Nodes, equal to 1. By which Rule that Equation comes out $48''$, nearly.

Both these Equations add to the mean Motion, whilst the Sun passes from the Quadrature, with the Apfides and Nodes, to their Conjunctions, and subtract from it, in its Passage from the Conjunctions to the Quadratures. The GIVEN QUANTITY of this last, like the former Equation, is to be increased or diminished in the Ratio of the Cubes of the Sun's Distance from the Earth inversely.

XVII. THE SECOND and GREATEST EQUATION of the MOON'S APOGEE.

By the same Theory of Gravity, our Author deduces the second and greatest Equation of the Apogee. We have already spoken of it in explaining the Motion of the Apfides from 7, 8, and 9 Corollaries of Prop. 66. B. 1. Its greatest Quantity, from the Theory of Gravity, as well as from Observation, is nearly $12^{\circ} 18'$, and at all Times it is to that Quantity, as the double Distance of the Sun from the Apogee is to the Radius.

XVIII. HORROX'S IMPROVEMENT of the LUNAR THEORY.

Horrox, as Sir Isaac observes, was the first Astronomer, who advanced the Theory of the Moon moving in an Ellipse, round the Earth placed in its lower Focus. Dr. Halley improved this Notion, by placing the Center of the Ellipse in an Epicycle, whose Center uniformly revolved round the Earth. The Motion in this Epicycle, expresses the fore-mentioned Inequalities, in the Progress and Regress of the Apfides, and the Change in the Quantity of the Eccentricity.

Let the mean Distance of the Moon from the Earth, says our Author, be divided into 100000 Parts; and let T (Fig. 3) represent the Earth; TC, the Moon's mean Eccentricity = 5505 such Parts. Produce TC to B, so as CB may be the Sine of the greatest semi-annual Equation of the Apogee, $12^{\circ} 18'$ to the Radius TC. Then, the Circle described upon the Center C, with the Radius CB, will be the EPICYCLE mentioned, in which, the Center of the Moon's Orbit is placed, and revolved to the Left, according to the Order of the Letters, B, D, A. Set off the Angle BCD, equal to twice the annual Argument, or the double Distance of the Sun's true Place from the Moon's Apogee, once equated. Then, CTD will be the second Equation of the Moon's Apogee for the Time given; TD the Eccentricity of the Orbit, tending to the Place of the Apogee, now twice equated; and CD, or CB, the Difference between the mean and the greatest Eccentricity = 1173, of the above Parts. N. B. The Radius of the Epicycle CD, or CB, as seen from the Moon, at her mean Distance from the Earth, subtends an Angle of $40' 20''$, = the Semi-Equation of Evection. How this Angle may be obtained, at once, without computing the Angle CTD, or the Eccentricity




Fig. III.

2. *A Reference.*

TD, is fully explained in the Letter to Lord MACCLESFIELD, on the Parallaetic Angle, to which the Reader is referred. Or you may see it performed farther on, P. 371.

But, the Moon's mean Place, the Place of its Apogee, its Eccentricity, the longer Axis of its Orbit, 200000, being given, the Moon's true Place in her Orbit, with her Distance from the Earth, are thence determined, by the known Method of solving the Equation of her Center.

3. *Changes of the Moon's Orbit according to the different Distances of the Sun from the Earth.*

In the Computations of the Equations of the Apogee, and Change of Eccentricity of the Orbit, which takes Place from the different Action of the Sun upon it, according to its different Aspects, in respect of the lunar Apfides, no Notice has hitherto been taken of the Diminution or Excess of solar Force, which affects these Changes in the Orbit, according as the Sun is farther from, or nearer to, the Earth. But, as the Sun acts with greater or less Power on the lunar Orbit, according as the Earth is nearest to its Perihelion, or Aphelion, and that, in the triplicate Ratio of its Distance, inversely. It is necessary, that the Excess, or Diminution, of the Sun's Action, upon the Orbit, should take Place in these Computations. — In Order to determine this Equation, Sir Isaac thus proceeds.

XIX. THE SECOND EQUATION of the MOON'S CENTER.

In the Perihelion of the $\odot$, when the solar Force is greatest, and where there is no solar Equat. the Center of D's Orbit moves faster about the Center C, (see Fig. 3) than in the Aphel. and that in the reciprocal tripl. Ratio of $\odot$'s Dist. fr. the

COMMENT on the MOON'S THEORY.

Earth. But, because, in Fig. III. the Equation of the Sun's Center, at other Distances, is included in the annual Argument, $\frac{1}{2}$ BCD; the Center of the Orbit thence moves faster, in its Epicycle BDA, in a reciprocal duplicate Ratio of those Distances from the Earth. But that it might move yet faster, round the Center C, in the reciprocal simple Ratio of those Distances; from the Point D, the first Center of the Moon's Orbit, set off the Point F, the second Center of the Moon's Orbit, making DF, to DC, as twice the Eccentricity of the Earth's Orbit to the Sun's mean Distance, and the Sun's mean diurnal Motion from the Moon's Apogee, to his diurnal mean Motion from his own Apogee, conjunctly: Or, as $33\frac{1}{2}$ to 1000, and $52' 27'' 16'''$ to $59' 8'' 10'''$, conjunctly, or 3 to 100.

And let the Angle CDF be equal to the Complement of the Sun's true Anomaly to 360° , for the Time given. The Line DF will be then 35,2 such Parts as TC contains 5505, or mean Distance of the Moon from the Earth 100000; and will subtend an Angle, seen, at a mean Distance, from the Moon, at Right Angles with that mean Distance, = $1' 12'' 30'''$, half the greatest Equation; the Double of which, $2' 25''$, according to our Author, is the Greatest of this Equation, which he calls the second Equation of the Moon's Center. Which Equation may be had, at all other Times, either by subtracting twice the annual Argument from the Sum of the Moon's equated and Sun's mean Anomaly, or else by adding the Distance of the Moon equated from the Sun, to the Distance of the Apogee of the Moon equated from the Apogee of the Sun, for the Argument; and saying, as the Radius is to the Sine of the Argument thus found, so is $2' 25''$ to the second Equation of the Moon's Center. To be added if the Argument is less than a Semi-Circle, but to be subtracted if greater. So shall you have (concludes our Author) the Moon's true Orbit Place in the Syzigies: in which Computation the Variation-Equations are excluded.

But this wants some farther Explanation.

In the preceding Fig. III. D representing the first Center of the Moon's Orbit, in the Epicycle, according to the double annual Argument BCD; F a new Center, translated out of that Epicycle, by the Excess of the Sun's Action upon it; it is evident, that the correct Equation of the Apogee is no longer the Angle CTD, but the Angle CTF; and the Eccentricity of the Orbit, to the Time given, will be represented by the Line TF, which, in this Situation, is greater than the former Eccentricity, TD. It appears also, that, the Angle FDC being the Complement of the Sun's Anomaly to 360° , the Point F, in the translated System, must, in the last Semi-circle of the Sun's Anomaly, when the Sun's Equation adds to the mean Motion, be in Antecedentia to the Point D, as in the Figure before us; where the Angle of the Sun's double Distance from the Moon's Apogee is no longer represented by DCB, but by FCB, less than the former by the Angle FCD. So the first Center of the Moon's Orbit, D, in the untranslated Epicycle, will, by doubling the annual Argument, (in which the Sun's Equation is twice included) for $\angle$ BCD, be carried about, out of the Apfides, in the inverse duplicate Ratio of the Sun's Distance; and will therefore require a Translation to F, to move round C in the simple Ratio of that Distance, inversely, corresponding with our Author's Theory, in this Respect: which should be the Acceleration of the second Center F, of the Orbit, in the reciprocal triplicate Ratio, only, (exclusive of the increased Acceleration by the redoubled Sun's Equation) every where, as in the Apfides; carried in a certain Curve, about the Center C, in the translated System, by the Means aforesaid.

BUT our Author, in this Computation, has given only the Equation which arises from the disturbing Force, taken in the simple Ratio of the Sun's Distance, inversely, the Sun's Equation, he says, involved in the Argumentum annum, supply the rest. But, this can only be understood where that Equation takes Place. For, in the Perihelion, and Aphelion, of the Earth, or of $\odot$, and 6 Signs, of the Earth's Anomaly, where the Excess, or Diminution, of the solar Force, is at its greatest Quantity, there the Line DF should be according to the triplicate Ratio of the disturbing Force. If we suppose the Center of the Ellipse to be carried about in the reciprocal duplicate Ratio of the Sun's Distance, the Equation, thence resulting, would be $2' 44''$; or, as Sir Isaac has determined it, DF would be to DC, as $33\frac{1}{2}$ to 1000, and $DF = 39,7$. But, to obtain the Proportion of each in the simple Ratio of the Sun's Distance, he lessens this Proportion, in the Ratio of the Sun's diurnal Motion from his own Apogee to his diurnal Motion from the Apogee of the Moon, or in the Ratio of $59' 8''$ to $52' 27''$, which gives $DF = 35,2$ and an Equation of $2' 25''$, as before. NOW, in order to obtain this Equation, in the triplicate Ratio of the Sun's Distance, in the Apfides; DF, must be taken to DC, as 50,85 to 1000, which Proportion will give DF 59,63, and the Equation of the Motion, in the translated Epicycle, according to the inverse triplicate Ratio, = $4' 6''$, which is the true Quantity in the Perihelion or Aphelion of the Earth.

In order to reduce THIS EQUATION, at all Times, to its true Quantity, we must consider it as consisting of two Parts. LET us call these Parts P, and Q: The first of these being the Quantity $2' 25''$, the Second of $1' 41''$. The Argument for the Equation P, will be the double annual Argument subtracted from the Sum of the Moon's equated and Sun's mean Anomaly. The Argument for the Equation Q, will be the Sun's Anomaly added to the double annual Argument, and subtracted from the Moon's Anomaly also. Both diminish the Place of the Moon, if their respective Arguments are more than six Signs, and increase its Longitude, if they are less. The Sum, or Difference, of these two Quantities, will give the Equation assigned, with the proper Sign connected; and the Center of P's Orbit will then be, every where, carried in a Curve about the Center C, nearly, as the Cube of the Sun's Distance from the Earth, inversely

1. Greatest 2d Equation of the Moon's Center determined.

2. Explanation of the Cause of the 2d Equation of the Moon's Center.

3. A Correction of the common 2d Equation of the Moon's Center.

4. The true Relation of this Equation at all Times.

COMMENT on the MOON'S THEORY.

XX. EVECTION-EQUATION of the MOON.

1. Another Method of computing the Quantities of the same Equation.

2. Correction of the Evection-Equation.

3. A further Correction of the Evection-Equation.

OR if, according to the shorter Method, laid down in the Letter to Lord MACCLESFIELD, referred to before, you compute, at once, the EVECTION-EQUATION, the respective Arguments, for both Parts of this Equation, will be for P, the Angle of Evection + Sun's Anomaly; and for Q, the Angle of Evection — Sun's Anomaly.

But, the EVECTION-EQUATION still wants a farther Correction, not considered by this great Author; which arises from the Obliquity of the Moon's Orbit with the Earth's Orbit. From this Position of the Plane of the lunar Orbit, it appears, that the Quantity of the solar Force, by which the Motion of the Moon is disturbed, as to Longitude, must be diminished: That Force requiring to be resolved into two Forces; one parallel, the other perpendicular, to the Direction thereof. It is plain, that the latter Force cannot affect the Motion as to Longitude; but only the former. So that, that Force shall be to the whole Force, in the triplicate Ratio of the Cosines of the Angles of Inclination. THE EVECTION-EQUATION diminished in that Ratio, will be $1^{\circ} 19' 41''$, which taken from $1^{\circ} 20' 41''$ Evection-Equation, the Equation thence resulting will be $1'$; according to Tables.

XXI. PRINCIPLES of the THEORY agree with OBSERVATION.

1. The Correctness of our Tables.

2. The Design of our Tables.

THUS, from our great Geometrician's Principles of the lunar Theory, the lunar Equations may be deduced, *a priori*, with a Degree of Exactness almost equal to Observation. By constructing Tables from the foregoing Data, the Moon's Longitude may be found, at all Times, within 2 Minutes of the Truth, and seldom with an Error of more than Half a Minute. This Theory, as Sir Isaac gave it the World, errs in no Part of the Orbit above 6 Minutes of a Degree; and the Method of Calculus, to be established upon the foregoing Revival of it, shews how little Alteration is necessary, to the making Tables from thence to serve the important Purpose of Longitude at Sea. That what is here offered, as an Amendment, or Correction of the THEORY, is as much intended to reflect Honour upon this great Man, as to render THIS WORK of general Use to Navigation.

XXII. The SUN and MOON'S RADICAL PLACES.

FOR the Establishment of astronomical Tables, according to this Theory, by the Assistance of Mr. Flamsteed, the then Astronomer Royal, Sir Isaac Newton fixed the following Longitudes of the Sun and Moon, as the correctest at the Royal Observatory, at Greenwich, to the last Day of December at Noon, old Style, 1700, or to the Beginning of 1701: viz. The mean Longitude of the Sun, $15^{\circ} 20' 43' 40''$: Of his Apogee, $7^{\circ} 44' 30''$. The mean Longitude of the Moon, $15^{\circ} 21' 0''$. Of her Apogee $8^{\circ} 20' 0''$; and of her ascending Node $27^{\circ} 24' 20''$: and the Difference of Meridians, between the Royal Observatories of Greenwich, and Paris, is given, from Observation, at $9' 20''$. On which our eminent Author observed, that the mean Motion of the Moon, and its Apogee, were not then discovered with sufficient Accuracy.

XXIII. CORRECTION and IMPROVEMENT of the Lunar NUMBERS and THEORY.

THE Observations of later Astronomers, particularly of Flamsteed's Successors, Dr. Halley, and Dr. Bradley, have corrected these Numbers to the wanted Accuracy. Dr. Bradley, one of the most accurate Observers that any Nation can boast of, has confirmed, by his Observations, an Opinion, started at first by Dr. Halley, and since pursued by Mr. Mayer, of Gottingen, and other foreign Astronomers, that the mean Motion of the Moon is more accelerated in her Orbit now than in former Ages. The Quantity of this Acceleration has been lately fixed at our Royal Observatory, for the Year 1750, at $12''$, for one hundred Years before and after that Era, and that the Increase of Acceleration, from that Time, is as the Square of the Centuries, and centesimal Parts of Centuries, before or after that Era. The mean Longitudes of the Sun, of his Apogee, and the Moon, and of her Apogee, and Nodes, are also made correct by later Observations; so that, now, in comparing very ancient Observations with our Tables, formed by these Corrections, the Errors are not greater, than the Rudeness and Inaccuracy of Observations, in those early Ages; will account for.

NOTE, There is this RULE (P. 299. Scholium to foregoing Propositions, B. III. Newton's Prin. transf. by Motte.) The Motion of the Sun being in the reciprocal duplicate Ratio of his Distance from the Earth, with Eccentricity $16 \frac{1}{2}$ to mean-Dist. 1000, generates an Equation $1^{\circ} 56' 20''$; but moving in the reciprocal triplicate Ratio generates an Equat. as 2 to 3, of the former; viz. $(2 : 3 :: 1^{\circ} 56' 20'' :) 2^{\circ} 54' 30''$ Equat. The Equations generated by the Sun's Mot. being directly as the Indices of the reciprocal generating Powers of his Distances from the Earth: the Revolutions being performed in the same Time; but if in different Time, the Equation before found will be more or less, as Time of 1st to Time of 2d Revolution; according to Machin, P. 38, Laws of Moon's Mot. — But whether this holds true in all Orbits and Indices of reciprocal generating Powers, is to be doubted, or proved; taking the same or like Position, in different Orbits; or taking their greatest Equations, respectively.

COMMENT on the MOON'S THEORY.

XXIV. LUNAR PROPERTIES and CONCLUSIONS.

IN the *Corollaries*, deduced from the 37th *Proportion*, Book III. relative to the MOTION of the TIDES, Sir Isaac Newton gives the following *Computations*.

1. That the Density of the Moon is to the Density of the Earth as 4891 to 4000, or as 11 to 9.
2. That, since the true Diameter of the Moon is to the Diameter of the Earth, as 100 to 365, the Mass of Matter in the Moon will be to the Mass of Matter in the Earth, as 1 to 39,788.
3. The accelerative Gravity, on the Surface of the Moon, will be 3 Times less than the accelerative Gravity on the Surface of the Earth.
4. That the Distance of the Moon's Center, from the Center of the Earth, will be to the Distance of the same from the Center of Gravity, as 40 to 39.
5. The mean Distance of the Moon from the Center of the Earth, is 60 Semidiameters, and one tenth, in the Syzigies; but, in the Quadratures, is 61 Semidiameters subtracting one 30th Part; which is in Proportion of 69 to 70.
6. The horizontal Parallax, in the Moon's Syzigies, Sir Isaac gives at $57' 20''$, under the Equator. For, from the Earth's being an oblate Spheroid, this Quantity diminishes, as that is receded from; and the Parallaxes in the Latitudes of 30° 38° 45° 52° 60° 90° are . . $57' 16''$ $57' 14''$ $57' 12''$ $57' 10''$ $57' 8''$ $57' 4''$.
7. The mean Semidiameter of the Moon, as seen from the Earth, is, at all Times, to the Moon's horizontal Parallax, as $15' 45''$ to $57' 20''$; or as .275 to 1, nearly. And its horary Motion, in Syzigies, Sir Isaac settles at $33' 32'' 32'''$; in the Quadratures at $32' 12''$. The greatest horizontal Parallax, at about $61' 10''$; the least, at $53' 30''$; the true horizontal Parallax being to the Mean, in the simple Ratio of the Moon's Distance from the Earth, inversely; and the true horary Motion of the Moon, to its mean Motion, in the duplicate Ratio of the true and mean horizontal Parallaxes.

S. C.

OF DETERMINING THE EQUATION OF EVECTION, referred to in P. 368, of this WORK, as determined at P. 25 and 26, in a PRINTED LETTER to the EARL of MACCLESFIELD.

THIS EVECTION-EQUATION, is the Difference between the mean and true central Equation of the Moon at all Times. —

THUS, the greatest Equation of the Moon's Center = $7^\circ 39' 20''$ Sun in *Apsides* D.

The least Equation thereof = $4^\circ 57' 58''$ Sun in their Quadrature.

Hence, the greatest *Evection-Equation* is the Difference between the greatest mean, and the greatest and least true central Equation, = $1^\circ 20' 41''$, when the Eccentricity = 55050, and mean Distance of the Moon from the Earth = 1000000.

Difference $2^\circ 41' 22''$
Half $1^\circ 20' 41''$ *Evection-Equation*.
6 18 39 *Mean central Equation*.

At all other Times, (when the Eccentricity is more or less than 55050, of such Parts as the mean Dist. from the Earth = 1000000) the Sine of this *Evection-Equation*, or of the Dif. between the present mean and true central Equation of the Moon, is to the Sine of the greatest *Evection*, $1^\circ 20' 41''$, nearly, as the Sine of the Excess of the Moon's mean Anomaly, equated by $\frac{1}{2}$ mean central Equation, according to its Sign, above twice the Sun from the Moon's Apogee, is to Radius. — Adding this Equation to the Moon's Longitude when the Argument is more than six Signs, and subtracting it therefrom when the Argument is less than six Signs.

Thus, As Radius	Logarithms.
To S. Arg. Evect. $2^\circ 50' = 65^\circ$. .	9.9572757
So S. $1^\circ 20' 41''$ Gr. Evecton . . .	8.3704700
To S. $1^\circ 13' 7''$ Present Evecton, nearly.	8.3277457

Or, As Radius	Logarithms.
To S. Arg. 65°	9.9572757
So S. $1^\circ 20' 26''$ Gr. Evecton . . .	8.3691223
To S. $1^\circ 12' 54''$	8.3263980

Present Evecton nearly, according to our Tables.

OTHERWISE. Like solving the central Equation of the Moon, to a given Eccentricity. Imagine another Ellipsis, whose Semitransverse = the mean Dist. of the Moon from the Earth = 1000000, and each Focus thereof distant from the Center = the Radius of the Epicycle (see P. 368. Fig. III.) = Half of the Dif. of the greatest and least Eccentricity of D's Orbit, = 1173; and the Arg. of Evecton similar to the mean Anomaly of the Moon, revolving about the upper Focus. Then there is given, the Angle of Argument at the upper Focus, whose Supplement is included by two Sides of a Triangle, one the Distance of the Foci, and the other = the Transverse Diameter, both issuing from the upper Focus, whose Sum and Diff. are as the Aphelion to the Perihelion Distance, to find the less Angle opposite, by Case 5 Oblique Plane Triangles. Which Angle being doubled = the *Evection-Equation* sought, very nearly.

Thus, As Aphelion Dist.	Logarithms.	
To Perihelion Dist.	4.9949354	} Const. Log. 9.9898110 To be added to the Tangent of Half the Evecton-Equation.
So Tan. $\frac{1}{2}$ Arg. = Tang. $\frac{1}{2}$ Sum opposite $\angle$ s in the given Δ $32^\circ 30' 0''$	4.9948756	
To Tan of Half their Dif.	9.8041873	
	9.7939983	
Less Angle . . . the Dif.	36 22	
Doubled	$1^\circ 12' 44''$	

$1^\circ 12' 44''$ Evecton-Equation required, nearly, as before.

If greater Accuracy is required, solve the elliptic Equation correctly, by the general Method, P. 96.
D. Cowper directs this Equation to be increased or diminished, in the simple Ratio of the Moon's horizontal Parallax, and then according to the Diff. of the Cubes of the Sun's Distance inversely, for giving the correct central Equation, when this 2d is connected with the first Part.

The MOON'S LONGITUDE computed from the TABLES, to a 100 PLACES, and compared with the respective OBSERVATIONS. C.

Eclipses, and whose Observations	Years of Christ.	Equal Time of Moon's trans- fiting the Meridian.					Annual Argument. s o ' "	D à O. s o ' "	Moon's Longitude observed.				Error of Com- putation. ' "
		M	D	H	M	S			s	o	'	"	
Moon Ecl.	1628	January	10	9	26	31	1 21 26	6 0 0	♌	0	27	16	+ 0 42
Sun . . .	1661	March	19	23	16	11	7 0 21	0 0 33	♍	11	4	9	+ 0 5
Sun . . .	1676	May	31	22	23	30	0 23 3	0 0 17	♎	21	26	17	- 0 4
Moon . .	1681	August	18	15	19	0	8 6 29	6 0 0	♏	6	14	33	+ 0 34
Moon . .	1682	February	11	11	13	19	1 13 36	6 0 0	♐	3	47	33	+ 0 5
Moon . .	1685	November	30	10	30	35	5 25 0	6 0 0	♑	19	39	39	+ 0 43
Flam- stead.	1689	November	16	11	47	20	0 0 1	6 0 45	♒	6	12	43	+ 1 3
	1690	April	18	16	23	50	4 16 4	8 2 0	♓	11	43	52	+ 1 44
	1690	April	20	18	2	32	4 17 50	8 25 5	♈	6	30	55	- 0 28
	1715	April	21	20	2	40	6 21 27	11 29 8	♉	11	20	17	- 0 2
	1722	November	27	1	21	22	3 16 44	11 29 40	♊	15	53	5	+ 1 58
	1725	December	5	9	8	5	11 22 19	4 21 24	♋	15	42	12	+ 1 1
			8	12	19	26	11 25 11	6 4 46	♌	2	58	38	+ 0 29
	1726	January	6	12	8	23	0 21 20	5 28 25	♍	26	7	46	- 0 50
			13	18	0	45	0 27 51	9 1 48	♎	6	54	59	- 0 16
		March	6	12	10	8	2 13 58	5 29 29	♏	26	42	2	- 0 15
			12	16	56	19	2 19 24	8 13 13	♐	16	46	53	- 1 10
		April	5	12	20	22	3 10 9	6 5 30	♑	2	15	10	- 0 4
		May	9	16	4	15	4 9 29	7 25 54	♒	26	11	14	+ 1 26
			13	19	3	0	4 13 0	9 12 17	♓	15	49	35	- 1 25
		June	7	15	33	34	5 4 7	7 18 13	♈	16	18	50	+ 1 35
		July	3	12	43	4	5 26 3	6 7 44	♉	0	0	32	+ 0 28
			10	17	44	7	6 2 10	8 28 56	♊	27	54	1	- 0 19
		August	12	21	20	21	7 0 26	10 19 17	♋	19	39	22	- 0 25
			22	5	7	25	7 8 27	2 21 50	♌	1	40	23	+ 1 18
		September	4	15	13	50	7 20 3	7 23 2	♍	16	21	33	- 0 59
		December	17	3	56	48	10 22 5	1 25 56	♎	3	20	1	- 0 15
	1727	March	23	10	17	43	1 17 53	5 1 23	♏	14	35	27	- 0 47
		May	22	11	3	24	3 9 29	5 18 47	♐	0	12	2	+ 1 3
		June	29	17	39	0	4 11 56	8 26 43	♑	14	58	28	- 0 24
		September	18	11	34	42	6 21 12	5 26 21	♒	2	25	41	+ 0 5
			26	18	10	22	6 28 27	9 2 2	♓	16	32	9	+ 0 14
	1728	March	13	11	25	5	11 29 9	5 19 51	♈	24	6	20	+ 1 13
			24	21	17	28	0 9 7	10 15 12	♉	0	30	42	- 1 16
		May	8	8	41	11	1 17 26	4 11 50	♊	10	7	18	+ 1 44
		June	10	12	17	1	2 15 36	6 3 19	♋	4	6	2	+ 1 17
		October	14	18	13	17	6 4 18	9 2 5	♌	5	11	23	+ 0 50
	1729	May	7	17	47	34	0 6 0	8 22 50	♍	21	8	49	- 1 48
			24	6	6	48	0 20 4	2 29 58	♎	13	47	44	- 0 57
		September	28	13	51	44	4 9 7	7 2 55	♏	20	11	31	+ 0 39
		October	2	17	11	53	4 12 46	8 18 13	♐	9	21	12	+ 1 25
			3	18	1	35	4 13 41	8 29 53	♑	21	46	46	+ 1 50
			6	20	23	20	4 16 24	10 6 33	♒	0	52	10	+ 1 5
		November	2	18	15	6	5 10 19	9 3 13	♓	5	11	23	+ 0 50
	1730	April	1	20	58	50	9 24 15	10 12 16	♈	4	50	54	- 0 30
		November	20	17	34	18	4 15 31	8 23 37	♉	3	31	30	- 0 41
	1731	January	4	6	15	8	5 25 40	3 7 38	♊	2	45	27	+ 1 11
			19	17	52	20	6 9 36	9 0 39	♋	11	44	5	- 0 32
		February	2	5	49	14	6 21 39	2 29 52	♌	24	17	15	- 0 29
		April	10	11	39	16	8 20 49	5 26 45	♍	28	6	40	+ 0 2
	1732	January	4	14	56	0	4 15 5	7 12 51	♎	8	45	28	+ 0 39
		February	29	12	13	30	6 5 0	6 2 57	♏	24	47	30	+ 0 45
		September	19	8	55	22	11 29 13	4 15 53	♐	23	10	18	- 0 10
			22	11	37	28	0 1 57	5 29 2	♑	10	8	27	- 0 22
			28	17	27	31	0 7 25	8 22 42	♒	9	56	59	+ 0 13

Dr. Halley.

The MOON'S LONGITUDE computed from the TABLES (continued) and compared with the respective OBSERVATIONS. C.

Eclipses. and whose Observations	Years of Christ.	Equal Time of Moon's tran- siting the Meridian.					Annual Argument.	D à O.	Moon's Longitude observed.				Error of Com- putation.
		M	D	H	M	S	s o /	s o /	s	o	'	"	' "
Dr. Halley.	1732	November	20	9	40	0	1 24 21	6 0 0	11	10	3	17	- 1 12
	1733	April	11	7	28	48	6 0 35	3 20 27	Ω	22	37	32	+ 0 28
			17	11	47	54	6 5 55	5 28 17	m	6	58	50	+ 0 34
			25	18	19	41	6 13 1	9 2 18	W	18	55	9	- 1 14
			26	19	8	56	6 13 54	9 15 34	X	3	1	6	- 0 54
		May	13	9	0	33	6 28 4	4 17 39	Y	20	43	41	- 0 31
			27	20	27	53	7 10 21	10 9 30	Z	26	26	16	+ 0 11
		September	9	9	24	57	10 9 37	4 23 8	α	20	28	34	- 1 20
			14	13	53	12	10 14 8	7 2 10	β	5	43	15	- 0 10
			18	17	48	55	10 17 46	8 28 36	γ	5	33	34	+ 0 27
			21	20	25	14	10 20 29	10 7 42	δ	17	11	25	+ 0 44
	1734	April	12	16	33	43	4 20 53	8 5 28	ε	9	36	4	- 1 12
		December	22	6	11	42	0 1 12	3 5 0	ζ	16	52	31	+ 0 23
	1735	January	3	17	24	15	0 12 27	8 21 41	η	16	43	41	- 0 35
			22	7	37	0	0 29 10	3 25 13	θ	8	12	52	- 0 59
			25	10	38	20	1 1 58	5 8 8	ι	24	26	42	+ 0 42
		September	20	13	10	20	7 27 13	6 0 0	κ	8	19	48	+ 0 32
	1736	March	15	12	6	43	1 5 20	6 0 0	λ	6	36	27	+ 0 21
		September	8	14	52	0	6 7 7	6 0 0	μ	27	20	57	+ 0 28
			23	4	34	6	6 19 29	11 29 49	ν	11	28	43	+ 0 32
Dr. Bradley.	1737	February	18	2	37	53	11 2 10	0 0 5	ξ	11	11	23	- 0 56
	1743	September	22	12	11	36	9 3 53	6 4 44	ο	15	15	0	- 0 7
		October	19	9	59	38	9 27 36	5 1 56	π	8	34	14	- 0 8
		November	9	3	14	33	10 16 0	1 17 25	ρ	16	6	0	- 0 44
		December	13	6	25	48	11 16 43	3 6 48	σ	9	20	15	+ 0 17
			19	12	1	32	11 22 30	5 29 26	τ	8	18	51	+ 0 9
			26	18	21	40	11 28 52	9 6 9	υ	22	30	40	- 0 32
	1744	March	23	17	38	42	2 16 43	8 21 48	φ	6	24	1	- 0 36
		April	10	8	4	38	3 2 20	3 26 12	χ	27	45	41	- 0 11
		May	13	10	44	0	4 0 45	5 13 31	ψ	17	1	57	+ 0 4
			23	18	47	21	4 8 48	9 9 49	ω	22	47	0	- 1 10
			24	19	28	40	4 10 0	9 21 33	Α	5	34	17	- 1 18
			26	20	57	3	4 12 13	10 16 9	Β	2	20	31	- 0 44
		June	2	2	55	31	4 17 31	1 9 28	Γ	2	50	24	- 1 32
			5	5	38	23	4 20 9	3 2 8	Δ	17	51	59	- 1 32
			9	8	41	9	4 23 40	4 13 50	Ε	12	46	17	- 0 43
			18	16	2	51	5 1 38	7 26 26	Ζ	5	41	19	+ 1 2
		July	12	11	42	4	11 29 25	5 23 45	Η	24	52	5	- 1 17
			13	12	32	40	5 22 36	6 4 46	Θ	7	10	8	- 0 0
		August	19	17	56	17	6 24 26	9 2 47	Ι	10	42	18	- 0 34
Moon Ecl. Sun . . . Moon . .		September	4	4	55	23	7 5 12	2 18 0	Κ	8	32	0	+ 0 34
			15	15	51	15	7 17 46	8 1 33	Λ	6	20	56	- 1 10
	1747	February	13	17	17	0	9 10 53	6 0 1	Μ	6	17	7	+ 0 38
	1748	July	14	0	15	48	0 10 22	0 0 24	Ν	3	9	37	+ 0 11
	1750	June	8	8	58	30	8 18 18	5 29 58	Ξ	28	17	24	- 1 26

☞ Number of Observations 100; Difference of the Errors, in Seconds of a Degree, 4428", whence the mean Difference of Error, $\pm 44'' 17'''$.

But the $\left\{ \begin{array}{l} \text{Negative} \\ \text{Affirmative} \end{array} \right\}$ Errors $\left\{ \begin{array}{l} = - 37' 15'' \\ = + 36 33 \end{array} \right\} = \pm 73' 48'' \text{ Diff.}$

Whence, the Sum is negative . . . - 42"

N. B. The 100 lunar Places, in Pages 372 and 373, were computed before the *Acceleration-Table*, P. 374, for correcting the Moon's mean Motions, and *Table*, P. 375, for correcting the VI. and VII. Equations, were made; which Places, and that in *Example*, P. 54, are computed independent of those correcting *Tables*.

Total.

Total CORRECTION of the Mean MOTION of the MOON.
According to Correction of the Moon's MEAN MOTION and ACCELERATION; for any Year of a given Century
before and since CHRIST. C.

Centuries before and after Christ.	Correction of M. Motion D, for given Centu- ries, 10 ^s 7° 52' 45", Mot. for 1 Century.	Acceleration, as the Square of the Centuries. For 1620 and 1820, + 10".	Total Correction of M. Mot. D, the Sum or Dif. of Correction, M. Mot. and Acceleration.	Total Diffe- rences in a Cen- tury.	EXAMPLES of the CONSTRUCTION of this TABLE.
B. C.	0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50	0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50	0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50	0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50	EXAMPLE I. Radical Year 1720 B. C. + 1000 Centuries 27,20 × 2' 20" = 1° 3' 28" 10" × 27,20 □ = 739,84 × 10" = Acceleration = + 2° 3' 18" Correction of M. Mot. — 1 3 28 Total Cor. M. Mot. + 0 59 50
A. C.	100 200 300 400 500 600 700 800 900 1000 1100 1200 1300 1400 1500 1600 1700 1720 1750 1800 1900 2000	100 200 300 400 500 600 700 800 900 1000 1100 1200 1300 1400 1500 1600 1700 1720 1750 1800 1900 2000	100 200 300 400 500 600 700 800 900 1000 1100 1200 1300 1400 1500 1600 1700 1720 1750 1800 1900 2000	100 200 300 400 500 600 700 800 900 1000 1100 1200 1300 1400 1500 1600 1700 1720 1750 1800 1900 2000	EXAMPLE II. Radical Year 1720 A. C. — 900 Centuries 8,20 × 2' 20" = 19' 8" 10" × 8,20 □ = 67,24 × 10" = Acceleration = + 0° 11' 12" Correction of M. Mot. — 19 8 Total Cor. M. Mot. — 7 56
					EXAMPLE III. Radical Year — 1720 A. C. 2000 Centuries 2,80 × 2' 20" = 6' 32" 10" × 2,80 □ = 7,84 × 10" = Acceleration = + 1' 18" Correction of M. Mot. + 6 32 Total Cor. M. Mot. + 7 50

☞ The Correction of mean Motion of the Moon is according to 2' 20" for each Century, from the radical Year : — before, and + after 1720. And the Acceleration is according to 10" for each square Century, from the radical Year : + before and after 1720. Whence, by connecting those Quantities, the total Correction of mean Motion, and total Difference in each Century, are determined, as above.

N. B. THE Acceleration of the Moon's Apogee is the same as that of the Moon; but with a contrary Sign, or —, from the radical Year; and consequently, the Acceleration of the Moon's Anomaly is double the Acceleration of the Moon; with the same Sign, or +, from the radical Year, 1720. The Acceleration of the Moon's Node, from the same radical Year 1720, is the Half + $\frac{1}{2}$ of that Half, of the Moon's Acceleration, with the same Sign, or +, from the radical Year.

Acceleration D.	Acceleration D's An.	Acceleration D's Ap.	Acceleration D's N.
THUS, + A.	+ 2A.	— A.	+ $\frac{1}{2}$ A + $\frac{\frac{1}{2} A}{5}$.

THE above Acceleration-Table is added to be examined by future Observations; that it may be determined whether the Moon's mean Motion is quicker in this Age than our Tables of mean Motion represent: as many Observations of lunar Eclipses, and other Observations, without the Syzygies, seem to prove an Acceleration, different to what is remarked in P. 55.

A TABLE

A TABLE for correcting the Sum of the VIth and VIIth EQUATIONS of the MOON, at the Beginning of the Tables. C.

Argument. D à ☉.	Dec. Xrs.	Lo. Log.	Argument. D à ☉.	EXAMPLES of the USE of the Decimal Multipliers and Logistical Logarithms.	
0° 6' 00"	,59	2291	0° 12' 6"	EXAMPLE I.	
5	,59	2291	25	+ 4' 6"	Lo. Log. 1.1654
10	,60	2218	20	D à ☉, 1°	0.1612
15	,62	2076	15		
20	,64	1938	10		
25	,66	1805	5		
1 7 0	,69	1612	0 11 5	2' 50"	1.3266
5	,72	1427	25	EXAMPLE II.	
10	,75	1249	20	- 1' 10"	Lo. Log. 1.7112
15	,79	1024	15	D à ☉, 1° 15'	0.1024
20	,83	0809	10		
25	,86	0655	5	- 0' 55"	1.8136
2 8 0	,89	0506	0 10 4		
5	,92	0362	25	RULE. Multiply the Sum of the VI. and VII. Equations, connected with its proper Sign, by	
10	,95	0223	20	the decimal Xr. correspondent to D à ☉, for the Product of the required Equation. Or add the	
15	,97	0132	15	Lo. Log. correspondent to D à ☉, to the Lo. Log. of the Sum of the VI. and VII. Equations, for	
20	,99	0044	10	Lo. Log. of the required Equation, to which prefix the Sign of the Sum of Equations.	
25	1,00	0000	5		
3 9 0	1,00	0000	0 9 3		

VI. Equation, + 2' 25"

VII. Equation, + 1 41

Sum, + 4 6

In Seconds, + 246"

D à ☉ . . 1° . . 69 Xr.

2214

1476

169,74 = + 2' 50" Equation.

VI. Equation, - 2' 0"

VII. Equation, + 0 50

Sum, - 1 10

In Seconds, - 70"

D à ☉ . . 1° 15' . . 79 Xr.

630

490

55,30 = - 55" Equation.

A COMPARISON of the Number, Arguments, and GREATEST EQUATIONS of the MOON.

According to our Royal Astronomer.				According to Mayer.				According to Sir I. Newton.			
N.	Arguments.	Sig. & Deg.	Greatest Equations D.	No.	Args. correspond.	Sig. & Deg.	Greatest Eq. D.	No.	Arguments and Greatest Eq.		
I.	☉'s M. Anom.	{ 3° 00'	D - 11' 25" Ap. + 26'	I.	☉'s M. Anom.	{ 3° 20'	D - 11' 20"	1.	Arg. ☉ M. An. - 11° 50'		
a	denotes Annual.	{ 9 0	= 19' 38" + 6' 22"	a		{ 8 29			Apogee D . . + 19 43		
II.	☉ à ☽'s Ap.	{ 1 15	+ . . 5' 59" =	X.	D à ☉ - M. An	{ 1 15	± . . 3' 45"	2.	Arg. ☉ à ☽ Ap. 3' 34" ☉ Ap.		
s	denotes Semiannual.	{ 10 15	3' 45" + 2' 14"	s	reversed Arg.	{ 10 15			☉ in ☽'s An. 3' 56" ☽ Per.		
III.	☉ à ☽	{ 1 15	+ . . 0' 47"	IX.	☽ à ☉	{ 1 15	± . . 0' 47"	3.	Arg. . . ☉ à ☽		
s		{ 10 15		s	reversed Arg.	{ 10 15			☽ - 1° 15' - 0' 47" D		
IV.	Mean Evection	{ 3 0	± . . 1' 0"	VII.	2. D à ☽	{ 3 0	± . . 0' 58"		ants { 10 15 + 0' 47" D		
s	+ 2. ☉ à ☽	{ 9 0		s	- ☽ An.	{ 9 0			- to 3° + to 6° - to 9° + to 12°		
V.	Mean Evection	{ 3 0	± . . 1' 25"	VI.	2. D à ☉	{ 3 0	± . . 1' 30"				
s	+ 2. ☽ M. An.	{ 9 0		s	+ ☽ An.	{ 9 0					
VI.	Mean Evection	{ 3 0	± . . 2' 25"	IV.	2. D à ☉	{ 3 0	± . . 1' 48"	5.	Arg. D à ☉ + ☽ Ap. à ☉ Ap		
s	+ ☉'s M. An.	{ 9 0		s	+ ☉ An.	{ 9 0			3° ± 2' 25"		
VII.	Mean Evection.	{ 3 0	± . . 1' 41"	V.	2. D à ☉ - ☉ An.	{ 3 0	± . . 1' 12"		+ to 6° - to 12° Arg.		
s	- ☉'s M. An.	{ 9 0		s	- ☽ An.	{ 9 0					
VIII.	☽'s corrected An.	{ 3 4	+ 6° 18' 38"	XI.	☽ corrected An.	{ 3 4	± 6° 18' 44"	4.	Arg. ☽'s An. fr. 1st and 2d		
a		{ 8 26		a		{ 8 26			Eq. Ap. ☽ Central Eq. accordg		
IX.	Tr. Evcc. or M. Evcc.	{ 3 1	+ 1° 20' 26"	XII.	2. D à ☉ cor.	{ 3 1	± 1° 20' 43"		to Eccentricity ☽ Orb. includg		
a	+ 1/2 Eq. ☽'s Cent.	{ 8 29		s	☽ An.	{ 8 29			sur 8 and 9; or Mayer's 11 and		
X.	☽ eq. à ☉ tr.	{ 1 14	+ . 36' 50" ☽ falc.	XIII.	☽ eq. à ☉ true.	{ 1 14	+ 39' 3" ☽ fal.	6.	Arg. D à ☉ ☽ Fr. 33' 14"		
s		{ 4 14	- . 39 22, gibbous.	s		{ 4 15	- 41 41, gibb.		In ☽'s An. To 37 11		
								7.	Doubtful to Newton.		

No. For the MOON'S Latitude.

I.	D in Orb à ☽.	{ 3° 00'	+ 5° 8' 28"	I.	D in Orb à ☽.	{ 3° 00'	+ 5° 9' 8"	☽ M. Evcc. - 2. D +
II.	D à ☉ - D à ☽	{ 3 0	± . . 8 52	II.	2. D à ☉ - D à ☽	{ 3 0	± . . 8 50	2. ☉ + ☽ An. + any Com.

No. For the Moon's PARALLAX.

I.	M. An. ☽ equated.	6° 0	+ . 60' 25"	I.	☽ cor. An.	6 0	+ . 60' 26"	Because 2. ☽ - 2. ☉ - ☽ An.
II.	Mean Evection	0 0	- . . . 42"	II.	2. D à ☉ - ☽ An.	0 0	- . . . 38	= Mean Evcc. = ☽ M. An.
III.	☽ eq. à ☉	3 0	- . . 1 0	III.	☽ eq. à ☉	6 0	- . . . 28	- 2. ☉ à ☽ Apogee.

Mayer's II. III. and VIII. Equations have these Arguments and Greatest Equations.

Arguments.	Sig. Deg.	Gr. Equat.	Arguments transformed.
II. 2. D à ☉ + ☉ M. An.	{ 3° 00'	+ 0' 54"	Or Arg. Mean Evection + Moon's M. Anomaly + Sun's M. Anomaly.
III. 2. D à ☉ - ☉ M. An.	{ 3 0	+ 1 2	Or Arg. Mean Evection + Moon's M. Anomaly - Sun's M. Anomaly.
IV. ☽ M. An. - ☉ M. An.	{ 3 0	± 0 40	Or Arg. Mean Evection + Sun's M. Pl. + Sun's Ap. - 2. Moon's Ap.

OF DETERMINING the MAXIMA, and CONSTRUCTION of the LUNAR EQUATIONS at the Beginning of this WORK.

I. FOR the greatest ANNUAL EQUATION of the MOON.

Put $Q = S.$ Sun's Equat. at greatest; $s =$ the D 's of $\odot$'s Period; $m =$ Moon's Period to $\star$; $n = D$'s Synodic Period to Sun; Then,

$$\frac{3^{mm}}{2,10611n} \times Q = \text{Sine greatest annual Eq. } D = 11' 25''.$$

$3 \dots \dots \dots$	0.4771212	$2,1061 \dots$	0.3234790
$m^2 = 27^d, 3216 \text{ squared}$	2.8730122	$s = 365^d, 2565$	2.5625979
$Q = S. 10' 55' 50''$	8.5274775	$n = 29^d, 53$	1.4702634

Sum, 11.8776109 Sum, 4.3563403
 $- 4.3563403$

Gr. an. Eq. D $11' 25''$ S. 7.5212706 Dif.

For gr. Equation D 's Apogee, or Ω .

Put $p = m.$ periodic Time D 's Ap. or Ω , and $q =$ Time of Synodic Revolution of the Sun to Ap. or Ω . Then

$$\frac{39}{2,2551p} \times Q = S. \text{ gr. Eq. Ap. And } \frac{3,1669q}{2p} \times Q = S. \text{ gr. Eq. } \Omega.$$

$3 \dots \dots \dots$	0.4771212	$2,2551 \dots$	0.3531658
$q = 411^d, 7863$	2.6146718	$p = 3231^d, 357$	3.5093850
$Q = S. 10' 55' 50''$	8.5274775		

Sum, 11.6192705 Sum, 3.8625508
 $- 3.8625508$

Gr. Eq. Ap. $19' 38''$ Dif. S. 7.7567197
 $+ 6 22$ Correction of Erection added.

26 0

Again, 3.1669 0.5006344 $2 \dots \dots$ 0.3010300
 $q = 346^d, 6197$ 2.5398533 $p = 6798^d, 203$ 3.8323941
 $Q = S. 10' 55' 50''$ 8.5274775

Sum, 11.5679652
 $- 4.1334241$

G. Eq. Ω $9' 21''$ Dif. S. 7.4345411

For present Equations D , Apogee, and Ω .

As $\left\{ \begin{array}{l} \text{gr. Eq. } \odot \\ 10' 55' 50'' \end{array} \right\}$	to $\left\{ \begin{array}{l} \text{gr. an. Eq. } D \\ \text{gr. Eq. } D \text{ Ap.} \\ \text{gr. Eq. } D \text{ } \Omega \end{array} \right\}$	so $\left\{ \begin{array}{l} \text{Pref. Eq. } \odot \\ \text{Eq. } \odot \end{array} \right\}$	to $\left\{ \begin{array}{l} \text{Pref. Eq. } D \\ \text{Pre. Eq. Ap.} \\ \text{Pref. Eq. } \Omega \end{array} \right\}$
Sir Isaac Newton determines	Gr. Eq. D	Ap.	Ω .
Mr. Machin from his Principles	$11' 49''$	$19' 43''$	$9' 24''$
	$12 5$	$21 57$	$8 56$

When Sir I. Newton had found the ann. Eq. D , he proportions,
 As m. diur. Mot. $\odot$ to M. diur. Mot. Ap. $6' 44'' 4''$ so $\odot$ Eq. $10' 56' 20''$
 $\odot$ $59' 18'' 19''$ to M. diur. Mot. Ω $3 10 38$ so $\odot$ $1' 58 10$
 to gr. Eq. Ap. $19' 43''$ respectively, $\left\{ \begin{array}{l} 10' 53' 45'' \\ 19' 38'' \end{array} \right\}$
 to gr. Eq. Ω $9 24$ as above. $\left\{ \begin{array}{l} 2' 54' 30'' \\ 2' 54' 30'' \end{array} \right\}$
 The $\odot$ adds in 1st 6 Signs. sub. in the last 6 Signs.
 Eq. of $\odot$ adds $\odot$'s M. An. but adds Sun's mean Anomaly
 being the Argument.

H. FOR the greatest first SEMI-ANNUAL EQUATION D .

As synodic Time $\odot$ to D 's Ap. $411^d, 786$ squared, co.	4.7706568
To synodic Time D to $\odot$ $29^d, 53$ squared	2.9405268
So S. gr. 2d Equat. Moon's Ap. $12' 18'$	9.3284416
To S. gr. first Semi-an. Equat. req. $3' 46''$ in Octants $\odot$	7.0396252

Or, in the same Ratio,

As Sq. M. diur. Mot. D to $\odot$, $12^d 11' 27'' = 43887''$ sq. co.	Logarithms, 0.7153282
To Sq. M. diur. Mot. $\odot$ to D 's Ap. $52' 27'' = 3147''$ sq.	6.957934
So Sine gr. 2d Equat. Ap. $12' 18'$	9.3284416

To S. $3' 46''$ as before, in Octants	7.0395632
---------------------------------------	-------------

Now, in Octants this Equation is increased and diminished in the reciprocal triplicate Ratio of the Sun's Distance from the Earth; and is $3' 34''$ when least, and $3' 56''$ when greatest. — But when the Moon's Apogee is out of the Octants,

As Rad. : to Arg. or 2. $\odot$ to D 's Ap. (or, according to Newton, twice Moon's Apogee from nearest Syzygy) :: $3' 46''$, or other greatest Equation in Octants : present Equation, required.

This Equation adds to mean Motion in the Passage of the Moon's Apogee, from the Quadrature of the Sun to the Syzygy; but subtracts in its Passage from the Syzygy to the Quadrature.

III. FOR the greatest second SEMI-ANNUAL EQUATION.

As Radius	10.0000000
To Sine of mean Inclination D 's Orb $5' 8' 28''$	8.9523314
So S. Half Dif. between gr. and least Incl. $8' 52''$	7.4114360
To S. $47'' 40''$, greatest Equation, required	6.3638374
[in Octants of $\odot$ to Ω ; in Per. $45''$ in Ap. $49''$]	

Being reciprocally, as Cube of the Sun's Distance from the Earth.

In other Positions,

As Rad. : S. Arg. 2. $\odot$ to Ω (or, according to Newton, to S. twice Ω , from nearest Syzygy) :: S. greatest Equation $47''$, or other Equation in Octants : S. present Equation, required, out of Octants of the $\odot$ to Ω , which is the Argument of this Equation.

IV. FOR the greatest of THIS EQUATION.

As Radius	10.0000000
To Cube Cos. mean Inclination of D 's Orb, $5' 8' 28''$	9.9947467
So Sine of Erection-Equation $10' 20' 40''$	8.3703804
To Sine of $10' 19' 42''$	8.3651271

Dif. Greatest Equation, required, $58''$ By Tab. 1'
 As Rad. : S. Arg. mean Erection $+ 2. \odot$ to Ω :: $58''$ or $1'$: present Equation.

V. FOR the greatest of THIS EQUATION.

As Sum of Forces 6.50 355.450 co.	7.4467847
To 3, Dif. solar Force D Per. and Apog. 1718	9.2350232
So Radius	10.0000000
To S. $1' 39''$, greatest Equation, required	6.6818079
As Rad. : S. Arg. Erection $+ 2 D$ M. Anom. :: $1' 39''$: present Equation.	

*VI. and VII. FOR the Sum of these GREATEST EQUATIONS.

As Semi-Transv. Earth's Orbit 100000 co.	5.0000000
To 3 Times Eccentricity $1685 = 5055$	3.7037212
So S. Erection $10' 20' 40''$	8.3703804
To S. Sum Equations $4' 5''$ required	7.0741016

$10' 55' 50''$ Sun's Equation,
 $57 55$ Half.

So $2 53 45$ Sum,

* As $100000 D$'s mean Dist. fr. $\odot$: to $2 DF$ (see P. 369, l. 12. fr. lat.)
 $= 11026 ::$ so Rad. : to Tang. $4' 6''$, = $\angle$ subtended by $2 DF$, seen from the D , at a mean Dist. and Perp. thereto; or seen at D , in a parallel Position to DF , from the upper Focus of D 's Orbit. See Fig. III, foregoing; and Fig. IV, further on.

Of DETERMINING the MAXIMA, and CONSTRUCTION of the LUNAR EQUATIONS at the Beginning of this WORK.

VI. FOR the greatest of THIS EQUATION.

As Sem. Tran. Earth's Orbit	100000 co.	Logarithms.
To twice Eccen. 1685 = 3370		5.0000000
And 59 to 52		3.5276299
So S. Evection	10 20' 40"	9.9451513
		8.3703804
To S. this gr. Equation	2' 24" required	6.8431616
Sum Equations	4 5	

Dif. 1 41 greatest VII. Equation.

VI. Rad. : S. Arg. Evec. + Sun's M. An. :: 2' 24" : present Equation.

VII. Rad. : S. Arg. Evec. — Sun's M. An. :: 1' 41" : present Equation.

N. B. The Tables shew, by the Arguments, when the Equations add or subtract.

VIII. GREATEST EQUATION, 6° 18' 38", and Construction, is seen in P. 48.

IX. GREATEST EQUATION, 1° 20' 40", and Construction, are seen P. 373. See also P. 362.

X. First and principal GREATEST VARIATION-EQUATION deduced, P. 361 and 362.

N. B. The greatest second Variation-Equation is to Rad. : as 2 to 344, nearly. 342 to 344 is as the least and greatest Distance of the Moon from the Sun in Conjunction and Opposition. — Hence the greatest second Variation = 2'.

And as Rad. : S. Arg. $\odot$ à $\odot$:: 2' : present second Variation-Equation. The present principal and also second Variation form the Compound Variation-Equation; connected together with their proper Signs.

The Tables in general shew when the Equations are to be added or subtracted.

COMPOUND EQUATIONS explained.

I. Compound annual Equations of the Moon's Apogee adds 6' 22" to the greatest Quantity for correcting the Evection.

II. or 1st Semi-annual. — The EVECTION, at a Mean, at each Apis, is $\pm 4' 28'' \frac{1}{2}$; the Half 2' 14". The mean Semi-annual, at greatest, being 3' 45" + 2' 14" = 5' 59", as by Tables. — But the synodic Time of the Sun's Motion from the Moon's Apogee in the lower Semi-circle of her Orb, being less than in the higher Semi-circle, this Equation is diminished also. Hence arises another Equation, which, like the second Variation, is greatest in Quadratures of the Sun with the Apisides, and amounts to about 30". It subtracts in the first Semi-Circle, and adds to the Moon's mean Motion in the last Semi-Circle.

For MOON's Mean LATITUDE.

Const. As Rad. : S. 5° 8' 28" :: S. $\odot$ à $\odot$: S. present Lat. $\odot$.

The mean horizontal Parallax is determined according to the Arg. $\odot$'s M. Anom. or equated by 10 first Equations $\odot$, and compounded with some few Seconds; then reduced to the true horizontal Parallax by Tab. whose Argument is mean Evection (like the mean central and Evection Equations) and farther corrected by Tab. whose Argument is $\odot$'s true Place from $\odot$. See P. 53.

FOR COMPARISON of COMPUTATION, Our Astronomical Tables are readily reduced to the Astronomia Arcana, or GREENWICH TABLES, of the Mean PLACES of the SUN and MOON, by the following EQUATIONS.

EQUATION of the SUN's Mean Places.

M. Pl. $\odot$	M. An. $\odot$	Ap. $\odot$	Use the same Signs for
+ 0' 17"	+ 1' 19"	- 1' 2"	Time back as forward.

EQUATION $\odot$'s M. Motion for 100 Years forward from 1700.

Eq. M. Mo. $\odot$	Eq. Mo. An. $\odot$	Eq. Mo. $\odot$ Ap.	Use contrary Signs for
0' 0"	- 5' 50"	+ 5' 50"	Time back.

N. B. The same Signs are used for Time back, as for Time forward, in mean Places of $\odot$ and $\odot$.

But contrary Signs for Time back to those for Time forward in Motion $\odot$ and $\odot$ equated. — Cut off 2 Figures to Right, of Seconds for 100 Years for the Seconds for 1 Year and multiply by the No. of Years from 1700 for the whole Equation, in Seconds, with its proper Sign.

The Mean of the Dif. Maxima, in Ostants, thence arising, will be 5' 38", compound Equation. The Construction, As Rad. : S. 2° à $\odot$'s Ap. :: 5' 38" : present Equation. Again, As Rad. : S. $\odot$ à $\odot$'s Ap. :: 30" to another Equation.

X. The mean Variation being 35' 10" + 2' 14", and Correction of mean Anomaly according to variable Eccentricity = 1' 6", added, make 38' 30", compounded Equation. But this Excess of Reduction, between the greatest and least, being 24", and always Subtraction of Variation, the mean tabular Variation is 38' 6" $\pm$ 9", further Correction of Evection, according to the Moon's Anomaly, as the Sun is in the lower, or higher Apis, gives 38' 15" in the 1st and 4th Quadrant; and 37' 57", in 2d and 3d Quadrant.

The Construction of these Equations is,

Const. Log. 9.9903357 1.4 Quad. + Log. Tan. 45° = Log. Tan. 9.9904115 2.3 Quad.

4th Proportional, which taken from 45° leaves 38' 15" } Arg. Moon à Quadrature of the Sun.

Again, As Rad. : S. $\odot$ à $\odot$:: 2' : to present second Equation, as before.

The Equation in the Ostants for correcting the Variation according to the Moon's Anomaly, is as before 1' 39". But, by Observation, in the Quadrature and Syzygies, amounts to 1' 10" only; somewhat correspondent to the Excess of Newton's second Variation. In order to accommodate the Tables to this Correction, the Evection is diminished 15", and this Equation 14". In the Quadratures and Syzygies, the Sum of this Diminution is 29", and reduces this Equation to 1' 10". In the Ostants, the Diminution is additional to this Equation, and raises it again to its Maximum 1' 39".

The Compound-Variation is compounded of the principal, and second Variation, and also the Half of the Correction of Evection.

CONSTRUCTION of the REDUCTION-EQUATION.

As Rad. : to Cosine of the mean Inclination of the Moon's Orb :: Tangent of 45° : to a Tangent of an Angle, which, subtracted from 45° leaves the greatest mean Reduction, required.

As Radius 10.0000000

To Cosine of mean Inclination of Moon's Orb. 5° 8' 28" 9.9982493

So Tangent 45° 10.0000000

To Tang. 44 53 4 9.9982493

Dif. 6' 56" Reduction, required.

As Rad. : S. 6' 56" :: S. doub. Arg. : S. present mean Reduc. Equat.

For EQUATION of MOON's mean to her true LATITUDE.

Const. As Rad. : S. 8' 52" :: S. Arg. : S. present Equation.

EQUATION of the MOON's Mean Places.

M. Pl. $\odot$	M. Pl. An. $\odot$	M. Pl. Ap. $\odot$	M. Pl. $\odot$
- 0' 49"	+ 0' 54"	- 1' 43"	+ 1' 50"

EQUATION $\odot$'s Motion for 100 Years forward from 1700.

M. Mot. $\odot$	M. Mo. $\odot$ An.	M. Mo. $\odot$ Ap.	M. Mo. $\odot$
+ 3' 5"	0' 0"	+ 3' 5"	+ 3' 5"

Acceleration of $\odot$'s Mot. for 100 Years forward or back.


+ 12" | + 24" | - 12" | + 8"

The Quantities of Acceleration are as the Square of Centuries or Parts of Centuries from 1750.

EXPLANATION

EXPLANATION of NEWTON'S LUNAR THEORY. See Fig. IV.

FIG. IV.



AP, The *Apfides* of the Moon's M. Orbit, or *transverse Diameter* = 2000000 Parts, answerable to the mean E. centricity = 55050 Parts = TC, or CB. T represents the Earth, in the lower Focus of the Moon's mean elliptic Orbit, PLHAHP, round which the Moon's true Motion is performed. B represents the upper Focus of the same Orbit, round which the equal Motion of the Moon is nearly performed. HH, the conjugate Diameter; LL, the Latus rectum; RDNOR, the Epicycle of Halley, (whose Center is at C) in which the Center of the Moon's Orbit is supposed to revolve to the Left.

Parts.
Therefore, TN = least Eccentricity Moon's Orb = 43319
TR = greatest Eccentricity . . . = 66782
TC being = mean Eccentricity . . . = 55050
Hence, NC (= CD) = CR = TR - TC = 11732
DF = 596

NOW, the Point D is moved to the Left in Halley's Epicycle, so as the Angle RCD may be always equal to twice Sun's Apogee equated; and CTD = ATQ, second Equation of the Moon's Apogee.

But Newton, for further Improvement of the lunar Theory, translates D to F, so that $\angle CDF$ = Complement of the Sun's true Anomaly to 360° , and DF to DC, as twice the Eccentricity of the Earth's Orbit to the Sun's mean Distance from the Earth, and the Sun's mean diurnal Motion from the Moon's Apogee to the Sun's mean diurnal Motion from its own Apogee, conjunctly, viz. as $33\frac{1}{2}$ to 1000, and $52' 27'' 16'''$ to $59' 8'' 10'''$, conjunctly, or as 3 to 100, nearly; whereby DF = 352 Parts of Eccentricity, according to Newton. But, if CD, the Epicycle's Radius, be considered to DF as 10000 to 508, (for a farther Correction of the Theory) then, by Proportion, DF = 596,5 Parts of Eccentricity. And this DF, at the Distance of the Center of the Moon's Orb TC, subtends the Angle at the Earth, which the Translation of the Center of her Orbit from D to F generates in its Motion.

The Double of which Line DF, in a parallel Position, at the Distance of the upper Focus, B, of the Moon's Orbit from the Earth, seen at double Dist. of the lower Focus, subtends, at the Earth, the same Angle as DF did before, seen at the lower Focus, which that Translation from D to F generates in the Motion about this upper Focus. But, at the Distance of the Moon from the Earth, 2DF at the upper Focus, in a parallel Position to the first Line DF, subtends an Angle at the Moon, which the said Translation generates in the Moon's Motion. This Angle is called 2d Equation of the Moon's Center, which, at a mean Distance of the Moon from the Earth, is nearly as the Sine of the Angle comprehended by Lines drawn from F and D to the Moon; and amounts to $2' 25''$ when greatest, according to Newton; 2DF being then at right $\angle$ s with the Moon's mean Distance. But DF being taken = 596,5 instead of Newton's 352, the greatest Equation will then amount to $4' 6''$, according to our Tables. And the Argument for finding this Equation at all Times, (according to Newton) is by adding the Distance of the Moon to the Moon's Apogee from the Sun's Apogee.

Then, As Rad. : S. Argument thus found :: $2' 25''$: present second Equation of the Moon's Center. To be added when the Argument is less than a Semi-Circle, and subtracted when greater. And thus, by Newton's placing the Center of the Moon's Orb at F, to revolve round Halley's moving Point at D, it is made to describe a certain curve Line, of variable Radius FC, about the Center C, with a Velocity almost reciprocally as the Cube of the Sun's Distance from the Earth; according to the Laws of Nature, observed.

TB D, Angle at upper Focus.

AB D, the mean Anomaly of the Moon.

BC D, the eccentric Anomaly of the Moon.

BT D, the true Anomaly of the Moon.

RCD = twice Sun's Ap. = 2, An. Argument.

TD or TF = present Eccentricity of the Moon's Orb.

CTD or CTF = second Equation of the Moon's Apogee.

QQ, the present Position of the Apfides or Apogee.

C D or C F, the Semi-Angle of Evection, at greatest (when CD D or CF D = 90°) = $40' 20''$ per fer . DCD or DCF Argu-

MB D, Angle of Correction of the Moon's mean Anomaly, and $\angle ABM$, the Moon's corrected mean Anomaly, according to Bullialdus. S. C.

DIFFERENT OBLIQUITIES of the ECLIPTIC as observed in different AGES of the WORLD, by eminent ASTRONOMERS.									
Yrs bef. Chr.	Astronomers.	Obliquity.	Yrs since	Astronomers.	Obliquity.	Yrs sin. Ch.	Astronomers.	Obliquity.	
		° ' "	Christ.		° ' "	1490	Dominicus Maria Novaras	23 29 0	
280	Aristarchus	23 51 0	380	Mabumed Eben Ga-	23 35 0	1514	Ferrariensis	23 29 0	
270	Eratosthenes	23 51 0		ber	23 35 0	1572	Venerus	23 28 0	
240	Hipparchus	23 51 0	911	Ababet Eben Corra	23 33 30	1670	Tycho Brahe	23 31 0	
Since Chr.			992	Abu Mabumed Al	23 22 21	1760	Hevelius	23 30 20	
140	Ptolemy	23 51 20		Cogandi	23 22 21	1760	P. Mengoli	23 28 24	
825	Benimula	23 35 0	1269	Cojab Nasirodni	23 30 0	1673	Mr. Flamstead	23 29 0	
827	Almamun	23 35 0	1363	Eben Sbatir	23 31 0	1700	Dr. Halley	23 29 0	
828	Iabia Eben A-	23 35 0	1437	Uleg Fieg	23 30 17	1750	Dr. Bradley	23 28 30	
	bumanfar	23 35 0	1460	Regiomontanus	23 29 0				


N. B. The principal Difference of the above Observations has partly arisen from the Imperfection of the Instruments with which they were taken, the different Skillfulness of Observers, and their imperfect Knowledge of Parallaxes and Refractions. See Philosoph. Transactions, No. 163. And Marcus Manilius.

EXPLANATION

EXPLANATION of MACHIN'S LUNAR THEORY. See Fig. V.

FIG. V.

THE CIRCLE ABC represents the Moon's Orbit about the Earth, placed in the Center T, at a mean Distance, the Moon having no other Gravity than to the Earth. The Diameter ATC divides the Part of the Moon's Orbit ABC towards the Sun from that Part ADC, opposite to the Sun; the Diameter BTD is the Line of the Moon's Conjunction with or Opposition to the Sun. The Fig. LMNO is an Ellipsis, whose Center is carried round the Earth in the Orbit AEBFCGDH: Its longer Axis LN, always parallel to BD, joining the Earth and Sun, TS, is double its shorter Axis MO. While the Fig. LMNO is carried to the Left, from A to E, B, F, C, G, D, H, &c. the Moon revolves contrary, from M to P, N, Q, O, R, L, V, M, describing equal Areas in equal Times about the Center of this elliptic Epicycle, so as to perform a Revolution therein, in the same Time that the Center of the said elliptic Epicycle performs its Revolution, in the circular Orbit.



The Moon is always in the remoter Extremity of its shorter Axis in M, and O, when in her Quarters from the Sun; and in the nearer Extremity of its longer Axis, at the Time of New and Full Moon. The shorter Semi-Axis of this Ellipsis MA, is to the Distance of its Center from the Earth, AT, in the duplicate Proportion of the Moon's periodical Time about the Earth to the Sun's periodical Time; being as 47 to 8400, nearly.

The Figure MPNQORLVM described by this compound Motion of the Moon in the elliptic Epicycle, while the Center thereof is carried round the Earth, very nearly represents the Form of the Moon's Orbit; supposing it without Eccentricity, and that its Plane was coincident with the Plane of the Ecliptic; and also that the Sun continued in the same Place, during the Moon's whole Revolution about the Earth.

OF MACHIN'S THEORY of the MOON'S MOTION.

PROPOSITION. A Body being deflected by two Forces, tending constantly to two fixed Points, will describe, by Lines drawn from those fixed Points to the Body, equal Solids in equal Times, about the Line joining the said fixed Points.

This general Law of centripetal Forces, according to Kepler, is, that all Bodies describe equal Areas in equal Times, about the single Center of their respective Revolutions; but as this Law has been proved defective, by Sir Isaac Newton, when a Body gravitates to two different Bodies at the same Time, as every Satellite gravitates towards its Primary, while it gravitates also towards the Sun, Mr. Machin, to supply the Defect of Kepler's Law, of Gravitation to one Center only, has explained the Motions of the Moon, and of the other Satellites, by the Law of the above Proposition, according to a Gravity tending to two different Centers at once.

COROLLARY. If a Body gravitates towards two fixed Centers, be supposed, for given small Intervals of Time, moving in a Plane passing through one of those fixed Centers, the Inclination of the said Plane to the Line joining both the Centers, will vary according to the Area described: That is, if the Area be augmented, the Inclination will be diminished; and if the Area be diminished, the Inclination of the said Plane will be augmented; that the Solids described, according to the Proposition, may be equal.

THIS Corollary, being rightly applied, will serve to explain the Variation of the Inclination of the Plane of the Moon's Orbit with the Plane of the Ecliptic; without entering into the intricate Calculations contained in the Corollaries to 34 Prop. of the III. Book of the Principia, to determine the greatest Quantity of Variation, when the Line of Nodes is in Quadrature with the Sun; which, in particular Numbers, is found to be $2^{\circ} 43''$, or $2^{\circ} 45''$, according to Newton.

But in this Case before us, we have a plain and general Rule, from the foregoing Premises; namely, that the greatest Variation, in the said Position of the Moon's Orbit, is to the mean Inclination of the Plane thereof, as the Difference of the greatest and least Areas, described in the same Time, by the Moon about the Earth, when in the Conjunction and in the Quadratures, to the mean Area.

Hence, if S be to L, as the Sun's to the Moon's Period; then the greatest is to the least Area, as $\sqrt{SS+3LL}$ to S, or as $S + \frac{3LL}{2S}$ to S, nearly. So that the Difference of the Areas, is to the mean Area, as $\frac{3LL}{2S}$ to $SS + \frac{3LL}{2S}$; in which Proportion is the Variation of the Inclination of the Plane of the Moon's Orbit, at this Time, to the mean Inclination thereof; agreeing, nearly, with Sir Isaac Newton's Computation.

OF MACHIN'S THEORY of the MOON'S MOTION.

MOREOVER, there is a *general Method*, as Mr. Machin observes, for assigning the *Laws* of the Motion of a Body to and from the Center, abstractedly considered from its Motion about a Center. And this Motion, to and from the Center, is called by *Kepler*, the *libratory Motion* of that Body, distinct from its *circular Motion*: The Knowledge of which Motion Mr. Machin thinks necessary for defining the *Laws* of the Revolution of a Body, in Respect of the Motion of the *Apsides* of its Orbit; which Revolution, from one *Apsis* to the other, is performed in the Time of the whole *libratory Motion* of the Body; the *Apsides* of the Orbit being the extreme Parts thereof, in which only the *libratory Motion*, to and from the Center, ceases.

According to *this Theory*, the Motion of a Body round the Center of Revolution, is not considered as a continued Deflection from a *straight Line*; but as compounded of a *circulatory Motion* round the Center, and of a *rectilinear Motion* to and from the Center of Revolution. To each of which Motions, Mr. Machin assigns its proper *Equant*.

The general Property of an *Equant* for any Motion, is a *Curve-Line* described about the Center thereof, whose Rays are reciprocally in the *subduplicate Velocity* at the Center, or *Velocity of Revolution*.

The *EQUANT* for the *libratory Motion*, is a *curve-lined Figure*, the Areas of which are proportional to the Times wherein the several Spaces of Libration are performed; which Figure is determined by the Law of Gravity, towards the Center, being known. For the *libratory Force*, to accelerate or retard the Motion, to or from the Center, is the Difference between the Gravity of the Body towards the Center, and the *centrifugal Force* arising from the *circulatory Motion*; the latter being always subject to one Rule: Because, in all the *Revolutions* about a Center, in any Curve, whether described by a *centripetal Force*, or not, the *centrifugal Force* will be directly, in the duplicate Proportion of the Area described in a given Time, and reciprocally, in the triplicate Proportion of the Distance; being an immediate Consequence of a known Proposition of *Huygens*. And the like Proportion also holds as to the *centripetal Force*, in all *circular Motions* whatsoever, from a known Prop. in *Sir Isaac Newton*. And what is true of the *centripetal Force* in Circles, is universally so of the other Force, in Orbits generated of any Form.

Moreover, that by knowing the Gravity of the Body, the other Force being always known, the Difference thereof (being the *absolute Force* to move the Body to and from the Center) will be also known. And, from thence, the Velocity of the Motion, and the Space described, in a given Time, may be found; and also the *Equant* described.

Example. If the Gravity be reciprocally as the Square of the Distance, the *Equant* for the *libratory Motion*, will be found to be an *Ellipsis*, similar to the Orbit, whose longer Axis is double the *Eccentricity*. And the Center of the *libratory Motion*, that is the Place where it is *swiftest*, will be in the Focus of that Orbit. The Time of Libration, through the several Spaces, will be measured by the *Sectors* of the said *Ellipsis*, similar to those described by the Body round the Focus of the Orbit. And the Period of the *libratory Motion* will be the same with the Period of the Revolution.

In any other Law of Gravity, the *Equant* for the *libratory Motion*, will be either of a Form different from the Orbit, or if of the same Form, will not be alike divided.

The *Equant*, for the *libratory Motion* of the Moon, is a *Curve* of the third Kind, whose Equation is of four Dimensions; but may be described by an *Ellipsis*; the Center of Libration not being in the Focus.

From this Method of resolving the Moon's Motion, it is not difficult to shew the *general Causes* of the Change of *Eccentricity*, and the *Inequality* of the Motion of the Moon's *Apogee*. For, when the Line of *Apsides* is moving towards the Sun, it is easy to shew, that since the external Force in the *Apsides* is then *centrifugal*, it will contribute to lengthen then the Space and Time of Libration. By lengthening the Space, it augments the *Eccentricity*; and by lengthening the Time of Libration it also prolongs the Time of the Revolution to the *Apsis*; causing what is improperly called a *progressive Motion* of the *Apsis*. But when the Line of *Apsis* is moving to the *Quadratures*, the external Force in the *Apsides*, is then *centripetal*, which will contribute to shorten the Space and Time of the Libration. And, by shortening the Space, the *Eccentricity* will thereby be lessened; and by shortening the Time of Libration, the Time of Revolution to the Axis will thereby be

contracted; causing what is improperly called a *retrograde Motion* of the *Apsis*.

REMARKS. The Motion of the Moon, in the before described *elliptic Epicycle*, only shews the *Elongation* of the Moon from the Center of that *Epicycle*; but does not describe an accurate *Ellipsis*, according to the *Newtonian Theory*.

When the Moon is in that Half of its Orbit next to the Sun, it then being nearer than the Earth to the Sun, it will have a greater Gravity than the Earth to the Sun; which Excess of Gravity, according to the *Newtonian Theory*, consists of two Parts; one acting in the Line PE, parallel to that which joins the Earth and Sun; and the other acting in the Line EC, directed to the Earth. And these two Forces being compounded into one, constitute a Force directed in the Line PE, (see Fig. V.) which is in Proportion to the Force of Gravity, as that Line PE, is to TE, nearly. Hence, as there is a Force constantly impelling the Moon somewhere towards the Point e; this Force is supposed to inflect the Motion into a *Curve-Line* about that Point; for the same Reason as the Gravity of it to the Earth is supposed to inflect its Motion into a *Curve-Line* about the Earth: Not that the Moon can actually have so many distinct Motions; but that one simple Motion of the Moon round the Sun is supposed to result from a Composition of these several Impulses, or Motions.

If AE represent the Equation of the Sun's Center, S the Sun's Period; P, the *synodical Period* of the Moon to the Sun; L, the Moon's Period to the Stars; Then, $\frac{3LL}{2PS} \times AE =$ the annual Equation of the Moon's mean Motion.

If AE represent the Equation of the Sun's Center; P, the mean periodical Time of the *Apogee* or *Node*; S, the mean *synodical Time* of the Sun's Revolution to the *Apogee* or *Node*: Then will

$\frac{3S}{2P} \times AE =$ the annual Equation of the *Apogee* or *Node*, according as S and P are defined.

According to which Rules, the Equation of the Sun's Center to $\left\{ \begin{array}{l} \text{ann. Equat. } \Delta \\ \text{ann. Equat. } \Delta \text{'s Ap.} \\ \text{ann. Equat. } \Delta \text{'s Node} \end{array} \right\}$ will be in a given Proportion, respectively.

The above general Propositions and Proportions, *mutatis mutandis*, serve for any other Satellite, as well as for the Moon.


Mr. Machin shews a Method of correcting the Angle of mean Anomaly at the upper Focus (by an *elliptic Equant*, see P. 82.) so as to give the true Anomaly, according to the Proportions of *Ward's Hypothesis*.

A METHOD of finding the DIFFERENCE between the Angle of a Planet at the upper Focus (according to *Ward's Hypothesis*) and the true Angle thereof from *Aphelion*. According to the late *Abraham De Moivre*, F. R. S. See his *Analytica*.

IF EPD be an *Ellipsis*; C, the Center; S, F, the Foci; DE, OY, the two Axes; and a Planet revolves about the Center of Force in the Focus S, and you make $\angle EFQ$ the mean Anomaly: $\angle$ right Angles :: as the Area PSE :: to the Area DOPE of the *Semi-Ellipsis*.

You may find PFQ = the Difference of the Angles EFP, EFQ, as follows.

Let CD = r, CO = m, SP = x, FP = y, &c. Then, by the Method of Fluxions, $CP = \sqrt{rr + mm - mx}$.



OF MACHIN'S THEORY of the MOON'S MOTION.

THE Angle of the Moon's *Elongation* from the Center, represented by $\angle TP$ is the *Variation* or Reflexion of the Moon; the Properties of which appear from Figure V.

The *Variation-Equation* is the Sine of the double Distance of the Moon from the Quadrature or Conjunction with the Sun; it being the Difference of the two Angles $\angle TA$ and $\angle PTA$, next the Sun, whose Tangents, by Construction of the Figure, are in a given Proportion.

The *Variation-Equation* is, all other Circumstances being alike, in the duplicate Proportion of the synodical Time of the Moon's Revolution to the Sun. For the *Variation* is in Proportion to the mean Diameter of the *Epicycle*, in the duplicate Proportion of the synodical Time of Revolution.

The GREATEST VARIATION is an Angle whose Sine is to Radius as the Difference of the greatest and least Distances TM and TN next the Sun, that is $3AM$ to their Sum.

According to which Proportion of the Lines described, this Rule makes the *Elongation* near 29 Minutes, which would be the *Variation*, supposing the Moon performed its Revolution to the Sun in the Time of its Revolution about the Earth. But this *Elongation* of 29' being increased in the duplicate Proportion of the synodical to the periodical Time of Revolution, it will give nearly 34' of *Variation*.

What is said of the *Epicycle* is upon Supposition, that the Earth's Orbit round the Sun is a Circle. But if the Eccentricity of the elliptic annual Orbit be considered, the mean Diameters of the *Epicycle* must increase or decrease in the reciprocal triplicate Proportion of the Sun's Distance.

Mr. Machin next shews a general Method of finding the *Inequality* by an *Equant*, in any Revolution, when the hourly Motion, or Velocity of the revolving Body varies, in Respect of its Aspect to some other revolving Body.

That, in any Revolution, the mean Motion and *Inequality* are to be assigned by Means of a curvilinear Figure, wherein equal Areas are described about the Center in equal Times; the Property of which Figure is, that the Rays from the Center, are always in the reciprocal subduplicate Proportion of the hourly Velocity about the Center.

He gives an Instance, in a Figure called the elliptic *Equant*, of the Sun's periodical Revolution to the Node, to the periodical Time of the Sun's Revolution, as the Area of the elliptic *Equant* to the Area of the circumscribing Circle.

That, if a Circle be described, about the Center of the curvilinear Figure, and equal to its Area, it will cut the Figure in the Place where the Sun has the mean Motion from the Node.

That, if any circular Sector be made, which shall contain an Area, equal to the Sector in the Figure, drawn through a Perpendicular's Intersection of the Fig. let fall from the Extreme of the Arch of the circular Sector, on the Diameter, the Angle of the circular Sector will be the Angle of mean Motion of the Sun from the Node.

That the Difference between the Angle of the circular Sector, representing the Sun's mean Motion from the Node, and the Angle of the Figure's Sector, representing the true Motion of the Sun from the Node, will be the Equation of the Sun's Motion from the Node.

That by the same Rule, an Equation may be determined of the Moon from the Sun; or Moon from the Apogee or Node; by a curvilinear Figure described about the Center; the Relation of which Areas shall represent the mean and true Motions; whence the *Inequality* or Equation of these Motions will be shewn.

That, as in every Revolution, there is a certain Figure proper to shew this *Inequality* of Motion, such a Figure may therefore be properly called the *Equant* for that Motion or Revolution.

That, in every Revolution, where the *Equant* has the same Property, the *Inequalities* or Equations will vary according to the same Rule.

EXAMPLE. If the *Equant* be an Ellipsis about the Center, as in the Figure of the Motion of the Sun from the Node.

1. The mean Motion in the whole Revolution will be a geometrical mean Proportional between the greatest Motion, in the Extremity of the shorter Axis, and the least Motion, in the Extremity of the longer Axis; because the Radius of the Circle, which is equal to an Ellipsis, is a mean Proportional between the two Semi-axes.

2. That the Tangents of the Angles of the mean and true Motion are in the given Proportion of the two Axes of the Ellipses; being as the respective Ordinates to those Motions.

3. The Sine of the Angle of the greatest Inequality in the *Orbits*, is to the Radius as Half the Sum of the Axes to Half their Difference.

That the *Equant* is an Ellipsis about the Center, in every Motion where the Excess of Velocity about the Center, above the least Velocity, is always in the duplicate Proportion of the Sine of the Angle of the true Motion from where the Velocity about the Center is least.

Whence it follows that these Motions are represented by an elliptic *Equant*, viz. The monthly Motion of the Moon from the Node; the annual Motion of the Sun from the Node, and of the Moon from the Sun; accelerated or retarded, according to the Change of the Area described about the Earth; (as in Newton's 26 Prop. III. Book.) Also the annual Motion of the Sun from the Moon's Apogee.

Thus, he instances, in the Motion of the Nodes, that the Node is in its swiftest retrograde Motion, when the Sun and Moon are in Conjunction or Opposition, and in Quadrature with the Line of Nodes.

According to Sir Isaac Newton's Method, (at the End of the 13 Prop. of the III. Book) the Force of the Sun to produce a Motion in the Node, at this Time, is equal to 3 Times the mean solar Force; that is, by Construction of the elliptic *Epicycle*, in Fig. V. equal to a Force which is to the Force of Gravity, as $3AM$ to AT , or 3 Times the shorter Semi-axis of the elliptic *Epicycle* to the Distance of the Center thereof from the Center of the Earth.

But, if the Moon revolves in an elliptic *Epicycle*, as before described, the Force, to make a Motion in the Node mentioned, will be to the Force of Gravity at the Time, as $3BN$ to BT , or thrice the longer Semi-axis of the elliptic *Epicycle*, to the shorter Semi-axis of the Moon's Orbit; being double the former Force. But, according to Sir I. NEWTON's METHOD, the Motion of the Node, at this Time, is to the Moon's Motion, as the solar Force to create a Motion in the Node is to the Force of Gravity. And, if the Moon be conceived revolving in a Circle, with the Velocity of its Motion from the Node, at this Time, when the Node is swiftest, and the Plane of the said Circle be supposed to have a Rotation upon an Axis perpendicular to the Plane of the *Ecliptic*, and the contrary Way to the Moon's Motion, so as to produce the Motion of the Node; leaving the Moon to move with its own Motion about the Earth; the Force to make a Motion in the Node seems to be the Difference of the former Forces to retain it with the Velocity of its Motion in the moveable and immoveable Planes. But the Velocity of Bodies, revolving in Circles, are in the subduplicate Proportion of the central Forces.

The Motion of the Moon from the Node, at this Time, when the Node moves swiftest, is to the Motion of the Moon, in the subduplicate Proportion of the Sum of the Forces to the Force of Gravity, or as the Sum of TB and $3BN$ to TB ; that is, as the Sum of the shorter Semi-axis of the Moon's Orb and 4 Times the longer Semi-axis of the elliptic *Epicycle*, to the Sum of the said shorter Semi-axis and once the longer Semi-axis of the said *Epicycle*.

This would be the greatest Motion of the Node, provided that the Plane of the Moon's Orbit was nearly coincident with the Plane of the *Ecliptic*; but the Inclination thereof being considered, the motive Force for the Node must be diminished in the Proportion of the Cosine of Inclination to Radius.

The Distance TB , before-mentioned, being equal to 8400, and $3BN$ being 282, the Inclination of the Plane, in this Position, is $4^\circ 59' 35''$; the Cosine of which to Radius is as 525 to 527 nearly; therefore the Force of Gravity is to the motive Force of the Node, thus diminished, in the compound Proportion of 8400 to 283 and 527 to 525; being nearly as 4216 to 141. So that the greatest Motion of the Moon from the Node, is to the Motion of the Moon, in the subduplicate Proportion of 4357 to 4216, or nearly as 613 to 603. According to which Computation, the greatest hourly Motion of the Node comes out $32'' 47'''$; but, by Sir Isaac Newton's Method, it amounts to $33'' 10''' \frac{1}{2}$.

This is the swiftest retrograde Motion of the Node when the Line of the Nodes is in Quadrature with the Sun, and the Moon is in the greatest Latitude, in Conjunction or Opposition to the Sun,

OF MACHIN'S THEORY of the MOON'S MOTION.

But the *Equant* for the Motion of the Moon from the Node in this Month, when the *Line* of the Nodes is in the *Quadrature* with the Sun, is an *Ellipsis* about the Center; and therefore the mean Motion, in this Month, will be found by the following RULE.

The mean Motion of the Moon from the Node, in that Month when the *Line* of the Nodes is in a *Quadrature* with the Sun, is a geometrical mean Proportional between the greatest Motion of the Moon from the Node, and the Motion of the Moon.

This Motion will be to the mean Motion of the Moon, in the subduplicate Proportion of 613 to 603, nearly as 1221 to 1211. That the Motion of the Node, in this Month, will be to the Motion of the Moon, as 10 to 1211; making the mean hourly Motion $16'' 19''' \frac{1}{4}$. But, according to Sir I. Newton, it amounts to $16'' 35'''$; yet by Corrections that he pursues, it reduces to $16'' 16''' \frac{2}{3}$.

The EQUANT for the annual Motion of the Sun from the Node being also an *Ellipsis*, it follows that

The mean Motion of the Sun from the Node is a geometrical mean Proportional, between the Motion of the Sun and the mean Motion of the Sun from the Node, in the Month when the *Line* of the Nodes is in a *Quadrature* with the Sun.

The mean Motion of the Node, when the *Line* of Nodes is in *Quadrature* with the Sun, is already shewn to be, to the Moon's mean Motion, as 10 to 1211; and the Motion of the Sun is to the Motion of the Moon, as 160 to 2139; whence, the Motion of the Node and Motion of the Sun will be as 1549 to 1395. Therefore, by the said Rule, the Sun's mean Motion from the Node, is to the Sun's mean Motion, in the subduplicate Proportion of 1549 to 1395, being nearly as 98 to 93, corresponding with Observation; there being 98 Revolutions of the Sun to the Node in 93 Revolutions of the Sun. — The subduplicate Proportion taken more nearly, is 941 to 893, which will give $19^\circ 21' 3''$, for the Motion of the Node from the fixed Stars, in a syderal Year. The Motion observed is $19^\circ 21' 22''$.

If the Computation from the Rule had been more exactly made, the annual Motion produced had been $19^\circ 21' 7'' \frac{1}{2}$, which is $14''$ less than the Motion observed by Astronomers.

The foregoing Difference of Motion, probably, may arise from the Sun's *Parallax*; and if so, may furnish the best and most certain Method of determining the true Distance of the Sun. For the Sun's Force being something more on that Half of the Moon's Orb which is towards the Sun, than it is on the other Half, the *elliptic Epicycle* must be accordingly larger in the first Case than in the latter. And by Computation, the mean Motion of the Node arising (after Consideration is taken of this Difference) is more than the mean Motion from the mean Magnitude of the *Epicycle*, by near $2''$ in a Year, for every Minute of the *parallactic Angle* of the Moon's Orbit. And, by the best Computation, this Difference of $14''$, in the annual Motion of the Node, will arise from about $8''$ of *Parallax*, making the Sun's Distance above 25000 Semidiameters of the Earth.

As the *Equant*, for the Motion of the Node, in that Month when the *Line* of the Nodes is in *Quadrature* with the Sun, is an *Ellipsis*; so, in any other Month, it is also an *Ellipsis*: the Motion of the Node being direct and retrograde by Turns, in the Moon's passing from the *Quadrature* with the Sun to the Place of its Node, and from thence to its *Quadrature* aforesaid.

The Inclination of the Plane of the Moon's Orbit to the Plane of the *Ecliptic* is also shewn by these *elliptic Equants*, (see P. 25. *Laws of the Moon's Motion*, at the End of the *Principia*, translated by Motte.)

The Motion of the Moon from the Sun, accelerated and retarded by the Increment of the Area described by the Moon about the Earth (according to 26. Prop. III. Book, *Principia*) is also reduced to an *elliptic Equant*, by taking the shorter to the longer Axis, in the subduplicate or 4th Root, Proportion of the Force of the Moon's Gravity to the Earth to the said Force added to 3 Times the mean solar Force; that is, as TA (in Fig. V.) is the first 3 mean Proportionals between TA and TA + 3AM.

In this Proportion is the Area described by the Moon about the Earth, when in *Quadrature* with the Sun, to the mean Area: Or, as the mean Area to the Area described, in the *Syzygies*. Hence, the

greatest Area in the *Syzygies* is to the least in the *Quadratures*, in the subduplicate Proportion of TA + 3AM to TA, or as $\sqrt{8541}$ to $\sqrt{8400}$.

But this is on *Supposition*, that the Moon revolves to the Sun in the same Time that it revolves about the Earth; which nearly agrees with Sir I. Newton's Computation in the before-cited Proposition 26. B. III.

In like Manner, an *elliptic Equant* may be constructed to shew the mean Motion of the Moon's Apogee, according to the Rules given by Sir Isaac Newton (in *Corollaries* of 45. Prop. 1. Book) for stating the Motion of the Apogee; viz. by taking the greatest retrograde Motion of the Apogee from the Force of the Sun upon the Moon in the *Quadratures*; and the greatest direct Motion, from the Force of the Sun upon the Moon when in *Conjunction* and *Opposition*; each, according to his Rule delivered in the 2d *Corollary* to the said Proposition. And, if an *Ellipsis* be made, whose Axes are in a subduplicate Proportion of the Moon's Motion from the Apogee, when in the said swiftest direct and retrograde Motions, the said *Ellipsis* will be nearly the *Equant* for the Motion of the Moon from the Apogee, and be nearly in the Form of that above, for the Increment of the Area. But the Motion of the Apogee, by this Method, will be found to be no more than $10^\circ 37' 22''$, in the Revolution of the Moon from *Apsis* to *Apsis*, which should be $3^\circ 4' 7'' \frac{1}{2}$, according to Observation. — By which, there is more Force necessary to account for the Motion of the Moon's Apogee than what arises from the Variation of the Moon's Gravity to the Sun, in its Revolution about the Earth.

But, if the Cause of this Motion be supposed to arise from the Variation of the Moon's Gravity to the Earth, as it revolves round, in the *elliptic Epicycle*, this Difference of Force, nearly the double of the former, will be sufficient to account for the Motion; but not so exactly as might be expected.

Mr. Machin proceeds to lay down a Rule, (founded on new Principles) which is, that The mean Motion of the Moon from its Apogee, is to the mean Motion of the Moon, in the subduplicate Proportion of the Dist. between twice the mean Semidiameters of the *Epicycle*, and the Moon's mean Dist. to her mean Dist. See Fig. V; where AM = 47, AN = 94, and AT = 8400.

EXAMPLE. Half the shorter Axis of the *Epicycle*, is $23 \frac{1}{2}$, deducted from AT (= 8400) = mean Distance from the Center of the Moon's Orb, 8376 $\frac{1}{2}$; the Sum of the Semiaxes of the *Epicycle*, 141; the Difference of the mean Distance 8376 $\frac{1}{2}$, and Sum of the said Semiaxes, of the *Epicycle*, = 8235 $\frac{1}{2}$. Wherefore the Motion of the Moon from the Apogee is to the Motion of the Moon, as the Subduplicate of 8235 $\frac{1}{2}$ to 8376 $\frac{1}{2}$, as 1877 to 1893; so that there ought to be 16 Revolutions of the Apogee in 1893 Revolutions of the Moon; agreeing, to great Precision, with the modern Numbers observed. Whence the mean Motion of the Apogee, in a syderal Year, will be $40^\circ 40' 40'' \frac{1}{2}$; nearly agreeing with the Numbers from Newton's Theory, $40^\circ 40' 43''$, as before observed in the Comment.

The mean Motion of the Apogee being stated, according to the physical Rules aforesaid, the following Rule serves for determining the Variation of Eccentricity of the Moon's Orbit.

RULE. The least is to the mean Eccentricity in the duplicate Proportion of the Sun's mean Motion from the Apogee of the Moon's Orbit, to the Sun's mean Motion. Or in the duplicate Proportion of the periodical Time of the Sun's Revolution to the mean periodical Time of its Revolution to the Moon's Apogee.

By the foregoing Rule for the mean Motion of the Apogee, there are 16 Revolutions of the Apogee in 1893 Revolutions of the Moon; but there being 254 Revolutions of the Moon in 19 Revolutions of the Sun; there must be about 7 Revolutions of the Apogee in about 62 Revolutions of the Sun; or rather about 20 in 177.

That the Period of the Sun to the Stars, and of the Sun to the Moon's Apogee, are, nearly, as 157 to 177; the Duplicate of which Proportion is, that of 107 to 136; which, according to the Rule, should be the Proportion of the least to the mean Eccentricity.

Hence,

OF MACHIN'S THEORY of the MOON'S MOTION.

Hence, by this Rule, the mean Eccentricity (or Half the Sum of the greatest and least) ought to be the Difference of the mean from the least, (or Half the Difference of greatest and least) as 136 to 29.

The mean Eccentricity according to Flamsteed or Horrox, is .055236; to mean Distance of Sun 1; Half Difference between greatest and least .011617; being nearly as $135\frac{1}{2}$ to $28\frac{1}{2}$.

According to Newton, the mean Eccentricity .05505 to 1, the mean Distance or Semi-Transverse of the Earth's Orbit; Half the Difference of greatest and least .01173; nearly as $135\frac{1}{2}$ to $28\frac{1}{2}$, very nearly as above.

The Rule here laid down is only to be used, for determining the variable Eccentricity, when the Eccentricity of the Orbit is small.

The greatest and least Eccentricity being known, the Equant for the Motion of the Sun from the Apogee will be an Ellipsis; whose greater and less Axis are equal to the greatest and least Eccentricities. And therefore by the Property of such an Equant,

The Sine of the greatest Equation of the Apogee will be to Radius, as the Difference of the Axes of the Equant is their Sum; that is, as the Difference of the greatest and least Eccentricities to their Sum.

EXAMPLE. The Difference to the Sum (before determined) as 29 to 136 :: Radius : $120^{\circ} 18' 40''$, as before determined in the Comment.

Hence also, Mr. Machin gives this Rule, for determining the Equation of the Apogee in any Aspect of the Sun to the Apogee.

The Tangent of the Angle of the mean Distance of the Sun from the Moon's Apogee is to the Tangent of the Angle of the true Distance of the Sun from the said Apogee, as the greatest is to the least Eccentricity, or as 165 to 107, nearly.

Mr. Machin then proceeds to shew that a Curve may be described about a Center, to represent the Inequality of any Motion proposed, whose Rays are reciprocally in the subduplicate Proportion of the Velocity about the Center. That, where the Inequality about the Center is but small, there is no Need of great Exactness in the Quadrature of the Curve for shewing what the Equation is. Whence all the small annual Equations (he shews) of the Moon's Motion, arising from the different Distances of the Sun, at different Times of the Year, are reduced to one Rule, near enough for the Purpose.

That since the Sun's Force to produce these annual Inequalities, is reciprocally in the triplicate Proportion of his Distances, the Rays of the Equant, for such a Motion, will be in the sesquuplicate Proportion of that Distance, &c. as before described; who determineth the small Equations of the Moon, Apogee, and Node, accordingly, as has been noted.

He proceeds to an Example of the Use of the Equant, in finding the Equations of the Moon's Center, by a complicated Figure, (see the End of Newton's Principia, translated by Motte) but as we have already given a Sketch of this Rule in P. 82. we need not repeat it here.

— What we have introduced is to give the Reader the completer Idea of physical Astronomy; which Sir Isaac Newton first promoted; and has since been improved by several mathematical Astronomers, De la Caille, &c.

Mr. T. S. F. R. S. the celebrated Author of the miscellaneous Tracts, distinguished for his eminent Abilities, has given a Correction of Ward's Hypothesis, (at P. 53 of the said Tracts) by Means of one, two, three, or more Equations, representing that Mr. Machin has given one of his three Equations defective; injudiciously asserting, that the Inequality in the Motion of a Planet about the upper Focus consists of three Parts; "As if (says the Tract Author) the Nature of the Subject admitted of just that Number, and no more; whereas the Parts or Equations arising in the Consideration, are without Number; being the Terms of an infinite Series, wherein the Eccentricity is the converging Quantity. Sir Isaac Newton has given two Terms of this Series, which are right; but the new Equation added by the Commentator (Mr. Machin) is not so; the Sign thereof, the Coefficient, and the Law by which it increases and decreases, being all different from what they ought to be."

Moreover, (in P. 54 of the said Tracts) the Tract Author has criticized the same Commentator for giving the Ratio of the Diameters of the Moon's Orbit (setting aside the Consideration of Eccentricity) as 59 to 60; Sir Isaac Newton having otherwise physically proved it to be as 69 to 70; with which the Tract Author's physical Computation wonderfully agrees.

OF DE LA CAILLE'S THEORY, and PHYSICAL RULES, of the MOON'S MOTION.
(See this celebrated AUTHOR'S ASTRONOMY for the DEMONSTRATION.)

I. THE different Inequalities in the Motion of the Moon are produced by the different Modifications its central Force to the Sun occasions in its central Force to the Earth.

II. The Moon's perturbate Force is always the third Side of a right lined Triangle, one Side whereof is the Radius of the Moon's Orbit, and the other Side is the triple of the Cosine of the Moon's Distance to the nearest Syzygy.

III. Therefore the Force accelerating or retarding the Moon's Motion in its Orbit, is always as $\frac{3}{2}$ of the Sine of twice the Moon's Distance from the Line of the Syzygies.

IV. And the Force diminishing or increasing the Moon's central Force or Gravity, respectively to the Earth, is as the Sum, or as the Difference of Half the Radius, and of $\frac{3}{2}$ of the Cosine of twice the Moon's Distance from the Line of the Syzygies.

V. All the Inequalities happening to the Moon's Motion, in its Passage from one Syzygy to its next Quadrature, are gradually restored in the Moon's Passage from this Quadrature to the next Syzygy; afterwards the like Inequalities begin again in the same Order from this Syzygy to the next Quadrature, and are again restored in the Passage to the following Syzygy, and so on.

VI. When the Moon is in the Syzygies, its perturbate Force diminishes the central Force, respectively to the Earth, and increases it when in the Quadratures: But the Diminution in the Syzygies, is double the Augmentation in the Quadratures.

VII. There are four Points in the Moon's Orbit, each about $54^{\circ} 44'$ from either Side of the Syzygies, where the Gravitation of the Moon to the Sun makes no Alteration in its central Force in respect to the Earth.

VIII. The Variation of the Moon's central Force respectively to the Earth, tends to flatten its Orbit towards the Syzygies, and to lengthen it towards the Quadratures; so that the Moon's Orbit, at first circular, is now become oval or elliptic; its greater Axis being in the Line of Quadratures, and less Axis in that of the Syzygies, and the Earth in one of the Foci.

IX. The Velocity of the Moon must decrease from the Syzygies to the Quadratures; and accelerate from the Quadratures to the Syzygies.

X. The Quantity whereby the Moon's Velocity is diminished in going from the Syzygy to the Quadrature, or accelerated from the Quadrature to the Syzygy, must increase from the Syzygy to the next Octant; and decrease in the same Proportion, from that Octant to the following Quadrature: It being always in the Ratio of the Sine of twice the Moon's Distance from the Syzygy. This Inequality is called the Moon's Variation.

OF DE LA CAILLE's THEORY, and PHYSICAL RULES, of the MOON's MOTION.

XI. In the Syzygies and Quadratures, a Ray from the Moon to the Earth will describe Areas in Proportion to the Time; in every other Point of the Orbit, the Areas are less in Proportion to the Times, as the Moon is nearer its Ostants.

The Moon's Orbit has been supposed a Circle; though it is in Fact an Ellipsis; in one of whose Foci the Earth is placed. Whence the explained Inequalities may cause the Ellipsis to be more convex in the Quadratures, and more rectilinear in the Syzygies. Which Inequalities must contribute to make the Moon recede from the Earth in the Quadratures, and approach it in the Syzygies; though the Moon's elliptic Orbit is not very eccentric. And since the Moon's central Force diminishes in the Syzygies, and increases in the Quadrature, it is evident (from what is demonstrated in De la Caille's Astronomy) it must sometimes be too great, and sometimes too little, to be inversely as the Square of the Moon's Distance from the Earth.

The Difference in the Moon's Distances, the Variation of the Eccentricities of the Earth and Moon, the Inclination of the Plane of the Moon's Orbit to that of the Ecliptic, constitute, and are the Cause of, five very considerable periodical Inequalities in the Moon's Motion.

1. An Inequality in the Time of several of the Moon's successive periodic Revolutions.

2. The Motion of the Line of the Apfides.

3. A continual Change in the Eccentricity of the Moon's Orbit.

4. The Retrogradation of the Moon's Nodes.

5. The Change in the Inclination of the Moon's Orbit.

XII. The Times of the Moon's periodic Revolutions are longer when the Earth is in its Perihelion than when in its Aphelion.

XIII. When the Moon is in the Syzygies, the Line of its Apfides advances directly, according to the Order of Signs; but moves retrograde when the Moon is in the Quadratures: It advances with the greatest Velocity, when concurring with the Line of Syzygies; and retrogrades with the greatest Velocity when concurring with the Line of Quadratures; the Sum of the Quantities it gains in several Revolutions, exceeding the Quantities it retrogrades during these Revolutions, causes an entire Revolution of the Line of Apfides, according to the Order of Signs, in about 19 Years.

XIV. The Eccentricity of the Moon's Orbit varying every Instant, is greatest when the Line of Apfides coincides with that of the Syzygies, and least when the Line of Apfides coincides with that of the Quadratures.

XV. When the Moon is in the Quadratures, the Nodes of its Orbit are always stationary; but commonly retrograde when the Moon is in the Syzygies. In the Interval from one Quadrature to the other, they move the most retrograde when the Line of the Nodes concurs with that of the Quadratures; and least, when it concurs with that of the Syzygies.

XVI. In every Revolution of the Moon, the Angle of Inclination of its Orbit to the Plane of the Ecliptic, increases in the Interval from a Syzygy to the next Quadrature; and decreases in the Interval from the Quadrature to the following Syzygy. It is greatest in the Quadrature when the Line of the Nodes concurs with that of the Syzygies; and least, in the Syzygy when the Line of the Nodes concurs with that of the Quadratures.

XVII. The Increase of the central Force, in the Quadratures, is to the Force acting upon the Moon's Plane, as the Cube of Radius, is to the triple Sine of the Moon's Distance from the Quadrature, multiplied by the Sine of the Nodes Distance from the Syzygy, and by the Sine of the Inclination of the Moon's Orbit.

XVIII. In the Passage from the Syzygy to the following Quadrature the Moon's Node moves retrograde, and the Inclination of the Orbit increases.

XIX. The Moon's Node moves retrograde, and the Inclination of the Orbit decreases, in the Passage from the Quadrature to the following Syzygy.

XX. In general, The Moon's Node moves according to the Order of Signs, and is stationary only when the Moon is in Quadrature, or when without Latitude. In all other Cases, it moves retrograde with a Velocity increased in the Proportion to the Moon's Proximity to the Syzygy, and as its Latitude is greater.

XXI. The Inclination of the Moon's Orbit, evidently changes four Times in every Revolution, twice increasing and twice diminishing. It is greatest when the Line of Nodes concurs with that of the Quadratures; and least when the Line of the Nodes coincides with that of the Syzygies.

XXII. The Moon's perturbate Force, and consequently all its Inequalities, are somewhat greater towards the Conjunctions than towards the Oppositions.

XXIII. In the Quadratures, the Increase of the Moon's Gravity to the Earth is always as the Moon's Distance from the Earth divided by the Cube of the Earth's Distance from the Sun. Or (the Distances being inversely as the apparent Diameters) as the Cube of the Sun's Diameter seen from the Earth, divided by the Moon's Diameter seen from the Earth.

XXIV. What holds true in the Moon's Inequalities is likewise applicable to the Satellites of Jupiter and Saturn. But because of the great Distance of those Planets from the Sun, the Inequalities of their Satellites must be very small; and, are sensible only in those nearest the Planet, on Account of their Proximity.

XXV. The Combination of the Moon's Gravity towards the Earth and the Sun, must evidently produce the same Effects, whether the Earth be supposed to describe an Ellipsis about the Sun, fixed in one of its Foci, or whether the Sun describes an Ellipsis about the Earth, fixed in one of its Foci: For these Effects depend upon the Moon's Position and Distance in respect of the Earth and Sun; and therefore whether the Earth or the Sun moves, their Situations will be the same to one another; and consequently to the Moon.

REMARK. To settle the Moon's Theory, and construct Tables for computing her true Longitude and Latitude, as seen from the Earth, it will be necessary to determine, by astronomical Observation, the Nineteen Elements, following.

1. The Moon's mean periodical Time.
2. The mean periodical Time of the Moon's Apogee.
3. The mean periodical Time of the ascending Node.
4. An Epocha of the Moon's mean Place.
5. An Epocha of the mean Place of the Apogee.
6. An Epocha of the mean Place of the Nodes.

These Epochas on the Noon of 31st December, 1745, N. S. mean Time; were; for the Moon $0^{\circ} 17' 33'' 30''$; for the Apogee $0^{\circ} 8' 12' 49''$; for the Node $11^{\circ} 27' 37' 0''$.

7. The greatest Equation of the Moon's mean Place.
8. That of the mean Place of the Moon's Apogee.
9. That of the mean Place of the Moon's Node.

These three Inequalities serve to compensate the Inequalities of the periodical Revolutions, according as the Earth is in its Perihelion or Aphelion; and are proportional to the Equation of the Sun's Center. The first, for the Moon, is $11' 50''$; for the Apogee $20' 0''$; and for the Node $9' 30''$.

10. The greatest Equation of the Moon's Apogee, which is $12' 18''$.
11. The mean Eccentricity of the Moon's Orbit; which is 5505 such Parts as the Moon's mean Dist. from the Earth contains 100000.
12. The Relation between the greatest and least Eccentricities; being as 6677 to 4333.

By these Relations the Equation of the Moon's Center is found in all Cases.

13. The Moon's mean Variation in Ostants, being $35' 10''$.
14. The Relation between the greatest and least Variation; being from $37' 25''$ to $33' 4''$.
15. The greatest second Equat. of Moon's Node; being $1^{\circ} 29' 40''$.
16. The mean Inclination of the Moon's Orbit; being about $5^{\circ} 8' 30''$.
17. The Proportion of the greatest to the least Inclination being as $5^{\circ} 17' 30''$ to $4^{\circ} 59' 30''$.
18. The mean Parallax of the Moon, being $57' 5''$.
19. The Ratio of the Parallax to the Moon's Diameter, being as $55' 40''$ to $30'$, nearly.

Besides which Elements there are others giving small Equations. —

We refer the Reader to our Tables and Comment for the Explanation and Use of all these Elements.

MEAN TIME of the FIRST MEAN NEW MOON, in JANUARY, from 1700 to 1800, O. S. With the Places of the Sun's Mean Anomaly, Annual Argument, and Sun from the Moon's Node, corresponding.

Years of Christ	First M. New Moon in Jan.	Sun's Mean Anomaly.	Sun from Moon's Apo.	Sun from ☉	Years of Christ	First M. New Moon in Jan.	Sun's Mean Anomaly.	Sun from Moon's Apo.	Sun from ☉
d h m f	s o ' "	s o ' "	s o ' "	s o ' "	d h m f	s o ' "	s o ' "	s o ' "	s o ' "
1700	9 14 42 44	6 21 43 53	0 0 52 40	4 13 10 32	1752	4 18 47 25	6 16 28 39	1 11 4 20	1 24 19 53
1	27 12 15 23	7 10 6 5	11 6 29 47	4 21 53 35	53	22 16 20 4	7 4 50 50	0 16 41 26	3 3 2 55
2	16 21 4 0	6 29 21 56	9 16 17 52	5 29 56 22	54	12 1 8 41	6 24 6 41	10 26 29 31	3 11 5 42
3	6 5 52 37	6 18 37 48	7 26 5 57	6 7 59 9	55	1 9 57 17	6 13 22 32	9 6 17 36	3 19 8 29
1704	25 3 25 16	7 6 59 59	7 1 43 3	7 16 42 30	1756	20 7 29 57	7 1 44 44	8 11 54 42	4 27 51 30
5	13 12 13 53	6 26 15 50	5 11 31 8	7 24 44 57	57	8 16 18 34	6 21 0 36	6 21 42 48	3 5 54 18
6	2 21 2 30	6 15 31 4	3 21 19 13	8 2 47 44	58	27 13 51 14	7 9 22 47	5 27 19 54	6 14 37 19
7	21 18 35 9	7 3 53 52	2 26 56 19	9 11 30 45	59	16 22 39 50	6 28 38 38	4 7 7 59	6 22 40 6
1708	11 3 23 46	6 23 9 43	1 6 44 24	9 19 33 32	1760	6 7 28 27	6 17 54 29	2 16 56 4	7 0 42 53
9	29 0 56 26	7 11 31 55	0 12 21 30	10 28 16 34	61	24 5 1 6	7 6 16 40	1 22 33 10	8 9 25 54
10	18 9 45 2	7 0 47 46	10 22 9 35	11 6 19 21	62	13 13 49 43	6 25 32 32	0 2 21 14	8 17 28 41
11	7 18 33 39	6 20 3 38	9 1 57 40	11 14 22 9	63	2 22 38 20	6 14 48 23	10 12 9 19	8 25 31 28
1712	26 6 6 18	7 8 25 48	8 7 34 46	0 23 5 10	1764	21 20 10 59	7 3 10 34	9 17 46 25	10 4 14 29
13	15 0 54 55	6 27 41 40	6 17 22 51	1 1 7 56	65	10 4 59 36	6 22 26 26	7 27 34 31	10 12 17 17
14	4 9 43 32	6 16 57 30	4 27 10 56	1 9 10 44	66	29 2 34 15	7 10 48 37	7 3 11 37	11 21 0 19
15	23 7 16 11	7 5 19 43	4 2 48 2	2 17 53 44	67	18 11 20 52	7 0 4 28	5 12 59 42	11 29 3 6
1716	12 16 4 48	6 24 35 33	2 12 36 7	2 25 56 32	1768	7 20 9 29	6 19 20 20	3 22 47 47	0 7 5 53
17	1 0 53 30	6 13 51 26	0 22 24 12	3 3 59 20	69	25 17 42 8	7 7 42 30	2 28 24 53	1 15 48 54
18	19 22 26 4	7 2 13 37	11 28 1 18	4 12 42 20	70	15 2 30 45	6 26 58 21	1 8 12 58	1 23 51 41
19	9 7 14 41	6 21 29 28	10 7 49 23	4 20 45 8	71	4 11 19 21	6 16 14 12	11 18 1 3	2 1 54 28
1720	28 4 47 20	7 9 51 39	9 13 26 29	5 29 28 9	1772	23 8 52 1	7 4 36 23	10 23 38 9	3 10 37 29
21	16 13 35 57	6 29 7 31	7 23 14 34	6 7 30 57	73	11 17 40 38	6 23 52 15	9 3 26 14	3 18 40 17
22	5 22 24 33	6 18 23 22	6 3 2 39	6 15 33 43	74	1 2 29 14	6 13 8 8	7 13 14 19	3 26 43 5
23	24 19 57 13	7 6 45 33	5 8 39 45	7 24 16 44	75	20 0 1 54	7 1 30 19	6 18 51 25	5 5 26 6
1724	14 4 45 50	6 26 1 24	3 18 27 50	8 2 19 31	1776	9 8 50 31	6 20 46 10	4 28 39 30	5 13 28 53
25	2 13 34 26	6 15 7 15	1 28 15 55	8 10 22 20	77	27 6 23 10	7 9 8 20	4 4 16 36	6 22 11 54
26	21 11 7 6	7 3 39 26	1 3 53 1	9 19 5 20	78	16 15 11 47	6 28 24 11	2 14 4 41	7 0 14 41
27	10 19 55 43	6 22 55 19	11 13 41 6	9 27 8 7	79	6 0 0 23	6 17 40 2	0 23 52 46	7 8 17 28
1728	0 4 44 19	6 12 11 10	9 23 29 12	10 5 10 54	1780	24 21 33 3	7 6 2 14	11 29 29 52	8 17 0 29
29	18 2 16 59	7 0 33 21	8 29 6 17	11 13 53 56	81	13 6 21 40	6 25 18 6	10 9 17 57	8 25 3 16
30	7 11 5 35	6 19 49 21	7 8 54 22	11 21 56 42	82	2 15 10 16	6 14 33 56	8 19 6 1	9 3 6 3
31	26 8 38 15	7 8 11 23	6 14 31 28	1 0 39 44	83	21 12 42 56	7 2 56 7	7 24 43 7	10 11 49 4
1732	45 17 26 52	6 27 27 13	4 24 19 33	1 8 42 31	1784	10 21 31 32	6 22 11 58	6 4 31 12	10 19 51 51
33	4 2 15 28	6 16 43 5	3 4 7 38	1 16 45 19	85	28 19 4 12	7 10 34 10	5 10 8 18	11 28 34 53
34	22 23 48 8	7 5 5 17	2 9 44 44	1 25 28 20	86	18 3 52 49	6 29 50 2	3 19 56 23	0 6 37 40
35	12 8 36 44	6 24 21 8	0 19 32 49	3 3 31 7	87	7 12 41 25	6 19 5 53	1 29 44 29	0 14 40 27
1736	1 17 25 21	6 13 36 59	10 29 20 54	3 11 33 54	1788	26 10 14 5	7 7 28 4	1 5 21 34	1 23 23 29
37	19 14 58 1	7 1 59 10	10 4 58 0	4 20 16 56	89	14 19 2 42	6 26 43 55	11 15 9 40	2 1 26 16
38	8 23 46 37	6 21 15 1	8 14 46 5	4 28 19 43	90	4 3 51 18	6 15 59 46	9 24 57 45	2 9 29 3
39	27 21 19 17	7 9 37 12	7 20 23 11	6 7 2 44	91	23 1 23 58	7 4 21 57	9 0 34 50	3 18 12 5
1740	17 6 7 54	6 28 53 4	6 0 11 16	6 15 5 31	1792	12 10 12 34	6 23 37 46	7 10 22 56	3 26 14 52
41	5 14 56 30	6 18 8 56	4 9 59 22	6 23 8 19	93	0 19 1 11	6 12 53 41	5 20 11 1	4 4 17 40
42	24 12 29 10	7 6 31 7	3 15 36 27	8 1 51 20	94	19 16 33 51	7 1 15 52	4 25 48 6	5 13 0 41
43	13 21 17 46	6 25 46 58	1 25 24 32	8 9 54 7	95	9 1 22 27	6 20 31 43	3 5 36 12	5 21 3 28
1744	3 6 6 23	6 15 2 49	0 5 12 37	8 17 56 55	1796	27 22 55 7	7 8 53 54	2 11 13 17	6 29 46 29
45	21 3 39 3	7 3 25 1	11 10 49 43	9 26 39 56	97	16 7 43 43	6 26 9 44	0 21 1 23	7 7 49 16
46	10 12 27 39	6 22 40 52	9 20 37 48	10 4 42 43	98	5 16 32 20	6 17 25 36	11 0 49 28	7 15 52 3
47	29 10 0 19	7 11 3 3	8 26 14 54	11 13 25 44	99	24 14 5 0	7 5 47 47	10 6 26 34	8 24 35 5
1748	18 18 48 55	7 0 18 54	7 6 2 59	11 21 28 31	1800	13 22 53 36	6 25 3 38	8 16 14 39	9 2 37 52
49	7 3 37 32	6 19 34 45	5 15 51 4	11 29 31 18					
50	26 1 10 12	7 7 56 56	4 21 28 10	1 8 14 19					
51	15 9 58 48	6 27 12 48	3 1 16 15	1 16 17 6					

For Convenience of Computation, the Computer must reduce the New and Full Moons of the Old, to the Days of New Style, by adding the Number of Days Difference of Style.

Number of DAYS in the MONTHLY REVOLUTIONS of the MOON with the SUN, from January. With the Motion of the Sun's Mean Anomaly, Annual Argument, and Sun from Moon's Node, corresponding.

No. Days.	Ms.	+ Days from January D.	Motion Sun's M. Anomaly.	Mo. Sun from Moon's Apog.	Motion Sun from ☉	No. Days.	Ms.	+ Days from January D.	Motion Sun's M. Anomaly.	Mo. Sun from Moon's Apog.	Motion Sun from ☉
d h m s	s o ' "	s o ' "	s o ' "	s o ' "	s o ' "	d h m s	s o ' "	s o ' "	s o ' "	s o ' "	s o ' "
29 Jan	29 12 44 3	0 29 6 19	0 25 49 0	1 0 40 14	206 Jul	25 17 8 21	6 23 44 15	6 0 43 3	7 4 41 38		
59 Feb	28 1 28 6	1 28 12 39	1 21 38 1	2 1 20 28	236 Aug	24 5 52 24	7 22 50 34	6 26 32 3	8 5 21 51		
88 Mar	29 14 12 9	2 27 18 58	2 17 27 1	3 2 0 42	265 Sep	22 18 36 27	8 21 56 53	7 22 21 4	9 6 2 5		
118 Apr	28 2 56 12	3 26 25 17	3 13 16 2	4 2 40 56	295 Oct	22 7 20 30	9 21 3 13	8 18 10 4	10 6 42 19		
147 May	27 15 40 15	4 25 31 37	4 9 5 2	5 3 21 10	324 Nov	20 20 4 34	10 20 9 32	9 13 59 5	11 7 22 33		
177 Jun	26 4 24 18	5 24 37 56	5 4 54 3	6 4 1 24	354 Dec	20 8 48 37	11 19 15 51	10 9 48 5	0 8 2 47		
In Leap-Year, take out one Day less, after February, than the tabular Number; but take out the same Motion.						Half Revol.					
						14 18 22 2	0 14 33 10	0 12 54 30	0 15 20 7		

N. B. The above is the only Table wherein a Day's Motion, at the Year's Beginning is not advanced, in Leap-Years; being constructed according to the Old Method of Computation requiring it to be so.

Subtracting for Half a Revolution from the Time and Places in January O. S. will give the Time and Places for the preceding Full Moon; i. e. When the Days on which a New Moon happens exceed 15. Or the above Table is easily reduced to New Style. But the Examination of ancient Eclipses is according to the above Table by Old Style.

R EVOLUTIONS of the MOON to the SUN, next after Julian CENTURIES completed.																	
Synodic Months.	Julian Centuries.	Time of 1st New Moon after Jul. Cent. completed.				Sun's Mean Anomaly.				Sun from Moon's Apogee.				Sun from Node.			
		D	H	M	S	s	o	i	"	s	o	i	"	s	o	i	"
1237	100	4	8	10	52	0	3	19	44	8	15	21	59	4	19	27	17
2474	200	8	16	21	45	0	6	39	29	5	0	43	58	9	8	54	35
3711	300	13	0	32	37	0	9	59	13	1	16	5	57	1	28	21	52
4948	400	17	8	43	29	0	13	18	58	10	1	27	56	6	17	49	10
6185	500	21	16	54	22	0	16	38	42	6	16	49	54	11	7	16	27
7422	600	26	1	5	14	0	19	58	27	3	2	11	53	3	26	43	45
8658	700	0	20	32	3	11	24	11	52	10	21	44	52	7	15	30	48
9895	800	5	4	42	56	11	27	31	37	7	7	6	51	0	4	58	6
11132	900	9	12	53	48	0	0	51	21	3	22	28	50	4	24	25	23
12369	1000	13	21	4	40	0	4	11	6	0	7	50	49	9	13	52	40
13606	1100	18	5	15	33	0	7	30	50	8	23	12	48	2	3	19	58
14843	1200	22	13	26	25	0	10	50	35	5	8	34	47	6	22	47	15
16080	1300	26	21	37	17	0	14	10	19	1	23	56	46	11	12	14	33
17316	1400	1	17	4	6	11	18	23	44	9	13	29	44	3	1	1	36
18553	1500	6	1	14	59	11	21	43	29	5	28	51	43	7	20	28	54
19790	1600	10	9	25	51	11	25	3	13	2	14	13	42	0	9	56	11
21027	1700	14	17	36	44	11	28	22	58	10	29	35	41	4	29	23	28
22264	1800	19	1	47	36	0	1	42	42	7	14	57	40	9	18	50	46
23501	1900	23	9	58	28	0	5	2	27	4	0	19	39	2	8	18	3
24738	2000	27	18	9	21	0	8	22	11	0	15	41	38	6	27	45	21
25974	2100	2	13	36	10	11	12	35	36	8	5	14	35	10	16	32	24
27211	2200	6	21	47	2	11	15	55	20	4	20	36	34	3	5	59	41
28448	2300	11	5	57	54	11	19	15	5	1	5	58	33	7	25	26	59
29685	2400	15	14	8	47	11	22	34	49	9	21	20	32	0	14	54	16
30922	2500	19	22	19	39	11	25	54	34	6	6	42	31	5	4	21	34
32159	2600	24	6	30	31	11	29	14	18	2	22	4	29	9	23	48	51
33396	2700	28	14	41	24	0	2	34	3	11	7	26	28	2	13	16	9
34632	2800	3	10	8	13	11	6	47	28	6	26	59	27	6	2	3	12
35869	2900	7	18	19	5	11	10	7	12	3	12	21	26	10	21	30	29
37106	3000	12	2	29	57	11	13	26	57	11	27	43	25	3	10	57	47
38343	3100	16	10	40	50	11	16	46	41	8	17	5	24	8	0	25	4
39580	3200	20	18	51	42	11	20	6	26	4	28	27	23	9	19	52	22

ECLIPSE-REVOLUTIONS and other PERIODS of the MOON to the SUN.																	
Synodic Months.	Periods in Julian Years and Parts.				Sun's Mean Anomaly.				Sun from Moon's Apogee.				Sun from Node.				
	Julian Years.	D	H	M	S	s	o	i	"	s	o	i	"	s	o	i	"
Pl. Pe. 111	8	355	21	29	38	11	20	41	42	11	15	39	48	5	14	25	47
1. 223	18	11	7	43	20	0	10	29	42	11	27	8	37	11	29	31	48
235	19	0	16	31	56	11	29	45	33	10	6	56	42	0	7	34	35
251	20	107	4	16	45	3	15	26	42	0	0	0	49	4	18	18	18
270	21	303	6	13	43	9	28	16	49	4	10	31	58	0	1	2	42
364	29	157	3	14	30	5	4	21	3	1	7	18	39	6	4	4	32
2. 446	36	21	15	26	40	0	20	59	24	11	24	17	14	11	29	3	35
534	43	64	8	3	8	2	2	15	42	3	15	9	53	5	28	4	2
587	47	168	10	57	50	5	14	50	45	1	4	27	15	0	3	36	19
669	54	32	23	10	0	1	1	29	6	11	21	25	51	11	28	35	23
767	62	4	23	6	59	0	3	48	38	0	1	28	34	4	4	18	8
810	65	178	18	41	10	5	25	20	27	0	1	35	53	0	3	8	7
4. 892	72	43	6	53	20	1	11	58	48	11	18	34	28	11	28	7	10
980	79	85	23	29	49	2	23	15	7	3	10	27	8	5	27	7	36
5. 1115	90	54	14	36	40	1	22	28	30	11	15	43	5	11	27	38	58
1203	97	96	7	13	9	3	3	45	50	3	7	35	43	5	26	39	24
6. 1338	108	65	22	20	0	2	2	58	12	11	12	51	42	11	27	10	45
2148	173	243	17	1	10	7	28	18	40	0	14	27	36	0	0	18	52
2301	186	13	21	20	57	0	11	25	52	0	4	25	43	0	12	54	25
2371	191	255	0	44	31	8	8	48	22	0	11	36	13	11	29	50	39
2641	213	192	6	58	14	6	7	15	11	4	22	8	10	0	0	53	22
2817	227	276	16	11	11	8	29	47	46	0	5	53	28	11	28	54	14
3040	245	286	23	54	31	9	10	17	28	0	3	2	5	11	28	26	2
3621	292	277	6	28	3	9	0	30	17	8	2	35	17	5	28	0	58
3709	299	319	23	4	31	10	11	46	35	11	24	27	56	11	27	1	24
4267	344	361	0	44	53	11	22	34	0	0	0	13	59	6	31	11	0
4601	371	363	5	57	51	11	23	45	23	11	13	2	24	11	25	8	34
5059	409	8	6	13	8	0	4	0	46	9	17	7	44	0	2	14	56
5458	441	102	23	9	25	3	7	2	57	4	28	1	38	11	29	47	37
6444	521	0	3	3	31	11	24	56	0	1	13	22	46	0	0	56	36
6890	557	21	18	30	12	0	15	55	24	1	7	40	2	0	0	0	11
15928	1287	287	6	1	34	9	0	9	40	2	29	47	40	0	0	19	14
22325	1804	359	10	34	42	11	7	8	19	11	29	47	6	11	29	44	56

N.B. 1 0 29^d.530590852 Sig. 9701786292 Sig. 8605595917 Sig. 1,02235124
 716 57 324^d.21^h.40^m.23^s. 10^d.19^h.26^m.13^s. 4^d.4^h.49^m.12^s. 0^d.0^h.6^m.16^s.
 &c. And other Periods are also had by compounding the above-Periods.

Plinian PERIOD of Solar ECLIPSES. According to New Stile.							Plinian PERIOD of Lunar ECLIPSES. According to New Stile.										
Years of Chr. N. S.	Mid. Ecl. of ☉, at Lond. apt. Time.				Sun's M. Anom.	Sun's D's Ap.	Lat. ☉ fr. Sun.	N. or S. asc. or def.	Years of Chr. N. S.	Mid. Ecl. of ☾, at Lond. apt. Time.				Sun's M. Anom.	Sun's D's Ap.	Lat. ☾ fr. Sun.	N. or S. asc. or def.
	M	D	H	M	s o /	s o /	' "			M	D	H	M	s o /	s o /	' "	
1701	Febru.	7	11	6	7 10 3	11 7 30	35 20	N.A.	1701	Febru.	22	11	29	7 24 51	11 20 54	47 55	N.D.
	August	3	21	30	1 4 56	4 10 40	26 22	S.D.		August	18	1	32	1 18 54	4 22 45	48 24	S.A.
1702	Janua.	27	13	42	6 29 4	9 16 50	3 35	S.A.	1703	Janua.	2	18	59	6 4 23	7 13 35	35 43	S.D.
	July	24	9	39	0 24 20	2 20 50	18 14	N.D.		June	28	13	10	11 28 36	0 18 1	17 43	N.A.
1703	Janua.	16	23	27	6 18 22	7 26 23	43 50	S.A.		Decem.	22	18	30	5 23 16	5 22 44	2 1	N.D.
	July	13	14	36½	0 13 27	1 0 47	61 0	N.D.	1704	June	17	6	24	11 18 12	10 28 25	28 14	S.A.
	Decem.	8	3	44	5 8 52	5 9 33	79 13	N.A.		Decem.	10	19	10	5 12 12	4 1 56	40 3	N.D.
1704	Janua.	6	14	15	6 7 53	6 6 6	84 20	S.A.	1706	April	27	13	33	9 27 44	6 23 22	40 10	N.A.
	June	2	1	7	11 3 13	10 15 30	51 33	S.D.		Oct.	21	6	56	3 21 54	11 24 58	41 17	S.D.
	Nov.	26	17	38	4 28 20	3 19 16	38 37	N.A.	1707	April	16	13	38	9 16 38	5 2 59	0 52	S.A.
1705	May	22	7	55	10 22 23	8 25 29	8 26	S.D.		Oct.	10	22	25	3 11 27	10 4 56	1 41	N.D.
	Nov.	16	1	20	4 17 33	1 28 48	1 29	S.A.	1708	April	4	17	38	9 5 42	3 12 40	42 13	S.A.
1706	May	11	21	34	10 11 51	7 5 40	36 12	N.D.		Sept.	29	8	59	3 0 47	8 14 46	43 52	N.D.
	Nov.	5	2	19	4 6 30	2 8 6	40 21	S.A.	1710	Feb.	13	10	49½	7 15 38	11 6 28	34 0	N.A.
1707	May	1	14	19	10 1 27	5 16 30	80 24	N.D.		August	8	22	0	1 9 33	4 8 34	31 42	S.D.
	Sept.	25	11	4	2 26 12	9 20 43	81 33	N.A.	1711	Feb.	3	0	32½	7 5 6	9 16 14	7 36	S.A.
	Oct.	25	2	16	3 25 25	10 17 26	78 36	S.A.		July	29	5	52	0 28 47	2 18 35	11 48	N.D.
1708	Mar.	21	18	53	8 21 57	3 0 30	33 21	S.D.	1712	Janua.	23	7	47	6 24 18	7 25 43	47 21	S.A.
	Sept.	13	21	0	2 15 30	8 1 17	39 43	N.A.		July	17	20	10½	3 18 16	0 28 52	56 40	N.D.
1709	Mar.	11	0	17	8 11 5	1 10 5	7 26	N.D.	1713	June	8	6	30	11 9 1	10 14 2	50 45	S.D.
	Sept.	3	12	36	2 5 3	6 11 26	4 30	S.A.		Decem.	1	15	16	5 2 50	3 17 18	43 10	N.A.
1710	Feb.	28	0	14	7 29 59	11 19 27	46 31	N.D.	1714	May	28	19	5	10 28 26	8 24 37	6 14	S.D.
	August	24	5	16	1 24 38	4 21 40	49 15	S.A.		Nov.	21	1	4	4 22 9	1 26 52	2 42	N.A.
1711	Feb.	17	1	38	7 18 56	9 28 48	84 45	N.D.	1715	May	18	0	31	10 17 35	7 4 32	36 41	N.D.
	July	15	7	20	0 15 2	2 6 45	61 10	N.A.		Nov.	10	16	0	4 11 40	0 6 41	39 13	S.A.
1712	Janua.	7	22	1	6 9 6	7 11 49	38 58	S.D.	1717	Mar.	26	15	15	8 26 24	2 28 21	37 57	S.D.
	July	3	10	34½	0 4 5	0 16 39	18 43	N.A.		Sept.	20	5	56	2 21 27	8 0 26	37 25	N.A.
	Decem.	27	13	27	5 28 39	5 21 34	2 35	N.D.	1718	Mar.	16	3	58	8 15 50	1 8 15	3 41	N.D.
1713	June	22	11	18	11 23 1	10 16 27	23 40	S.A.		Sept.	9	7	52	2 10 26	6 10 4	3 39	S.A.
	Decem.	17	4	2	5 18 9	4 1 17	43 49	N.D.	Besides these 29 lunar and 41 solar Eclipses in this Plinian Period, of 18y 11d 7h 43m 20s, there were 5 solar Ones of no Use, viz in Mar. 1707, Jan. 1711, Oct. 1714, Aug. 1718, O. S. arising but to few Digits near the S. Pole. The 5th was on May 13, 1714, N. S. but 1 Digit in Quantity, near the N. Pole. So that the New Moon in May, 1732, was not eclipsed. But as, by the shifting of ☾'s Node, the Earth and Moon's Shadows are also shifted from falling on each other of those Bodies, so as to fall into the Expanse, the Tab. of Eclipse Periods, or No. of synodic Revolutions of ☾ to the ☉ (when the Sun comes near the Moon's Node) in P. 386, serves to shew when Eclipses will happen again, after a given No of Years, whether in Plinian or other Periods. For each Plinian Period there is an Equation of Time-Table, and Diff. Moon's Lat. below.								
1714	June	11	16	44	11 12 8	9 6 26	66 25	S.A.									
	Decem.	6	13	28	5 7 26	2 10 49	83 8	N.D.									
1715	May	2	21	35½	10 2 36	6 21 36	43 25	N.A.									
	Oct.	26	20	55	3 27 5	11 23 31	42 37	S.D.									
1716	April	21	14	7	9 22 15	5 1 49	1 19	S.A.									
	Oct.	14	22	6	3 16 2	10 2 56	3 2	S.D.									
1717	April	11	4	35	9 11 44	3 11 54	45 5	S.A.									
	Oct.	4	6	21	3 5 16	8 12 40	38 30	N.D.									
1718	Mar.	1	19	30	8 1 41	0 25 34	74 30	N.A.									
	Mar.	11	12	5	9 0 57	1 21 42	86 7	S.A.									
	Sept.	23	20	38	2 24 45	6 22 41	81 0	N.D.									

EQUATION of TIME and of MOON's Latitude, N. or S. ascending or descending, after one Plinian Period.

Argument. Sun's Mean Anomaly.												Argument. Sun from Moon's Apogee, 1 eqd.									
M. An.	Time.	Dif. Moon's Lat.	M. An.	Time.	Dif. Moon's Lat.	M. An.	Time.	Dif. Moon's Lat.	M. An.	Time.	Dif. Moon's Lat.	Sun's Ap. Time.	Sun's Ap. Time.	Sun's Ap. Time.	Sun's Ap. Time.	Sun's Ap. Time.	Sun's Ap. Time.	Sun's Ap. Time.	Sun's Ap. Time.	Sun's Ap. Time.	Sun's Ap. Time.
s o m s	' "	' "	s o m s	' "	' "	s o m s	' "	' "	s o m s	' "	' "	s o m s	' "	' "	s o m s	' "	' "	s o m s	' "	' "	s o m s
0 0 45 2	4 27	3 5	7 19	2 7	6 5	46 19	0 24	9 5	9 7	2 50	0 0 29 7	3 5	3 42	6 5	25 3	9 5	0 22	0 0 45 2	4 27	3 5	7 19
5 44 30	4 26	10	11 20	1 56	10	45 22	0 26	10	12 58	3 0	10 28 44	10 5	59	10	24 50	10	2 0	5 44 30	4 26	10	11 20
10 43 44	4 24	15	15 13	1 46	15	44 2	0 29	15	16 42	3 10	15 28 12	15 8	6	15	24 30	15	4 31	10 43 44	4 24	15	15 13
15 42 36	4 21	20	19 4	1 36	20	42 27	0 34	20	20 18	3 20	20 27 18	20 10	10	20	24 7	20	6 57	15 42 36	4 21	20	19 4
20 41 9	4 16	25	22 46	1 26	25	40 26	0 39	25	23 46	3 29	25 26 12	25 12	8	25	23 30	25	9 23	20 41 9	4 16	25	22 46
25 39 25	4 11	4	26 17	1 16	7	37 59	0 46	10	26 58	3 38	1 0 24 47	4	0 13 59	7	0 22 41	10	0 11 45	25 39 25	4 11	4	26 17
30 37 23	4 6	5	29 39	1 7	5	35 22	0 53	5	29 57	3 46	5 23 19	5	15 44	5	21 42	5	14 4	30 37 23	4 6	5	29 39
35 35 4	4 0	10	32 47	0 59	10	32 28	1 0	10	32 48	3 54	10 21 38	10	17 18	10	20 38	10	16 22	35 35 4	4 0	10	32 47
40 32 30	3 53	15	35 39	0 52	15	29 18	1 8	15	35 21	4 0	15 19 40	15	18 40	15	19 27	15	18 33	40 32 30	3 53	15	35 39
45 29 41	3 45	20	38 18	0 45	20	25 55	1 17	20	37 38	4 6	20 17 36	20	20 0	20	18 10	20	20 30	45 29 41	3 45	20	38 18
50 26 37	3 37	25	40 40	0 39	25	22 24	1 26	25	39 35	4 12	25 15 25	25	11 10	25	16 35	25	22 21	50 26 37	3 37	25	40 40
55 23 21	3 28	3	42 37	0 34	3	18 40	1 36	11	41 17	4 17	3 0 13 11	5	0 22 11	8	0 15 0	11	0 23 59	55 23 21	3 28	3	42 37
60 19 55	3 19	5	44 11	0 30	5	14 50	1 46	5	42 42	4 21	5 10 48	5	22 57	5	13 15	5	25 29	60 19 55	3 19	5	44 11
65 16 20	3 9	10	45 30	0 26	10	10 53	1 57	10	43 50	4 24	10 8 21	10	23 41	10	11 21	10	26 45	65 16 20	3 9	10	45 30
70 12 32	2 59	15	46 25	0 24	15	6 55	2 7	15	44 35	4 26	15 5 53	15	24 19	15	9 19	15	27 36	70 12 32	2 59	15	46 25
75 8 43	2 49	20	46 58	0 23	20	2 51	2 18	20	45 2	4 27	20 3 25	20	24 39	20	7 12	20	28 27	75 8 43	2 49	20	46 58
80 4 48	2 38	25	47 7	0 23	25	1 11	2 29	25	45 13	4 28	25 1 0	25	24 59	25	5 0	25	28 54	80 4 48	2 38	25	47 7
85 0 45	2 28	3	46 54	0 23	3	5 13	2 40	12	45 2	4 27	3 0 1 20	6	0 25 51	9	0 2 42	12	0 29 7	85 0 45	2 28	3	46 54

N. B. The mean Dif. of Sun's ☉ in a Plinian Period being [Decim.] 118,984329 wanting [Decim.] .470130, of a Degree of a Revolution if that Decimal be multiplied into the Number of Plinian Periods since any Eclipse, the Product will be the Degrees of Sun's Dif. ☉, less than the Dif. when the Eclipse happened. Which Dif. taken from the Sun's 1st Dif. ☉, will shew the Sun's present Dif. ☉. And if within 18 or 120 (the Limits of solar and lunar Eclipses, respectively) an Eclipse of one of those Luminaries will respectively happen again; otherwise not.

RADICAL Mean PLACES and MOTIONS of the MOON from the SUN, &c. for determining the Mean TIME of the mean, and then the true NEW and FULL MOONS, with (or without) Acceleration of Mean Motion. According to Old Style.

Years of Christ.	☿ à ☽	☿'s M. An.	☿ à ☽'s Ap.	☿ à ☽	Years of Christ.	☿ à ☽	☿'s M. An.	☿ à ☽'s Ap.	☿ à ☽
s o i "	s o i "	s o i "	s o i "	s o i "	s o i "	s o i "	s o i "	s o i "	s o i "
1652	0 6 48 7	6 13 41 55	4 26 11 38	9 5 27 23	1704	2 5 41 4	6 13 12 17	6 10 36 39	6 21 37 42
53	4 16 25 30	6 13 26 34	3 15 17 28	0 24 32 46	5	6 15 18 27	6 12 56 57	4 19 42 29	7 10 43 5
54	8 26 2 54	6 13 11 14	2 4 23 18	10 13 38 16	6	10 24 55 50	6 12 41 36	8 18 48 19	7 29 48 29
55	1 5 40 17	6 12 55 54	0 13 29 7	11 2 43 33	7	8 4 33 14	6 12 26 15	2 7 54 8	8 18 53 52
1656	5 27 29 7	6 13 39 38	11 13 27 24	11 22 51 15	1708	7 26 22 3	6 13 10 1	0 27 52 26	9 9 1 35
57	10 7 6 30	6 13 24 18	10 2 33 14	0 11 56 38	9	0 5 59 26	6 12 54 41	11 16 58 16	9 28 6 58
58	2 16 43 53	6 13 8 57	8 21 39 4	1 1 2 2	10	4 15 36 49	6 12 39 20	10 6 4 6	10 17 12 22
59	6 26 21 17	6 12 53 37	7 10 44 53	1 20 7 25	11	8 25 14 13	6 12 24 0	8 25 9 55	11 6 17 45
1660	11 18 10 6	6 13 37 22	6 0 43 11	2 10 15 8	1712	1 17 3 3	6 13 7 44	7 15 8 12	11 26 25 27
61	3 27 47 29	6 13 22 2	4 19 49 1	0 29 20 31	13	5 26 40 26	6 12 52 24	6 4 14 2	0 15 26 50
62	8 7 24 53	6 13 6 41	3 8 54 51	3 18 25 55	14	10 6 17 49	6 12 57 3	4 23 19 52	1 4 36 14
63	0 17 2 17	6 12 51 21	1 28 0 40	4 7 31 18	15	2 15 55 13	6 12 21 43	3 12 25 41	1 23 41 37
1664	5 8 51 6	6 13 35 5	0 17 58 57	4 27 39 0	1716	7 7 44 3	6 13 5 27	2 2 23 58	2 13 49 19
65	9 18 28 29	6 13 19 45	0 7 4 47	5 16 44 23	17	11 17 21 26	6 12 50 7	0 21 29 48	3 2 54 42
66	1 28 5 52	6 13 4 24	9 26 10 37	5 35 49 47	18	3 26 58 49	6 12 34 46	11 10 35 38	3 22 0 6
67	6 7 43 16	6 12 49 4	8 15 16 26	6 24 55 10	19	8 6 36 13	6 12 19 26	9 29 41 27	4 11 5 29
1668	10 29 32 6	6 13 32 48	7 5 14 43	7 15 2 52	1720	0 28 25 3	6 13 3 10	8 19 39 44	5 1 13 11
69	3 9 9 29	6 13 17 28	5 24 20 33	8 4 8 15	21	5 8 2 26	6 12 47 50	7 8 45 34	5 20 18 34
70	7 18 46 52	6 13 2 7	4 13 26 23	8 23 13 39	22	9 17 39 49	6 12 32 29	5 27 51 24	6 9 23 58
71	11 28 24 16	6 12 46 47	3 2 32 12	9 12 19 2	23	1 27 17 13	6 12 17 9	4 16 57 13	6 28 29 21
1672	4 20 13 6	6 13 30 31	1 22 30 29	10 2 26 44	1724	6 19 6 2	6 13 0 54	3 6 55 31	7 18 37 4
73	8 29 50 29	6 13 15 11	0 11 36 19	10 21 32 7	25	10 28 43 25	6 12 45 34	1 26 1 21	8 7 42 27
74	1 9 27 52	6 12 59 50	11 0 42 9	11 10 37 31	26	3 8 20 48	6 12 30 13	0 5 7 11	8 26 47 51
75	5 19 5 16	6 12 44 30	9 19 47 58	11 29 42 54	27	7 17 58 12	6 12 14 53	11 4 13 0	9 15 53 14
1676	10 10 54 5	6 13 28 15	8 9 46 16	0 19 50 37	1728	0 9 47 2	6 12 58 37	9 24 11 17	10 6 0 56
77	2 20 31 28	6 13 12 55	6 28 52 6	1 8 56 0	29	4 19 24 25	6 12 43 17	8 13 17 7	10 25 6 19
78	7 0 8 51	6 12 57 34	5 17 57 56	1 28 1 24	30	8 29 1 48	6 12 27 56	7 2 22 57	11 14 17 43
79	11 9 46 15	6 12 42 14	4 7 3 45	2 17 6 47	31	4 21 48 49	6 12 12 36	5 21 28 46	0 3 17 6
1680	4 1 35 5	6 13 25 58	2 7 2 2	3 7 14 29	1732	6 0 28 2	6 12 56 20	4 11 27 3	0 23 24 48
81	8 11 12 28	6 13 10 38	0 26 7 52	3 26 19 52	33	10 10 5 25	6 12 41 0	3 0 32 53	1 12 30 11
82	0 20 49 51	6 12 55 17	11 15 13 42	4 15 25 16	34	2 19 42 48	6 12 25 39	1 19 38 43	2 1 35 35
83	5 0 27 15	6 12 39 57	10 4 19 31	5 4 30 39	35	6 29 20 12	6 12 10 19	0 8 44 32	2 20 46 58
1684	9 22 16 4	6 13 23 42	8 24 17 49	5 24 38 22	1736	11 21 5 1	6 12 54 4	10 28 42 50	3 10 48 41
85	2 1 53 27	6 13 8 22	7 13 23 39	6 13 43 45	37	4 0 46 24	6 12 38 44	9 17 48 40	3 29 54 4
86	6 11 30 50	6 12 53 1	6 2 29 29	7 2 49 9	38	8 10 23 47	6 12 23 23	8 6 54 30	4 18 59 28
87	10 21 8 14	6 12 37 41	4 21 35 18	7 21 54 32	39	0 20 1 11	6 12 8 3	6 26 0 19	5 8 4 51
1688	3 12 57 4	6 13 21 25	3 11 33 35	8 12 2 14	1340	5 11 50 1	6 12 51 47	5 15 58 36	5 28 12 33
89	7 22 34 27	6 13 6 5	2 0 39 25	9 1 7 37	41	9 21 27 24	6 12 36 27	4 5 4 26	6 17 17 56
90	0 2 11 50	6 12 50 44	0 19 45 15	9 20 13 1	42	2 1 4 47	6 12 21 6	2 24 10 16	7 6 23 20
91	4 11 49 14	6 12 35 24	11 8 51 4	10 9 18 24	43	6 10 42 11	6 12 5 46	1 13 16 5	7 25 28 43
1692	9 3 38 4	6 13 19 8	9 28 49 21	10 29 26 6	1744	11 2 31 1	6 12 49 30	0 3 14 22	8 15 36 25
93	1 13 15 27	6 13 3 48	8 17 55 11	11 18 31 29	45	3 12 8 24	6 12 34 10	10 22 20 12	9 4 41 48
94	5 22 52 50	6 12 48 27	7 7 1 2	0 7 36 53	46	7 25 45 47	6 12 18 49	9 11 26 2	9 23 47 12
95	10 2 30 14	6 12 33 7	5 26 6 50	0 26 42 16	47	0 1 23 11	6 12 3 29	8 0 31 51	10 12 52 35
1696	2 24 19 4	6 13 16 51	4 16 5 7	1 16 49 58	1748	4 23 12 1	6 12 47 13	6 20 30 8	11 3 0 17
97	7 3 56 27	6 13 1 31	3 5 10 57	2 5 55 21	49	9 2 49 24	6 12 31 53	5 9 35 58	11 22 5 40
98	11 13 33 50	6 12 46 10	1 24 16 47	2 55 0 45	50	1 12 26 47	6 12 16 32	3 28 41 48	0 11 11 4
99	3 23 11 14	6 12 30 50	0 13 22 36	3 14 6 8	51	5 22 4 11	6 12 1 12	2 17 47 37	1 0 16 27
1700	8 15 0 4	6 13 14 34	11 23 20 53	4 4 13 50	1752	10 13 53 0	6 12 44 57	1 7 45 55	1 20 24 10
1	0 24 37 27	6 12 59 14	10 12 26 43	4 23 19 13	For Convenience of Computation, the Computer must reduce the New and Full Moons of the Old to the Day of New Style, by adding the Number of Days Difference of the Style.				
2	5 4 14 50	6 12 43 53	9 1 32 33	5 12 24 37					
3	9 13 52 14	6 12 28 33	7 20 38 22	6 1 30 0					
Mean MOTION of the MOON from the SUN, &c. for Julian CENTURIES.									
Julian Cent.	Mot. ☿ à ☽	Mot. ☿ An.	M. ☿ à Ap.	M. ☿ à ☽	Julian Cent.	Mot. ☿ à ☽	M. ☿ An.	M. ☿ à Ap.	M. ☿ à ☽
s o i "	s o i "	s o i "	s o i "	s o i "	s o i "	s o i "	s o i "	s o i "	s o i "
100	10 7 4 53	11 29 3 2	8 11 34 17	4 14 56 47	1300	1 2 3 29	11 17 39 26	1 0 25 41	10 14 18 11
200	8 14 9 46	11 28 6 4	4 23 8 34	8 29 53 34	1400	11 9 8 22	11 16 42 28	9 11 59 58	2 29 14 58
300	6 21 14 39	11 27 9 6	1 4 42 51	1 14 50 21	1500	9 16 13 15	11 15 45 30	5 23 34 15	7 14 11 45
400	4 28 19 32	11 26 12 8	9 16 17 8	5 29 47 8	1600	7 23 18 8	11 14 48 32	4 14 24 14	11 28 42 48
500	3 5 24 25	11 25 15 10	5 27 51 25	10 14 43 55	1700	5 19 57 12	11 13 51 34	10 16 42 49	4 14 5 19
600	1 12 29 18	11 24 18 12	2 9 25 42	2 29 40 42	1800	2 12 52 19	11 12 54 36	9 7 42 48	1 13 52 27
700	11 19 34 11	11 23 21 14	10 20 59 59	7 14 37 29	1900	0 21 37 40	11 11 0 40	11 21 25 40	5 28 5 40
800	8 3 43 57	11 22 24 16	3 14 8 33	4 14 31 3	2000	10 28 42 33	11 10 3 42	8 2 59 57	1 13 52 27
900	6 10 43 50	11 21 27 18	4 18 51 24	5 29 21 24	2100	9 5 47 26	11 9 6 44	4 14 24 14	2 28 49 14

Mean MOTION of the MOON from the SUN, &c. for MONTHS.

Mths.	Mot. D à Sun	Mot. $\odot$ An.	M. $\odot$ à D Ap.	M. Sun à $\odot$
Jan.	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0
31 Feb.	0 17 54 48	1 0 33 13	0 27 6 5	1 2 11 48
59 Mar.	0 11 29 15 15	1 28 9 1	1 22 34 48	2 2 16 39
90 April	0 0 17 10 3	2 28 42 15	2 18 40 54	3 3 28 27
120 May	0 0 22 53 23	3 28 16 20	3 14 54 32	4 4 37 57
151 June	0 1 10 48 10	4 28 49 33	4 12 0 36	5 6 49 44
181 July	0 1 16 31 31	5 28 23 38	5 8 14 14	6 7 59 14
212 Aug.	0 2 4 26 19	6 28 56 50	6 5 20 19	7 10 11 1
243 Sept.	0 2 22 21 7	7 29 30 3	7 2 26 24	8 12 22 47
273 Oct.	0 2 28 4 27	8 29 4 8	7 28 40 2	9 13 32 18
304 Nov.	0 3 15 59 15	9 29 37 22	8 25 46 7	10 15 44 6
334 Dec.	0 3 21 42 36	10 29 11 26	9 21 59 45	11 16 53 35

Mean MOTION of the MOON from the SUN, &c. for DAYS

Days.	Mot. D à Sun	M. Sun's An.	M. $\odot$ à D Ap.	Mo. Sun à $\odot$
1	0 12 11 27	0 0 59 8	0 0 52 27	0 1 2 19
2	0 24 23 53	0 1 58 17	0 1 44 55	0 2 4 38
3	1 6 34 20	0 2 57 25	0 2 37 22	0 3 6 57
4	1 18 45 47	0 3 56 32	0 3 29 49	0 4 9 16
5	2 0 57 13	0 4 55 41	0 4 22 16	0 5 11 35
6	2 13 8 40	0 5 54 49	0 5 14 44	0 6 13 54
7	2 25 20 7	0 6 53 57	0 6 7 11	0 7 16 13
8	3 7 31 34	0 7 53 6	0 6 59 38	0 8 18 32
9	3 19 43 0	0 8 52 14	0 7 52 5	0 9 20 51
10	4 1 54 27	0 9 51 21	0 8 44 33	0 10 23 10
11	4 14 5 54	0 10 50 30	0 9 37 0	0 11 25 29
12	4 26 17 20	0 11 49 38	0 10 29 27	0 12 27 48
13	5 8 28 47	0 12 48 46	0 11 21 54	0 13 30 7
14	5 20 40 14	0 13 47 55	0 12 14 22	0 14 32 26
15	6 2 51 40	0 14 47 3	0 13 6 49	0 15 34 44
16	6 15 3 7	0 15 46 11	0 13 59 16	0 16 37 3
17	6 27 14 34	0 16 45 19	0 14 51 44	0 17 39 22
18	7 9 26 0	0 17 44 27	0 15 44 11	0 18 41 41
19	7 21 37 27	0 18 43 35	0 16 36 38	0 19 44 0
20	8 3 48 54	0 19 42 44	0 17 29 5	0 20 46 19
21	8 16 0 21	0 20 41 52	0 18 21 33	0 21 48 38
22	8 28 11 47	0 21 40 50	0 19 14 0	0 23 50 57
23	9 10 23 14	0 22 40 8	0 20 6 27	0 24 52 16
24	9 22 34 41	0 23 39 16	0 20 58 54	0 25 55 35
25	10 4 46 7	0 24 38 24	0 21 51 22	0 26 57 54
26	10 16 57 34	0 25 37 33	0 22 43 49	0 27 0 13
27	10 29 9 1	0 26 36 41	0 23 36 16	0 28 2 32
28	11 11 20 27	0 27 35 48	0 24 28 43	0 29 4 51
29	11 23 31 54	0 28 34 57	0 25 21 11	1 0 7 10
30	0 5 43 21	0 29 34 5	0 26 13 38	1 1 9 29
31	0 17 54 48	1 0 33 13	0 27 6 5	1 2 11 48

In Jan. and Feb. Leap-Year, take out for a Day sooner.

USE of the FOREGOING TABLES of MOON from the Sun.

COLLECT the Mean Places (from the Radical Places and Mean Motions connected) for the Beginning of that Month, in which the Mean Time of the Mean or true New or Full Moon is required. — Then connect therewith the total Quantity of Correction of M. Motion and Acceleration of the Moon, (from Tab. p. 374 foregoing) and the Result will give the corrected or true Mean Place of D à $\odot$, at the Beginning of the said Month; connecting also the Variation of the D 's Ap. and $\odot$, corresponding to the said total Correction, (of contrary Signs) with the Places of $\odot$ à D 's Ap. and $\odot$ à $\odot$ (according to p. 374) the Place of the Sun's M. An. being permanent, as its Motion is esteem'd to be invariable. — Now deduct the final Place of

D à Sun from 12^h (if New Moon is sought) or from 6^h or 18^h, (if Full Moon is sought) and then take from the Remainder, in Mot. D à Sun, for as many Days, Hours, Minutes and Seconds as can be deducted: For which Days, Hours, &c. you must add the correspondent Motions to the Sun's An. Sun à D Ap. and Moon à Sun. By which final Places, you must obtain and subjoin to the Time of Mean Syzygy, the Time, with its proper Sign, answering, in Mot. D à Sun, to the Sum of the Sun's Equation, and of the Sum of the Moon's annual, 6th, and Elliptic Equation, in Syzygies (taken out by the proper Arguments, and changing the Sign of the Sum of D 's Equations) for the true Time of the New or Full Moon required. See Examples farther on.

Mean MOTION of D from $\odot$, &c. for Hours, Minutes, and Seconds.

H	Mot. D à Sun	M. Sun's An.	M. $\odot$ à D Ap.	M. Sun à $\odot$
M	0 1 1 1	0 1 1 1	0 1 1 1	0 1 1 1
S	1 1 1 1	1 1 1 1	1 1 1 1	1 1 1 1
1	0 30 29	0 2 28	0 2 11	0 2 36
2	1 0 57	0 4 56	0 4 22	0 5 12
3	1 31 26	0 7 24	0 6 33	0 7 47
4	2 1 54	0 9 51	0 8 44	0 10 23
5	2 32 23	0 12 19	0 10 56	0 12 59
6	3 2 52	0 14 47	0 13 7	0 15 35
7	3 33 20	0 17 15	0 15 18	0 18 10
8	4 3 49	0 19 43	0 17 29	0 20 46
9	4 34 17	0 22 11	0 19 40	0 23 22
10	5 4 46	0 24 38	0 21 51	0 25 58
11	5 35 15	0 27 6	0 24 2	0 28 33
12	6 5 43	0 29 34	0 26 13	0 31 9
13	6 36 12	0 32 2	0 28 25	0 33 45
14	7 6 41	0 34 30	0 30 36	0 36 21
15	7 39 9	0 36 58	0 32 47	0 38 56
16	8 7 38	0 39 26	0 34 58	0 41 32
17	8 38 6	0 41 53	0 37 9	0 44 8
18	9 8 35	0 44 21	0 39 20	0 46 44
19	9 39 4	0 46 49	0 41 31	0 49 20
20	10 9 32	0 49 17	0 43 42	0 51 55
21	10 40 1	0 51 45	0 45 53	0 54 31
22	11 10 29	0 54 13	0 48 5	0 57 7
23	11 40 58	0 56 40	0 50 16	0 59 43
24	12 11 27	0 59 8	0 52 27	1 2 18
25	12 41 55	1 1 36	0 54 38	1 4 54
26	13 12 24	1 4 4	0 56 49	1 7 30
27	13 42 53	1 6 32	0 59 0	1 10 6
28	14 13 21	1 9 0	1 1 11	1 12 41
29	14 43 50	1 11 28	1 3 22	1 15 17
30	15 14 18	1 13 55	1 5 33	1 17 53
31	15 44 47	1 16 23	1 7 45	1 20 29
32	16 15 16	1 18 51	1 9 56	1 23 5
33	16 45 44	1 21 19	1 12 7	1 25 40
34	17 16 13	1 23 47	1 14 18	1 28 16
35	17 46 41	1 26 15	1 16 29	1 30 52
36	18 17 10	1 28 42	1 18 40	1 33 28
37	18 49 39	1 31 10	1 20 51	1 36 3
38	19 18 7	1 33 38	1 23 2	1 38 39
39	19 48 36	1 36 6	1 25 14	1 41 15
40	20 19 4	1 38 34	1 27 25	1 43 51
41	20 49 33	1 41 2	1 29 36	1 46 26
42	21 20 2	1 43 30	1 31 47	1 49 2
43	21 50 30	1 45 57	1 33 58	1 51 38
44	22 20 59	1 48 25	1 36 9	1 54 14
45	22 51 28	1 50 53	1 38 20	1 56 49
46	23 21 56	1 53 21	1 40 31	1 59 25
47	23 52 25	1 55 49	1 42 42	2 2 1
48	24 22 53	1 58 17	1 44 54	2 4 37
49	24 53 22	2 0 44	1 47 5	2 7 13
50	25 23 51	2 3 12	1 49 16	2 9 48
51	25 54 19	2 5 40	1 51 27	2 12 24
52	26 24 48	2 8 8	1 53 38	2 15 0
53	26 55 16	2 10 36	1 55 49	2 17 36
54	27 25 45	2 13 4	1 58 0	2 20 11
55	27 56 14	2 15 32	2 0 11	2 22 47
56	28 26 42	2 17 59	2 2 23	2 25 23
57	28 57 11	2 20 27	2 4 34	2 27 59
58	29 27 40	2 22 55	2 6 45	2 30 34
59	29 58 8	2 25 23	2 8 56	2 33 10
60	30 28 37	2 27 51	2 11 7	2 35 46

EQUATION

EQUATION of the MOON'S CENTER (and II. Equation, Supplemental, compounded) at New and Full MOON. To be used with the 1st and 6th Supplemental Equations of the MOON, (the 6th to be doubled) p. 61, 62 and 74, and Sun's Equation, p. 12.

Argument. SUN from the Moon's Apogee 1 equated, at New; and + Six Signs at Full MOON.

Sun à D Ap.	Sig. 0	Diff.	Sig. 1.	Diff.	Sig. 2.	Diff.	Sig. 3.	Diff.	Sig. 4.	Diff.	Sig. 5.	Diff.	Sun à D Ap.
0	0 0 0	1 11	0 0 0	1 11	0 0 0	1 11	0 0 0	1 11	0 0 0	1 11	0 0 0	1 11	0
0	0 0 0	5 36	2 38 53	4 40	4 27 45	2 21	4 57 46	0 26	4 8 20	2 49	2 19 27	4 27	30
1	0 5 36	5 36	2 43 33	4 36	4 30 6	2 15	4 57 20	0 31	4 5 31	2 53	2 15 7	4 22	29
2	0 11 12	5 35	2 48 9	4 33	4 32 21	2 10	4 56 49	0 37	4 2 38	2 57	2 10 45	4 23	28
3	0 16 47	5 35	2 52 42	4 29	4 34 31	2 4	4 56 12	0 43	3 59 41	3 2	2 6 22	4 25	27
4	0 22 23	5 35	2 57 11	4 26	4 36 35	1 59	4 55 29	0 47	3 56 39	3 4	2 1 57	4 26	26
5	0 27 58	5 35	3 1 37	4 22	4 38 34	1 53	4 54 42	0 52	3 53 35	3 7	1 57 31	4 29	25
6	0 33 33	5 33	3 5 59	4 17	4 40 27	1 48	4 53 50	0 58	3 50 28	3 12	1 53 2	4 31	24
7	0 39 6	5 32	3 10 16	4 14	4 42 15	1 43	4 52 52	1 4	3 47 16	3 15	1 48 31	4 32	23
8	0 44 38	5 31	3 14 30	4 10	4 43 58	1 37	4 51 48	1 8	3 44 1	3 19	1 43 59	4 33	22
9	0 50 9	5 31	3 18 40	4 5	4 45 35	1 30	4 50 40	1 14	3 40 42	3 24	1 39 26	4 34	21
10	0 55 40	5 29	3 22 45	4 0	4 47 5	1 25	4 49 26	1 19	3 37 18	3 26	1 34 52	4 36	20
11	1 1 9	5 28	3 26 45	3 56	4 48 30	1 20	4 48 7	1 23	3 33 52	3 30	1 30 16	4 38	19
12	1 6 37	5 27	3 30 41	3 52	4 49 50	1 15	4 46 44	1 28	3 30 22	3 33	1 25 38	4 40	18
13	1 12 4	5 26	3 34 33	3 47	4 51 5	1 8	4 45 16	1 34	3 26 49	3 35	1 20 58	4 41	17
14	1 17 30	5 23	3 38 20	3 42	4 52 13	1 2	4 43 42	1 38	3 23 14	3 38	1 16 18	4 42	16
15	1 22 53	5 20	3 42 2	3 38	4 53 15	0 58	4 42 4	1 43	3 19 36	3 42	1 11 37	4 42	15
16	1 28 13	5 18	3 45 40	3 34	4 54 13	0 52	4 40 21	1 48	3 15 54	3 44	1 6 55	4 42	14
17	1 33 31	5 17	3 49 14	3 29	4 55 5	0 46	4 38 33	1 52	3 12 10	3 48	1 2 13	4 43	13
18	1 38 48	5 15	3 52 43	3 23	4 55 51	0 41	4 36 41	1 57	3 8 22	3 51	0 57 30	4 45	12
19	1 44 3	5 13	3 56 6	3 18	4 56 32	0 35	4 34 44	2 2	3 4 31	3 53	0 52 45	4 45	11
20	1 49 16	5 11	3 59 24	3 13	4 57 7	0 29	4 32 42	2 7	3 0 38	3 56	0 48 0	4 46	10
21	1 54 27	5 9	4 2 37	3 9	4 57 36	0 23	4 30 35	2 11	2 56 42	3 58	0 43 14	4 46	9
22	1 59 36	5 5	4 5 46	3 3	4 57 59	0 17	4 28 24	2 15	2 52 44	4 1	0 38 28	4 47	8
23	2 4 41	5 2	4 8 49	2 58	4 58 16	0 13	4 26 9	2 19	2 48 43	4 4	0 33 41	4 49	7
24	2 9 43	5 0	4 11 47	2 53	4 58 29	0 6	4 23 50	2 25	2 44 39	4 7	0 28 52	4 48	6
25	2 14 43	4 58	4 14 40	2 47	4 58 35	0 1	4 21 25	2 29	2 40 32	4 8	0 24 4	4 48	5
26	2 19 41	4 54	4 17 27	2 41	4 58 36	0 5	4 18 56	2 33	2 36 24	4 10	0 19 16	4 49	4
27	2 24 35	4 49	4 20 8	2 38	4 58 31	0 9	4 16 23	2 37	2 32 14	4 13	0 14 27	4 49	3
28	2 29 24	4 46	4 22 46	2 33	4 58 22	0 15	4 13 46	2 41	2 28 1	4 16	0 9 38	4 49	2
29	2 34 10	4 43	4 25 19	2 26	4 58 7	0 21	4 11 5	2 45	2 23 45	4 18	0 4 49	4 49	1
30	2 38 53	4 43	4 27 45		4 57 46		4 8 20		2 19 27		0 0 0	4 49	0
Sun à D Ap.	Sig. 11	Diff.	Sig. 10	Diff.	Sig. 9.	Diff.	Sig. 8.	Diff.	Sig. 7.	Diff.	Sig. 6.	Diff.	Sun à D Ap.

The MOON'S LATITUDE, INCLINATION and REDUCTION to the ECLIPTIC, at New and Full MOON.

Argument. Sun's Distance from the MOON's mean NODE, or Argument of Latitude.

Sun from Node.	Sig. 0 N. Sig. 6 S. Latitude.	Diff.	Reduction.	Diff.	Inclination, Way to Ecliptic.	Diff.	Sig. 1 N. Sig. 7 S. Latitude.	Diff.	Reduction.	Diff.	Sig. 2 N. Sig. 8 S. Latitude.	Diff.	Reduction.	Diff.	Sun from Node.
0	0 0 0	1 11	0 0 0	1 11	0 0 0	1 11	0 0 0	1 11	0 0 0	1 11	0 0 0	1 11	0 0 0	1 11	0
0	0 0 0	5 15	0 0	14	5 17 20	0 3	2 30 12	4 36	6 1	7	4 19 39	2 34	5 59	8	30
1	0 5 15	5 15	0 14	15	5 17 17	0 9	2 34 42	4 27	6 8	6	4 22 13	2 29	5 51	8	29
2	0 10 30	5 14	0 29	14	5 17 8	0 14	2 39 9	4 25	6 14	6	4 24 42	2 24	5 43	8	28
3	0 15 44	5 15	0 43	15	5 16 54	0 21	2 43 34	4 22	6 20	6	4 27 6	2 19	5 35	9	27
4	0 20 59	5 14	0 58	14	5 16 33	0 27	2 47 56	4 19	6 26	5	4 29 25	2 15	5 26	8	26
5	0 26 13	5 13	1 12	14	5 16 6	0 33	2 52 15	4 15	6 31	5	4 31 40	2 9	5 17	9	25
6	0 31 26	5 13	1 26	15	5 15 33	0 38	2 56 30	4 12	6 36	4	4 33 49	2 4	5 8	10	24
7	0 36 39	5 12	1 41	14	5 14 55	0 44	3 0 42	4 9	6 40	4	4 35 53	1 59	4 58	10	23
8	0 41 51	5 11	1 55	14	5 14 11	0 49	3 4 51	4 5	6 43	3	4 37 52	1 55	4 48	11	22
9	0 47 2	5 11	2 9	13	5 13 22	0 55	3 8 56	4 2	6 46	3	4 39 47	1 50	4 37	11	21
10	0 52 13	5 10	2 22	13	5 12 27	1 11	3 12 58	3 58	6 49	3	4 41 37	1 44	4 26	11	20
11	0 57 23	5 8	2 36	14	5 11 26	1 17	3 16 56	3 55	6 52	2	4 43 21	1 39	4 15	12	19
12	1 2 31	5 7	2 49	13	5 10 19	1 7	3 20 51	3 51	6 54	1	4 45 0	1 34	4 3	12	18
13	1 7 38	5 6	3 3	13	5 9 6	1 3	3 24 42	3 47	6 55	1	4 46 34	1 29	3 51	12	17
14	1 12 44	5 5	3 16	13	5 7 47	1 25	3 28 29	3 43	6 56	0	4 48 3	1 23	3 39	12	16
15	1 17 49	5 3	3 29	12	5 6 22	1 30	3 32 12	3 39	6 56	1	4 49 26	1 18	3 27	12	15
16	1 22 52	5 2	3 41	12	5 4 52	1 36	3 35 51	3 35	6 55	1	4 50 44	1 13	3 14	13	14
17	1 27 54	5 0	3 53	12	5 3 16	2 41	3 39 26	3 31	6 54	1	4 51 57	1 7	3 1	13	13
18	1 32 54	4 58	4 5	12	5 1 35	1 46	3 42 57	3 27	6 53	2	4 53 4	1 2	2 48	13	12
19	1 37 52	4 57	4 17	11	4 59 49	1 53	3 46 24	3 24	6 51	2	4 54 6	0 57	2 35	13	11
20	1 42 49	4 55	4 28	11	4 57 56	2 3	3 49 48	3 19	6 49	3	4 55 3	0 52	2 21	13	10
21	1 47 44	4 53	4 39	11	4 55 58	2 8	3 53 7	3 15	6 46	3	4 55 55	0 47	2 8	13	9
22	1 52 37	4 50	4 50	10	4 53 55	2 14	3 56 22	3 11	6 43	4	4 56 42	0 41	1 54	14	8
23	1 57 27	4 48	5 0	10	4 51 47	2 19	3 59 33	3 6	6 39	4	4 57 23	0 35	1 40	14	7
24	2 2 15	4 45	5 10	10	4 49 33	2 24	4 2 39	3 1	6 35	5	4 57 58	0 29	1 26	14	6
25	2 7 0	4 43	5 20	9	4 47 14	2 29	4 5 40	2 56	6 30	5	4 58 27	0 24	1 12	14	5
26	2 11 43	4 41	5 29	9	4 44 50	2 30	4 8 36	2 52	6 25	6	4 58 51	0 19	0 58	15	4
27	2 16 24	4 38	5 38	8	4 42 20	2 35	4 11 28	2 48	6 19	6	4 59 10	0 14	0 43	15	3
28	2 21 2	4 37	5 46	8	4 39 45	2 40	4 14 16	2 44	6 13	7	4 59 24	0 8	0 29	15	2
29	2 25 38	4 34	5 54	7	4 37 5	2 46	4 17 0	2 39	6 6	7	4 59 32	0 3	0 14	15	1
30	2 30 12	4 34	6 1	7	4 34 19		4 19 39		6 59		4 59 35		0 0	14	0
Sun from Node.	Latitude	Diff.	Reduction.	Diff.	Inclination, Way to Ecliptic.	Diff.	Latitude	Diff.	Reduction.	Diff.	Latitude	Diff.	Reduction.	Diff.	Sun from Node.

THE true Hourly MOTIONS, DIAMETERS, and PARALLAXES of the SUN and MOON, in ECLIPSES.

N. B. In Eclipses of the Moon, add 6 Signs to the Arg. Annum, or $\odot$ à $\downarrow$ Ap. & equated.

Sun's M. Anomaly	Sun's true Hourly Motion.	Sun's Diameter.	Sun's M. Anomaly	Sun from Moon's Apogee.	M's true Hourly Motion.	Moon's Horizont. Diameter.	Moon's Horizont. Parallax.	Sun from Moon's Apogee.	Sun & Moon's Altitud.	Sun's Paral- lax.	Increase of Sun & M's Diameter.
0	0	2 23	31 38	0	12	29 25	53 28	0	12	0	0
5	2 23	31 38	25	5	29 36	29 25	53 29	25	2	11	1
10	2 23	31 38	20	10	29 39	29 26	53 31	20	4	11	2
15	2 23	31 39	15	15	29 39	29 28	53 35	15	6	11	3
20	2 23	31 40	10	20	29 45	29 30	53 41	10	8	11	4
25	2 23	31 41	5	25	29 53	29 34	53 48	5	10	11	5
1	0	2 24	31 42	0	11	29 40	53 57	0	12	11	6
5	2 24	31 43	25	5	30 11	29 46	54 7	25	14	11	7
10	2 24	31 45	20	10	30 22	29 51	54 18	20	16	11	8
15	2 24	31 47	15	15	30 36	29 58	54 32	15	18	11	9
20	2 25	31 49	10	20	30 50	30 5	54 45	10	20	11	10
25	2 25	31 51	5	25	31 6	30 14	55 0	5	22	11	10
2	0	2 25	31 53	0	10	31 23	55 16	0	24	10	11
5	2 26	31 56	25	5	31 42	30 32	55 33	25	26	10	12
10	2 26	31 59	20	10	32 1	30 42	55 51	20	28	10	13
15	2 26	32 1	15	15	32 23	30 52	56 10	15	30	10	14
20	2 27	32 4	10	20	32 45	31 3	56 30	10	32	10	15
25	2 27	32 7	5	25	33 8	31 14	56 49	5	34	9	16
3	0	2 28	32 10	0	9	33 32	57 9	0	36	9	16
5	2 28	32 13	25	5	33 56	31 36	57 29	25	39	9	18
10	2 28	32 15	20	10	34 21	31 47	57 49	20	42	9	19
15	2 29	32 18	15	15	34 45	31 58	58 9	15	45	8	20
20	2 29	32 21	10	20	35 8	32 9	58 28	10	48	7	21
25	2 30	32 23	5	25	35 31	32 20	58 47	5	51	7	22
4	0	2 30	32 26	0	8	35 54	59 5	0	54	6	23
5	2 31	32 29	25	5	36 14	32 39	59 23	25	57	6	23
10	2 31	32 31	20	10	36 34	32 48	59 40	20	60	5	24
15	2 31	32 33	15	15	36 53	32 57	59 55	15	63	5	25
20	2 32	32 35	10	20	37 10	33 5	60 10	10	66	4	26
25	2 32	32 37	5	25	37 24	33 12	60 23	5	69	4	26
5	0	2 32	32 39	0	7	37 39	60 34	0	72	3	27
5	2 32	32 40	25	5	37 50	33 24	60 44	25	75	3	27
10	2 33	32 41	20	10	38 0	33 28	60 52	20	78	2	27
15	2 33	32 42	15	15	38 6	33 31	60 58	15	81	1	28
20	2 33	32 43	10	20	38 14	33 34	61 3	10	84	0	28
25	2 33	32 43	5	25	38 17	33 36	61 6	5	87	0	28
6	0	2 33	32 43	0	6	38 18	61 7	0	90	0	29

The TIMES between the true Ecliptic Conjunction or Opposition of the Sun and Moon, and the Middle of Eclipse, or greatest Obscuration, at the nearest Dist. of their Centers. Half of which Times, with the same Signs, will be nearly the Difference between the Orbit Syzygy and said Middle of the Eclipse.

D à 8.		True Hourly MOTION of the MOON from the SUN.										D à 8.	
Sig. o. 6.		27'	28'	29'	30'	31'	32'	33'	34'	35'	36'	Sig. 5. 11.	
0	om os	om os	om os	om os	om os	om os	om os	om os	om os	om os	om os	0	
1	1 14	1 11	1 8	1 6	1 4	1 2	1 0	0 58	0 55	0 52	0 49	30	
2	2 28	2 22	2 17	2 12	2 7	2 3	1 59	1 55	1 52	1 48	1 45	29	
3	3 41	3 33	3 25	3 17	3 10	3 4	2 58	2 52	2 47	2 42	2 36	28	
4	4 54	4 43	4 32	4 23	4 13	4 5	3 57	3 49	3 42	3 36	3 29	27	
5	6 7	5 53	5 40	5 28	5 16	5 5	4 55	4 46	4 38	4 29	4 20	26	
6	7 19	7 2	6 46	6 32	6 18	6 5	5 54	5 43	5 32	5 22	5 12	25	
7	8 30	8 10	7 52	7 36	7 20	7 5	6 52	6 39	6 26	6 15	6 4	24	
8	9 41	9 18	8 58	8 39	8 21	8 4	7 49	7 34	7 20	7 7	6 24	23	
9	10 51	10 26	10 3	9 41	9 22	9 3	8 45	8 28	8 13	7 58	7 42	22	
10	12 0	11 32	11 7	10 43	10 21	10 1	9 41	9 23	9 5	8 49	8 32	21	
11	13 8	12 38	12 10	11 44	11 20	10 57	10 36	10 16	9 57	9 39	9 21	20	
12	14 16	13 43	13 13	12 44	12 18	11 53	11 30	11 8	10 48	10 28	10 9	19	
13	15 22	14 46	14 14	13 43	13 15	12 48	12 23	11 59	11 38	11 16	10 54	18	
14	16 26	15 48	15 13	14 40	14 10	13 42	13 15	12 50	12 27	12 4	11 49	17	
15	17 29	16 49	16 11	15 37	15 4	14 34	14 6	13 39	13 14	12 50	12 24	16	
16	18 30	17 48	17 9	16 32	15 58	15 26	14 56	14 27	14 1	13 35	13 8	15	
17	19 30	18 46	18 5	17 26	16 50	16 16	15 44	15 15	14 47	14 20	13 52	14	
												13	

N. B. As hourly Mot. $\downarrow$ à $\odot$: 60 Min. of Time :: Reduction to Ecliptic in Min. and Sec. of Deg. : Reduc. in Min. and Sec. of Time, from the Orbit to the Ecliptic Syzygy, and also to the Middle of the Eclipse. — Which being contrarily situated from the Orbit Syzygy, double the said Time will be nearly the Reduction from Ecliptic Conjunction or Opposition to Middle of Eclipse; having the same Signs with the Reduction Equation to the Middle of Eclipse from the Orbit Syzygy. And therefore the Reduction from Middle of the Eclipse to the Orbit Syzygy, or from the Orbit to the Ecliptic Syzygy, by Half the whole Equation, must be with a contrary Sign.

The

The ANGLES which the Way of the MOON from the SUN makes with a Circle of LATITUDE:
In ECLIPSES of those LUMINARIES.

D & Q.		True hourly MOTION of the MOON from the SUN.										D & Q.	
Sig. 0.6.		27'	28'	29'	30'	31'	32'	33'	34'	35'	36'	Sig. 5. 11.	
0	84 14	84 15	84 16	84 17	84 18	84 18	84 19	84 20	84 20	84 21	84 21	30	
1	84 14	84 15	84 16	84 17	84 18	84 18	84 19	84 20	84 20	84 21	84 21	29	
2	84 14	84 15	84 16	84 17	84 18	84 18	84 19	84 20	84 20	84 21	84 21	28	
3	84 14	84 15	84 16	84 17	84 18	84 18	84 19	84 20	84 20	84 21	84 21	27	
4	84 15	84 16	84 17	84 18	84 18	84 19	84 20	84 21	84 21	84 22	84 22	26	
5	84 15	84 16	84 17	84 18	84 19	84 20	84 21	84 21	84 22	84 22	84 23	25	
6	84 16	84 17	84 18	84 19	84 20	84 21	84 22	84 22	84 23	84 23	84 24	24	
7	84 17	84 18	84 19	84 20	84 21	84 22	84 23	84 23	84 24	84 24	84 25	23	
8	84 18	84 19	84 20	84 21	84 22	84 23	84 24	84 24	84 25	84 26	84 26	22	
9	84 19	84 20	84 21	84 22	84 23	84 24	84 25	84 25	84 26	84 27	84 27	21	
10	84 20	84 21	84 22	84 23	84 24	84 25	84 26	84 26	84 27	84 28	84 28	20	
11	84 21	84 22	84 23	84 24	84 25	84 26	84 27	84 27	84 28	84 29	84 29	19	
12	84 22	84 23	84 24	84 25	84 26	84 27	84 28	84 28	84 29	84 30	84 30	18	
13	84 24	84 25	84 26	84 27	84 28	84 29	84 30	84 30	84 31	84 32	84 32	17	
14	84 25	84 26	84 27	84 28	84 29	84 30	84 31	84 31	84 32	84 33	84 33	16	
15	84 27	84 28	84 29	84 30	84 31	84 32	84 33	84 33	84 34	84 35	84 35	15	
16	84 29	84 30	84 31	84 32	84 33	84 34	84 35	84 35	84 36	84 37	84 37	14	
17	84 31	84 32	84 33	84 34	84 35	84 36	84 37	84 37	84 38	84 39	84 39	13	

The Complement of these Angles to 90° shews the Angles that the Way of the Moon from the Sun makes with the Ecliptic.

EXAMPLE. I. To find the mean Time of the mean and true Orbit New Moon for the Month of May, in the 585th Year current before Christ?

By the Nature of Construction of the Tables, 1 Year current before Christ is represented by 0, 101 before Christ, by 100, in the Tables, &c. each current Year before Christ by 1 less in the Tables, as we have noticed in our foregoing Improvements of Chronology; consequently the 585th Year current before Christ is represented by 584 in the Tables. And, since 584 before, + 1716 since Christ = 23 complete Centuries, or 2300 Years, by the Tables, from the Beginning of the 585th Year current before Christ, to the Beginning of the 1716th Year current since Christ. Therefore, by Tab. P. 385.

Dates, &c.	1716 since Chr. O.S. Jan.	Sub. 2300 Yrs Mot.	Rem. 585 bef. Chr. Jan.	May Biff. 1 Day less.	M. Time Mean new D, May	Eq. T. bet. M. and tr. D .. +	M. Time tr. new D req. May	in 585 Yr. cur. bef. Chr. O.S.	By Tab. P. 390, D Lat. N.	1716 since Chr. O.S. Jan.	Sub. 2300 Yrs Mot.	Rem. 585 bef. Chr. Jan.	May Biff. 1 Day less.	M. Time Mean new D, May	Eq. T. bet. M. and tr. D .. +	M. Time tr. new D req. May	in 585 Yr. cur. bef. Chr. O.S.	By Tab. P. 390, D Lat. N.	1716 since Chr. O.S. Jan.	Sub. 2300 Yrs Mot.	Rem. 585 bef. Chr. Jan.	May Biff. 1 Day less.	M. Time Mean new D, May	Eq. T. bet. M. and tr. D .. +	M. Time tr. new D req. May	in 585 Yr. cur. bef. Chr. O.S.	By Tab. P. 390, D Lat. N.
	12 16 4 48	11 5 57 54	1 10 6 54	26 15 40 15	28 1 47 9	.. 2 13 23	28 4 0 32	.. 20 0	.. 20 0	12 16 4 48	11 5 57 54	1 10 6 54	26 15 40 15	28 1 47 9	.. 2 13 23	28 4 0 32	.. 20 0	.. 20 0	12 16 4 48	11 5 57 54	1 10 6 54	26 15 40 15	28 1 47 9	.. 2 13 23	28 4 0 32	.. 20 0	.. 20 0
	6 24 35 33	11 19 15 5	7 5 20 28	4 25 31 37	0 0 52 5	6 15 9 13	Arg. 6. Equat. Supplemental, P. 74.	5 15 42 35	Eq. Sun - 1 43	6 24 35 33	11 19 15 5	7 5 20 28	4 25 31 37	0 0 52 5	6 15 9 13	Arg. 6. Equat. Supplemental, P. 74.	5 15 42 35	Eq. Sun - 1 43	6 24 35 33	11 19 15 5	7 5 20 28	4 25 31 37	0 0 52 5	6 15 9 13	Arg. 6. Equat. Supplemental, P. 74.	5 15 42 35	Eq. Sun - 1 43
	2 12 36 6	1 5 58 33	1 6 37 33	4 9 5 2	5 15 42 35	Eq. Ap. cont. + 17	5 15 42 35	Eq. Sun - 1 43	5 15 42 35	2 12 36 6	1 5 58 33	1 6 37 33	4 9 5 2	5 15 42 35	Eq. Ap. cont. + 17	5 15 42 35	Eq. Sun - 1 43	5 15 42 35	2 12 36 6	1 5 58 33	1 6 37 33	4 9 5 2	5 15 42 35	Eq. Ap. cont. + 17	5 15 42 35	Eq. Sun - 1 43	5 15 42 35
	2 25 56 32	7 25 26 59	7 0 29 33	5 3 21 10	0 3 50 43	Eq. Q. cont. - 8	0 3 50 35	Eq. Sun - 1 43	0 3 48 52	2 25 56 32	7 25 26 59	7 0 29 33	5 3 21 10	0 3 50 43	Eq. Q. cont. - 8	0 3 50 35	Eq. Sun - 1 43	0 3 48 52	2 25 56 32	7 25 26 59	7 0 29 33	5 3 21 10	0 3 50 43	Eq. Q. cont. - 8	0 3 50 35	Eq. Sun - 1 43	0 3 48 52
	6 Equat. doubled, P. 74.	Equat. at N. and F. D. ?	P. 390. . . .	Sum Equat. D . . .	Equat. Sun. P. 12. . .	Sum Eq. = M. Dist. D à C. +	at true Conj. . . Hours 2 . .	Minutes 13 . . .	Seconds 23 . . .	6 Equat. doubled, P. 74.	Equat. at N. and F. D. ?	P. 390. . . .	Sum Equat. D . . .	Equat. Sun. P. 12. . .	Sum Eq. = M. Dist. D à C. +	at true Conj. . . Hours 2 . .	Minutes 13 . . .	Seconds 23 . . .	6 Equat. doubled, P. 74.	Equat. at N. and F. D. ?	P. 390. . . .	Sum Equat. D . . .	Equat. Sun. P. 12. . .	Sum Eq. = M. Dist. D à C. +	at true Conj. . . Hours 2 . .	Minutes 13 . . .	Seconds 23 . . .

* At this Time a great Eclipse of the Sun happened, that put an End to the Battle then fighting between the Medes and Lydians, on which Peace ensued. — This Eclipse happened above 2 Hours later at the Meridian where the Battle was fought, near the River Halys, the Boundary betwixt Media and Lydia, in Asia Minor, in the 4th Year of the 48th Olympiad, according to Herodotus, as predicted by Thales, of Miletus, or Sardis, in Asia. — But Ferguson's Astr. P. 168, will have this Eclipse to be in 603 Year bef. Chr. May 18, 8h 30m Forenoon, (London-Meridian) 18 Yrs sooner. The Sun was then above 4 Degrees from the Moon's Node.

The Dominical Letters for this Year (Bisextile) being FE, according to Table of Chronology, P. 148, May 28, in the 585th Year current, before Christ, the Day on which this Eclipse happened was on a Wednesday. In which Year the Cycle of the Sun was 13, Golden Number 6, and Roman Indiction 4; being in the 4129th Year of the Julian Period. And 1716, since Christ, being in the 6429th Year of the Julian Period, the Difference = 2300 Years aforesaid.

Mr. C. Brent, (in P. 341, Compend. Astron.) makes the 28th of May, on which this Eclipse happened, to be on a Monday, instead of on Wednesday, (observed also in Ferguson's Astronomy) which we should not have here noticed, had he not disputed (in Lond. Mag. for April, 1760) the Truth, by pretending "It was 585 Years complete, before Christ, (i. e. 586 current) then not Leap-Year; when G was the Dominical Letter." That "Otherwise the Eclipse (according to Newton's Chronology) could not have been on May 28." Whereas there was no Eclipse in the 586th Year current before Christ, on May 28, when the Dominical Letter was G and Cycle of Sun 12, (his Rule, P. 341, Compend. Astr. for finding the Sun's Cycle, not true) so it must be on the 28th of May, in the 585th Year current, before Christ, on a Wednesday, according to Newton's Chronology when this Eclipse happened; as the Computation proves above; and as his own Computation proves (P. 340. Comp. Astr.) of 2320 Years completed back from May 1, 1736, since Christ, to May 1, 585, before Christ; who has the same Mistake, P. 345, 346, Compend. Astr. making the 12th of March, in the 4th Year before Christ, on Sunday, the Dominical Letters BA (though not Leap-Year) for the 12th of March, in the 5th Year, before Christ.

For Years and Months, since Christ, you must add or subtract the Years and Months Motion forward or backward, to and from the Years, Time over and above, and Places, answering to January in the Eighteenth Century, P. 385.

EXAMPLE

EXAMPLE II. To find the true Places of the Sun and Moon at the Time of the foregoing Eclipse of the Sun, according to the Moon's true Orbit-Conjunction?

RULE. Compute the Sun's Apogee to the Time of the mean new Moon, and annex the other mean Places, foregoing, to that Time. Then add the Motion answerable to the Equation of Time, if that adds; or subtract, if that subtracts; for the mean Places, at the true Syzygy. Then add the Sun's Apogee and Anomaly for the Sun's mean Place, from which taking the Sun's Ap. and Sun from Moon's Node, there will remain the Places of the Moon's Apogee and Node. And the Sun's Apogee added to Moon's Apogee will give the Moon's Anomaly. The Sum of which and Sun's Ap. = M. Pl. $\odot$; with which connect the proper Equations as below for the Moon's true Place. The Sun's true Place being had from the Sun's Equation connected with his mean Place, before found.

The METHOD of COMPUTATION below is intirely NEW.

1716 since Ch. May 28	Sun's Apog.	Sun's M. An.	Sun's $\odot$'s Ap.	Sun's $\odot$.	$\odot$ to $\odot$.
Sub. 2300 Years Motion —	3° 80' 18"	s o / "	s o / "	s o / "	s o / "
585 bef. Ch. May 28 ^d 1 ^h 47 ^m 9 ^s	1 28 42 48	o o 52 5	5 15 42 35	o 3 50 43	o o o o
Add for 12 Equations. } M. 13 S. 23		. . . 4 56	. . . 4 22	. . . 5 12	o 1 o 57
585, May 28 ^d 4 ^h 0 ^m 32 ^s Past, — 1 21		 33	 29	 35	6 48
	Sun's M. An.	o o 57 34	5 15 47 26	o 3 56 30	o 1 7 45
	Sun's Ap. +	1 28 42 48	1 29 40 22	1 29 40 22	5 15 47 26
	Sun's M. Place	1 29 40 22	Ap. 8 13 52 56	8 1 25 43 52	5 16 55 11
	Sun's Equation	— 1 54	— 1 28 42 48	⊖ 1 29 39 9	8 11 52 56
	Sun's true Pl.	1 29 38 28	6 15 10 8	o 3 55 17	2 o 48 7
	$\odot$ tr. Orb Pl.	1 29 39 9	Arg. 6 Eq ⁿ . P. 74.	[Arg. $\odot$'s Lat.	
	Diff.	+ 41	5 15 45 32	1 Eq. P. 61 . . . + o o 11	
			Arg. cent. Eq. past	6 Eq. doubled, P. 74. — o 1 15	
			Conjunction, = 1 ^m	Eq. at New and Full — 1 7 54	
			21 ^s to be deducted.		
				Moon's true Orb. Place 1 29 39 9	
				Moon's Latitude . . . 20 34	N.

PROPOSITION.

The Sum of the Moon's Equations in Syzygies connected by a contrary Sign with the Sun's Equation, and its proper Sign, will be the Distance of Time, nearly, in mean Motion, of the Moon from the Sun, on this Side, or beyond, a true Conjunction or Opposition of the Sun and Moon, from the Time of a mean Syzygy.

DEMONSTRATION. Let $\odot$ and $\odot$ represent the Sun and Moon's mean Places, respectively, at the Time of a true Orbit-Conjunction or Opposition of the Moon and Sun. $\odot - \odot$ will be the mean Distance of the Moon from the Sun at that Time. Let $\pm S$ denote the Sun's Equation, and $\pm M$ denote the Sum of the Moon's Equations, at the Time of a mean Syzygy. Now because there is but small Difference between the Equations for the Sun and Moon's mean Places, at the Time of a mean and true Syzygy (especially when the Interval is small.) Therefore, $\odot \pm M \text{ min. } \odot \pm S = o$, or 6 Signs, nearly at the true Syzygy. Consequently, Mean Distance = $\odot - \odot = \pm S \mp M + o$ or 6 Signs, by Transposition. That is, the mean Distance of the Moon from the Sun, at the true Syzygy nearly = Sum of the Sun's Equation S , with its proper Sign, and the Moon's Equation M , with a contrary Sign, $+ o$, or 6 Signs, at mean Conjunction or Opposition, respectively. And therefore, that Sum reduced to Time, in mean Motion of the Moon from the Sun, reckoned from o , or 6^s $\odot$ from $\odot$ at the mean Syzygy, and that Interval of Time added or deducted to or from the Time* of mean Syzygy, according as the Moon's true Place is short of, or beyond, the Sun's true Place, or its Opposite, as the Sign of those Equations in one Sum directs, will give the mean Time of the true Syzygy, required, very nearly: And the more nearly the less the Interval happens to be; because then the Moon and Sun's Equations at mean and true Syzygy are the nearer alike. Q. E. D.

If greater Accuracy be required, the Equations of the Sun and Moon may be again found from the Arguments to the last near Time of the true Syzygy, and the Difference between the Sum of those, and the Sum of the former Equations taken, and reduced to Time, in Motion of the Moon from the Sun, and added to, or subtracted from the last near Time, according as the Sign of the Difference of the Sum of the former, taken from the Sum of the latter of those Equations, directs, will give the correcter Time of the true Syzygy, and so on. Or the Sun and Moon's true Places may be found to the nearly true Syzygy, and the Difference of those Places, reduced to Time, in mean Motion $\odot$ fr. $\odot$, and added to, or subtracted from, the Time of the nearly true Syzygy, according as the Moon's true Place is behind or before the Sun's true Place, will give the correcter Time, required.

PRECEPTS for finding the TIMES and PLACES of ORBIT NEW and FULL MOONS.

TO find the mean TIME of the nearly true ORBIT CONJUNCTION or OPPOSITION of the SUN and MOON?

From Table, (P. 385) of the 1st mean New Moon in January, according to Old Style, collect the Days, and Parts of a Day, answering to the Year and Month (taking out a Day less in Leap-Year after February) and collect also the Degrees and Parts of a Degree of the Sun's mean Anomaly, mean Distance of the Sun from the Moon's Apogee, and Node; which are the same for Leap as for common Years. Or, if the Time of the next following or foregoing Full Moon, in any Month, is required, you must add or subtract, to or from the Time, and Places, of the mean New Moon, respectively, the Time and Motion for Half a synodical Month, to be found at the Bottom of the Months.

From Table, I. Equation $\odot$, (P. 61 and 62) take out the Equations of the Moon, Apogee, and Node, answering to the Sun's mean Anomaly, which apply to the former Places, with a contrary Sign to that directed in the Tables, and you will have the Sun's Distance from the Apogee and Node, once equated. Add to these, or subtract therefrom, (according to its Title) the Sun's Equation, and you will have the true Dist. $\odot$ from $\odot$'s Apogee, and true Dist. $\odot$ from $\odot$ 1. equated.

The mean Distance of the Sun from the Moon's Apogee 1 equated, being subtracted from the Sun's mean Anomaly, the Remainder will be the Argument of the VI. Equation supplemental, P. 74, at New Moon; but at Full Moon, this Argument, and also that of the Sun from the Moon's Apogee, for the central Equation in Syzygies, must be increased by 6 Signs added to each; because the Arg. 6 Equat. at Opposition, = $\odot$ to $\odot + \odot$ Ap. — $\odot$ Ap. = 6^s + $\odot$ Ap. — $\odot$ Ap. And Arg. central Equation, at Opposition, = $\odot$ Anom. + 6 Signs = $\odot$ from $\odot$'s Ap. + 6 Signs.

In P. 61 and 62, find 1st, or annual Equation of the Moon, with a contrary Sign, according to the Sun's mean Anomaly. In P. 74, find 6. Equation supplemental, answering to the proper Argument; which Equation you must double, and connect its contrary Sign. In P. 390, find the central Equation, at New or Full, from its proper Argument, with a contrary Sign, the Sum of these 3 Equations with their contrary Signs, (or, which amounts to the same Thing, the Sum of these 3 proper lunar Equations with a contrary Sign) connected with the Equation of the Sun's Center (which you must also find) with its proper Sign, will give the mean Distance of the Moon from the Sun, very nearly, at a true Conjunction or Opposition; which, reduced to Time, (by Tab. P. 389, of M. Mot. $\odot$ fr. $\odot$) will be nearly the true Equation of mean Time, with its proper Sign, for the Interval of Time, to be added to, or subtracted from, the Time of the mean New or Full Moon, at o or 6 Signs mean Dist. $\odot$ to $\odot$, for the mean Time of the nearly true Orbit Conjunction or Opposition, required.

* In P. A S Halley's Astron. Tables, his Precept-Writer applies the Excess of the Sun's Equation, and Sum of the Moon's Equations, (reduced to Time, in mean Mot. $\odot$ to $\odot$) to the Time of the mean for that of the true Syzygy, without considering the Sign of the greater, and therefore errs in his Account of the true Syzygy preceding the mean, and in the mean preceding the true.

PRECEPTS for finding the TIMES and PLACES of ORBIT NEW and FULL MOONS.

The Place of the ☾ subtracted from the Sun's true Place, in Conjunction, but from its opposite Place in Opposition, will give the *Argument* of the Moon's Latitude, in Syzygies; by which, and *Tab. P. 390*, the Moon's Latitude is found, and her Place in the Ecliptic, by *Reduction* from her Orbit-Place; both which Orbit and Ecliptic Places in the Syzygies are nearly the same.

But the Time between Ecliptic Conjunction or Opposition, and the Middle of Eclipses, is found by *Tab. P. 391*, Half of which Time is the Difference between an Orbit Syzygy and Middle of an Eclipse.

To determine the Sun and Moon's true PLACES, correspondent to the mean Time of the near or true Orbit CONJUNCTION and OPPOSITION?

Compute the Sun's Apogee (from *Tables* at the Beginning of this *Work*) to the Time of the mean New or Full Moon, and annex the other mean Places already found to that Time; with all which Places (of Sun's M. Anom. Sun from Moon's Ap. and ☾, and Moon from Sun, or 6° or 6s) connect their respective Motions for the Equation of Time, or Interval, between the mean and nearly true Orbit Syzygy, according to the proper Sign. Then add the Sun's Apogee and mean Anom. ☽, for the Sun's mean Place, from which taking the Sun from the Moon's Apogee, and also Sun from Moon's ☾, there will remain the mean Places of the Moon's Apogee and Node. And the Sun from the Moon's Apogee, added to ☽ from ☾, will give the Moon's mean Anomaly. The Sum of which Moon's Anomaly, and Moon's Apogee (before found) will give the mean Place of the Moon in her Orbit, with which if you now connect her proper Equations 1, 6, and the central in Syzygies, taken out from the true or equated Arguments, (in *P. 61, 74, and 390*) you will have the Moon's nearly true Place in her Orbit.

The Sun's true Place being here also had from the Equation of his Center, the Distance of the Moon's from the Sun's true Place, i. e. the Remainder after the Sun is taken from the Moon's Place (being a small Arc) turned into Time (by *Tab. mean Mot. ☽ à ☾*) and added to or subtracted from the former Time of the Orbit Syzygy, according as the Moon's true Place is, by the Length of that Arc, behind or before the Sun's Place, will give the correcter Time of the Orbit Syzygy, required.

N. B. In Orbit Opposition, or Full Moon, you must add 6 Signs to the Sun from the Moon's Apogee, for the Argument of the Moon's central Equation at that Time; and also 6 Signs must be added for the Argument of the 6. Equation, in Opposition.

REMARK. In Dr. Halley's *Analytic Tables*, for computing the Moon's Place, at her Orbit-Conjunction or Opposition, (than which our common *Tables* are not less expeditious) the 1st or annual Equation of the Moon, Apogee, and ☾, being connected or compounded with the Moon's mean Motion, Apogee, and ☾, answering to the Sun's mean Anomaly.

And in Imitation of Dr. Halley, we have also as usefully compounded the 1st semiannual Equation of the Moon, with the Moon's central Equation in Syzygies; having both the same Arg. annum. (See *P. 390*) Which central Equation is computed by finding the mean from the true Anomaly of the Moon, and taking their Difference.

EXAMPLE III. To find the mean Time of the mean and nearly true New Moon, or Orbit-Conjunction, in the Month of April 1764, since Christ New Style?

☞ 11 Days must either be added to the Day of Old Style, in January, for the Day of New Style, correspondent, among the Radixes; or else a Month must be so taken, that when 11 Days are added to its Days corresponding, the Sum may come out in April, or the Month wherein the New or Full Moon is required, the nearest possible to the first Day of that Month.

Dates, &c.	1st m. n. ☽ Jan.	Sun's M. Anom.	Sun à Moon's Ap.	Sun à ☾.	Equations of the Moon with contrary Signs.
By <i>P. 385</i> . 1764 fin. Ch. O. S.	21 20 10 59	7 3 10 34	9 17 46 25	10 4 14 29	1 Equat. ☽, <i>P. 61</i> . + 0° 11' 48"
Biff. Feb. (adding 11 Ds) Mar.	10 1 28 6	1 28 12 39	1 21 38 1	2 1 20 28	6 Eq. doubled, <i>P. 74</i> . + 0 4 28
Reduces to Apr. N. S.	0 21 39 5	9 1 23 13	11 9 24 26	0 5 34 57	Eq. at N. ☽, <i>P.</i> . - 1 44 2
Equat. bet. M. and true ☽	+ 55 2	- 11 9 4 27	1 Eq. cont. - 19 59	1 Eq. cont. + 9 29	- 1 27 46
M. Time Orbit N. ☽, April, 1764, N. S. required.	0 22 34 7	9 22 18 46	11 9 4 27	0 5 44 26	Sun's Equat. <i>P. 12</i> . - 1 55 43
According to <i>Ferguson's Astr.</i> <i>P. 198</i>	0 22 15 0	Arg. 6. Equat. supplemental, <i>P. 74</i> . at New Moon.	☾ Eq. + 1 55 43	☾ Eq. + 1 55 43	Sum, mean Dist. ☽ ☽ à ☾, in Syzygy. } + 27 57
Diff.	- 19 7		11 11 0 10	0 7 40 9	Minutes 55 . . 27 56
			True Arg. central Eq. at New Moon.	True Sun à ☾, 1 eqd. the Sun eclipsed considerably.	Seconds 2 . . 1
					Equation or Interval of Time, [+.

EXAMPLE IV. To find the true Orbit Places of the Sun and Moon, at the Time of the foregoing Eclipse of the Sun, according to the Moon's nearly true Conjunction?

	Sun's Apogee.	Sun's M. Anom.	Sun à ☽'s Ap.	Sun à ☾.	☽ à ☾.
1764, March 31, N. S.	3 8 49 18	9 1 23 13	11 9 24 26	0 5 34 57	0 0 0 0
21 ^h 39 ^m 5 ^s		2 16	2 0	2 23	27 57
Add for Eq. M. 55 S. 2					
22 34 7	Sun's M. An.	9 1 25 29	11 9 26 26	0 5 37 20	0 0 27 57
Short + 18	Sun's Ap. +	3 8 49 18	0 10 14 47	0 10 14 47	11 9 26 26
					+
1764, Apr. 0 ^d 22 34 25	Sun's M. Pl.	0 10 14 47	Ap. 1 0 48 21	☾ 0 4 37 27	11 9 54 23
Nearer Time.	Sun's Equat.	+ 1 55 42	- 3 8 49 18	☾ 0 12 10 29	1 0 48 21
					☽'s M. Anom.
	Sun's true Pl.	0 12 10 29	9 21 59 3	0 7 33 2	0 10 42 44
	☽ true Orb. Pl.	0 12 10 20	1 Eq. Ap. + 19 59	[Arg. ☽'s Lat. 1 Eq. prop. <i>P. 61</i> . 6 Eq. doubled, <i>P. 74</i> . Cent. Eq. at N. ☽.	☽'s M. Pl.
		Diff. - 9	9 22 19 2		- 11 48
		= 18 ^s short of Conjunction.	Arg. 6. Equat. 11 11 2 9		- 4 28
			Arg. central Eq.	☽'s true Orb. Pl.	+ 1 43 52
				0 12 10 20	Reduc. - 1' 48".
				Moon's Lat. 39 31	N.

☞ If you find the Sun and Moon's true Places to the Time of mean Syzygy, and reduce what the Moon is short of, or beyond, a Conjunction or Opposition into Time, in mean Mot. ☽ à ☾, then this Time being added to or subtracted from the Time of mean Syzygy, it will give you nearly the Time of true Syzygy, required.

EXAMPLE

EXAMPLE V. To find the mean Time of the mean and nearly true Opposition or Full Moon, in the Month of April, 1764?

1764.	1st m. n. D Apr.	Sun's M. Anom.	Sun à Moon's Ap.	Sun à Q.	Equations of the Moon with contrary Signs.
From above, April, N. S.	0 21 39 5	9 1 23 13	11 9 24 26	0 5 34 57	1 Equat. D, P. 61 + 0 11' 19"
1/2 Rev. or synod. Month, +	14 18 22 2	0 14 33 10	0 12 54 30	0 15 20 7	6 Eq. doubd, P. 74. — 0 2 24
Mean Opposition, April	15 16 1 7	9 15 56 23	11 22 18 56	0 20 55 4	CenEq. at Full, p. 390 + 0 30 5
Eq. bet. M. and true Oppn	+ 4 45 27	— 11 21 59 50	1 Eq. cont. — 19 6	1 Eq. cont. + 9 5	
M. Time true Opposition, Apr.		9 23 56 33	11 21 59 50	0 21 4 9	Sum, + 39 0
1764, N. S. required . .	15 20 46 34	+ 6	⊙ Eq. + 1 44 57	⊙ Eq. + 1 44 57	Sun's Equat. + 1 44 57
According to Ferguson's Afr.		3 23 56 33	11 23 44 47	0 22 49 6	Sum, M. Dif. D à ⊙, + 2 23 57
P. 199.	15 20 56 0	Arg. 6. Equat. P.	+ 6	True Sun à Q. Sun	[in Syzygy.
Error	+ 19 26	74. at Full D.	5 23 44 47	being above 12° from	Hours 4 . . 2 1 54
			Arg. central Equat.	the Q, here happens	22 3
			at Full Moon.	no Eclipse of the D,	Minutes 45 . . 21 50
				as near the opposite	Seconds 27 . . 13
				Q.	Equat. or Interval of Time, +.

N. B. You may take out the 1st, 6th, and central Equations of D as they are, and then connect the Sum thereof, changing the Sign, with the Equation of the Sun's Center, for the mean Dist. D à ⊙, to be turned into Time, and used as above, connected, in its proper Sign, with the Time of the mean Syzygy.

EXAMPLE VI. To find the true Places of the Sun and Moon, at the Time of the foregoing nearly true Orbit Opposition of the Sun and Moon?

1764, Apr. 15, N. S.	Sun's Apogee.	Sun's M. Anom.	Sun à D's Apogee.	Sun à Q.	D à Sun.
16h 1m 7s	3 8 49 20	9 15 56 23	11 22 18 56	0 20 55 4	0 0 0 0
Add Equat. {	4	 9 51	8 44	10 23	0 2 1 54
. . 45 27	By Tab. P. 389	 1 52	1 39	1 58	23 5
20 46 34	Sun's M. An	9 16 8 6	11 22 29 19	0 21 7 25	0 2 24 59
Short + 2 45	Sun's Apogee	3 8 49 20	0 24 57 26	0 24 57 26	11 22 29 19
1764, Apr. 15d 20 49 19	Sun's M. Pl.	0 24 57 26	Ap. 1 2 28 7	⊙ 0 3 50 1	11 24 54 18
Nearer Time.	Sun's Equat.	+ 1 50 36	1. Eq. + 19 6	⊙ 0 26 48 2	1 2 28 7
	Sun's true Pl	0 26 48 2	1 2 47 13	6 22 58 1	0 27 22 25
	D tr. Orb. Pl.	0 26 46 38	— 3 8 49 20	[Arg. Lat.	D's M. Anom.
		— 1 24	9 23 57 53	1 Equat. D . . .	D's M. Pl.
		= 2m 45s short	+ 6	6 Eq. D doubled	. . . — 11 19
		of Opposition.	3 23 57 53	Eq. Center at Full	+ 4 24
			Arg. 6. Equat.	True Orb. Pl. D	— 28 52
			11 24 0 49	Moon's Lat.	0 26 46 38
			+ 6	Sun à D Ap.	.. 1 57 17
			5 24 0 49		Reduc. — 5'.
			Arg. cent. Equat. at		S.
			Full Moon.		

N. B. The Arg. 6. Equat. = { D's Ap. 1. eqd. — ⊙'s Ap. at New;
but + 6s at Full Moon.

Arg. Central Eq. = { Pl. Sun — D's Apog. 1. equated, at New;
but + 6s at Full Moon.

EXAMPLE VII. To find the Mean of Time of the Moon's mean and nearly true Orbit Conjunction with the Sun, (by Table, P. 383.) for the Month of May in the 585th Year current before Christ, according to the Moon's Acceleration, in Tab. P. 374?

See Precept in Tab. P. 389.	D à Sun.	Sun's M. Anom.	Sun à Moon's Ap.	Sun à Q.	By this Method* the Acceleration and Correction of M. Mo. D, Ap. and Q, is connected for the Begining of any Year, fin. Chr.
1716 since Christ, O. S.	7s 7° 44' 3"	6s 13° 5' 27"	2s 20 23' 58"	2s 13° 49' 19"	
2300 Years Motion —	7 12 52 19	11 8 9 46	0 26 8 31	7 13 46 1	
585 bef. Chr. current	11 24 51 44	7 4 55 41	1 6 15 27	7 0 3 18	Accel. Ap. and Q connected
Accelerat. & Cor. M. Mot. P. 374.	⊕ + 42 10*		D Ap. + 1 28 10*	Q — 44 54*	[with a contrary Sign.
Corrected . .	11 25 33 54	7 4 55 41	1 7 43 37	6 29 18 24	
May . .	0 22 53 23	3 28 16 20	3 14 54 32	4 4 37 57	
	0 48 27 17	11 3 12 1	4 22 38 9	11 3 56 21	The Mot. of Sun's M. An.
	12	0 27 36 47	0 24 29 35	0 29 5 53	Sun à D Ap. and Q are added
Remains . .	11 11 32 43	0 0 48 48	5 17 7 44	0 3 2 14	for Ds, Hrs. Min. &c. as below.
Days 28 . .	11 11 20 27	5 17 8 0	1 Eq. Ap. con. + 16	1 Eq. Q con. — 7	Days 28, Ho. 0, M. 24, S. 10.
	12 16	6 13 40 48	5 17 8 0	0 3 2 7	Mean Places correspondent.
Hours 0, Minutes 24	. . . 12 11	Arg. 6. Eq. P. 74.	Sun's Eq. — 1 36	Sun's Eq. — 1 36	Equations of Moon below.
Seconds 10	 5		1 17 6 24	0 3 0 31	1 Eq. D prop. P. 61. + 0 0' 10"
In 585, May 28d 0h 24m 10s	M. Time M. New Moon.		True Arg. cent. Eq. at	True Sun from Node,	6 Eq. D doub. P. 74. — 0 1 4
Eq. Time + 2 0 8	the Interval bet. M. and tr. N. D.		New Moon.	1 equated.	Cen. Eq. N. D p. 390 — 1 1 43
In 585, May 28d 2 24 18	M. Time true Orbit N. D, reqd.			Sun greatly eclipsed.	+ cont. . . Sum — 1 2 37
bef. Chr. 4 0 32	Computed in Ex. I. P. 392.				Sun's Equat. — 0 1 36
					+ Sig. D Eq. ?
					changed, Sum + 1 1 1
					Mean Dist. D à Sun, in Syzygy.
					Hours 2 . . 1 0 57
					Min. 0, Sec. 8 4
					Equat. or Interval of Time, +.

Diff. 1 36 14 sooner in 2300 Years Julian.

To which Time the Sun and Moon's true Places may be computed, as in Example II. Or you may first take the Reduc. D to Ecliptic, by Tab. P. 392. doubling the ' and " for Min. and Sec. Reduc. Time; with a contrary Sign.

EXAMPLE VIII. To find the mean Time of the mean and nearly true Orbit Opposition, or Full Moon, in March 1764; without Acceleration and Correction of mean Motion. See Tables, P. 388 and 389.

	Moon á Sun.	Sun's M. Anom.	Sun á Moon's Ap.	Sun á Node.	
1664 since Cbr. O. S.	5 8 51 6	6 13 35 5	0 17 58 57	4 27 39 0	
+100 Years Mot. . .	10 7 4 53	11 29 3 2	8 11 34 17	4 14 56 47	
March	11 29 15 15	1 28 9 1	1 22 34 48	2 2 16 39	
1764 March	3 15 11 14	8 10 47 8	10 22 8 2	11 14 52 26	Days 6, Motion:
	6	0 5 54 49	0 5 14 44	0 6 13 54	
Moon fr. Mean Opposit.	2 14 48 46	0 7 24	0 6 33	0 7 47	Hours 3.
N. S. 17. O.S. Days 6	2 13 8 40	0 42	0 37	0 44	Minutes 17, Seconds 4.
Hours 3	1 40 6	8 16 50 3	10 27 29 56	11 21 14 51	Places correspond. to Δ á Sun = 6s.
	1 31 26	10 27 10 23	1 Eq. con. — 19 33	1 Eq. con. + 9 17	Equations of the Moon below.
Minutes 17	8 40	9 19 39 40	10 27 10 23	11 21 24 8	1 Eq. Δ , P. 62. — 0° 11' 33"
	 8 38	+6	Sun's Eq. +1 53 18	Sun's Eq. +1 53 18	6 Eq. doubled, P. 74. + 0 4 33
Seconds 4	 2	3 19 39 40	10 29 3 41	11 23 17 26	Cent. Eq. Δ at Full — 2 23 29
		Arg. 6. Eq. P. 74.	+ 6	True Sun á Node.	+ contr. . . Sum — 2 30 29
d h m s				Where Moon 7° from	+ Sig. Δ Eq. changed, Sum + 4 23 47
In 1764, March 17 3 17 4, M. Time M. Full Moon.			4 29 3 41	opposite 8°. There-	Mean Dist. Moon á Sun, in Syzygy.
Equat. Time + 8 39 20 Interval bet. M. and tr. Opp.			Arg. Cent. Eq. at Full	fore Δ eclipsed being	Hours 8 . . 4 3 49
			Moon.	less than 12° á 8°.	Minutes 39 . . 19 48
In 1764, March 17 11 56 24 N. S. M. Time true Orbit					Seconds 20 . . . 10
[Opposition, required.]					

In *Ferguson's Astronomy*, (and also in a *New Work* bearing his Name we have seen at the Observatory at *Greenwich*, in *Manuscript*, examining the Times of *antient Eclipses*, referred to *Dr. Bradley's* Inspection and Approbation) we find a *similar Method* to that above, for determining the mean Time of the mean and true Orbit Conjunction and Opposition of the Sun and Moon. But this Author (*erroneously*) gives the Sum of the Moon's Equations with a *contrary Sign*, and of the Sun's Equation with its *proper Sign*, reduced to Time in mean Motion of the Moon (as his *three lunar Equation Tables and Sun's Equation Table*, there reduced to Time, evince) instead of reducing those Equations to Time (as we have demonstrated they should be reduced) in mean Motion of the Moon from the Sun. Who therefore is greatly wide of Truth in his computed Times of Lunations and Eclipses, as follows.

According to *Ferguson's Astronomy*.

	d	h	m	
1761 May 4 5 4 New Moon.				
18 10 24 Full Moon.				Δ Eclipse.
1764 March 31 22 15 New Moon.				
April 1 10 16 $\odot$ Eclipse.				
April 15 20 56 Full Moon.				
1801 March 13 9 13 Mean New.				
424 March 9 20 29 New Moon.				
721 March 8 7 36 Full Moon.				

According to *De la Caille's Astronomy*.

	d	h	m	s	
May 4 ^d 5 ^h 37 ^m 0 ^s New Moon.					
18 10 23 0 Full Moon. Δ Ecl.					
March 31 22 29 51 New Moon.					
April 1 10 43 0 $\odot$ Eclipse.					
April 15 22 38 0 Full Moon.					

For the Meridian of *Greenwich*.

If *Dr. Bradley* has not detected the foregoing Mistake in Judgment, we here observe it, that *Mr. Ferguson's Computations*, printed, or in *Manuscript*, may receive a general Revise and Correction; or new Edition of the whole Work.

* * Mr. *Ferguson* deducts a Day in his 3 and 4th Examples above; though he proposes to avoid that Inconvenience, by beginning his Computations at *March*, instead of *January*, where they will admit of it; which appear no Advantage to us; but rather the contrary. — He says, P. 201, of his *Astronomy*, "Our Saviour was born in a Leap Year," though the 1st Year current of Christ (at the Beginning of which our Saviour was born) was the 1st Year after Leap-Year, or after the Leap Year which was the 1st current before Christ; or why is 1 Year directed to be deducted out of the Years before Christ (before a Division of the Remainder by 4) to shew the 1, 2, 3, or 0 Year since Leap-Year, or when it is Bissextile; according as 3, 2, 1, or 0, remains? *Ferguson right*

PRECEPTS for finding the TIMES and PLACES of Ecliptic NEW and FULL MOONS.

To find the mean TIME of the true Ecliptic CONJUNCTION or OPPOSITION of the SUN and MOON?

RULE. To the mean Time of a mean, or that of a nearly true New or Full Moon (found by the foregoing Rules) compute the Sun and Moon's true Places, in the Ecliptic. And if they are *exactly* the same, or opposite, (*which seldom happens*) the Time of the true New or Full Moon is discovered. If they differ, note that Difference, and compute, again, the Sun and Moon's Places in the Ecliptic, for an Hour before or after the Time, for which you have their Places already computed; whereby you will gain the true hourly Motion of the Moon from the Sun in the Ecliptic.

Now say, As the hourly Motion of the Moon from the Sun is to 1 Hour or 60^m of Time :: so is the Distance of the Moon from the Sun, at mean or nearly true New or Full Moon (or Difference of their Places before noted) : to the Interval of Time between the mean, or near Ecliptic New or Full Moon. Which Time being added to, or subtracted from, the Time of the mean, or nearly true Conjunction or Opposition, according as the Moon's computed Place is short of, or beyond the Sun's computed Place, or its Opposite, at that Time, and you will have the mean Time (nearly) of the true Ecliptic Conjunction or Opposition of the Sun and Moon, required. When greater Exactness is wanted, compute again the Sun and Moon's true Places to the last found Time of Conjunction and Opposition, and if they are precisely the same, or opposite, you have your Desire; if otherwise, proportion for the Difference, to get the exact Time of Ecliptic Conjunction or Opposition.

* A true Ecliptic Conjunction of the Sun and Moon happens when the Sun's true Place is the same with the Moon's true Ecliptic Place. And a true Ecliptic Opposition happens when the Sun's true Place is directly opposite to the Moon's true Ecliptic Place.

EXAMPLE

PRECEPTS for finding the TIMES and PLACES of Ecliptic NEW and FULL MOONS.

EXAMPLE 2. To find the Sun and Moon's true Places, at their Ecliptic Opposition, in the Year 1764, March 17^d 11^h 53^m nearly. See Example VIII. P. 396.

Sun's true Place at that Time by first Tables . . . 11^h 27^m 53^s 58"
Moon's true Place by 1, 8, 9, 10, and 11 Equat. C. 5 27 38 46

Moon á Sun at that Time 5 29 44 48
Moon short of Opposition, or 6 Signs 15 12
Sun's true Place for an Hour later 11 27 56 27
Moon's true Place for that Time 5 28 16 35

Moon á Sun at that Time 6 0 20 8
Moon past an Opposition or 6^s 20 8
Hourly Motion of the Sun 2 29
Hourly Motion of the Moon 37 49
Hourly Motion of the Moon á Sun 35 20

Say 35' 20" :: 60^m :: 15' 12" : 25^m 49^s

March 17^d 11^h 53 0 +

Hence, the tr. Ecliptic Conj. will be 1764, Mar. 17^d 12^h 18^m 49^s nearly.

By M. De la Caille's Ephem. 1764, Mar. 17^d 12^h 2^m 16^s

The Time of this Ecliptic Opposition may be more nearly computed by using all the Equation Tables of the Moon. C.

EXAMPLE X. To find the foregoing Ecliptic Opposition of the Sun and Moon, on March 17, by the Supplemental Equation Tables D, of Mr. Mayer, P. 97, &c.

Sun's true Place, 1764, Mar. 17^d 11^h 53^m, as before, 11^h 27^m 53^s 58"
D tr. Ecliptic Pl. by 1, 11, 12, 13, and 14 Eq. Mayer, 5 27 42 18

Moon á Sun at that Time 5 29 48 20
Short of Ecliptic Opposition or 6 Signs by 11 40
Sun's true Place for an Hour later as before 11 27 56 27
Moon's true Place for that Time 5 28 19 49

Moon á Sun at that Time 6 0 23 22
Moon past Opposition or 6^s 23 22
Hourly Motion of the Sun 2 29
Hourly Motion of the Moon 37 31
Hourly Motion of the Moon á Sun 35 2

Say 35' 2" : 60^m :: 11' 40" : 19^m 59^s

March 17^d 11^h 53 0 +

Hence the Eclipt. Conj. March 17^d 12^h 12^m 59^s in 1764.

By M. De la Caille's Eph. March 17 12 2 16 Greenwich.

Eclipse 10 43 Dif.

The Time of this Ecliptic Opposition of the Sun and Moon may be also more nearly computed by using all M. Mayer's Eq. Tables, P. 97, &c.

TO determine the NUMBER of ECLIPSES in any YEAR, and on what DAYS of the MONTH they happen?

By Table, P. 385, take out the Place of the Sun from the Node for the Beginning and Month-day of any Year in the 18th Century, which you may reduce to the Beginning and Month-day of any other Year, in a remote Century, by adding or deducting the Days, and Motion of the Sun from Node for the Century, or Centuries distant, forward or back of the Radical Days and Place of the Sun from the Node. Then add the Motion of the Sun from the Node for Months forward in that Year to make the Sun from the Node, at New and Full, come within the Limits of a solar or lunar Eclipse, for as many Months as possible, (which is done by Inspection of Motion of Sun from Node) viz. within 18° Sun from Node, at New; and 12°, at Full Moon.

EXAMPLE XI. To determine the Number of Eclipses for the Year 1764, and on what Days they happen in that Year?

By Tab. P. 385.

	d	h	m	s	Sun á Node.
O. S. New D, 1764 Jan.	21	20	10	59	10 4 14 29
N. S. Add 11					
January 29	12	44	3		1 0 40 14
1/2 Revolution	14	18	22	2	0 15 20 7
Full. March 17	3	17	14		11 20 14 50
1/2 Revolution +	14	18	22	2	0 15 20 7
New. March 31	21	39	6		0 5 34 57

All the ECLIPSES that can happen, in 1764.

The same may, as readily, be performed for any other Year, forward or back of the present Year.

Moon 10° short of South Node. Eclipse D. I.

Sun 50¹/₂ past N. Node. Eclipse Sun. II.

	d	h	m	s	Sun á Node.
Again, O. S. 1764, January	21	20	10	59	10 4 14 29
N. S. Add 11					
August, Biff. 23	5	52	24		8 5 21 51

New. September 25 2 3 23
1/2 Revolution — 14 18 22 2

Full. September 10 7 41 21

When the Sun from the Moon's Node is just without, or near the Limits of a solar or lunar Eclipse, you must examine all Circumstances closely to avoid missing an Eclipse.

Sun 90¹/₂ past S. Node. Eclipse Sun. IV.

Moon 6° short of S. Node. Eclipse D. III.

PRECEPTS for computing ECLIPSES of the SUN and MOON.

WHAT AN ECLIPSE OF THE MOON IS.

All dark Bodies, being exposed to the Sun, cast a Shadow behind them, by intercepting the Sun's Rays, and hindering the opposite Space from being enlightened. And the Earth, being a dark Body, casts a Shadow into the Space opposite to the Sun, wherein if the Moon should come, in her Course through her Orbit, she must be deprived of the Sun's Light and be so disqualified from enlightening the Earth, by Reflection of her borrowed Light, as to occasion that Appearance which is called an Eclipse of the Moon.

There is, besides the totally dark Shadow, which is a conic Space totally deprived of the Sun's Light, an ambient Space which is but partly deprived of his Light; that is the Space whereon the Sun's whole Body does not shine, but the Rays coming from some Part of his Body only enter and enlighten that Space more or less, according as the Place is nearer to or farther from the totally dark Shadow: This partially dark Shadow is called the Penumbra.

PRECEPTS for computing ECLIPSES of the SUN and MOON.

TO compute an ECLIPSE of the MOON.

Precept 1. At the Time of true Opposition of the Sun and Moon (found as before) find the Angle which the Way of the Moon from the Sun makes with a Circle of Latitude (by Tab. P. 392) from the Argument of Latitude, and hourly Motion of the Moon from the Sun. Then say,

As Rad. : S. Moon's Lat. :: S. said $\angle$ Moon to Sun : S. nearest Approach of the Centers of the Moon and Earth's Shadow.

Prec. 2. From the Table (P. 391) of the hourly Motions and Semidiameters of the Sun, with the Argument of the Sun's mean Anomaly, take out his Semidiameter. And, from the same Table, with the proper Argument in Eclipses, take out the Moon's horizontal Parallax and Semidiameter.

Then from the Sum of the horizontal Parallaxes of the Sun and Moon, subtract the Semidiameter of the Sun, and the Remainder will be the apparent Semidiameter of the Earth's Shadow. — To be increased by 50" for the Earth's Atmosphere.

Prec. 3. To the Semidiameter of the Earth's Shadow, thus increased, add the Semidiameter of the Moon; and if the Sum of which is more than the nearest Approach of their Centers, (found as before) the Moon will be eclipsed at that Time; but otherwise she will not.

Prec. 4. The Moon being found eclipsed; from the Sum of the Semidiameters of the Moon and Earth's Shadow, deduct the nearest Approach of their Centers, and the Remainder will be the Part deficient. Now say,

As Semidiam. of Moon : 6 Digits or 360' :: Part deficient : Digits eclipsed.

Prec. 5. From Tab. (P. 391) with Arg. Lat. and hourly Mot. Moon fr. Sun, take out the Time between the true Opposition and the mean Eclipse, or greatest Obscuration; which being added to, or subtracted from, the Time of the true Ecliptic Opposition, according as the Moon is yet short of, or past, the next Node, will give you the Time of the greatest Obscuration.

Prec. 6. From the Square of the Sum of the Semidiameters of the Moon and Earth's Shadow, in Seconds, deduct the Square of the nearest Approach of their Centers in Seconds of a Degree, the Square Root of what remains will be Motion of Semiduration, in Seconds of a Degree.

Prec. 7. If the Eclipse be discovered to be above 12 Digits (or total) from the Square of the Difference of the Semidiameters of the Moon and Earth's Shadow, in Seconds of a Degree, deduct the Square of the nearest Approach of their Centers, in Seconds of a Degree, the Square Root of what remains will be the Motion of Semiduration, in total Darkness, in Seconds of a Degree. Now say,

As hourly Mot. Moon to Sun in her Orb (reserved in finding the true Opposition) : is to 1 Hour or 60^m Time :: so is the Motion of Semiduration : to the Time of Semiduration.

And :: so is the Motion of Semiduration in total Darkness : to the Time of Semiduration in total Darkness. Which being respectively subtracted from, and added to, the Time of the greatest Obscuration, will give you the Time of the Beginning and End of the Moon's Eclipse, and also of the total Darkness.

WHAT AN ECLIPSE OF THE SUN IS.

The Appearance, called an Eclipse of the Sun, might properly be called an Eclipse of the Earth; the Earth and its Inhabitants at that Time being only deprived of the Sun's Light, while the Sun himself suffers no Defect of Light. If we consider a Spectator placed at the Moon in the Time of what is called an Eclipse of the Sun, he will behold the Moon's totally dark Shadow, and her Penumbra, pass over the Earth's Disk or Face, like as an Inhabitant of the Earth sees the Earth's dark Shadow, and her Penumbra, pass over the Moon's Disk, or Face, in an Eclipse of the Moon. Where it is to be observed, that the Sun is seen eclipsed when only the Moon's Penumbra falls upon the Earth; whereas the Moon is never seen, or said to be, eclipsed, but when she falls into the totally dark Shadow of the Earth. And while the Earth diurnally revolves about its Axis, a Spectator at the Moon will see Countries, Cities, and Towns, move upon the Earth's Disk or Face, from West to East, or from Right to Left. That as such Spectator at the Moon, at the Time of a Sun's Eclipse, sees only the Earth's enlightened Hemisphere, he will see Places emerge from behind the Western Edge of the Earth's Disk, and move Eastward, when the Inhabitants of such Places of the Earth see the Sun rise and move towards the West. And when such Spectator in the Moon sees the Edge of her Penumbra touch any Place upon the Earth's Disk, the Inhabitants of that Place on the Earth will see the Sun begin to be eclipsed.

TO compute a general and central ECLIPSE of the SUN.

Precept 1. At the Time of the true Conjunction of the Sun and Moon (found as before directed) from the horizontal Parallax of the Moon, subtract the horizontal Parallax of the Sun, the Remainder will be the Semidiameter of the Earth's Disk. Likewise, to the Semidiameter of the Sun add the Semidiameter of the Moon, the Sum of which will be the Semidiameter of the Moon's Penumbra.

Prec. 2. To the Semidiameter of the Earth's Disk, add the Semidiameter of the Moon's Penumbra, and if the Sum exceeds the nearest Approach of their Centers (found like the nearest Approach of the Centers of the Moon and Earth's Shadow, in an Eclipse of the Moon) the Sun will be seen eclipsed in some Part of the Earth; but otherwise no Eclipse of the Sun can then happen.

Prec. 3. The Sun being found eclipsed in some Part of the Earth, the Beginning, Middle, and End of the general and also central Eclipse (with Respect to the whole Earth) may be found in like Manner as is before directed, for finding the Beginning, Middle, and End of an Eclipse of the Moon.

But to find the Places on the Earth where the principal or particular Appearances of an Eclipse of the Sun happen, may be done by an orthographic Projection of the Sphere, or such as would be seen by a Spectator placed at a very great Distance, in a right Line; joining the Centers of the Sun and Earth.

The Beginning, greatest Obscuration, and End of an Eclipse of the Sun, in any particular Place on the Earth, are also determined by an analytic Computation from converging Series, grounded on a geometrical Projection; (See Duntborne's Practical Astronomy of the Moon, inscribed to Dr. Long of Cambridge) being a PROLIX METHOD.

TO compute a particular or partial ECLIPSE of the SUN, by PARALLAXES, according to the COMMON METHOD

Precept 1. To the apparent Time of the true Conjunction of the Sun and Moon, (found as before directed) compute the Parallax of the Moon from the Sun, in Longitude.

Prec. 2. To Half an Hour, an Hour or more (as the said Parallax is small or great) before or after the true Conjunction, according as the Moon is in the eastern or western Quadrant of the Ecliptic, compute again the Parallax of the Moon from the Sun in Longitude; by which Means (from the true) you will have the visible half hourly, hourly, &c. Motion of the Moon from the Sun. Now say,

As visible Motion of Moon to Sun : is to Time taken :: so is the Parallax of the Moon from the Sun in Longitude, at the Time of true Conjunction : to the Interval of the true and visible Conjunction. — Which added to, or subtracted from, the Time of the true Conjunction, according as the Moon is in the eastern or western Quadrant of the Ecliptic, will give you the Time of the visible Conjunction of the Sun and Moon.

Prec. 3. To which Time, compute again the true Places of the Sun and Moon, with the Parallax of the Moon from the Sun in Longitude and Latitude, and so find her apparent or visible Place, in Respect of the Sun.

Prec. 4. To an Hour before the Time of visible Conjunction, compute the visible Longitude of the Moon from the Sun, in Antecedence; with her visible Latitude; and also to an Hour after the visible Conjunction, compute the visible Longitude of the Moon from the Sun in Consequence; with her visible Latitude. Now say,

As Sine of the visible Motion of the Moon from the Sun in Longitude in those two Hours : is to Radius :: so is the Tangent of the Difference

of visible Latitudes : to the Tangent of the Angle of the visible Way
of the Moon with the Ecliptic at the Time of the visible Conjunction.

And, as Radius : is to Sine of the visible Latitude of the Moon at the Time of the visible Conjunction :: so is the Sine of the Angle of the visible Way of the Moon with the Ecliptic at that Time ; to the Sine of the Motion, seen from the visible Conjunction to the greatest Obscuration.

And :: *so is the Cosine of that Angle : to the Sine of the visible Distance of the Centers of the Sun and Moon at the Time of the greatest Obscuration, that is, their nearest visible Distance.*

If this Distance be less than the Semidiameter of the Moon's Penumbra (found as before directed) the Sun will be eclipsed in that Place of the Earth; but otherwise not.

If the Sun be found eclipsed, from the Semidiameter of the *Penumbra* subtract the nearest *visible* Distance of the Centers of the Sun and Moon, the *Remainder* will be the Part *deficient*. — From which the *Digits* eclipsed may be found, as before *directed* in P. 398.

Prece. 5. Now say, As the visible Motion of the Moon from the Sun from an Hour before to an Hour after the Time of Conjunction : is to 2 Hours, or 120^m :: so is the Motion seen from the visible Conjunction to the greatest Obscuration : to the Interval in Time. — Which being added to, or subtracted from, the Time of the visible Conjunction (as the Moon's visible Latitude is decreasing or increasing) gives the Time of the greatest Obscuration.

Proc. 6. From the *Square* of the Semidiameter of the Penumbra, in Seconds, subtract the *Square* of the nearest visible Distance of the Centers of the Sun and Moon in Seconds, the *Square Root* of the Remainder will be the visible Motion of Semiduration in Seconds of a Degree.


Præc. 7. Now say, *As Sine of the visible Motion of the Moon à Sun in Longitude for an Hour before the visible Conjunction : is to Radius :: so is the Tangent of the Difference of visible Latitude in that Time : to the Tangent of the Angle of the visible Way of the Moon with the Ecliptic, from the Beginning of the Eclipse to the visible Conjunction.*

And, As Radius : is to the Sine of the visible Motion of Semiduration
:: so is the Cosine of that Angle : to the Sine of the visible Motion in
Longitude from the Beginning of the Eclipse to the greatest Obscuration.

PREC. 8. Likewise, As the visible Motion of the Moon & Sun in Longitude, in the Hour before the visible Conjunction: is to 1 Hour or 60^m :: so is the Motion in Longitude, seen from the Beginning of the Eclipse to the greatest Obscuration: to the Time of Incidence — Which subtracted from the Time of greatest Obscuration, gives the Time of the Beginning of the Eclipse.

☞ By the like Proportions (from the visible Longitude of the Moon from the Sun in Consequence, with her visible Latitude for an Hour after the visible Conjunction) the Time of Emergence and the End of Eclipse may be determined; with sufficient Labour and Attention.

IN the annexed Fig. the Circle ABCD represents the Earth's enlightened Disk, BED the *Ecliptic* thereon projected, or *orthographically*, as would be seen by a *Spectator* placed at a great Distance in a right Line joining the Centers of the *Sun* and *Earth*. EC the *Ecliptic* A and SIK the *Ar* projected on th



the *Ecliptic Axis*, C its Pole, P the Projection of the Earth's Pole; and SIK the Path of the Moon's *Shadow*, or her Way from the Sun projected on the *Disk*.

NOW, in the right-angled *spherical* Triangle, CMP, we have given (besides the right Angle at M) the Side CP, the Distance of the Poles of Ecliptic and Equator, and $\angle PCM =$ Sun's Distance from the equinoctial Point; from whence the Side CM, the Measure of the $\angle CEM$ (the Inclination of the Earth's and Ecliptic's Axes) is determined.

Here it is to be noted, that when the Sun is in the ascending Signs ♈, ♉, ♊, ♋, ♌, ♍, the Point P, the Pole of the Equator in the Projection falls to the Right of the Ecliptic Axis; but when the Sun is in the descending Signs, ♎, ♏, ♐, ♑, ♒, ♓, the Point P falls to the Left of the Ecliptic Axis; contrary to the Way it falls when the Projection is supposed to be made on a Plane parallel to the Earth's Disk, at the Moon's Orbit.

The $\angle$ CEI, made by a *Perpendicular* to the Path of the Moon's *Shadow*, with the *Axis* of the *Ecliptic* is determined to be the *Complement* of the *Angle*, which the *Way* of the *Moon* from the *Sun* makes with a *Circle of Latitude*. But you must observe that this *Perpendicular* falls to the *Left* of the *Ecliptic Axis* when the *Moon* is nearer to her *ascending* than *descending* Node; but to the *Right* thereof when she is nearer to her *descending* than *ascending* Node.

The $\angle$ s CEM and CEI, being, as before, determined, the $\angle$ IEM (their Sum or Difference) the Place F on the Earth's Surface, first touched by the *Penumbra*, and where the rising Sun begins to be eclipsed in his *superior* Limb, are likewise determined, as follows.

Draw the Meridian PF; then, in the right angled Plane Triangle SIE, are given the Sides SE, the Sum of the Semidiameters of the Earth's Disk and Moon's Penumra, and IE, the nearest Approach of their Centers; from whence the $\angle$ SEI is determined; and by adding or deducting the $\angle$ IEM known, the $\angle$ SEM will be also known.

NOW, in the *spherical Triangle* PMF, on the Earth's Surface, right angled at M, the Sides PM, equal to the Sun's Declination, and FM, the Measure of the $\angle$ SEM, are given. Hence, you may find the Side PF, the Complement of *Latitude* of the Place F, and the $\angle$ MPF; which $\angle$, or its Supplement to 180° (as the elevated Pole, in the Place F, is darkened or enlightened) is the Angle contained between the Meridian of the Place F, and the Meridian of the Place where the Sun is, at that Time, *vertical*. But the Longitude of the Place where the Sun is *vertical* is known by the Time; and therefore the Longitude of the Place F, will be known. — And having determined both *Latitude* and *Longitude*, the Place itself will be known.

By the like Method you may find the Place on the Earth, where the Center of the *Perumbra* enters the Earth's *Disk*, and the Sun rises *centrally* eclipsed; as likewise those Places where the *central Eclipse*, and *Penumbra* both quit the Earth.

And, by the same Method you may find the *Place G*, upon the Earth's Surface, under the *Penumbra's* Center; at the Time of the true *Conjunction*, when the Sun is centrally eclipsed in the *nonagesimal* Degree, or highest Point of the *Ecliptic*.

For, if EC, the *Disk's* Semidiameter, be made *Radius*, then GE, the Moon's Latitude, will be the *Sine* of the Sun's Distance from the *Vertex* of the Place G. Therefore, in the *spherical* Triangle GEP, on the Earth's Surface, the *Sides* PE, the Sun's Distance from the Pole, and GE, his Distance from the *Vertex*, and the $\angle$ GEP are given; from whence are found the Side GP, the Complement of Latitude of the Place G, and the $\angle$ GPE, the Difference of Meridians of the Place G, and where the Sun is then vertical; from which the Place G itself will, as before, be also known.

In like Manner, the Place of the Earth may be found, under the Center of the *Penumbra* at any given Instant of Time.

The Place H, upon the Earth's Surface, where the Sun is *centrally eclipsed* in the Meridian, may be found by only resolving the right angled Triangle EIH, wherein are given all the Angles, and the Side IE, to

To find the PLACES, on the EARTH, of the principal APPEARANCES of Solar ECLIPSES, by PROJECTION.

find the other two Sides; viz. IH divided by the Moon's hourly Motion from the Sun will give the Difference of the Time from the Middle of the Eclipse, when the Center of the Shadow is upon the UNIVERSAL MERIDIAN, or Line HPE, wherein the Earth's Axis is projected. It is called Universal Meridian because when a Spectator placed in the Moon observes any Place, on the Earth's Surface comes to touch this Line (by the Earth's diurnal Rotation) it will be Mid-day to the Inhabitants of that Place. Hence, the Longitude of the Place H will be known; being equal to the Longitude of the Place where the Sun is vertical, when the Center of the Shade is upon the said universal Meridian: And the other Side HE is the Sine of the Sun's Distance from the Vertex of the Place H (making the Semidiameter of the Disk, Radius) whereby the Latitude of that Place will be also known.

The Beginning, greatest Obscuration, and End, of an Eclipse of the Sun, for any particular Place of the Earth, may be determined by algebraic Computation, and infinite Series, grounded on a geometrical Projection, similar to the foregoing (as may be seen in Dunthorne's Astronomy.) But, as this Method is very tedious and prolix, and fitter for Speculation than Practice, we therefore omit it. And we leave it to the Computer's PRACTICE and EXPERIENCE to apply the plain and easy RULES we have delivered, for computing ECLIPSES of the SUN and MOON in general. See P. 260, for the Application.

NUMBER of ECLIPSES of the SUN and MOON that can happen in ONE YEAR.

THE Number of Eclipses of the Sun and Moon in any Year cannot be less than two nor above seven. The usual Number is four but very seldom above six.

See P. 397, for finding the Number of Eclipses in any Year.

If at the first New Moon in January, before the 11th Day, (the Number of Days Difference between the lunar and solar Year) the Sun be near 18 Degrees short of the Node, the Sun will be then eclipsed, and the next Full Moon, and also following New Moon, will be eclipsed; viz. 2 Eclipses of the Sun and 1 of the Moon will happen before the 9th of February, that Year. And as many will happen in the 6th Lunation after, in the same Order, before August 5, viz. 1 of the Sun, 1 of the Moon, and another of the Sun, in 177 Days from the former; being the Time of 6 mean Lunations. And 1 Eclipse more of the Sun will happen at the End of 12 Lunations from the 1st, being the greatest Number of Eclipses, viz. 5 of the Sun, and 2 of the Moon, that can happen in a Year, including 13 New Moons.

	Ecl. Sun. New. Sun à Q.	Ecl. Moon. Next Full. Sun à Q.	Ecl. Sun. Next New. Sun à Q.
Before Jan. 11	11° 12' 30"	11° 27' 05"	0° 13' 10"
EX. 3 Eclipses 6 Lunations +	6 4 1	6 4 1	6 4 1
3 more Eclipses 6 Lunations +	5 16 31 6 4 1	6 1 51	6 17 11
1 more Eclipse	11 20 32	In all 7 Eclipses.	

But if on the first Full Moon before the 11th of January, the Sun from the Node be near 18 Degrees, as at the first New Moon, in that Month, and changing Eclipses of the Sun to those of the Moon, in the above Example, then there can only happen 2 Eclipses of the Sun, in that Year, being the least Number. And two Eclipses of the Sun can only happen in a Year, when, at the first New Moon in January, the Sun is very near the Moon's Node; so that the mean or different Number of Eclipses happen (between the greatest and least Numbers, 7 and 2) when at the first New Moon in January, the Sun à Q is in the middle or at different Distance, betwixt the Sun's eclipsed Limits, of 0 and 18 Degrees, Sun à Q, as appears from above.

HISTORICAL ECLIPSES of the SUN and MOON. To be examined. From RICCIOLUS.

Bef. Chr. 754 July 5	But April 21, by an old Calendar, an Eclipse of the Sun. Rome began to be built.
721 Mar. 19	Total Eclipse of the Moon. The Assyrian Empire ended, and Babylonian established.
585 May 28	Eclipse of the Sun, predicted by Thales. Peace ensued betwixt the Medes and Lydians.
523 July 16	Eclipse of the Moon. Followed by the Death of Cambyfes.
502 Nov. 19	Eclipse of the Moon. Followed by the Slaughter of the Sabines, and Death of Valerius Publicola.
491 Apr. 25	Eclipse of the Moon. Followed by a great Famine at Rome, and Beginning of the Peloponnesian War.*
463 Apr. 30	Eclipse of the Sun. War with Persians, falling off from Egyptians.
431 Aug. 3	Total Eclipse of the Sun. A Comet and Plague at Athens.†

* The Beginning of this War is here antedated 60 Years.

† This Eclipse was in the 1st Year of the Peloponnesian War, according to Ferguson's Astronomy.

Mr. Ferguson, in P. 168 of his Astronomy, not arguing from the Distance and Motion of the Sun from the Node, at the Time of and for Conjunction and Opposition, when Eclipses of Sun and Moon happen (but from the Time of the Sun's Passage by the Node) assigns not the true Reason for the Number of Eclipses that can happen in a Year. But he makes us ample Amends in his accurate Draughts, and Account of other Subjects.

Of the PLINIAN PERIOD of ECLIPSES, and their RETURNS.

THE Period of Eclipses of 223 synodical Months (discovered by the Chaldeans, and called by them Saros, and also called the Plinian Period, as used by Pliny) consists of 18 Years, 11 Days, 7 Hours, 43 Minutes, and 20 Seconds, when 4 Leap-Years are contained; but when 5 Leap-Years intervene, it consists of 1 Day less, or of 18 Years, 10 Days, 7 Hours, 43 Min. and 20 Seconds; very necessary to be noted in its Use.

The Addition of the odd Hours in this Period may shift a solar Eclipse into Night, and a lunar one into Day; so as both to become invisible, after a Period or two. The Sun à Q, at the Conclusion of any Number of Periods, will shew whether an Eclipse of the Sun or Moon returns, within the Limits of 18 or 12 Degrees respectively; which does not agree with Mr. Ferguson's Account of no Eclipse of the Sun repeated, after 500 Years; nor yet with his 12 Thousand Years Revolution, on their Return from travelling in the Expanse, during that Interval. For 470130 of a Deg. is what Sun à Q goes back in one Period; by which dividing 180, the Quotient = 38,287 the Number of Periods, when the Moon's Shadow quits the Earth; which X'd by 18 Yrs, 422 in one Period = 705 Yrs, 32, &c. Yrs. (and not 500) when the Moon's Shadow quits the Earth, and travels in the Expanse. Now, 3600 ÷ 180 = 20; therefore 705,32 X 20 = 14106 Years, the Period of Returns of solar Eclipses; and not 12 Thousand, according to Mr. Ferguson. — Again, 120 ÷ 470130 = 25,5248 Periods, when the Earth's Shadow quits the Moon, which X'd by 18 Yrs, 422 = 470 Yrs, 217, when the Earth's Shadow wholly travels in the Expanse. — But, 3600 ÷ 120 = 30, which X'd into 470,217 = 14106 Yrs, as before, the Period of lunar Eclipses, travelling in the Expanse.

The falling back of the Sun and Moon from the Q in every Period, is the Cause of these Eclipses about the ascending Node happening more southerly, at each Return; and of those about the descending Q happening more northerly till they wear off the Earth and Moon's Surfaces, and travel in the Expanse.

Of Antient ECLIPSES.

THE present Astronomical Tables are found to disagree with the Times of antient Eclipses, on Account of the Moon's Acceleration of mean Motion, requiring a considerable Addition to her antient mean Places, as we have illustrated. Therefore, the Eclipse of the Sun predicted by Thales, in the 4th Year of the 48 Olympiad, putting an End to the Battle betwixt the Medes and Lydians, might very well correspond with the Time of that Eclipse by Computation; without going so far back with Mr. Ferguson, as a Plinian or Chaldean Period, to make the Time, Appearance, and Event, agree. On which Principle of Acceleration, and no other, the Times of antient Eclipses, in general, are to be examined, and reconciled with Observation.

PLURALITY of ECLIPSES.

MORE Eclipses happen of the Sun than of the Moon; because the Sun's eclipsed Limits are greater than that of the Moon's, as 180 to 120 Sun from Q.

But more visible Eclipses of the Moon than of the Sun happen; because Eclipses of the Moon are alike seen from all Parts of the Earth placed next the Moon. But Eclipses of the Sun are only visible to that small Part of the Earth (placed next the Sun) whereon the Moon's Shadow falls. And at the farthest Distances of the Moon from the Earth, her Shadow terminates before it reaches the Earth, at New Moon; and the Sun's greatest Eclipse is then called annular, with a bright Ring appearing round the Moon's Edge.

ANGLE of the HOUR CIRCLE (passing through the SUN or MOON) with the ECLIPTIC. For finding the Parallax Angle of the Sun or Moon, in ECLIPSES; and also in all Latitudes of the Moon.

Argument. Right Ascension of the SUN or MOON.

R. A Sun or D	6 Sig. 180 Deg.	1 Sig. 30 Deg.	2 Sig. 60 Deg.	60 Deg.	R. A Sun or D											
	HourCirc.	Dif.	HourCirc.	Dif.	HourCirc.	Dif.										
o	o	i	''	i	''	o	i	''	i	''	o					
0	66	31	30	0	14	69	49	10	12	55	78	30	40	21	12	30
1	66	31	44	0	41	70	2	5	13	17	78	51	52	21	25	29
2	66	32	25	1	8	70	15	22	13	37	79	13	17	21	34	28
3	66	33	33	1	35	70	28	59	14	0	79	34	51	21	44	27
4	66	35	8	2	3	70	42	59	14	20	79	56	35	21	54	26
5	66	37	11	2	30	70	57	19	14	40	80	18	29	22	3	25
6	66	39	41	2	56	71	11	59	15	0	80	40	32	22	12	24
7	66	42	37	3	24	71	26	59	15	20	81	2	44	22	20	23
8	66	46	1	3	51	71	42	19	15	40	81	25	4	22	29	22
9	66	49	52	4	17	71	57	59	15	59	81	47	33	22	37	21
10	66	54	9	4	44	72	13	58	16	17	82	10	10	22	44	20
11	66	58	53	5	10	72	30	15	16	35	82	32	54	22	51	19
12	67	4	3	5	37	72	46	50	16	54	82	55	45	22	58	18
13	67	9	40	6	4	73	3	44	17	14	83	18	43	23	3	17
14	67	15	44	6	29	73	20	58	17	25	83	41	46	23	10	16
15	67	22	13	6	55	73	38	23	17	45	84	4	56	23	15	15
16	67	29	8	7	21	73	56	8	18	2	84	28	11	23	21	14
17	67	36	29	7	46	74	14	10	18	18	84	51	32	23	25	13
18	67	44	15	8	11	74	32	28	18	34	85	14	57	23	30	12
19	67	52	26	8	36	74	51	2	18	48	85	38	27	23	34	11
20	68	1	2	9	2	75	9	50	19	6	86	2	1	23	37	10
21	68	10	4	9	26	75	28	56	19	17	86	25	38	23	41	9
22	68	19	30	9	50	75	48	13	19	32	86	49	19	23	44	8
23	68	29	20	10	14	76	7	45	19	47	87	13	3	23	46	7
24	68	39	34	10	38	76	27	32	20	0	87	36	49	23	49	6
25	68	50	12	11	2	76	47	32	20	12	88	0	38	23	50	5
26	69	1	14	11	25	77	7	45	20	26	88	24	28	23	51	4
27	69	12	39	11	47	77	28	11	20	38	88	48	19	23	54	3
28	69	24	26	12	11	77	48	49	20	50	89	12	13	23	53	2
29	69	36	37	12	33	78	9	39	21	1	89	36	6	23	54	1
30	69	49	10			78	30	40			90	0	0			0
R. A Sun or D	5 Sig. 150 Deg.	4 Sig. 120 Deg.	3 Sig. 90 Deg.	R. A Sun or D												
	11	10	9													

Construction. As Radius : S. Ecliptic Obliquity :: Cos. R. A. from next Equin. Point, :: Cos. $\angle$ Hour Circle with the Ecliptic.

Example. To find the Angle of Hour Circle with the Ecliptic to 65° R. A. from the next equinoctial Point?

As Radius	Logarithms.
To S 23° 28' 30"	9.6002636
So Cos. 65°	9.6259483
To Cos. 80° 18' 29"	9.2262119
$\angle$ Hour Circle required.	

Visible Conjunction of Sun and Moon, as seen from the Earth's Superficies, always differs from the Time of true Conjunction, as seen from the Earth's Center, except when they are conjoined in the nonagesimal Degree; when the true and visible or apparent Time is the same. — Hence, if the true Conjunction falls to the Eastward of the nonagesimal Degree, the Moon's Place is advanced by her Parallax of Longitude, when the visible Conjunction is before the true. But if the true Conjunction falls to the Westward of the nonagesimal Degree, the Moon's Place is retarded by the Parallax in Longitude, and the visible or apparent Conjunction of Sun and Moon, then follows the Time of the true Conjunction, very necessary to know in computing solar Eclipses.

Moon's Dist. fr. Ecliptic, or vis. Conj. to Orbit. Conj. or Midd. Ecl. ☉.

Ar. Vis. Lat. D.

Vis. Lat. D.	Dist. fr. Ecl. ☉.
0	0 5
1	1 10
2	2 15
3	3 20
4	4 25
5	5 30
6	6 35
7	7 40
8	8 45
9	9 50
10	10 55
11	11 55
12	12 55
13	13 55
14	14 55
15	15 55
16	16 55
17	17 55
18	18 55
19	19 55
20	20 55
21	21 55
22	22 55
23	23 55
24	24 55
25	25 55
26	26 55
27	27 55
28	28 55
29	29 55
30	30 55

Lat. D. Dist. above or below the Ecl. ☉.

USE.

As vis. hour-ly Mot : 60m :: Dist. fr. Mid. Ecl. in Mot. : Dist. in Minutes of Time.

PRINCIPLES of PROJECTION of a Solar ECLIPSE, Geometrically.

A Plane being conceived to touch the Moon's Orbit in a Point where a Line (at right Angles to the said Plane) intersecting the said Orbit, connects the Centers of the Earth and Sun; and an infinite Number of straight Lines being supposed to pass from the Sun's Center, through the said Plane of the Earth's Periphery, to the Axis thereof, and likewise to the Axis of the Ecliptic, and the Path of any Vertex, the said straight Lines will, orthographically, project the Earth's Disk, its Axis, the Axis of the Ecliptic, and Path of the Vertex, on the aforesaid Plane.

See Flamsteed's Doctrine of the Sphere, P. 27.

If the Sun be in the Equinoctial, the Paths of the Vertices (described by the Rotation of the Earth's Surface) will be projected in straight Lines upon the said Plane; but if the Sun has Declination, those Paths will be projected in Ellipses, upon the said Plane.

The Transverse-Diameter of the Ellipsis, representing any Path, is equal to double the right Line of the Distance of the said Vertex from the Pole; that is equal to twice the Cosine of the Place's Latitude. But the Conjugate is equal to the Difference of the right Sines of the Sum and Difference of the Distances of the Path and Sun from the Pole. That is equal to the Sine Compl. of the Sun's Declination added to the Co-latitude of the Place less the Right Sine of the Difference of the Complement of the Sun's Declination, and Co-latitude of the Place.

The Transverse Diameter lies at right Angles to the Earth's Axis, and the Conjugate coincides with it.

Of the MOON'S PARALLAXES.

1. If the Distance of the Moon from the Point ascending or descending of the Ecliptic, be less than her Altitude, she has then no Parallax in Latitude, which never happens but in such Latitudes where the Moon's Orb or Ecliptic are vertical Circles.

2. If the Moon's Distance from the Point ascending or descending be just 90°, then a vertical Circle intersects the Ecliptic at right Angles, and then is no Parallax of Longitude but only of Latitude.

3. If the Vertical passing through the Moon's Center fall on the Ecliptic, at oblique Angles, then there is Parallax in Longitude and Latitude.


4. To all Places on the Earth having above 28° 46' 20" North Latitude, the Moon's Parallax is Southward, and is depressed below her true Place, according as she is East or West of the 90°. — These Parallaxes are necessary to be known in the Computation of solar Eclipses.

EXAMPLES of computing the PARALLACTIC ANGLE, according to DIFFERENT METHODS.
By D. COWPER.

1717, September 17th 9^h 58^m 20^s, apparent Time, observed Emission of an Occultation of Aldebaran, at Crane-Court, London.

Sun's Longitude	2° 38' 0"
Right Ascension	182 24 57
Time from Noon in Degs.	149 35 0
R. Asc. Med. Caeli	331 59 57
Moon's Long. in Ecliptic	11 5 38 0
Latitude S.	4 38 33
Declination N.	16 42 15
Right Ascension	64 34 32
Angle at the Pole	92 34 35
Moon's Zenith-Distance	78 33 59
Angle of Position, at P's	39 21 26
Center, or $\angle PDL$	
Angle of Parall. Lat. and	
Circ. Decl. $\angle PDL$, the less	80 9 10
Or $\angle PDL$ Supplement the greater	99 50 50
Parallactic $\angle$, or $\angle PDL$	
$\angle PDL$, required.	60 29 24

P, the North Pole.
Z, the Zenith.
HH, the Horizon.
EE, the Equator.
EC, the Ecliptic.
LL, Parallel Latitude.
OO, Moon's Orbit.
D, Moon's Orbit-Place with 4° 38' S. Latitude.
PZ, Comp. Lat. 38° 28' 0".
PD, Moon's Distance from Pole 73° 17' 45".
ZD, Moon's Comp. Alt. 78° 33' 59".
 $\angle DPZ$, 92° 34' 35".
S descending Node D.



$\angle ZeC$, parallactic $\angle$ at the Ecliptic; being the Angle of a vertical ZD with the Ecliptic EC.
 $\angle ZDL$, Parallactic $\angle$ at the Moon; being the Angle of a vertical ZD with a Parallel of Latitude LL.

By the Universal METHOD.

Meridian Angle	69° 14' 26"
Altitude Med. Caeli	26 56 34
Moon's Azimuth from N. or $\angle PZD$	77 28 25
Parallactic $\angle ZeC$	60 38 14

To find the parallactic Angle at the Moon, or $\angle ZDL$, from the parallactic Angle at the Ecliptic $\angle ZeC$?

As Cos. D Lat. 4° 38' 33" : Rad. :: Cos. parallactic $\angle$ at D 60° 38' 14" above : Cos. 60° 31' 52", parallactic Angle at the Moon, required.

By the 90th DEGREE.

Medium Caeli in Ecliptic	29° 53' 59"
Altitude 90th Degree	33 26 10
Longitude 90th Degree	0 4 35 14
Moon from 90th Degree	2 1 2 46
Moon Zenith Distance	78 33 43
Parallactic $\angle ZeC$	60 32 2

According to D. COWPER's short METHOD.

1737, Feb. 18 th 2 ^h 25 ^m 9 ^s Ap. Time, Beginning. Sun Eclip. observed at Fleetstreet.	
Sun's Longitude	11° 6' 55"
R. Ascension	3 42 34 48
Time fr. Noon in Degrees	36 17 15
R. Asc. Med. Caeli	18 52 3
360°	
Moon's Ecliptic Long.	11 10 44
Latitude N.	0 37 40
Declination S.	6 48 11

D's Right Ascension 342° 23' 56"
 $\angle$ at the Pole 36 28 7
Moon's Zenith Dist. 66 7 41
 $\angle$ Position at D 23 52 54
Angle of Paral. Lat. and Circ. Decl. 67 41 5
Diff. parallactic Angle 43 48 11

According to UNIVERSAL METHOD.

For same Time.	
Meridian Angle	67° 51' 20"
Altitude M. Caeli	46 31 10
Moon's Azim. fr. S.	40 11 48
Parallactic Angle	43 48 20

According to 90th DEGREE.

For same Time.	
Med. Caeli in Ecl.	0° 20' 26" 4"
Altitude 90th Deg.	50 24 20
Longitude 90th Deg.	1 10 6 25
Moon fr. 90th Deg.	1 28 55 41
Moon's Zenith Dist.	66 7 56
Parallactic Angle	43 48 12

EXPLANATION of the foregoing DIFFERENT PARALLACTIC ANGLES.

When the Moon is in or near the Ecliptic, or near 90° or 180° , with Latitude, the parallactic Angle at the Ecliptic $\angle ZeC$ will be sufficient to answer the Purpose.

But if she be near 90° or 180° the parallactic Angle at the Moon must be used; the Difference between both which Angles being then considerable.

P, the N. Pole of the World.
E, Pole of the Ecliptic.
Z, the Zenith.
CN, Part of the Ecliptic.
CD, Lat. D South.
N, the Place of 90th Deg. Ecliptic.
LDL, a Parallel of Lat.
Pm, the Meridian.
 $\angle PmN$, the Meridian Angle.



ZE = Altitude 90th Deg or Dist. of Zenith fr. Pole of Ecliptic.
 $\angle CmZ$ also the Merid. $\angle$ Zenith, or $\angle CeD$, the parallactic $\angle$ at the Ecliptic.
 $\angle CDe$, the Compl. of parallactic $\angle$ at the Moon, that a Vertical ZD makes with a Circle of Latitude ED.

For, As Cos. Lat. CD : Rad. :: Cos. parallactic Angle at Ecliptic, CeD : Cos. parallactic Angle at the Moon, of ZDL, viz. to CDDe, required, according to the above Proportions by D. Cowper.—The true parallactic Angle, at all Times, being the Angle of a Vertical and Parallel of Latitude, or the Complement of the Angle that a Vertical makes with a Circle of Declination, viz. Complement of $\angle EDZ$ being $\angle eDL$; agreeing with Mr. Duntborne's Remark in his Astronomy, P. 63:

From whence follows this QUERE. If ED, a Circle of Latitude, cut the Ecliptic CN, and Parallel of Latitude LL, both at right Angles, why should not the alternate Angles $\angle ZeN$, $\angle ZDL$, respecting the same Parallels, cut by the vertical Circle ZD be the same, or the parallactic Angle at Ecliptic and at the Moon, be the same?

It is observed by one of our Correspondents, (a very judicious Mathematician and Astronomer) that he never met with any Author who taught the true Method of representing the Appearances of Eclipses, &c. nor ever saw them truly described in any Almanac or Ephemeris, for the British Dominions, except in the late Weaver's Ephemeris.

This Correspondent computed the parallactic Angle, and the Moon's visible Latitude, in several Positions, at the Beginning, Middle, and End of some Eclipses, for the Island of Antigua, that he might truly delineate the Manner of their Appearance there which (by Reason of the Ecliptic transiting the Zenith once in every 24 Hours) is quite different to what happens here in England, both with Respect to the Position of the Moon's Face, and the visible Curve she describes.

The solar Eclipse, happening the 18th of February, 1737, has been computed by the *operose Method* delivered in Leadbetter's *Uranoscopia*, from P. 147 to 153; viz. for finding the Meridian Angle, culminating Point, 90th Degree, &c. in the Moon's Orbit. And likewise that Eclipse was computed, for the same Place, with these Requisites in the *Ecliptic*; but the last Computation came much nearest to the Observation. And you may observe in the Examples of the Use of Dr. Halley's Tables, (d and d 2) that the 90th Degree, and *parallactic Angle* at the Moon, are both taken in the *Ecliptic*; though the Moon had almost 50 of Latitude.

It is likewise evident, that as the *Ecliptic* is a fixed Circle, (not changing its Inclination, and Place of Intersection with the Equinoctial, like the Moon's Orbit) in which the Sun constantly appears, and is that Circle to which all the Longitudes and Latitudes of the Planets are computed, and referred (being the Standard for determining the Moon's visible Longitude and Latitude) therefore the other Method of computing the *parallactic Angle* made by the Intersection of a Vertical and the Moon's Orbit, is rejected; as being both tedious and inaccurate.

PRECEPTS of lunar COMPUTATION.

TO find the Moon's true Place, from the mean Time given, when one of her Limbs passes the Meridian, her observed Right Ascension at that Time, (having no Parallax) and Distance of her lower (or upper) Limb from the Vertex.

In the Year 1725, December 5, the Western Limb of the Moon was observed by Dr. Halley to pass the Meridian of the Royal Observatory at Greenwich, at 9^h 8^m 5^s, mean Time, its observed Right Ascension being 42° 26' 15", and Distance of the lower Limb from the Vertex 34° 9' 15": Required from thence the Moon's true Place in the *Ecliptic*, and also her Latitude?

EXAMPLE.

By Halley's Tables of the Moon's *Ecliptic Longitude* 8 15° 42' 34" (See P. c. 1.) Her Latitude North 1 39 57

By same Tables. Moon's horizontal Parallax then 0 59 43
Which Parallax, at any Time, is also had fr. an *Ephem.*

Half + 1/2 of that Half is the Moon's horizontal Diameter 0 32 50
The observed Dist. of the Moon's lower Limb fr. Vertex 34 9 15
To which add Refraction + 36

Ap. Dist. D's lower Limb fr. Vertex, clear of Refraction 34 9 51

Now, As Radius	10.0000000	Com. Log.
To S. hor. Parallax 59' 43"	8.2397998	9021 Lo. Log.
So S. Dist. à Vert. 34° 9' 51"	9.7494008	2506 Ar. Com. L. S.

To S. D Par. in Alt. 33' 32" 7.9892006 2527 L. L.

This Parallax subtracted fr. ap. Dist. D's lower Limb 0 33 32

from the Vertex, clear of Refraction 33 36 19

Remains true Dist. Moon's lower Limb fr. Vertex 33 36 19

Subtr. from which the horizontal Sem. D 16 25

Rem. the true Dist. Moon's Center from Vertex 33 19 54

The Compl. of Lat. of Observatory, added to which 38 31 30

S. M. The true Dist. D's Center from N. Pole 71 51 24

Now, As Sine D's true Dist. à 71° 51' 24" co. 0.0221481 9778 Log. Sine

To Radius 10.0000000

So S. hor. Diam. D, 32' 50" 7.9800345 2617 L. L.

To S. D's hor. Diam. in R. A. 34' 33" 8.0021826 1.2395 L. L.

To the observed R. A. of the Moon's Western Limb 42° 26' 15"

Add the Moon's horizontal Semi-Diameter in R. A. 0 17 16

Sum, is the true R. A. of the Moon's Center 42 43 31

Now, From the Moon's R. A. 42° 43' 31" Quest. XXX.

And her Distance from the N. Pole 71 51 24 P. 228.

Her *Ecliptic Longitude* will be 8 15 42 12 by the

Latitude North 1 38 37 Observation.

Longitude } less than by Tables { 0' 22" the Error of Com-

Latitude } 1 20 putation.

The horizontal Diameter of the Moon is to her Diameter in Longi-

tude, as *Cosine* of the Moon's Latitude to Radius.

And her horizontal Diameter is to her Diameter in right Ascension,

as the *Cosine* of her Declination is to Radius.

The Seconds of a Degree (in Tab. P. 391) answering to the Moon's

Distance from the Vertex, and also to her Distance from her Apogee,

being added to the Moon's horizontal Diameter, will give her apparent

Diameter, which, increased in the respective foregoing Proportions,

will give her apparent Diameters in Longitude and right Ascension.

When the Moon's Eastern } Limb } touches the Meridian,
Western }
Subtract } her Semidiameter in right Ascension or Long. { from } her
add } her right Ascension or Longitude of the Limb observed, for the right Af-
cenion or Longitude of her Center. { to }

REMARK on Dr. Halley's TABLES. See the END of his Preface-
Writer's PREFACE.

"It appears from Dr. Halley's Papers, that both in computing the
"right Ascension of the Moon's Limb, and finding the Longitude of her
"Center from Observation, he constantly used the apparent, instead of the
"true, Semidiameter of the Moon; and consequently from the Beginning
"of the Table *Lunæ Meridianæ*, &c. to the latter End of the Year
"1725, from the New to the Full Moons, while the Western Limb was
"observed, the computed right Ascensions of that Limb are set down too
"backward; and during the Remainder of the Month, while the Eastern
"Limb was observed, the computed right Ascensions of that Limb are
"set down too forward.

"And from the Year 1725, the Longitudes of the Moon's Center de-
"duced from the Observations of the Western Limb are set down too for-
"ward, and those from the Observations of the Eastern Limb too back-
"ward.

"This Error is not considerable, and seldom exceeds 15"; therefore in
"common Cases it may be neglected. — But if any one undertakes to correct
"the Moon's mean Motion, or to new model the Equations, it will be ne-
"cessary to correct the Errors in the Abacus, or Table, by the Excess
"of the Moon's apparent Semidiameter in right Ascension, or Longitude,
"above her horizontal Semidiameter of the like Denomination."

To find, according to HALLEY'S METHOD, the Time when the
Moon's Eastern Limb passed the Meridian of Greenwich-Observatory, on
December 28, 1745?

EXAMPLE.

Take a *Plinian Period* back, which to December 18, 1727, is 18
Years, 10 Days, (because it contains 5 Leap-Years) on which Day,
in the *Tab. Lunæ Meridianæ*, the Moon's Eastern Limb was observed
to pass the Meridian at 13^h 45^m 21^s, mean Time.

But 18 Yrs. 10 Ds. want 7^h 43^m 20^s of a complete Period; in

which Time the Moon's mean Motion is 4° 14' 22"

And the Meridian passes over an Arc of the Equator 16^m 55^s

equal to this Arc of mean Motion, in 13^h 28^m 26^s

Subtract which therefore from the Time of Observa- Mean Time.

tion, and there remains 13^h 28^m 26^s

The near Time of the Passage of the Moon's Eastern Limb over the

Meridian of Greenwich Observatory, required.

But, the Moon's computed Long. to this Time is 6° 43' 8"

Her Latitude N. 3 49 24

In the last Column of Halley's Abacus, *Tab. Lunæ meridianæ*, the

Error of Computation for December 18, 1727 is — 3' 1"

Therefore, add which to the computed Longitude, and

you have the correct Longitude for the Time 6° 46' 9"

Hence, the right Ascension of the Moon is found 130 12 38

Which increased by her true Semidiameter in R. A. 17 18

Gives the R. A. of the Moon's Eastern Limb 130 29 56

To Sun's mean Long. at the given Time 288° 25' 2"

Add an Arc answerable to the above M. } 202 6 30

Time 13^h 28^m 26^s } Sum, is the right Ascension of the Meridian 130 31 32

Which right Ascension is past the Moon's Limb by an Arc of 1 36

The Supplemental Equation-Tables, at Beginning of this Work, used with the Eccentricity of the Moon's Orbit, or 2d Equation of her Apogee, are for Trial and Comparison of Computation, by other Methods; in Order for making Improvement in the Use and Number of lunar Equations, by different Methods.

This

Different METHODS of observing the LONGITUDE.

This Arc the Meridian passes over in about 6 $\frac{1}{2}$ Time, which subtracted from the Time given, would leave the Time of the Moon's Transit correct, over the Meridian sought, but for the Moon, increasing her right Ascension after her Limb passed the Meridian, and thereby diminishing her Distance from the same.

To allow for which add 4" to the Arc of Distance and it becomes

Which Arc the Meridian passes over in	00	1'	40"
Subtract which from the Time given	13 ^h	28 ^m	26 ^s
There remains the mean Time, correct, of the Moon's Eastern Limb passing Greenwich Meridian, according to Tables	13	28	19
Observed by Dr. Bradley	13	28	21

Of HALLEY'S METHOD of determining the LONGITUDE by OBSERVATION.

1. HE observes the mean Time of Approach of the Moon's Limb to a fixed Star.

2. The mean Time, and angular Distance of the Moon's Limb from a fixed Star, nearly in the same Parallel of Latitude.

3. The mean Time, and angular Distance of the Moon's Limb from the Sun's next Limb in her first and last Quarter.

All observed under a distant Meridian from Greenwich Observatory.

And by assuming a mean Time at Greenwich Meridian, to which his astronomical Tables are adjusted, (by repeated Trials) till it corresponds with the respective mean Time of the apparent Observations, made under a distant Meridian, he determines the Difference of Longitude: Being the Degrees answerable to the Difference between the assumed mean Time at Greenwich Observatory, and the mean Time under the distant Meridian, where and when the respective Phenomenon, or Appearance, is observed.

This he effects (as may be seen at the Beginning of his Tables) by computing the Sun's Longitude and the Moon's Longitude corrected, and Latitude to the assumed Time at Greenwich with the Moon's horizontal Parallax to that Time.

Then the mean Time, in Degrees, when the Observation was made, being added to the Sun's mean Longitude, at Greenwich, will give the right Ascension of the distant Meridian, nearly.

And by the Moon's Parallax, he gains the parallactic Angle: Whence from the Moon's true Distance from the Vertex (her apparent Semidiameter being given) he determines her Parallax in Altitude, Longitude, and Latitude, (according to Rules in P. 234) and thence her visible Distance and also of the Star, from the Vertex, (allowing for Refraction) and apparent Place of her Limb or Center, in Longitude; so as to correspond with the visible or apparent Contact, or angular Distance, of the Moon and Star; or Distance of the Moon and Sun first given.

Of DE LA CAILLE'S METHOD of determining the LONGITUDE, by OBSERVATION.

This celebrated great Astronomer, proceeds according to the Reverse of Dr. Halley's Method; from the apparent or visible to the true Distances of the Objects, observed.

He uses, for his Aid, l'Almanac nautique, wherein are computed the Distances of the Moon's Limb from a given fixed Star at Paris, to every Hour of the Night for each astronomical Day of the Year, under that Meridian; from whence the true angular Distances of the Moon from the same Star, to any other intermediate Time at Paris, are had by a single Proportion.

To which there is annexed the true Time of the Moon's Passage by the Meridian of Paris, and the Log. of her horizontal Parallax in Seconds of a Degree, at that Time.

Also the true Distance of the Moon's Limb from the Sun's next Limb are set down, for every Hour of the Day; with the Sun's Distance from each Pole, and Log. of the Moon's Parallax for each Day, at Noon, in the said Almanac nautique. — With the Logarithms for proportioning the Distances of the Moon and Star, or Moon and Sun, for Times between the Hours; with other auxiliary Tables, for facilitating the Operation; as may be seen in De la Caille's Ephemerides, from 1755 to 1765.

1. He finds the true Time of Observation under a distant Meridian from Paris, from Altitude of a Star, and Distance of the Eye from the Earth's Surface (allowing for Refraction) the true Distance of the Star from the Pole being also given, on a certain Day, when the true Time of its Passage by the Meridian of Paris is known, and to be nearly estimated for a distant Meridian, and the contrary.

2. He reduces the visible Altitude of the Moon and Star, observed under a distant Meridian, to the true Altitudes, at the Moment of their angular Distance taken.

3. He corrects the observed, or visible angular Distances of the Moon and Star to the true Distance; being an Error caused by Refraction.

4. He corrects the visible Distance observed to the true angular Distance of the Moon and Star, allowing for the Effect of the Moon's Parallax. — On which he remarks on a particular Case of the Moon and Star being in the same Vertical.

5. Then from the Result of these Operations, he finds the true Distance of the Moon and Star, as seen from the Earth's Center, for the Place of Observation, and true Time given; then proportioning for the true Time, to the same true Distance of the Moon and Star under Paris Meridian, the Difference of the Time of Observation, and Time at Paris, (when the Moon and Star have the same true Distance) reduced to Degrees, will be the Difference of Longitude, required; being a more direct, but a more tedious and difficult Method than Dr. Halley's; notwithstanding the Helps from the Almanac nautique, in proportioning the same Distance of the Moon and Star, or of the Moon and Sun, and a Table of Logarithms to the Distances from the Zenith; answering to Refractions, &c.

Of observing the LONGITUDE at SEA, by the MOON'S THEORY.

M. DE LA CAILLE takes Notice, in the Introduction to his EPHEMERIDES, from 1755 to 1765, that the Moon and Star's Altitudes, and their Distance from each other, are with Doubt, or Difficulty, taken near enough for Use. That, as the Moon's Motion is but 13° ± in 24 Hours, affording an Arch of about 2' of a Degree, moved over by the Moon in 4 Minutes of Time to a Degree Difference of Longitude, an Error of 3' of a Degree resulting from the Observation, and Computation of the Moon's true Place, under a distant Meridian, will create an Error of about 1 Degree and a Half in the Difference of observed Longitude. Who thinks we cannot be certain of the Longitude nearer than to 2 Degrees, on Account of the many contingent Obstacles in the Observation; however skillful the Observer and however accurately and well contrived his Instruments may be, with which he observes.

He also advances that in the several Circumstances entering into the Observation and Computation of the Longitude at Sea, that the Sum of the unavoidable Errors thereby are not less than 5' of a Degree. Who enumerates several Causes of Error, in taking an Observation of the Altitudes and Distance of the Moon and some Star, by a Quadrant of Reflection. 1. The defective Division of the Instrument. 2. The Difficulty of ascertaining the Parallelism of the Glasses in the Use of the Instrument. 3. The Difficulty of ascertaining the Contact of necessary Images and Parallelism, by the Diameters of Objects, in a Glass magnifying not more than 3 or 4 Times. 4. The Difficulty of adjusting the Images of Objects when not would measure their angular tr. Diff. 5. The Motion of the Ship interrupting our Attention to the Objects. 6. About 1' of a Degree Error, from the best lunar Tables, though rectified by Observation, in computing the Moon's true Place. Who farther adds, that Persons, having been at Sea, may have seen, that two Observers taking the Sun's Altitude at Noon, with a good Quadrant of Reflection, well rectified, shall differ 5, 6, 7, or 8 Minutes of a Degree, in rough Weather. That in the finest Weather and calmest Sea, no two Persons will agree in ascertaining the exact Latitude; but will differ at least, 3 or 4 Minutes of a Degree, which are the Limits assigned to Observations made at Sea, when the greatest Care is taken and the best Instruments are used in observing.

Of observing the LONGITUDE at SEA, by the MOON'S THEORY.

ON WHICH WE REMARK:

THAT if a Mean of the Latitudes, Altitudes, &c. taken by several Observers, of the same, at the same Time, be determined, it will come very near to Truth. For though it is possible one Observer may mistake, it does not follow that he must do so; because it is known, that several unavoidable Errors, lying a different Way, compensate rather than encrease one another. And by a Repetition of Observation, by one, or more Observers, using the same Methods, we may come near the Truth by taking a Mean of two or more Longitudes determined. And hence the Difference of Longitude, by the Moon's Place, observed under a distant Meridian, at a given or determined Time, compared with the corresponding Time, when the Moon has the same Place under a first Meridian, from correct Tables, will be very near the Truth, and extremely interesting for correcting a Ship's Longitude in long Voyages.

Therefore the Discovery of the Difference of Longitude, betwixt Greenwich and the Place of Observation, is Nothing more than the Difference of Time turned into Degrees, (allowing 15' of a Degree to 1 Minute of Time) betwixt the Times of the Moon having the same computed Place at Greenwich, as that observed under a distant Meridian; allowing for Parallax and Refraction, at the Place of Observation, for reducing the Moon's apparent to her true Place.

It is observed that the Moon's Altitude in the Night can be but imperfectly taken, though the Sky be then very clear; because the Horizon of the Sea is then seen but very confusedly; unless the Moon be very near the Horizon. This arises from the Moon's reflected Light upon the Surface of the Water, between the Eye and Horizon; the Sea and ill polished Glasses making the Effect unequal. A Train of Images are painted in the Vertical of the Moon, where Light hinders to distinguish the Horizon of the Sea beyond it; which however is always near the End of that Train, except when the Moon is near the Zenith. But when the Moon is very low, or near the Horizon, (between 5 and 15 Degrees Altitude) this reflected Light is confounded with the Horizon of the Sea, and then is the Time you may take her Altitude the most exactly and fit for the Purpose of the Longitude, according to De la Caille's Method.

A small Cloudiness or Thickness at the Horizon is reckoned another Obstacle to taking the Moon's true Altitude; because we thereby see the Extremity of the Horizon but very imperfectly, in considerable Altitudes of the Moon.

When the Moon passes to the Meridian more than 2 Hours after Sun-Rising, or before his Setting, her Light is then too feeble to determine, exactly, her Meridian Altitude. And when the Moon is covered with Clouds, in the Time of her Passage to the Meridian, there is Nothing to be done that Day. All which Inconveniences must be reduced to the proper Opportunity of observing the Longitude, or else they render the Result of the Observation extremely doubtful.

In this Method, besides the Moon's Altitude, it is necessary to have, as exact as possible, the Distance of the Moon from a near Star, that we may succeed without having 3 or 4' Error; and we must also have as exactly as possible the Altitude of the given Star, one of the brightest, not in the Vertical of the Moon, for avoiding the Inconveniences of its Reflection, that frequently occasions it to be seen on the Horizon of the Sea. In taking an Observation of its Altitude 7 or 8 Minutes of Error therein, will not be of much Concern in finding the Hour Angle, or Time of Observation. There are but few Cases wherein an Error in the Star's Altitude of 7 or 8 Minutes of a Degree will cause an Error of 15 Minutes of Time in the Hour Angle, or 3 Degrees and $\frac{1}{2}$ in the Longitude. But, there are a great Number of Cases where an Error of near 15 Minutes of a Degree in the Star's Altitude, will not make 1 Minute Error of Time in the Hour Angle, that we seek; and consequently not above 15 Minutes of a Degree Error in the Longitude or 5 or 6 English Leagues.

ACCOUNT of a PAMPHLET pretending a DISCOVERY of the LONGITUDE.

IN the Year 1710 there was a small Pamphlet, (which is fallen into our Hands) printed for D. Midwinter, at the three Crowns in St. Paul's Church-Yard, intitled A New and Easy METHOD to find the LONGITUDE at Sea. Which was from the Observation of the Altitudes of the Moon (though not of her Limb, but of her Center by

Gues) in the same Vertical with a fixed Star, taken by two Observers, with the Altitude of a second fixed Star, taken at the same Time, by a third Observer; from whence the Moon's Ecliptic Place and Latitude were determined, together with the Latitude of the Places and Time of Observation (but without any Allowance for Parallax and Refraction, whereby the Operation became altogether erroneous.) To which observed Time and Place of the Moon, the Time correspondent was also found in the Tables to the same Place, in a blundering Way, and thence, erroneously, the Difference of Longitude between the Meridian of the Tables, and Meridian of Observation.

OUR METHOD of determining the MOON'S true PLACE under a distant MERIDIAN.

By taking the Altitude of the Moon's upper or lower Limb, on the same Vertical with a noted fixed Star, whose Altitude is then taken (allowing for Parallax from an Ephemeris, and Refraction) at a given Time, in a given Latitude, you may, from thence, determine the Moon's true Longitude and Latitude, which will be of Use in determining the Distance of the Meridians, between the Place of Observation and that Meridian, for which correct astronomical Tables of the lunar Places are adapted. For the Difference of Time, in Degrees, between the Time of the Moon's true Place observed under a distant Meridian, and the Time of the same true Place computed from Tables suited to the first Meridian, will be the Difference of Longitude between those two Meridians.

But the Meridian Altitude of one of the Moon's Limbs, and Time when it touches the Meridian being taken, (allowing for Refraction and Parallax) and the Time before or after which a noted fixed Star transits the same Meridian, whose Meridian Altitude is truly ascertained, will give the Moon's true Place, as seen from the Earth's Center, from her visible Place seen from the Earth's Surface, the soonest of any Method (because there is no Parallax in right Ascension of the Moon, on the Meridian) and thence you will have the Difference of Longitude from the computed Time to the same true lunar Place.

EXAMPLE of computing the DIFFERENCE of LONGITUDE from GREENWICH.

By observing the MOON'S true PLACE under a distant Meridian.

Supposing that to the Westward of Greenwich, on July 14^d 1^h 20^m in the Morning, 1764, equal Time, the Moon's true Place (as seen from the Earth's Center) be found from her visible Place (as seen from the Earth's Surface) $\approx 4^{\circ} 39'$: Required, from thence, the Place's Longitude from Greenwich, where and when that Observation is made? By De la Caille's Eph. $\text{D's Pl. reduced to Greenwich.}$

1764, July 13, Noon $23^{\circ} 52' 28''$ Dif. $14^{\circ} 55'$, D's diurnal Mot.
14, Noon $7^{\circ} 23'$
 $23^{\circ} 52' 28''$ } Dif. $11^{\circ} 11'$. Say, As $14^{\circ} 55' : 24^h :: 11^{\circ} 11' :$
 $4^{\circ} 39'$ } 18^h 58^m , from the Noon of the 13th.

Hence it is 18^h past Noon, at Greenwich on the 13th astronomical Day of July, or at 6 in the Morning, on the 14th civil Day, 1764, when the Moon has Longitude $\approx 4^{\circ} 39'$, nearly.

But to determine the Difference of Longitude correctly, compute the Moon's true Place for 1764, July 13^d, 18^h, at Greenwich, and repeat the Operation, till the equal Time at Greenwich is found correspondent to $\approx 4^{\circ} 39'$ Long. D ; then the Difference between the computed Time at Greenwich and July 13^d 13^h 20^m Time of the Observation, under the distant Meridian, turned into Degrees, will give the correct Difference of West Longitude from Greenwich, required.

By first XI. Equa. marked C, and correcting Tab. 1764, July 13^d 18^h Greenwich, Moon's Place $\approx 3^{\circ} 59' 6''$
1764, July 13^d 13^h 20^m under dist. Merid. observed $\approx 4^{\circ} 39' 0''$

Moon's Pl. at Greenwich short of Place observed $39^{\circ} 54'$
1764, July 13^d 19^h, Greenwich, Moon's Pl. computed $\approx 4^{\circ} 33' 43''$
Hourly Motion of the Moon $34^{\circ} 37'$
Say, $34^{\circ} 37' : 1^h :: 39^{\circ} 54' : 1^h 9^m 10^s$ above 18^h , or
1764, 13^d 19^h 9^m 10^s Time at Greenwich } Moon's Pl. the same,
1764, 13 13 20 0 Time under distant } viz. $4^{\circ} 39'$.
Meridian

Dif. sooner $5^m 49^s 10$ under dist. Meridian.
Answering to $87^{\circ} 17' 30''$ W. Lon. fr. Greenwich, req. So for the Rest.

A NEW METHOD of making a LUNAR EPHEMERIS.

To estimate the MOON'S PLACE, and of making a lunar EPHEMERIS.

LET the Moon's Place be computed 4 Times in a Month, at least, (though 6 Times will be better) viz. on the Moon of the Syzygies, and Quadratures, being the extreme Limits of Motion.

If $4^{\circ} 10' 5''$ be added to the Moon's true Place, for any one Day, it will give her true Longitude for any other Day, within 12 or 15' at most, after 4326 Days (being 11 Years, 309 Days, or 12 Years wanting 56 Days) are elapsed.

Therefore, look into a correct Ephemeris for 12 Years before. And if you would know the Moon's true Place on the 1st of January, seek her Place for the 27th of February, 12 Years before; to which add $4^{\circ} 10' 5''$, and the Sum will be her true Longitude on the 1st Day of January, required, after 12 Years, within about 12 or 15' (at most) of her computed Place by good astronomical Tables. But on some Days of the Year, after 4326 Days elapsed, the Difference will be Nothing.

EXAMPLE.

Pl. Noon.	Lat.	D's Pl.
1752, Feb. 16, O.S. or 27, N.S. $\Omega 14^{\circ} 27'$	$4^{\circ} 52'$	De la Caille.
+ 4 1 5		Noon.
D's Pl. 1764, Jan. 1. Greenw. $\uparrow 15^{\circ} 32'$		$\uparrow 15^{\circ} 47'$
		$4^{\circ} 35'$

Proceeding thus, from Day to Day, with the Places found in an old Ephemeris, still adding $4^{\circ} 10' 5''$, and writing against the Sums, the respective Days answering thereto, successively, you will have a new Ephemeris of lunar Places, answering nearly to the Truth.

Parker's Ephemeris has been the best for this Purpose; because Dr. HALLEY condescended, annually, to assist that Ephemeridist in his Computations; especially in the lunar Places; as the late Mr. Henry Beighton, F. R. S. then Author of the Ladies Diary, observed in 1738, who was personally acquainted with both those Gentlemen.

As the Moon's true Place may differ from that computed, for the Days of the new Moon, Quarters, &c. some few Minutes, may proportionably be added to, or subtracted from, the computed Places, found after each Interval, according as the Case may require, which will determine the Moon's Place to a desired Accuracy, without a new Computation.

N. B. The foregoing Method (communicated by a Friend) of estimating the Moon's Place is proposed for Trial, and making a farther Improvement.

See Palladium 1758, P. 48, for a Method of determining the Moon's true Place after a Plinian Period (of 18 Years, 11 Days, &c.) and also her Latitude.

See also Palladium 1759, P. 68, for a Method of determining the Lunations and Moon's Rising and Setting, after a Plinian Period.

To find the near Time of the Moon's Rising, (or a Planet's) in a given Latitude?

Enter the Table of oblique Ascensions in Time, as in Leadbetter's II. Vol. of Astronomy, (proper to the Latitude of the Place) and with the Moon's or Planet's Longitude and Latitude, at the estimate Rising, take out the oblique Ascension in Time, from which deduct the Sun's right Ascension in Time at the estimate Rising, and the Remainder more or less 6 Hours, (according as it is less or more than 6 Hours) will be the near Time of Moon's (or a Planet's) Rising, in the Morning, required.

To find the near Time of the Moon's (or a Planet's) Setting, to a given Latitude?

Enter the Table of oblique Ascensions, in Time. (proper to the Latitude of the Place) with the opposite Sign and Degree of Longitude, and contrary Latitude to what the Moon (or Planet) has at the Time of Setting, and take out the oblique Descension in Time agreeing thereto, from which take the Sun's right Ascension in Time, and the Remainder, more or less 6 Hours, (according as it is less or more than 6 Hours) will be the Time of the Moon's (or Planet's) Setting after Noon, required.

Since the Difference of oblique Ascensions of 1 Degree (in our Latitudes) in a Sign of long Ascension, as $\overline{\alpha}$, Ω , $\overline{\pi}$, $\overline{\mu}$, $\overline{\eta}$, $\overline{\zeta}$, is but 6m of Time, at most, if the Moon's Longitude be within 10 or 12' of Truth, which it will seldom exceed, and for the most Part be less, it occasions but about 1m Error in the Time of Rising and Setting. And as for 3 or 4' of Latitude (which perhaps may sometimes happen) that will not cause above Half a Minute's Variation, at most, in her Rising.

But, if she be in the Signs of short Ascension, as in $\overline{\nu}$, $\overline{\omega}$, $\overline{\chi}$, $\overline{\psi}$, $\overline{\delta}$, $\overline{\epsilon}$, the Difference of oblique Ascension of 1 Degree, will not, sometimes, amount to 2 Minutes in Time. Consequently, should her Longitude vary as above, there will not (by this Method of Computation) be Half a Minute Difference (at most) from the true Rising and Setting.

In Computations of this Kind, it is necessary to be careful in reducing the Moon's Place and right Ascension, and Sun's right Ascension, for the Time given, to those for the estimate Time of her Rising and Setting; or (in repeating the Operation) for the last found Time.

In the foregoing Operations, Parallaxes and Refractions are not considered.

Of the MOTION of COMETS.

THE Reverend STEPHEN BOLTON, M. A. and Rector of Stalbridge, Dorsetshire, has obliged us with several of his curious Remarks and Improvements, on the Motion of Comets. The BASIS on which he founds his Improvements, and to which he refers all his Computations, depends on the angular Motion of the Earth round the Sun, from the first to the last Observation of the Comet. To exemplify which, he has given several Schemes of the angular Motion of the Earth round the Sun, in different Positions of different Comets observed.

His UNIVERSAL METHOD of subtracting or adding the Angles, found at the Sun, to and from other Angles which the Earth describes round the Sun, so as to correspond with the Angles observed in different Positions of the same Comet, or to reconcile an Agreement among one another, must infallibly prove the Data to be true, and when they may be depended upon.

Moreover, if the parabolic Trajectory of a Comet be found, very nearly, to answer the Observations at the Earth, it is very reasonable to conclude (he says) that the Velocity of that Comet, in its true elliptic Orbit, in and about the Perihelion, and Perihelion-Distance itself, will be very nearly the same, as in the fictitious parabolic Curve.

And, if so, then Nothing more is wanting (he says) but to find such Semi-Transverse to the Comet's elliptic Orbit, (or periodic Time, the one giving the other) as may exactly correspond with all the Observations of the Comet taken; and which (he says) may be effected by Trial.

See our easy and short Method of finding the true from the mean Anomaly of a Planet or Comet. P. 96.

In taking Observations of the Altitude of a Comet, and fixed Star, in the same Vertical, either above or below the Comet, Mr. Bolton projects the Light from a dark Lanthorn up a fine silver Wire playing in Water for a vertical Line.

The Altitude of the Comet, (allowing for Refraction) Time of Observation, and Latitude of the Place, being given, the Star's angular Distance from the Meridian is also given. Whence the Comet and Stars Azimuth from the North, the Star's Angle of Position, by a Vertical and Circle of Declination, the Star's Altitude, and consequently, the Difference of Altitudes, the Difference of right Ascension of Comet and Star, and Co-Declination of the Comet; and the Longitude and Latitude of that Comet, are found.

Two Observers, taking the Altitude of the Comet, and fixed Star, on the same Vertical, either above or below the Comet, at the same Time, on the same Azimuth, may spare the foregoing astronomical Operation.

EXAMPLE

EXAMPLE of lunar COMPUTATION. According to D. C.

By Halley's Tables.	SUN and MOON's PLACES computed. C.			ARGUMENTS of EQUATIONS and CONCLUSIONS.		
Sun's mean Longitude	s 6 12 52 52	s 6 12 52 52	s 6 10 56 46	Mean Anomaly of the D	s 6 1 5 22	
Sun's Equation	— 1 56 6	6 8 54 16	6 8 54 16	2. Mean Arg. Ann. or 2. mean Long.		
Sun's true Longitude	— 10 56 46	0 3 58 36	0 2 2 30	2. $\odot$ à D's Apogee	0 7 57 12	
Sun's mean Anomaly	3 4 41 37	An. Arg. to mean Long. Sun.	An. Arg. to true Long. Sun.	Mean Evection	5 23 8 10	
By Royal Astronomer.	D's Longitude.	D's Apogee.	D's Node.	1/2 Mean Equation of the D's Center + Diff. 6 Equations	+ 4 7	
1732	3 21 40 39	5 9 45 34	8 27 47 49	Arg. of Evection, correct, from Sun's mean Place	5 23 12 17	
September 22	8 11 44 41	.. 29 31 23	. 14 2 0	Evection-Equation to this Argument	— 0 9 40	
H. 11	... 6 2 21	... 3 4	... 1 27	Evection-Equation to Arg. Evac. (from $\odot$ true Pl.) 5 ^s 27 ^o 4' 29"	— 0 4 12	
37 ^m . . 28 ^s	... 20 34	... 11	... 5	Diff. = Equation of Evection from quadruplicate Ratio of Sun's Dist.	5 28	
Mean Motions	0 9 48 15	6 9 20 12	8 13 44 17	Newton's greatest second Equat. of D's Center	— 2 25	
Annual Equations	+ 11 23	— 25 56	+ 9 20	Greatest Equation of D's Center, from tripl. Ratio of Sun's Dist. resulting	3 3	
D 1. equated	0 9 59 38	6 8 54 16	8 13 53 37	According to Mayer, 3d + 4th Equation	3 0	
Apogee 1. equated —	6 8 54 16	D's Ap. 1. equated.	0 10 2 12	According to R. Astronomer, 6 + 7 Equation	4 6	
Mean Anomaly	6 1 5 22	6 1 5 22	3 26 8 35	These last Equations, viz. their Sum, are to the Evection-Equation, respectively at a Maximum, as 3. Eccentricity of the Earth's Orbit to its Semitransverse. (See P. 376.)		
6 Equations	+ 0 15	0 4 5 0	[1st Ar. Lat. 2. An. Arg.	The above is designed as a PROOF, that the Sum of the 6 and 7 Equations, at the Beginning OF THIS WORK, is = 4' 6", at a Maximum, deduced from the reciprocal triplicate Ratio of the Sun's Distance, agreeing with 5' 28" Equation, by Tables, according to the quadruplicate or 4th Power of the Sun's Distance inversely. — For, as the generated Equations are directly as the Indices of the reciprocal generating Powers of the Sun's Distance, Say,		
Anom. 7 Equat. Elliptic Equation	6 1 5 37	5 27 0 22	Evac. fr. $\odot$ tr. Pl. 1/2 M. Eq. D Cen. + [6 lefs Eq.	As 3 Index : 4' 6" Equat. :: 4 Index : 5' 28" Equat. by Tables, as above. Q. E. D.		
Anom. 8 Equat. Evection Equation	6 1 13 22	5 27 4 29	Arg. of Evac. fr. $\odot$ [tr. Pl.			
Anom. 9 equated Variation-Equation	6 1 9 10					
True Anomaly	6 1 7 56					
Apog. 1 equated +	6 8 54 16					
D's Orb. Long.	0 10 2 12					
Reduc. + Eq $\odot$ = 22"	+ 5 53					
D in Ecciptic Observed	10 8 5					
Error	— 0 22					
Omitting the 7th Equation according to Newton.	Error — 2' 2"					
By Halley's Tables, Error — 5' 25"						
		No. Argumts. Equations.				
		s 0 1 1 11 1 11				
		II. 0 2 2 0 26 ...				
		III. 9 26 53 ...				
		IV. 1 20 46 ...				
		V. 5 29 10 ...				
		VI. 9 1 42 2 25 ...				
		VII. 2 22 18 ...				
		An. 9. 6 ^s 10 ^o 9' 10 2 51 3 6				
		An. Ar. 0 2 2 30 — —2 51				
		D à $\odot$ 5 29 6 40 6 lefs Eq. +0 15				

To be published by Subscription.

Astronomia Perfecta, Or the Supplemental ASTRONOMER and NAVIGATOR, completing the Royal Astronomer and Navigator (lately published.)

Containing the Greenwich and Paris TABLES (lately acquired) of the Radical Mean Places and Motions of the SUN and MOON. CORRECT TABLES of the Radical Mean Places and Motions of the PLANETS and SATELLITES; and Equations for computing their respective true Places, and equal or true Time of their Appulses, Occultations, Transits, and ECLIPSES. Illustrated by Precepts and Examples. Also the Elements and Doctrine of COMETS. And many New Improvements in ASTRONOMY.

With an EPHEMERIS, for 6 Years from the Date of Publication,

Correcting all the Seamen's Calendars extant; and Tables fitted for observing the Longitude from the Royal Observatory at Greenwich.

By the AUTHOR of the ROYAL ASTRONOMER and NAVIGATOR.

First the Sun
 (A mighty Sphere) be fram'd; unlightsome first,
 Though of ethereal Mould: then form'd the Moon
 Globose, and every Magnitude of Stars;
 And sow'd with Stars, the Heav'n, thick as a Field.

Milt. Par. Lost. B. vii. l. 354.

EXAMPLE of COMPUTATION. According to Mr. MAYER's supplemental EQUATIONS. See P. 97.

To find the SUN and MOON's PLACES 1739, February 12^d 3^h 39^m 22^s $\frac{1}{2}$ true Time, or 3^h 54^m 10^s mean Time, for the Meridian of Paris OBSERVATORY.

	Sun's Longitude.	Apogee of the Sun.	☽'s Longitude.	Apog. ☽.	☽ ascend.
	s o ' "	s o ' "	s o ' "	s o ' "	s o ' "
1739	9 9 40 54	3 8 25 22	5 15 31 28	2 23 18 28	4 13 5 35
February 12	1 12 22 58	7	6 26 35 6	4 47 26	2 16 38
3 ^h	7 24	0	1 38 49	50	24
54 ^m 10 ^s	2 13	0	29 44	15	7
☉ Equation	10 22 13 29	3 8 25 29	0 14 15 8	2 28 6 59	2 17 9
	+ 1 21 9	7 13 48 0	☉ M. Anom.	9 16 8 9	4 10 48 26
				— 14 32	— 7 16
☉ Long. fr. Equ ^x .	10 23 34 38	Diam. Sun 3229	0 14 6 13	9 15 53 37	4 10 41 10
Precef. Equ ^x .	Correct Long. + 8	Hor. Mot. 231	+ 5 50 3	— 8 55	I. 8 9 50 40
Tr. Long. ☉ fr. Equ ^x .	10 23 34 46	98802	0 19 56 26	XI. 9 15 44 42	☽ à ☉ 1 26 57
	— 14	+ 3	35 24	☽ à ☉ 1 20 31 27	Doub. 3 23 54
☽ à ☉	1 20 40	98805	0 20 31 50	Doub. 3 11 2 54	II. 7 14 3
		☉ à ☽.	— 4 30	XII. 5 25 18 12	South Lat. 4 50 1
An. ☽	9 16 8	Lefs Equations.	0 20 27 20	+ 5 56 54	South Lat. 6 36
2. ☽ à ☉	3 11 20	— +	Long. ☽ fr. Eq.	— 0 6 41	South Lat. 4 56 37
Arg. I.	7 13 48 ...	8' 1"	Prec. Eq. — 14	+ 5 50 13	Or Tot. Lat. ☽ S.
II.	10 25 8 ...	...	0 20 27 6	XIII. 1 26 21 40	XI. 56 9
III.	7 27 32 ...	...	Tr. Long. ☽ fr. Eq.		XII. + 38
IV.	1 9 0 ...	...			XIII. — 10
V.	10 11 24 ...	0 54			Eq. Par. ☽ 56 37
VI.	0 7 28 ...	...			Hor. Diam. ☽ 30 53
VII.	6 20 46 ...	21			Equat. Precef. Equinox — 14
VIII.	2 2 20 ...	...			Long. Sun from Equinox 23° 34' 32"
IX.	5 17 14 ...	20			Long. ☽ from Equinox 20 27 6
X.	4 4 32 ...	3 7			Latitude ☽ South 4 56 37
	Total —	12 43			Long. ☽ observed 20 27 15
	+	3 48			Latitude observed 4 56 45
	—	8 55			Error in Longitude — 0 9
					Latitude — 0 8

Correction of the Longitude of the SUN and SUN's Distance from the EARTH. According to Mr. MAYER.

Argument. MOON from the SUN.

☽ à ☉.	Sig. 0.		Sig. 1.		Sig. 2.		Sig. 3.		Sig. 4.		Sig. 5.		☽ à ☉.
o	Long.	Diff.	Long.	Diff.	Long.	Diff.	Long.	Diff.	Long.	Diff.	Long.	Diff.	o
	+	+	+	+	+	+	+	—	+	—	+	—	
	"	Parts.	"	Parts.	"	Parts.	"	Parts.	"	Parts.	"	Parts.	
1	0	5	5	4	9	3	10	0	9	3	5	4	26
6	1	5	6	4	9	2	10	1	8	3	4	5	21
11	2	5	7	4	9	2	10	1	8	3	3	5	16
16	3	5	7	3	10	1	10	1	7	4	2	5	11
21	4	5	8	3	10	1	9	2	6	4	2	5	6
26	4	4	8	3	10	0	9	2	6	4	1	5	1
	—	+	—	+	—	+	—	—	—	—	—	—	
☽ à ☉.	Long.	Diff.	Long.	Diff.	Long.	Diff.	Long.	Diff.	Long.	Diff.	Long.	Diff.	☽ à ☉.
	Sig. 11.		Sig. 10.		Sig. 9.		Sig. 8.		Sig. 7.		Sig. 6.		

A TABLE and EXAMPLE *new modelled*, for USE, (the TRUTH of *Investigation* and *Construction* being proved.) According to MISCELLANEOUS TRACTS. Whereby you may reduce a COMET's LONGITUDE from the Perihelion in a *parabolic* Orbit, to the Longitude from the Perihelion in an *elliptic* Orbit; by the Difference of Motion in those Orbits.

Comet from Per.	Min. of Deg.	Corresp. common Log.	For Log. of Comet fr. Sun.	Comet from Per.	Min. of Deg.	Corresp. common Log.	For Log. of Comet fr. Sun.	Comet from Per.	Min. of Deg.	Corresp. common Log.	For Log. of Comet fr. Sun.	Comet from Per.	Min. of Deg.	Corresp. common Log.	For Log. of Comet fr. Sun.
0	—	—	—	0	—	—	—	0	+	+	—	0	+	+	—
1	30'	1.4770	5.8200	41	836'	2.9220	9.0006	81	152'	2.1827	9.4777	121	3547'	3.5499	9.7516
2	60	1.7770	6.4220	42	836	2.9229	9.0195	82	206	2.3139	9.4851	122	3675	3.5653	9.7602
3	90	1.9533	6.7735	43	836	2.9232	9.0378	83	261	2.4170	9.4924	123	3806	3.5805	9.7690
4	119	2.0778	7.0242	44	836	2.9228	9.0555	84	318	2.5021	9.4996	124	3941	3.5956	9.7781
5	149	2.1740	7.2178	45	835	2.9217	9.0727	85	376	2.5750	9.5067	125	4078	3.6105	9.7874
6	179	2.2521	7.2759	46	832	2.9200	9.0894	86	435	2.6386	9.5138	126	4220	3.6253	9.7971
7	208	2.3178	7.5095	47	827	2.9176	9.1056	87	496	2.6951	9.5207	127	4364	3.6399	9.8070
8	237	2.3745	7.6250	48	821	2.9145	9.1215	88	558	2.7470	9.5275	128	4513	3.6545	9.8172
9	266	2.4242	7.7269	49	814	2.9107	9.1369	89	622	2.7936	9.5342	129	4667	3.6690	9.8278
10	294	2.4683	7.8178	50	806	2.9061	9.1520	90	688	2.8373	9.5409	130	4825	3.6835	9.8387
11	322	2.5074	7.8998	51	796	2.9007	9.1666	91	754	2.8776	9.5475	131	4989	3.6980	9.8500
12	349	2.5431	7.9747	52	784	2.8945	9.1808	92	823	2.9152	9.5540	132	5158	3.7125	9.8618
13	376	2.5760	8.0437	53	772	2.8874	9.1947	93	892	2.9506	9.5605	133	5332	3.7269	9.8740
14	403	2.6058	8.1074	54	758	2.8795	9.2083	94	963	2.9837	9.5670	134	5512	3.7413	9.8866
15	430	2.6331	8.1666	55	743	2.8708	9.2215	95	1036	3.0154	9.5734	135	5698	3.7557	9.8997
16	455	2.6583	8.2217	56	726	2.8610	9.2344	96	1110	3.0455	9.5797	136	5890	3.7701	9.9132
17	480	2.6816	8.2735	57	708	2.8500	9.2469	97	1186	3.0742	9.5860	137	6088	3.7845	9.9273
18	505	2.7032	8.3222	58	688	2.8378	9.2592	98	1264	3.1016	9.5923	138	6297	3.7991	9.9419
19	529	2.7233	8.3682	59	667	2.8243	9.2712	99	1343	3.1279	9.5986	139	6513	3.8138	9.9571
20	552	2.7420	8.4116	60	645	2.8095	9.2829	100	1423	3.1531	9.6050	140	6739	3.8286	9.9729
21	575	2.7594	8.4528	61	621	2.7932	9.2943	101	1505	3.1774	9.6113	141	6974	3.8435	9.9893
22	596	2.7755	8.4921	62	596	2.7754	9.3055	102	1588	3.2008	9.6176	142	7219	3.8585	10.0064
23	617	2.7906	8.5295	63	570	2.7557	9.3163	103	1673	3.2236	9.6239	143	7475	3.8736	10.0242
24	637	2.8046	8.5652	64	542	2.7338	9.3270	104	1761	3.2457	9.6302	144	7743	3.8889	10.0427
25	657	2.8176	8.5994	65	512	2.7095	9.3374	105	1850	3.2671	9.6366	145	8024	3.9044	10.0620
26	676	2.8297	8.6320	66	482	2.6827	9.3477	106	1941	3.2879	9.6430	146	8316	3.9199	10.0822
27	694	2.8410	8.6634	67	450	2.6526	9.3576	107	2033	3.3081	9.6495	147	8630	3.9360	10.1032
28	710	2.8514	8.6934	68	416	2.6187	9.3674	108	2127	3.3278	9.6560	148	8962	3.9524	10.1251
29	726	2.8610	8.7224	69	381	2.5805	9.3769	109	2223	3.3469	9.6627	149	9312	3.9690	10.1481
30	741	2.8698	8.7502	70	344	2.5363	9.3862	110	2321	3.3656	9.6694	150	9681	3.9859	10.1720
31	755	2.8779	8.7770	71	306	2.4857	9.3954	111	2421	3.3840	9.6763	151	10072	4.0031	10.1970
32	768	2.8853	8.8030	72	267	2.4258	9.4044	112	2523	3.4019	9.6831	152	10491	4.0208	10.2231
33	780	2.8920	8.8280	73	226	2.3537	9.4132	113	2627	3.4195	9.6901	153	10940	4.0390	10.2504
34	791	2.8980	8.8521	74	184	2.2639	9.4218	114	2734	3.4368	9.6972	154	11410	4.0575	10.2792
35	801	2.9034	8.8755	75	140	2.1458	9.4303	115	2843	3.4538	9.7044	155	11929	4.0766	10.3094
36	809	2.9081	8.8980	76	95	1.9767	9.4386	116	2954	3.4704	9.7119	156	12483	4.0963	10.3412
37	817	2.9122	8.9198	77	48	1.6840	9.4468	117	3068	3.4868	9.7195	157	13086	4.1168	10.3746
38	823	2.9156	8.9409	78	+	+	9.4546	118	3184	3.5030	9.7272	158	13740	4.1380	10.4098
39	829	2.9184	8.9614	79	49	1.6910	9.4624	119	3302	3.5188	9.7352	159	14455	4.1600	10.4469
40	833	2.9205	8.9813	80	100	2.0006	9.4701	120	3422	3.5345	9.7433	160	15223	4.1825	10.4861

USE. In the 1st Column find the Comet's Longitude from the Perihelion the same as that given according to the Hypothesis of a *parabolic* Orbit (by *Halley's*, *De la Caille's*, or other Tables) against which Longitude, in the 3d Column, you have the Logarithm answering to a Number of Minutes in 2d Column. From which Logarithm subtract the Logarithm of the greater *Axe* divided by the *Perihelion Distance* of the Comet's elliptic Orbit, and the Remainder will be the Logarithm of a Number of Minutes to be added or subtracted to or from the said Longitude, as the Table above directs, for the Comet's true Longitude in the *elliptic* Orbit.

Likewise, from the Logarithm in the 4th Column, correspondent to the said Longitude, subtract the Logarithm of the said *Quotient*, and the Remainder, connected with 10 as *Index*, will be the Logarithm of a *Quantity* to be deducted, from the Logarithm of the Comet's Distance from the Sun, in the *parabolic Hypothesis*, for the Logarithm of the Comet's Distance from the SUN, according to the *elliptic* Orbit.

EXAMPLE. Let the greater *Axe* of the Ellipsis be = 35.727, and *Perihelion Distance* = 0.5825 (being the Orbit of the Comet seen in the Year 1682) and let the Longitude from *Perihelion* according to the *parabolic* Orbit, in *Dr. Halley's Table*, be 44° 3' 20", with the Log. 0.065838 of the Comet's Distance from the Sun, correspondent.

$$\text{The Log. of } \frac{35.727}{0.5825} = 1.7877 - \dots - 1.7877 - \\ 0.5825 \quad 2.9228 \text{ in Tab. } \dots 9.0555 \text{ against } 44^\circ \text{ Longitude.}$$

$$\text{Subtract } 13' 39'' = 13'.65 \dots 1.1351 \text{ Log. } \quad 7.2678 \\ \text{Com. à Perih. } 44^\circ 3' 20'' \text{ in par. Orb. } \quad - 10$$

$$\text{Rem. Long. } 43.49.41 \text{ from Perih. req. } \\ \text{in ellipt. Orb.}$$

$$\text{Or Log. } 3.2678 \dots \text{No. corresp. } .001853 - \\ \text{Log. Com. à } \odot \text{ in parab. Orb. } .065838$$

$$\text{Log. of Comet's Distance from the Sun, required } \dots 0.063985 \text{ Rem.} \\ \text{in ellipt. Orb.}$$

The foregoing Table not only reduces the Comet's Longitude from a *parabolic* to an *elliptic* Orbit, by the Difference of Motion in those two Orbits, but is farther useful, for nearly determining the *Species* of the elliptic Orbit which a *new* Comet describes, when the Observations made thereon differ essentially from the computed Places by the *parabolic Hypothesis*.

COMPUTATION of an OPPOSITION of the SUN and MOON, (with their Places.)
By D. COWPER.

D. COWPER, observing the Difference (in P. 397) between the Time of the Ecliptic Opposition of the Sun and Moon, computed by a few Equations, and the Time computed in M. De la Caille's Ephemerides, has computed (from all the lunar Equations, marked C.) the Time more correctly as follows, on March 17^d 12^h 3^m 36^s, in 1764, for Greenwich, D. C.

By De la Caille's Ephemerides 2 16 . . . for Greenwich.

1 20 Diff.

For, 1764, March 17^d 11^h 53^m, mean Time, Sun's true Place $\propto$ 27° 53' 58"
Moon's true Ecliptic Place at that Time 5 27 47 46

$\propto$ à $\odot$ short of an Opposition . . . 6 12
 $\odot$ à $\propto$ past an Opposition in Hour after . . . 28 53

Mot. $\propto$ à $\odot$ in an Hour, the Sum . . . 35 5

Now, 35' 5" : 60^m :: 6' 12" : 10^m 36^s to be added to the first Time, for Time of true Opposition.

March 17^d 11^h 53 0

1764, March 17 12 3 36 Mean Time of the true Opposition of the Sun and Moon, at Greenwich, D. C.

Moon's Lat. N. D. 0° 39' 1" } M. Anom. $\odot$ 8° 17' 11" 23" in 1764 March 17^d 11^h 53^m.
Horizontal Parallax 60 31 }

According to Mayer's Tables 1764, March 17^d 12^h 2^m . . . Sun's Longitude . . . 11° 27' 53' 58"
Moon's Ecliptic Longitude at that Time . . . 5 27 47 21

$\propto$ à $\odot$ short of Opposition . . . 6 37

Now, As 35' 5" hourly Mot. $\propto$ à $\odot$: 60^m :: 6' 37" short of Opposition : 11^m 19^s
March 17^d 12^h 2 0

Opposition according to Mayer . . . 1764, March 17 12 13 19 for Paris.

M. De la Caille 17 12 11 36 for Paris.

Moon's Lat. N. D. 0° 39' 20" }
Horizontal Parallax 60 57 }

1 43 Diff.

Correspondent to 1764, Mar. 17^d 3^m 59^s Mayer, and Mar. 17^d 2^m 16^s De la Caille, for Greenwich.

According to D. C.

1737, February 18^d 2^h 37^m 53^s Mean Time, Moon's Place computed in Syzygy $\propto$ 11° 7' 39"
Observed . . . $\propto$ 11 11 23 } By our supplemental
Equations, doubling
6th Equation = - 4' 50".

Moon's Lat. N. 0° 37' 57"

Error - 4 44
+ 2 25 for 6th Equation, single.

Error but - 2 19

The same computed by all the Moon's Equations C. (connecting the Correction of 6th and 7th Equations + 56")
 $\propto$ 11 11 23 the same as observed.

D. Cowper is doubtful whether the Acceleration of the Moon's mean Motion in this Century be a Correction or not, for Want of sufficient Observations. Who is of Opinion the Theory of Gravity requires, that the Apogee and Node should be accelerated too; but not in the Ratio as is given by Mr. Gaal Morris, (in the *Astronomia Arcana*, or *Greenwich Tables*, he has secretly printed) but in the Ratio of the annual Equations to each other, as 6, 10, and 5, very nearly; in which Proportion the Correction of the Moon's mean Motion is given in the foregoing Computation.

CONCLUSION, or LAND.

* * * The Plan of our WORK being already far extended beyond its original designed Limits, our Reader, it is hoped, will excuse us, for omitting the general geometrical Construction of SOLAR ECLIPSES, &c. to make Room for the foregoing METHOD of computing the LONGITUDE OF PLACES BY OBSERVATION OF THE MOON, and other more necessary Subjects.

What we have farther promised, and is expected from us, to complete our WORK, we refer the Readers for Satisfaction, to our *Astronomia Perfecta*, or SUPPLEMENTAL ASTRONOMER, and NAVIGATOR, (partly finished) to be published by Subscription, at Half a Guinea paid down for each complete Book, in Sheets, as soon as we find sufficient Encouragement: Works of this Nature having too few Readers, and are too expensive to be published otherwise; as we have already sufficiently experienced.

The Number of our Subscribers to this present Work are so few, that we think it rather a Diminution than a Reputation to Science to print so small a List. But we the less wonder at so few Encouragers of Science, as there have been so many Impositions, by Men taking Subscribers Money for Books, or engraved Performances, they were never qualified to execute, nor designed to deliver. Who therefore ought to be looked upon as Enemies to Science, Society, and their Country. And we are sorry to say that some Persons (who would be thought very honest) have traded with our Subscribers Money for Want of a Principle and Spirit to pay it, as they were ordered, for our Use, in carrying on this expensive Work: The original Plan of which being augmented from 30 to 53½ Sheets, in Quarto.

F I N I S.



